

Market Pricing for Matroid Rank Valuations

Kristóf Bérczi*

Naonori Kakimura[†]

Yusuke Kobayashi[‡]

July 20, 2020

Abstract

In this paper, we study the problem of maximizing social welfare in combinatorial markets through pricing schemes. We consider the existence of prices that are capable to achieve optimal social welfare without a central tie-breaking coordinator. In the case of two buyers with rank valuations, we give polynomial-time algorithms that always find such prices when one of the matroids is a simple partition matroid or both matroids are strongly base orderable. This result partially answers a question raised by Dütting and Végh in 2017. We further formalize a weighted variant of the conjecture of Dütting and Végh, and show that the weighted variant can be reduced to the unweighted one based on the weight-splitting theorem for weighted matroid intersection by Frank. We also show that a similar reduction technique works for M^2 -concave functions, or equivalently, gross substitutes functions.

1 Introduction

In this paper, we study the problem of maximizing social welfare in combinatorial markets through pricing schemes. Let us consider a combinatorial market consisting of indivisible goods and buyers, where each buyer has a valuation function that describes the buyer’s preferences over the subsets of items. The goal is to allocate the items to buyers in such a way that the social welfare, that is, the total sum of the buyers’ values, is maximized. Such an allocation can be found efficiently under reasonable assumptions on the valuations [32]. As an application of the Vickrey–Clarke–Groves (VCG) mechanism [6, 21, 37] for welfare maximization, the VCG auction is another illustrious example. However, the problem becomes much more intricate if the optimal welfare is ought to be achieved using simpler mechanisms employed in real world markets, such as pricing.

In a pricing scheme, the seller sets the item prices, and the *utility* of a buyer for a given bundle of items is defined as the value of the bundle with respect to the buyer’s valuation, minus the total price of the items in the bundle. Ideally, the prices are set in such a way that there exists an allocation of the items to buyers in which the market clears and everyone receives a bundle that maximizes her utility. A pair of pricing and allocation possessing these properties is called a *Walrasian equilibrium*¹, while we will refer to the price vector itself as *Walrasian pricing*. The

*MTA-ELTE Egerváry Research Group, Department of Operations Research, Eötvös Loránd University, Budapest, Hungary. Email: berkri@cs.elte.hu.

[†]Department of Mathematics, Faculty of Science and Technology, Keio University, Yokohama, Japan. Email: kakimura@math.keio.ac.jp.

[‡]Research Institute for Mathematical Sciences (RIMS), Kyoto University, Kyoto, Japan. Email: yusuke@kurims.kyoto.ac.jp.

¹Walrasian equilibrium is often called competitive pricing, or market equilibrium in the literature.

fundamental notion of Walrasian equilibrium first appeared in [38], and the definition immediately implies that the allocation in a Walrasian equilibrium maximizes social welfare. Therefore, the problem might seem to be settled for markets that admit such an equilibrium.

Cohen-Addad et al. [7] observed that Walrasian prices alone are not sufficient to coordinate the market. The reason is that ties among different bundles have to be broken up carefully by a central coordinator, in a manner consistent with the corresponding optimal allocation. However, in real markets, buyers walk into the shop in an arbitrary sequential order and choose an arbitrary best bundle for themselves without caring about social optimum. In their paper, it is shown that the absence of a tie-breaking rule may result in an arbitrary bad allocation.

To overcome these difficulties, Cohen-Addad et al. [7] introduced the notion of *dynamic pricing schemes*. In this setting, the seller is allowed to dynamically update the prices between buyer arrivals. Achieving optimal social welfare based on dynamic pricing would be clearly possible if the order in which buyers arrive was known in advance. Nevertheless, determining an optimal dynamic pricing scheme is highly non-trivial when the prices need to be set before getting access to the preferences of the next buyer.

The main open problem in [7] asked whether any market with *gross substitutes valuations* has a dynamic pricing scheme that achieves optimal social welfare. A market with gross substitutes valuations is known to be an important class of markets having Walrasian prices [25]. It is worth noting that the existence of an optimal scheme reduces to the existence of an appropriate initial price vector; an optimal allocation then can be determined by induction. For a formal definition, we refer the reader to [1].

As a starting step towards understanding the general case, Dütting and Végh [11] suggested to look at matroid rank functions as valuations, because a matroid rank function is a fundamental example of gross substitutes valuations. In particular, they proposed the following conjecture for the case of two buyers.² Here, a matroid with a ground set S and a base family \mathcal{B} is denoted by $M = (S, \mathcal{B})$ and we denote $p(X) := \sum_{s \in X} p(s)$ for $p : S \rightarrow \mathbb{R}$ and $X \subseteq S$.

Conjecture 1. *Let $M_1 = (S, \mathcal{B}_1)$ and $M_2 = (S, \mathcal{B}_2)$ be matroids with a common ground set S such that there exist disjoint bases $B_1 \in \mathcal{B}_1$ and $B_2 \in \mathcal{B}_2$ with $B_1 \cup B_2 = S$. Then, there exists a function $p : S \rightarrow \mathbb{R}$ (called a price vector) satisfying the following conditions.*

1. *For any $B_1 \in \arg \min_{X \in \mathcal{B}_1} p(X)$, it holds that $S \setminus B_1 \in \mathcal{B}_2$.*
2. *For any $B_2 \in \arg \min_{X \in \mathcal{B}_2} p(X)$, it holds that $S \setminus B_2 \in \mathcal{B}_1$.*

The requirements in the conjecture can be interpreted as follows. There are two buyers and each buyer $i \in \{1, 2\}$ wants to buy a set of items that forms a basis in \mathcal{B}_i . If buyer i comes to a shop first, then she chooses a cheapest set B_i in \mathcal{B}_i with an arbitrary tie-breaking rule. Regardless of the choice of B_i , the remaining set $S \setminus B_i$ is a desired set for the other buyer.

Actually, Conjecture 1 resolves the existence of a static pricing scheme for a two-buyer market with matroid rank valuations. That is, if Conjecture 1 is true, then the following conjecture is also true. See Lemma 10 for the details.

Conjecture 2. *Let $M_1 = (S, \mathcal{B}_1)$ and $M_2 = (S, \mathcal{B}_2)$ be matroids with rank functions r_1 and r_2 , respectively. Then, there exists a function $p : S \rightarrow \mathbb{R}$ satisfying the following conditions.*

²Dütting and Végh conjectured that the price vector p can be chosen to have all different values, that is, $p(s_1) \neq p(s_2)$ for $s_1 \neq s_2$. This difference is not essential, because we can apply a perturbation to p without affecting the requirements in Conjecture 1.

1. For any $B_1 \in \arg \max_{X \subseteq S} (r_1(X) - p(X))$ and for any $B_2 \in \arg \max_{Y \subseteq S \setminus B_1} (r_2(Y) - p(Y))$, $r_1(B_1) + r_2(B_2) = \max\{r_1(X) + r_2(Y) \mid X, Y \subseteq S, X \cap Y = \emptyset\}$.
2. For any $B_2 \in \arg \max_{Y \subseteq S} (r_2(Y) - p(Y))$ and for any $B_1 \in \arg \max_{X \subseteq S \setminus B_2} (r_1(X) - p(X))$, $r_1(B_1) + r_2(B_2) = \max\{r_1(X) + r_2(Y) \mid X, Y \subseteq S, X \cap Y = \emptyset\}$.

In the conjecture, if buyer i comes to a shop first, then she chooses an arbitrary bundle B_i that maximizes her utility $r_i - p$, and the second buyer chooses a best bundle in $S \setminus B_i$. The requirements mean that any choice of B_i results in an allocation maximizing the social welfare. Thus, whoever comes first, we can achieve the optimal social welfare.

Previous work The notion of Walrasian equilibrium dates back to 1874 [38], originally defined for divisible goods. In their analysis of the matching problem, Kelso and Crawford [25] introduced the so-called gross substitutes condition, and showed the existence of Walrasian prices for gross substitutes valuations. Gul and Stacchetti [22] later verified that, in a sense, this condition is necessary to ensure the existence of a Walrasian equilibrium.³

It was first observed by Cohen-Addad et al. [7] and Hsu et al. [23] that Walrasian prices are not sufficient to control the market as ties must be broken in a coordinated fashion that is consistent with maximizing social welfare. A natural idea for resolving this issue would be trying to find Walrasian prices where ties do not occur. However, Hsu et al. showed that minimal Walrasian prices always induce ties. Even more, Cohen-Addad et al. proved that no static prices can give more than 2/3 of the social welfare when buyers arrive sequentially. As a workaround, they proposed a dynamic pricing scheme for matching markets (i.e., unit-demand valuations), where the prices are updated between buyer-arrivals based upon the current inventory without knowing the identity of the next buyer. On the negative side, they presented a market with coverage valuations where Walrasian prices do exist, but no dynamic pricing scheme can achieve the optimal social welfare. Meanwhile, Hsu et al. showed that, under certain conditions, minimal Walrasian equilibrium prices induce low over-demand and high welfare. Recently, Berger et al. [1] considered markets beyond unit-demand valuations, and gave a characterization of all optimal allocations in multi-demand markets. Based on this, they provided a polynomial-time algorithm for finding optimal dynamic prices up to three multi-demand buyers.

To overcome the limitations of Walrasian equilibrium, Feldman et al. [16] proposed a relaxation called combinatorial Walrasian equilibrium in which the seller can partition the items into indivisible bundles prior to sale, and they provided an algorithm that determines bundle prices obtaining at least half of the optimal social welfare.

Another line of research concentrated on posted-price mechanisms in online settings. As alternatives to optimal auctions, Blumrosen and Holenstein [2] studied posted-price mechanisms and dynamic auctions in Bayesian settings under the objective of maximizing revenue. They gave a characterization of the optimal revenue for general distributions, and provided algorithms that achieve the optimal solution. Chawla et al. [3, 4] developed a theory of sequential posted-price mechanisms, and provided constant-factor approximation algorithms for several multi-dimensional multi-unit auction problems and generalizations to matroid feasibility constraints. In [15], Feldman et al. verified the existence of prices that, in expectation, achieve at least half of the optimal

³The simplest example of gross substitutes valuations are unit demand preferences, when each agent can enjoy at most one item. Gul and Stacchetti showed that gross substitutes preferences form the largest set containing unit demand preferences for which an existence theorem can be obtained.

social welfare for fractionally subadditive valuations, a class that includes all submodular functions. Dütting et al. [9, 10] provided a general framework for posted-price mechanisms in Bayesian settings. Chawla et al. [5] showed that static, anonymous bundle pricing mechanisms are useful when buyers’ preferences have complementarities. Ezra et al. [14] provided upper and lower bounds on the largest fraction of the optimal social welfare that can be guaranteed with static prices for several classes of valuations, such as submodular, XOS, or subadditive. A setting related to online bipartite matching, called the Max-Min Greedy matching, was considered in [12].

Our results In the present paper, we concentrate on combinatorial markets with two buyers having matroid rank valuations, where the matroid corresponding to buyer i is denoted by $M_i = (S, \mathcal{B}_i)$ for $i = 1, 2$. Since this setting is reduced to Conjecture 1, in which each buyer has to buy a set of items that forms a basis of a matroid, we focus on Conjecture 1.

While Conjecture 1 remains open in general, we give polynomial-time⁴ algorithms for two important special cases. In the first one, one of the matroids is a partition matroid. Although partition matroids have relatively simple structure, finding the proper price vector p is non-trivial even in this seemingly simple case.

Theorem 1. *Let M_1 be a partition matroid with partition classes of size at most 2 and with all-ones upper bound on the partition classes, and let M_2 be an arbitrary matroid. Then Conjectures 1 and 2 hold, and a price vector p satisfying the conditions can be computed in polynomial time.*

Next we consider strongly base orderable matroids, a class of matroids with distinctive structural properties. Roughly, in a strongly base orderable matroid, for any pair of bases, there exists a bijection between them satisfying a certain property (see Section 2 for the formal definition). Note that various matroids appearing in combinatorial and graph optimization problems belong to this class, such as partition, laminar, transversal matroids, or more generally, gammoids.

Theorem 2. *If both M_1 and M_2 are strongly base orderable, then Conjectures 1 and 2 hold. Furthermore, a price vector p satisfying the conditions can be computed in polynomial time if, for any pair of bases, the bijection between them can be computed in polynomial time.*

Another contribution of this paper is to show the equivalence between Conjecture 1 and its weighted counterpart as below.

Conjecture 3. *For $i \in \{1, 2\}$, let $M_i = (S, \mathcal{B}_i)$ be a matroid and $w_i : S \rightarrow \mathbb{R}$ be a weight function. Assume that there exist disjoint bases $B_1 \in \mathcal{B}_1$ and $B_2 \in \mathcal{B}_2$ with $B_1 \cup B_2 = S$. Then, there exists a function $p : S \rightarrow \mathbb{R}$ satisfying the following conditions.*

1. *For any $B_1 \in \arg \max_{X \in \mathcal{B}_1} (w_1(X) - p(X))$, we have that B_1 is a maximizer of $w_1(X) + w_2(S \setminus X)$ subject to $X \in \mathcal{B}_1$ and $S \setminus X \in \mathcal{B}_2$.*
2. *For any $B_2 \in \arg \max_{X \in \mathcal{B}_2} (w_2(X) - p(X))$, we have that B_2 is a maximizer of $w_1(S \setminus X) + w_2(X)$ subject to $S \setminus X \in \mathcal{B}_1$ and $X \in \mathcal{B}_2$.*

Clearly, Conjecture 1 is a special case of Conjecture 3; this follows easily by setting $w_1 \equiv w_2 \equiv 0$. Somewhat surprisingly, the reverse implication also holds for arbitrary matroids.

⁴In matroid algorithms, it is usually assumed that the matroids are accessed through independence oracles, and the complexity of an algorithm is measured by the number of oracle calls and other conventional elementary steps.

Theorem 3. *If Conjecture 1 is true, then Conjecture 3 is also true.*

More generally, we prove that Theorem 3 can be generalized to the case with gross substitutes valuations, i.e., M^\natural -concave functions. See Theorem 19 in Section 6 for the details.

Based on Theorem 3 and the properties of partition and strongly base orderable matroids, we have the following corollaries.

Corollary 4. *Let M_1 be a partition matroid with partition classes of size at most 2 and with all-ones upper bound on the partition classes, and let M_2 be an arbitrary matroid. Then Conjecture 3 holds, and a price vector p satisfying the conditions can be computed in polynomial time.*

Corollary 5. *If both M_1 and M_2 are strongly base orderable, then Conjecture 3 holds. Furthermore, a price vector p satisfying the conditions can be computed in polynomial time if, for any pair of bases, the bijection between them can be computed in polynomial time.*

Paper organization The rest of the paper is organized as follows. Basic definitions and notation are given in Section 2. Theorems 1 and 2 are proved in Sections 3 and 4, respectively. The connection between unweighted and weighted variants of the problem is discussed in Section 5. The reduction technique is extended to gross substitutes valuations in Section 6. We conclude the paper in Section 7.

2 Preliminaries

Basic notation The sets of reals, non-negative reals, integers, and non-negative integers are denoted by \mathbb{R} , \mathbb{R}_+ , \mathbb{Z} , and \mathbb{Z}_+ , respectively. Let S be a finite set. Given a subset $B \subseteq S$ and elements $x, y \in S$, we write $B - x + y$ for short to denote the set $(B \setminus \{x\}) \cup \{y\}$. The *symmetric difference* of two sets X and Y is $X \Delta Y := (X \setminus Y) \cup (Y \setminus X)$. For a function $f : S \rightarrow \mathbb{R}$, we use $f(X) := \sum_{x \in X} f(x)$. For two vectors $x, y \in \mathbb{R}^S$, we denote $x \cdot y := \sum_{s \in S} x(s)y(s)$.

Matroids and matroid intersection Matroids were introduced as an abstract generalization of linear independence in vector spaces [33, 39]. A *matroid* M is a pair (S, \mathcal{I}) where S is the *ground set* of the matroid and $\mathcal{I} \subseteq 2^S$ is the family of *independent sets* satisfying the *independence axioms*: (I1) $\emptyset \in \mathcal{I}$, (I2) $X \subseteq Y \in \mathcal{I} \Rightarrow X \in \mathcal{I}$, and (I3) $X, Y \in \mathcal{I}$, $|X| < |Y| \Rightarrow \exists e \in Y \setminus X$ s.t. $X + e \in \mathcal{I}$. A *loop* is an element that is non-independent on its own. The *rank* of a set $X \subseteq S$ is the maximum size of an independent set contained in X , and is denoted by $r(X)$. Here r is called the *rank function* of M . Maximal independent sets of M are called *bases* and their set is denoted by \mathcal{B} . Alternatively, matroids can be defined through the *basis axioms*: (B1) $\mathcal{B} \neq \emptyset$, and (B2) $B_1, B_2 \in \mathcal{B}, x \in B_1 \setminus B_2 \Rightarrow \exists y \in B_2 \setminus B_1$ s.t. $B_1 - x + y \in \mathcal{B}$. In this paper, a matroid is denoted by a pair (S, \mathcal{B}) , where S is a ground set and \mathcal{B} is a base family.

For a matroid $M = (S, \mathcal{B})$ and for $T \subseteq S$, *deleting* T gives a matroid M' on the ground set $S \setminus T$ such that a subset of $S \setminus T$ is independent in M' if and only if it is independent in M . For $T \subseteq S$, *contracting* T gives a matroid M' on the ground set $S \setminus T$ whose rank function is $r'(X) = r(X \cup T) - r(T)$, where r is the rank function of M . *Adding a parallel copy* of an element $s \in S$ gives a new matroid $M' = (S', \mathcal{B}')$ on ground set $S' = S + s'$ where $\mathcal{B}' = \{X \subseteq S' : \text{either } X \in \mathcal{B}, \text{ or } s \notin X, s' \in X \text{ and } X - s' + s \in \mathcal{B}\}$. The *direct sum* $M_1 \oplus M_2$ of matroids $M_1 = (S_1, \mathcal{B}_1)$ and $M_2 = (S_2, \mathcal{B}_2)$ on disjoint ground sets is a matroid $M = (S_1 \cup S_2, \mathcal{B})$ whose bases are the

disjoint unions of a basis M_1 and a basis of M_2 . The *sum* or *union* $M_1 + M_2$ of $M_1 = (S, \mathcal{B}_1)$ and $M_2 = (S, \mathcal{B}_2)$ on the same ground set is a matroid $M = (S, \mathcal{B})$ whose independent sets are the disjoint unions of an independent set of M_1 and an independent set of M_2 .

For a basis $B \in \mathcal{B}$, let us consider the bipartite graph $G = (S, E[B])$ defined by $E[B] := \{(x, y) \mid x \in B, y \in S \setminus B, B - x + y \in \mathcal{B}\}$. Krogdahl [26, 27, 28] verified the following statement (see also [35, Theorem 39.13]).

Theorem 6 (Krogdahl). *Let $M = (S, \mathcal{B})$ be a matroid and let $B \in \mathcal{B}$. Let $B' \subseteq S$ be such that $|B| = |B'|$ and $E[B]$ contains a unique perfect matching on $B \triangle B'$. Then $B' \in \mathcal{B}$.*

In the *weighted matroid intersection problem*, we are given two matroids $M_1 = (S, \mathcal{B}_1)$ and $M_2 = (S, \mathcal{B}_2)$ on the same ground set together with a weight function $w : S \rightarrow \mathbb{R}$, and the goal is to find a common basis maximizing $w(B)$, that is, $B \in \arg \max\{w(B) \mid B \in \mathcal{B}_1 \cap \mathcal{B}_2\}$. The celebrated weight-splitting theorem of Frank [17] gives a min-max relation for the weighted matroid intersection.

Theorem 7 (Frank). *The maximum w -weight of a common basis of $M_1 = (S, \mathcal{B}_1)$ and $M_2 = (S, \mathcal{B}_2)$ is equal to the minimum of $\max\{w_1(B) \mid B \in \mathcal{B}_1\} + \max\{w_2(B) \mid B \in \mathcal{B}_2\}$ subject to $w = w_1 + w_2$. In particular, for an optimal weight-splitting $w = w_1 + w_2$, it holds that $\arg \max\{w(B) \mid B \in \mathcal{B}_1 \cap \mathcal{B}_2\} = \arg \max\{w_1(B) \mid B \in \mathcal{B}_1\} \cap \arg \max\{w_2(B) \mid B \in \mathcal{B}_2\}$.*

A *k-uniform matroid* is a matroid $M = (S, \mathcal{B})$ where $\mathcal{B} = \{X \subseteq S \mid |X| = k\}$ for some $k \in \mathbb{Z}_+$. A *partition matroid* $M = (S, \mathcal{B})$ is the direct sum of uniform matroids, or in other words, $\mathcal{B} = \{X \subseteq S \mid |X \cap S_i| = k_i \text{ for } i = 1, \dots, q\}$ for some partition $S = S_1 \cup \dots \cup S_q$ of S and $k_i \in \mathbb{Z}_+$ for $i = 1, \dots, q$. Each S_i is called a *partition class*. In the paper, we will work with partition matroids satisfying $|S_i| \leq 2$ and $k_i = 1$ for $i = 1, \dots, q$.

For further details on matroids and the matroid intersection problem, we refer the reader to [34, 35].

Dual matroids The *dual* of a matroid $M = (S, \mathcal{B})$ is the matroid $M^* = (S, \mathcal{B}^*)$ where $\mathcal{B}^* = \{B^* \subseteq S \mid S \setminus B^* \in \mathcal{B}\}$. Given one of the standard oracles for M , the same oracle can be constructed for M^* as well.

We now rephrase Conjecture 1 by using dual matroids. Suppose that M_1 and M_2 are matroids as in Conjecture 1 and let $M_2^* = (S, \mathcal{B}_2^*)$ be the dual matroid of M_2 . Then, we can see that $S \setminus B_1 \in \mathcal{B}_2$ is equivalent to $B_1 \in \mathcal{B}_2^*$, and $B_2 \in \arg \min_{X \in \mathcal{B}_2} p(X)$ is equivalent to $S \setminus B_2 \in \arg \max_{X \in \mathcal{B}_2^*} p(X)$. Therefore, by replacing M_2 and $S \setminus B_2$ with M_2^* and B_2 , respectively, Conjecture 1 is equivalent to the following conjecture.

Conjecture 4. *Let $M_1 = (S, \mathcal{B}_1)$ and $M_2 = (S, \mathcal{B}_2)$ be matroids with a common ground set S such that there exists a common basis $B \in \mathcal{B}_1 \cap \mathcal{B}_2$. Then, there exists a function $p : S \rightarrow \mathbb{R}$ satisfying the following conditions.*

1. *For any $B_1 \in \arg \min_{X \in \mathcal{B}_1} p(X)$, it holds that $B_1 \in \mathcal{B}_2$.*
2. *For any $B_2 \in \arg \max_{X \in \mathcal{B}_2} p(X)$, it holds that $B_2 \in \mathcal{B}_1$.*

Conjecture 4 bears a lot of similarities with the problem of packing common bases in the intersection of two matroids. If M_1 and M_2 share two disjoint common bases then setting the prices low on one of them and high on the other gives a desired p . If S can be partitioned into two disjoint bases in both M_1 and M_2 , then the statement may be reminiscent of Rota's famous conjecture concerning rearrangements of bases [24].

Strongly base orderable matroids A matroid $M = (S, \mathcal{B})$ is *strongly base orderable* if for any two bases $B_1, B_2 \in \mathcal{B}$, there exists a bijection $f : B_1 \rightarrow B_2$ such that $(B_1 \setminus X) \cup f(X) \in \mathcal{B}$ for any $X \subseteq B_1$, where we denote $f(X) := \{f(e) \mid e \in X\}$. Davies and McDiarmid [8] observed the following (see also [35, Theorem 42.13]).

Theorem 8 (Davies and McDiarmid). *Let $M_1 = (S, \mathcal{B}_1)$ and $M_2 = (S, \mathcal{B}_2)$ be strongly base orderable matroids. If $X \subseteq S$ can be partitioned into k bases in both M_1 and M_2 , then X can be partitioned into k common bases. Furthermore, such k common bases can be computed in polynomial time if the bijection f can be computed in polynomial time for any pair of bases.*

The following technical lemma about strongly base orderable matroids will be used in the proof of Corollary 5.

Lemma 9. *Let $M = (S, \mathcal{B})$ be a strongly base orderable matroid, $q : S \rightarrow \mathbb{R}$ be a function, and define a matroid $\hat{M} = (S, \hat{\mathcal{B}})$ by $\hat{\mathcal{B}} = \arg \max_{X \in \mathcal{B}} q(X)$. Then \hat{M} is strongly base orderable.*

Proof. Let $B_1, B_2 \in \hat{\mathcal{B}}$. Since both B_1 and B_2 are bases of $M = (S, \mathcal{B})$, there exists a bijection $f : B_1 \rightarrow B_2$ such that $(B_1 \setminus X) \cup f(X) \in \mathcal{B}$ for any $X \subseteq B_1$. Since $q(B_1) \geq q((B_1 \setminus X) \cup f(X))$ for any $X \subseteq B_1$ by $B_1 \in \hat{\mathcal{B}}$, it holds that $q(X) \geq q(f(X))$. In particular, $q(x) \geq q(f(x))$ for any $x \in B_1$. Since $B_2 \in \hat{\mathcal{B}}$, we obtain $q(B_1) = q(B_2) = q(f(B_1))$, which shows that $q(x) = q(f(x))$ for any $x \in B_1$. Therefore, $q(B_1) = q((B_1 \setminus X) \cup f(X))$ for any $X \subseteq B_1$, and hence $(B_1 \setminus X) \cup f(X) \in \hat{\mathcal{B}}$. This shows that \hat{M} is strongly base orderable. \square

Market model In a combinatorial market, we are given a set S of *indivisible items* and a set J of *buyers*. Each buyer $i \in J$ has a *valuation function* $v_i : 2^S \rightarrow \mathbb{R}$ that describes the buyer's preferences over the subsets of items. Given *prices* $p : S \rightarrow \mathbb{R}$, the *utility* of buyer $i \in J$ for a subset $X \subseteq S$ is defined by $u_i(X) = v_i(X) - p(X)$. The buyers arrive in an undetermined order, and the next buyer always picks a subset of items that maximizes her utility. The goal is to set the prices in such a way that no matter which buyer arrives next, the final allocation of items maximizes the social welfare. In a dynamic pricing scheme, the prices can be updated between buyer arrivals based on the remaining sets of items and buyers.

We focus on the case of two buyers with matroid rank functions as valuations. Let $M_1 = (S, \mathcal{B}_1)$ and $M_2 = (S, \mathcal{B}_2)$ be matroids with rank functions r_1 and r_2 , respectively. The valuation of agent i is r_i for $i = 1, 2$. The valuations are accessed through one of the standard matroid oracles (e.g. independence or rank oracle). As described in the introduction, this setting can be reduced to the case in which each buyer always chooses a basis that maximizes her utility, that is, Conjecture 2 can be reduced to Conjecture 1.

Lemma 10. *If Conjecture 1 is true, then Conjecture 2 is also true.*

Proof. Let $M_1 = (S, \mathcal{B}_1)$ and $M_2 = (S, \mathcal{B}_2)$ be matroids as in Conjecture 2 and let $\hat{B}_1 \in \mathcal{B}_1$ and $\hat{B}_2 \in \mathcal{B}_2$ be a pair of bases that maximizes $|\hat{B}_1 \cup \hat{B}_2|$. For $i \in \{1, 2\}$, let M'_i be the matroid obtained from M_i by deleting $S \setminus (\hat{B}_1 \cup \hat{B}_2)$ and contracting $\hat{B}_1 \cap \hat{B}_2$. Then, $M'_1 = (S', \mathcal{B}'_1)$ and $M'_2 = (S', \mathcal{B}'_2)$ are matroids with a common ground set $S' := (\hat{B}_1 \cup \hat{B}_2) \setminus (\hat{B}_1 \cap \hat{B}_2)$ such that there exist disjoint bases $\hat{B}_1 \setminus \hat{B}_2 \in \mathcal{B}'_1$ and $\hat{B}_2 \setminus \hat{B}_1 \in \mathcal{B}'_2$ whose union is S' . Hence, by assuming that Conjecture 1 is true, there exists a price vector $p' : S' \rightarrow \mathbb{R}$ with the following conditions.

1. For any $B'_1 \in \arg \min_{X \in \mathcal{B}'_1} p'(X)$, it holds that $S' \setminus B'_1 \in \mathcal{B}'_2$.

2. For any $B'_2 \in \arg \min_{X \in \mathcal{B}'_2} p'(X)$, it holds that $S' \setminus B'_2 \in \mathcal{B}'_1$.

We observe that we can modify the price vector p' so that $0 < p'(s) < 1$ for every $s \in S'$, by replacing $p'(s)$ with $\alpha \cdot p'(s) + \beta$ for some $\alpha > 0$ and $\beta \in \mathbb{R}$. By using such a function p' , define $p : S \rightarrow \mathbb{R}$ by

$$p(s) = \begin{cases} p'(s) & \text{if } s \in S', \\ 0 & \text{if } s \in \hat{B}_1 \cap \hat{B}_2, \\ 1 & \text{if } s \in S \setminus (\hat{B}_1 \cup \hat{B}_2). \end{cases}$$

For $B_1 \in \arg \max_{X \subseteq S} (r_1(X) - p(X))$, the definition of p shows that $B_1 = B'_1 \cup (\hat{B}_1 \cap \hat{B}_2)$ for some $B'_1 \in \arg \min_{X \in \mathcal{B}'_1} p'(X)$. Since this implies $S' \setminus B'_1 \in \mathcal{B}'_2$, it holds that $S' \setminus B'_1$ is a maximal independent set of M_1 in $S \setminus B_1$ by the maximality of $|\hat{B}_1 \cup \hat{B}_2|$. Therefore, if $B_2 \in \arg \max_{Y \subseteq S \setminus B_1} (r_2(Y) - p(Y))$, then $B_2 = S' \setminus B'_1$ and hence

$$\begin{aligned} r_1(B_1) + r_2(B_2) &= |B'_1| + |\hat{B}_1 \cap \hat{B}_2| + |S' \setminus B'_1| = |\hat{B}_1 \cup \hat{B}_2| \\ &= \max\{r_1(X) + r_2(Y) \mid X, Y \subseteq S, X \cap Y = \emptyset\}, \end{aligned}$$

which shows the first requirement of Conjecture 2. The same argument works for $B_2 \in \arg \max_{X \subseteq S} (r_2(X) - p(X))$. Therefore, p satisfies the requirements in Conjecture 2. \square

Note that a pair of bases $\hat{B}_1 \in \mathcal{B}_1$ and $\hat{B}_2 \in \mathcal{B}_2$ maximizing $|\hat{B}_1 \cup \hat{B}_2|$ can be computed in polynomial time by applying a matroid intersection algorithm to M_1 and M_2^* . Note also that the price vector p obtained in the above proof is not necessarily a Walrasian price.

We can consider a weighted variant of Conjecture 1 in which we are given weight functions $w_1 : S \rightarrow \mathbb{R}$ and $w_2 : S \rightarrow \mathbb{R}$. For a buyer $i \in \{1, 2\}$ and for a basis $X \in \mathcal{B}_i$, the valuation $v_i(X)$ is defined as $w_i(X)$. Each buyer chooses a basis that maximizes her utility. Note that choosing a basis is a hard constraint, and hence we do not have to define $v_i(X)$ for $X \notin \mathcal{B}_i$. The goal is to find a price vector p that achieves the optimal social welfare $\max\{w_1(X) + w_2(S \setminus X) \mid X \in \mathcal{B}_1, S \setminus X \in \mathcal{B}_2\}$.

Recently, Berger et al. [1] investigated the existence of optimal dynamic pricing schemes for k -demand valuations. A valuation $v : 2^S \rightarrow \mathbb{R}_+$ is k -demand if $v(X) = \max\{\sum_{s \in Z} v(s) \mid Z \subseteq X, |Z| \leq k\}$. Although this setting is similar to our weighted variant for k -uniform matroids, our results do not directly generalize their work because of our assumption on the buyers' choices.

3 Partition matroids

The aim of this section is to prove the existence of a required price vector p for instances where M_1 is a partition matroid of special type.

Theorem 1. *Let M_1 be a partition matroid with partition classes of size at most 2 and with all-ones upper bound on the partition classes, and let M_2 be an arbitrary matroid. Then Conjectures 1 and 2 hold, and a price vector p satisfying the conditions can be computed in polynomial time.*

Proof. Since Conjectures 1 and 4 are equivalent by replacing M_2 with its dual M_2^* , we show Conjecture 4. Let $M_1 = (S, \mathcal{B}_1)$ be a partition matroid defined by partition $S = S_1 \cup \dots \cup S_q$ where $|S_i| \leq 2$ for $i = 1, \dots, q$, that is, $\mathcal{B}_1 = \{X \subseteq S \mid |X \cap S_i| = 1 \text{ for } i = 1, \dots, q\}$. Let $M_2 = (S, \mathcal{B}_2)$ be an arbitrary matroid such that M_1 and M_2 have a common basis.

Let $B_1 \in \mathcal{B}_1 \cap \mathcal{B}_2$ be an arbitrary common basis. Take another common basis $B_2 \in \mathcal{B}_1 \cap \mathcal{B}_2$ (possibly $B_2 = B_1$) such that $|B_1 \cap B_2|$ is minimized. We consider a bipartite digraph $D = (V, E)$ defined by

$$\begin{aligned} V &= (B_1 \cap B_2) \cup (S \setminus (B_1 \cup B_2)), \\ E &= \{(x, y) \mid x \in B_1 \cap B_2, y \in S \setminus (B_1 \cup B_2), B_1 - x + y \in \mathcal{B}_1\} \\ &\quad \cup \{(y, x) \mid x \in B_1 \cap B_2, y \in S \setminus (B_1 \cup B_2), B_2 - x + y \in \mathcal{B}_2\}. \end{aligned} \tag{1}$$

Claim 11. *The digraph D is acyclic.*

Proof. Let $x \in B_1 \cap B_2$ and $y \in S \setminus (B_1 \cup B_2)$. As M_1 is defined on a partition consisting of classes of size at most 2, $B_1 - x + y \in \mathcal{B}_1$ implies that $\{x, y\}$ is one of the partition classes. This implies that $B_1 - x + y \in \mathcal{B}_1$ if and only if $B_2 + x - y \in \mathcal{B}_1$.

Now suppose to the contrary that D contains a dicycle. Choose a dicycle C with the smallest number of vertices, which implies that C has no chord. Then, $B'_2 := B_2 \Delta V(C)$ is a common basis of M_1 and M_2 by the above observation and Theorem 6. Since $|B_1 \cap B'_2| < |B_1 \cap B_2|$, this contradicts that $|B_1 \cap B_2|$ is minimized. \square

Let $n = |S|$. We now consider a function $p : S \rightarrow \mathbb{R}$ satisfying the following: $p(x) := 0$ for $x \in B_1 \setminus B_2$, $p(x) := n + 1$ for $x \in B_2 \setminus B_1$, $p(x)$ are distinct values in $\{1, 2, \dots, n\}$ for $x \in V$, and $p(x) < p(y)$ for $(x, y) \in E$. Note that such a function exists by Claim 11, which can be found easily by the topological sorting. In what follows, we show that p satisfies that $\arg \min_{X \in \mathcal{B}_1} p(X) = \{B_1\}$ and $\arg \max_{X \in \mathcal{B}_2} p(X) = \{B_2\}$.

Claim 12. $\arg \min_{X \in \mathcal{B}_1} p(X) = \{B_1\}$ and $\arg \max_{X \in \mathcal{B}_2} p(X) = \{B_2\}$.

Proof. For a non-negative integer k , let $S_k := \{x \in S \mid p(x) \leq k\}$ and let I_k be a minimizer of $p(X)$ subject to X being a maximal independent set of M_1 and $X \subseteq S_k$. Note that I_k can be computed by a greedy algorithm. Since S_n contains a basis B_1 , the greedy algorithm chooses no element in $B_2 \setminus B_1$, which means that $I_k \cap (B_2 \setminus B_1) = \emptyset$ for every k . We also note that I_k is uniquely determined for each k , since $p(x)$'s are distinct for $x \in V$.

We show that $I_k = B_1 \cap S_k$ for every k by induction on k . Since $I_0 = B_1 \setminus B_2$, it is obvious that $I_0 = B_1 \cap S_0$. Fix $k \geq 1$ and assume that $I_{k-1} = B_1 \cap S_{k-1}$. Then, we have the following.

- If there exists $x \in B_1 \cap B_2$ with $p(x) = k$, then $I_k = I_{k-1} + x$, and hence $I_k = B_1 \cap S_k$.
- Suppose that there exists $y \in S \setminus (B_1 \cup B_2)$ with $p(y) = k$. We show that $I_{k-1} + y$ is not independent in M_1 . Suppose to the contrary that $I_{k-1} + y$ is independent. Then, there exists $x \in B_1 \setminus I_{k-1}$ such that $B_1 - x + y \in \mathcal{B}_1$, and hence $(x, y) \in E$. By the choice of p , we obtain $p(x) < p(y)$, i.e., $x \in S_{k-1}$. This contradicts $x \in B_1 \setminus I_{k-1}$, because $S_{k-1} \cap (B_1 \setminus I_{k-1}) = \emptyset$ by the induction hypothesis. Therefore, $I_{k-1} + y$ is not independent in M_1 , which shows that $I_k = I_{k-1}$ and $I_k = B_1 \cap S_k$.
- If there exists no $x \in V$ with $p(x) = k$, then $I_k = I_{k-1}$, and hence $I_k = B_1 \cap S_k$.

Therefore, $I_k = B_1 \cap S_k$ holds for every k by induction. This shows that $I_{n+1} = B_1 \cap S_{n+1} = B_1$, and hence $\arg \min_{X \in \mathcal{B}_1} p(X) = \{I_{n+1}\} = \{B_1\}$.

By a similar argument, we obtain $\arg \max_{X \in \mathcal{B}_2} p(X) = \{B_2\}$. \square

Since $B_1, B_2 \in \mathcal{B}_1 \cap \mathcal{B}_2$, this claim shows that p satisfies the requirements in Conjecture 4. Thus, Conjecture 4 holds, and hence Conjecture 1 also holds.

This together with Lemma 10 shows that Conjecture 2 also holds. Note that, in the proof of Lemma 10, we modify given matroids by deleting and contracting some elements, but this modification does not affect the assumption on M_1 . That is, if M_1 is a partition matroid with partition classes of size at most 2 and with all-ones upper bound on the partition classes, then the obtained matroid M'_1 is also a partition matroid of this type. \square

Remark 13. Note that in the proof of Theorem 1, we fixed the basis $B_1 \in \mathcal{B}_1$ arbitrarily. That is, for any $B_1 \in \mathcal{B}_1$, the optimal price vector p can be set in such a way that the maximum utility of the buyer corresponding to M_1 is attained on B_1 . It is not difficult to see that the analogous statement holds for any basis $B_2 \in \mathcal{B}_2$.

Remark 14. Even when \mathcal{B}_1 is a base family of a partition matroid as in Theorem 1, if \mathcal{B}_2 is an arbitrary set family of S , then the requirements in Conjecture 1 do not necessarily hold. To see this, suppose that $S = \{1, 2, 3, 4\}$, $\mathcal{B}_1 = \{\{1, 3\}, \{1, 4\}, \{2, 3\}, \{2, 4\}\}$, and $\mathcal{B}_2 = \{\{2, 4\}, \{1, 2\}, \{3, 4\}\}$. Then, $(B_1, B_2) = (\{1, 3\}, \{2, 4\})$ is a unique pair of disjoint sets such that $B_1 \in \mathcal{B}_1$, $B_2 \in \mathcal{B}_2$, and $B_1 \cup B_2 = S$. If p satisfies the requirements in Conjecture 1, then $p(1) < p(2)$ and $p(3) < p(4)$ hold by the first requirement and $p(4) < p(1)$ and $p(2) < p(3)$ hold by the second requirement. This shows that such p does not exist.

4 Strongly base orderable matroids

In this section, we show that Conjectures 1 and 2 hold for strongly base orderable matroids. The proof is based on a similar approach to that of Theorem 1. Nevertheless, there are small but crucial differences.

Theorem 2. *If both M_1 and M_2 are strongly base orderable, then Conjectures 1 and 2 hold. Furthermore, a price vector p satisfying the conditions can be computed in polynomial time if, for any pair of bases, the bijection between them can be computed in polynomial time.*

Proof. In order to show Conjecture 1, we first show Conjecture 4 under the assumption that M_1 and M_2 are strongly base orderable. Let $M_1 = (S, \mathcal{B}_1)$ and $M_2 = (S, \mathcal{B}_2)$ be strongly base orderable matroids that have a common basis. We take two common bases $B_1, B_2 \in \mathcal{B}_1 \cap \mathcal{B}_2$ (possibly $B_1 = B_2$) such that $|B_1 \cap B_2|$ is minimized. For each element $x \in S$, we add a parallel copy x' of x to the matroid M_i and denote the matroid thus obtained by $M_i^+ = (S \cup S', \mathcal{B}_i^+)$ for $i \in \{1, 2\}$. We denote $X' := \{x' \mid x \in X\}$ for $X \subseteq S$. Let $2M_i^+ = (S \cup S', 2\mathcal{B}_i^+)$ be the sum of two copies of M_i^+ . As M_i^+ clearly has two disjoint bases, we have $2\mathcal{B}_i^+ := \{Y_1 \cup Y_2 \mid Y_1 \text{ and } Y_2 \text{ are disjoint bases of } M_i^+\}$.

Claim 15. *For $i \in \{1, 2\}$, $2M_i^+$ is a strongly base orderable matroid.*

Proof. Fix $i \in \{1, 2\}$. We can easily see that M_i^+ is strongly base orderable. Suppose that we are given two bases $X_1, X_2 \in 2\mathcal{B}_i^+$, and suppose also that $X_1 = Y_1^1 \cup Y_1^2$ and $X_2 = Y_2^1 \cup Y_2^2$, where $Y_1^1, Y_1^2, Y_2^1, Y_2^2 \in \mathcal{B}_i^+$. Since M_i^+ is strongly base orderable, for $j \in \{1, 2\}$, there exists a bijection $f_j : Y_1^j \rightarrow Y_2^j$ such that $(Y_1^j \setminus X) \cup f_j(X) \in \mathcal{B}_i^+$ for any $X \subseteq Y_1^j$. Then, f_1 and f_2 naturally define a bijection $f : X_1 \rightarrow X_2$ such that $(X_1 \setminus X) \cup f(X) \in 2\mathcal{B}_i^+$ for any $X \subseteq X_1$. This shows that $2M_i^+$ is strongly base orderable. \square

Let $X_0 := (B_1 \cup B_2) \cup (B_1 \cap B_2)'$. Then, X_0 is a common basis of $2M_1^+$ and $2M_2^+$. We consider a bipartite digraph $D^+ = (V, E^+)$ defined by

$$\begin{aligned} V &= (B_1 \cap B_2) \cup (S \setminus (B_1 \cup B_2)), \\ E^+ &= \{(x, y) \mid x \in B_1 \cap B_2, y \in S \setminus (B_1 \cup B_2), X_0 - x + y \in 2\mathcal{B}_1^+\} \\ &\quad \cup \{(y, x) \mid x \in B_1 \cap B_2, y \in S \setminus (B_1 \cup B_2), X_0 - x + y \in 2\mathcal{B}_2^+\}. \end{aligned}$$

Claim 16. *The digraph D^+ is acyclic.*

Proof. Suppose to the contrary that D^+ contains a dicycle. Choose a dicycle C with the smallest number of vertices, which implies that C has no chord. Then, $X_0 \triangle V(C)$ is a common basis of $2M_1^+$ and $2M_2^+$ by Theorem 6. By Theorem 8 and Claim 15, $X_0 \triangle V(C)$ can be partitioned into two common bases of M_1^+ and M_2^+ . Let \tilde{B}_1 and \tilde{B}_2 be the sets in S corresponding to these common bases. Then, $\tilde{B}_1, \tilde{B}_2 \in \mathcal{B}_1 \cap \mathcal{B}_2$ and $|\tilde{B}_1 \cap \tilde{B}_2| < |B_1 \cap B_2|$. This contradicts that $|B_1 \cap B_2|$ is minimized. \square

We now consider the digraph $D = (V, E)$ defined by (1). For $x \in B_1 \cap B_2$ and $y \in S \setminus (B_1 \cup B_2)$, we observe that $B_1 - x + y \in \mathcal{B}_1$ implies $X_0 - x + y \in 2\mathcal{B}_1^+$ and $B_2 - x + y \in \mathcal{B}_2$ implies $X_0 - x + y \in 2\mathcal{B}_2^+$. This shows that D is a subgraph of D^+ , and hence D is acyclic by Claim 16. Therefore, we can find a function $p : S \rightarrow \mathbb{R}$ such that $p(x) := 0$ for $x \in B_1 \setminus B_2$, $p(x) := |S| + 1$ for $x \in B_2 \setminus B_1$, $p(x)$ are distinct values in $\{1, 2, \dots, |S|\}$ for $x \in V$, and $p(x) < p(y)$ for $(x, y) \in E$. Then, Claim 12 shows that $\arg \min_{X \in \mathcal{B}_1} p(X) = \{B_1\}$ and $\arg \max_{X \in \mathcal{B}_2} p(X) = \{B_2\}$. Since $B_1, B_2 \in \mathcal{B}_1 \cap \mathcal{B}_2$, p satisfies the requirements in Conjecture 4. Thus, Conjecture 4 holds.

This proof can be converted to a polynomial-time algorithm for computing p as follows. We first pick up two arbitrary common bases $B_1, B_2 \in \mathcal{B}_1 \cap \mathcal{B}_2$ and construct a digraph D^+ as above. If D^+ is acyclic, then we can find an appropriate function p . Otherwise, the proof of Claim 16 shows that we can find $\tilde{B}_1, \tilde{B}_2 \in \mathcal{B}_1 \cap \mathcal{B}_2$ with $|\tilde{B}_1 \cap \tilde{B}_2| < |B_1 \cap B_2|$. Then, we update $B_i \leftarrow \tilde{B}_i$ for $i \in \{1, 2\}$, construct D^+ , and repeat this procedure. Since $|B_1 \cap B_2|$ decreases monotonically, this procedure is executed at most $|S|$ times.

Recall that Conjectures 1 and 4 are equivalent by replacing M_2 with M_2^* . Since M_2 is strongly base orderable if and only if M_2^* is strongly base orderable, Conjecture 1 also holds for strongly base orderable matroids.

This together with Lemma 10 shows that Conjecture 2 also holds. We note that, if M_1 and M_2 are strongly base orderable matroids, then the matroids M_1' and M_2' obtained by deletion and contraction in the proof of Lemma 10 are also strongly base orderable. \square

Finally in this section, we show an application of Theorem 2 to bipartite matching, which is of independent interest. For a vertex v in a graph, let $\delta(v)$ denote the set of all the edges incident to v .

Corollary 17. *For a bipartite graph $G = (U, V; E)$ containing a perfect matching, there exists a weight function $w : E \rightarrow \mathbb{R}$ satisfying the following conditions.*

1. *For each $u \in U$, let e_u be a lightest edge in $\delta(u)$ with respect to w . Then, $\{e_u \mid u \in U\}$ is a perfect matching in G .*
2. *For each $v \in V$, let e_v be a heaviest edge in $\delta(v)$ with respect to w . Then, $\{e_v \mid v \in V\}$ is a perfect matching in G .*

Proof. Let $\mathcal{B}_1 = \{F \subseteq E \mid |F \cap \delta(u)| = 1 \text{ for any } u \in U\}$ and $\mathcal{B}_2 = \{F \subseteq E \mid |F \cap \delta(v)| = 1 \text{ for any } v \in V\}$. By definition, (E, \mathcal{B}_1) and (E, \mathcal{B}_2) are partition matroids, and hence they are strongly base orderable matroids. Since Conjecture 4 holds for strongly base orderable matroids and $\mathcal{B}_1 \cap \mathcal{B}_2$ is the set of perfect matchings in G , we obtain the corollary. \square

5 Reduction from the weighted case to the unweighted case

In this section, we show that the weighted problem can be reduced to the unweighted one, and prove Theorem 3.

Theorem 3. *If Conjecture 1 is true, then Conjecture 3 is also true.*

Proof. Since Conjectures 1 and 4 are equivalent, it suffices to show that Conjecture 3 is true by assuming that Conjecture 4 is true.

Suppose that we are given $M_i = (S, \mathcal{B}_i)$ and $w_i : S \rightarrow \mathbb{R}$ for $i \in \{1, 2\}$ as in Conjecture 3. We first consider the problem of finding a maximum weight common basis of M_1 and M_2^* with respect to $w_1 - w_2$, where $M_2^* = (S, \mathcal{B}_2^*)$ is the dual matroid of M_2 . By Theorem 7, there exist two functions $q_1 : S \rightarrow \mathbb{R}$ and $q_2 : S \rightarrow \mathbb{R}$ with $q_1 + q_2 = w_1 - w_2$ such that

$$\arg \max_{X \in \mathcal{B}_1 \cap \mathcal{B}_2^*} (w_1(X) - w_2(X)) = \left(\arg \max_{X \in \mathcal{B}_1} q_1(X) \right) \cap \left(\arg \max_{X \in \mathcal{B}_2^*} q_2(X) \right). \quad (2)$$

Define $\hat{\mathcal{B}}_1 = \arg \max_{X \in \mathcal{B}_1} q_1(X)$ and $\hat{\mathcal{B}}_2 = \arg \max_{X \in \mathcal{B}_2^*} q_2(X)$. Then, it is known that $\hat{M}_i = (S, \hat{\mathcal{B}}_i)$ is also a matroid for $i \in \{1, 2\}$ (see [13]). By (2), we obtain

$$\arg \max_{X \in \mathcal{B}_1 \cap \mathcal{B}_2^*} (w_1(X) - w_2(X)) = \hat{\mathcal{B}}_1 \cap \hat{\mathcal{B}}_2. \quad (3)$$

This together with $\mathcal{B}_1 \cap \mathcal{B}_2^* \neq \emptyset$ shows that $\hat{\mathcal{B}}_1 \cap \hat{\mathcal{B}}_2 \neq \emptyset$, and hence \hat{M}_1 and \hat{M}_2 satisfy the assumptions in Conjecture 4. Therefore, by assuming that Conjecture 4 is true, there exists a function $\hat{p} : S \rightarrow \mathbb{R}$ satisfying the following conditions.

- (a) For any $B_1 \in \arg \min_{X \in \hat{\mathcal{B}}_1} \hat{p}(X)$, it holds that $B_1 \in \hat{\mathcal{B}}_2$.
- (b) For any $B_2 \in \arg \max_{X \in \hat{\mathcal{B}}_2} \hat{p}(X)$, it holds that $B_2 \in \hat{\mathcal{B}}_1$.

Let $\delta := \min\{|q_i(X) - q_i(Y)| \mid i \in \{1, 2\}, X, Y \subseteq S, q_i(X) \neq q_i(Y)\}$ and let ε be a positive number such that $\varepsilon \cdot |\hat{p}(X)| < \delta/2$ for any $X \subseteq S$. We now show that $p := w_1 - q_1 + \varepsilon \cdot \hat{p}$ satisfies the requirements of Conjecture 3. Let B_1 be a set in $\arg \max_{X \in \mathcal{B}_1} (w_1(X) - p(X)) = \arg \max_{X \in \mathcal{B}_1} (q_1(X) - \varepsilon \cdot \hat{p}(X))$. Since $-\delta/2 < \varepsilon \cdot \hat{p}(X) < \delta/2$ for any $X \subseteq S$, we have that $B_1 \in \arg \max_{X \in \mathcal{B}_1} q_1(X) = \hat{\mathcal{B}}_1$ and $B_1 \in \arg \min_{X \in \hat{\mathcal{B}}_1} \hat{p}(X)$. Then (a) shows that $B_1 \in \hat{\mathcal{B}}_2$. Therefore,

$$B_1 \in \hat{\mathcal{B}}_1 \cap \hat{\mathcal{B}}_2 = \arg \max_{X \in \mathcal{B}_1 \cap \mathcal{B}_2^*} (w_1(X) - w_2(X)) = \arg \max_{X \in \mathcal{B}_1 \cap \mathcal{B}_2^*} (w_1(X) + w_2(S \setminus X))$$

holds by (3), which means that p satisfies the first requirement in Conjecture 3.

Similarly, let B_2 be a set in

$$\arg \max_{X \in \mathcal{B}_2} (w_2(X) - p(X)) = \arg \max_{X \in \mathcal{B}_2} (-q_2(X) - \varepsilon \cdot \hat{p}(X)) = \arg \max_{X \in \mathcal{B}_2} (q_2(S \setminus X) + \varepsilon \cdot \hat{p}(S \setminus X)).$$

This shows that $S \setminus B_2 \in \arg \max_{X \in \mathcal{B}_2^*} q_2(X) = \hat{\mathcal{B}}_2$ and $S \setminus B_2 \in \arg \max_{X \in \hat{\mathcal{B}}_2} \hat{p}(X)$. Then (b) shows that $S \setminus B_2 \in \hat{\mathcal{B}}_1$. Therefore,

$$S \setminus B_2 \in \hat{\mathcal{B}}_1 \cap \hat{\mathcal{B}}_2 = \arg \max_{X \in \mathcal{B}_1 \cap \mathcal{B}_2^*} (w_1(X) - w_2(X)) = \arg \max_{X \in \mathcal{B}_1 \cap \mathcal{B}_2^*} (w_1(X) + w_2(S \setminus X))$$

holds by (3), which means that p satisfies the second requirement in Conjecture 3. Therefore, Conjecture 3 is true if Conjecture 4 is true. \square

Remark 18. Algorithmically, if we can compute \hat{p} , then we can compute p efficiently as follows. First, we may assume that w_1 and w_2 are integral by multiplying by the common denominator. Then, we can take q_1 and q_2 so that they are integral [17]. Therefore, we have that $\delta \geq 1$, and hence $\varepsilon := 1/(1 + 2 \sum_{s \in S} |\hat{p}(s)|)$ satisfies the conditions in the proof. This shows that we can compute $p := w_1 - q_1 + \varepsilon \cdot \hat{p}$.

By Theorem 3, we obtain Corollaries 4 and 5 as follows. In the proof of Theorem 3, we consider Conjecture 4 for matroids $\hat{M}_i = (S, \hat{\mathcal{B}}_i)$, where $\hat{\mathcal{B}}_1 = \arg \max_{X \in \mathcal{B}_1} q_1(X)$ and $\hat{\mathcal{B}}_2 = \arg \max_{X \in \mathcal{B}_2^*} q_2(X)$. Observe that if M_1 is a partition matroid with partition classes of size at most 2 and with all-ones upper bound on the partition classes, then so is \hat{M}_1 . Furthermore, Lemma 9 shows that if M_i is strongly base orderable, then so is \hat{M}_i . Since Theorems 1 and 2 imply that Conjecture 4 also holds for these cases, we obtain Corollaries 4 and 5.

6 Gross substitutes valuations

In this section, we show that the reduction technique in Section 5 works also for M^\natural -concave functions, or equivalently, gross substitutes functions. M^\natural -concave functions are introduced by Murota and Shioura [31] and play a central role in the theory of discrete convex analysis. A function $f : \mathbb{Z}^S \rightarrow \mathbb{R} \cup \{-\infty\}$ is said to be M^\natural -concave if it satisfies the following exchange property:

(M^\natural -EXC) $\forall x, y \in \text{dom} f, \forall i \in \text{supp}^+(x - y), \exists j \in \text{supp}^-(x - y) \cup \{0\}$:

$$f(x) + f(y) \leq f(x - \chi_i + \chi_j) + f(y + \chi_i - \chi_j),$$

where $\text{dom} f = \{x \in \mathbb{Z}^S \mid f(x) > -\infty\}$, $\text{supp}^+(x) = \{i \in S \mid x(i) > 0\}$, $\text{supp}^-(x) = \{i \in S \mid x(i) < 0\}$ for $x \in \mathbb{Z}^S$, χ_i is the characteristic vector of $i \in S$, and χ_0 is the all-zero vector $\mathbf{0}$. When we consider a function f on $\{0, 1\}^S$, f can be regarded as a function on \mathbb{Z}^S by setting $f(x) = -\infty$ for $x \in \mathbb{Z}^S \setminus \{0, 1\}^S$. It is shown by Fujishige and Yang [?] that a function f on $\{0, 1\}^S$ is M^\natural -concave if and only if it is a gross substitutes function (see also [29, Theorem 6.34]). See survey papers [30, 36] for more details on M^\natural -concave functions and gross substitutes functions. For a set $Q \subseteq \mathbb{Z}^S$, we define a function f_Q on \mathbb{Z}^S by $f_Q(x) = 0$ if $x \in Q$ and $f_Q(x) = -\infty$ otherwise. We say that a set $Q \subseteq \mathbb{Z}^S$ is M^\natural -convex if f_Q is an M^\natural -concave function. It is known that a set is M^\natural -convex if and only if it is the set of integer points/vectors in an integral g-polymatroid [18, 19]. Let $\mathbf{1}$ denote the all-one vector in \mathbb{Z}^S .

We are interested in the existence of a pricing scheme for the two-buyer case with gross substitutes valuations (or equivalently, M^\natural -concave valuations), which is stated as follows.

Conjecture 5. For $i = 1, 2$, let $v_i : \{0, 1\}^S \rightarrow \mathbb{R} \cup \{-\infty\}$ be an M^\sharp -concave function. Then, there exists a vector $p \in \mathbb{R}^S$ satisfying the following conditions.

1. For any $x_1 \in \arg \max_{x \in \{0, 1\}^S} (v_1(x) - p \cdot x)$, it holds that $x_1 \in \arg \max_{x \in \{0, 1\}^S} (v_1(x) + v_2(\mathbf{1} - x))$.
2. For any $x_2 \in \arg \max_{x \in \{0, 1\}^S} (v_2(x) - p \cdot x)$, it holds that $x_2 \in \arg \max_{x \in \{0, 1\}^S} (v_1(\mathbf{1} - x) + v_2(x))$.

In Conjecture 5, a subset of S is represented by its characteristic vector. If buyer i comes to a shop first, then she chooses an arbitrary set x_i maximizing her utility $v_i(x) - p \cdot x$. Then, the second buyer takes the set of all the remaining items whose characteristic vector is $\mathbf{1} - x_i$. Conjecture 5 asserts that, regardless of the choice of x_i , this mechanism gives an allocation maximizing the social welfare.

As an unweighted version of this conjecture, we consider the following conjecture.

Conjecture 6. For $i = 1, 2$, let $Q_i \subseteq \{0, 1\}^S$ be an M^\sharp -convex set such that there exist $x_1 \in Q_1$ and $x_2 \in Q_2$ with $x_1 + x_2 = \mathbf{1}$. Then, there exists a vector $p \in \mathbb{R}^S$ satisfying the following conditions.

1. For any $x_1 \in \arg \min_{x \in Q_1} (p \cdot x)$, it holds that $\mathbf{1} - x_1 \in Q_2$.
2. For any $x_2 \in \arg \min_{x \in Q_2} (p \cdot x)$, it holds that $\mathbf{1} - x_2 \in Q_1$.

In Conjecture 6, each buyer i has an admissible set Q_i instead of a valuation. More precisely, each buyer i wants to buy a set of items whose characteristic vector x_i belongs to a given M^\sharp -convex set Q_i . We can easily see that Conjecture 6 is a special case of Conjecture 5, in which $v_i = f_{Q_i}$ for $i = 1, 2$. We now prove that the reverse implication also holds, which means that Conjecture 5 can be reduced to the unweighted case.

Theorem 19. If Conjecture 6 is true, then Conjecture 5 is also true.

Proof. Let $v_2^* : \{0, 1\}^S \rightarrow \mathbb{R} \cup \{-\infty\}$ be the function defined by $v_2^*(x) = v_2(\mathbf{1} - x)$ for $x \in \{0, 1\}^S$. Then, v_2^* is also an M^\sharp -concave function. Consider the problem of maximizing $v_1(x) + v_2^*(x)$ subject to $x \in \{0, 1\}^S$. By the M -convex intersection theorem (see [29, Theorem 8.17]), there exists a vector $q \in \mathbb{R}^S$ such that

$$\arg \max_{x \in \{0, 1\}^S} (v_1(x) + v_2^*(x)) = \left(\arg \max_{x \in \{0, 1\}^S} (v_1(x) - q \cdot x) \right) \cap \left(\arg \max_{x \in \{0, 1\}^S} (v_2^*(x) + q \cdot x) \right). \quad (4)$$

Define $Q_1 = \arg \max_{x \in \{0, 1\}^S} (v_1(x) - q \cdot x)$, $Q_2^* = \arg \max_{x \in \{0, 1\}^S} (v_2^*(x) + q \cdot x)$, and $Q_2 = \{\mathbf{1} - x \mid x \in Q_2^*\}$. Then, it is known that Q_1 and Q_2^* are M^\sharp -convex sets (see [29, Theorem 6.30(2)]), and so is Q_2 (see [29, Theorem 6.13(2)]). By (4), we obtain $\arg \max_{x \in \{0, 1\}^S} (v_1(x) + v_2^*(x)) = Q_1 \cap Q_2^*$. This shows that $Q_1 \cap Q_2^* \neq \emptyset$, and hence Q_1 and Q_2 satisfy the assumptions in Conjecture 6. Therefore, by assuming that Conjecture 6 is true, there exists a vector $\hat{p} \in \mathbb{R}^S$ satisfying the following conditions.

- (a) For any $x_1 \in \arg \min_{x \in Q_1} (\hat{p} \cdot x)$, it holds that $\mathbf{1} - x_1 \in Q_2$.
- (b) For any $x_2 \in \arg \min_{x \in Q_2} (\hat{p} \cdot x)$, it holds that $\mathbf{1} - x_2 \in Q_1$.

Then, by the same argument as the proof of Theorem 3, $p := q + \varepsilon \cdot \hat{p}$ satisfies the requirements in Conjecture 5, where ε is a sufficiently small positive number. \square

Remark 20. In a market model, it is common to assume that each valuation v_i is monotone and $v_i(\emptyset) = 0$. We note that these assumptions are not required in the proof of Theorem 19. In return for this, the obtained price vector p is not necessarily non-negative.

We note that Conjecture 1 is a special case of Conjecture 5, as the characteristic vector of all the bases of a matroid forms an M^{\sharp} -convex set. This relationship supports the importance of Conjecture 1.

7 Conclusion

We considered the existence of prices that are capable to achieve optimal social welfare without a central tie-breaking coordinator. Although such pricing looks similar to well-known Walrasian pricing, it is less understood even for two-buyer markets with gross substitute valuations. This paper focuses on two-buyer markets with rank valuations, and we gave polynomial-time algorithms that always find such prices when one of the matroids is a simple partition matroid or both matroids are strongly base orderable. This result partially answers a question of Dütting and Végh. We further formalized a weighted variant of the conjecture of Dütting and Végh, and showed that the weighted variant can be reduced to the unweighted one based on the weight-splitting theorem of Frank. We also showed that a similar reduction technique works for M^{\sharp} -concave functions, or equivalently, gross substitutes functions.

Acknowledgement Kristóf Bérczi was supported by the János Bolyai Research Fellowship of the Hungarian Academy of Sciences and by Projects no. NKFI-128673 and no. ED_18-1-2019-0030 provided from the National Research, Development and Innovation Fund of Hungary. Naonori Kakimura was supported by JSPS KAKENHI Grant Numbers JP17K00028 and JP18H05291, Japan. Yusuke Kobayashi was supported by JSPS KAKENHI grant numbers JP18H05291, JP19H05485, and JP20K11692, Japan. The work was supported by the Research Institute for Mathematical Sciences, an International Joint Usage/Research Center located in Kyoto University.

References

- [1] B. Berger, A. Eden, and M. Feldman. On the power and limits of dynamic pricing in combinatorial markets. *arXiv preprint arXiv:2002.06863*, 2020.
- [2] L. Blumrosen and T. Holenstein. Posted prices vs. negotiations: an asymptotic analysis. *EC*, 10:1386790–1386801, 2008.
- [3] S. Chawla, J. D. Hartline, D. L. Malec, and B. Sivan. Multi-parameter mechanism design and sequential posted pricing. In *Proceedings of the forty-second ACM symposium on Theory of computing*, pages 311–320, 2010.
- [4] S. Chawla, D. L. Malec, and B. Sivan. The power of randomness in bayesian optimal mechanism design. In *Proceedings of the 11th ACM conference on Electronic commerce*, pages 149–158, 2010.

- [5] S. Chawla, J. B. Miller, and Y. Teng. Pricing for online resource allocation: intervals and paths. In *Proceedings of the Thirtieth Annual ACM-SIAM Symposium on Discrete Algorithms*, pages 1962–1981. SIAM, 2019.
- [6] E. H. Clarke. Multipart pricing of public goods. *Public choice*, pages 17–33, 1971.
- [7] V. Cohen-Addad, A. Eden, M. Feldman, and A. Fiat. The invisible hand of dynamic market pricing. In *Proceedings of the 2016 ACM Conference on Economics and Computation*, pages 383–400, 2016.
- [8] J. Davies and C. McDiarmid. Disjoint common transversals and exchange structures. *Journal of the London Mathematical Society*, 2(1):55–62, 1976.
- [9] P. Dütting, M. Feldman, T. Kesselheim, and B. Lucier. Posted prices, smoothness, and combinatorial prophet inequalities. *arXiv preprint arXiv:1612.03161*, 2016.
- [10] P. Dütting, M. Feldman, T. Kesselheim, and B. Lucier. Prophet inequalities made easy: Stochastic optimization by pricing non-stochastic inputs. In *2017 IEEE 58th Annual Symposium on Foundations of Computer Science (FOCS)*, pages 540–551. IEEE, 2017.
- [11] P. Dütting and L. A. Végh. Private Communication, 2017.
- [12] A. Eden, U. Feige, and M. Feldman. Max-min greedy matching. *Approximation, Randomization, and Combinatorial Optimization. Algorithms and Techniques (APPROX/RANDOM 2019)*, 145:7, 2019.
- [13] J. Edmonds. Matroids and the greedy algorithm. *Mathematical programming*, 1(1):127–136, 1971.
- [14] T. Ezra, M. Feldman, T. Roughgarden, and W. Suksompong. Pricing multi-unit markets. In *International Conference on Web and Internet Economics*, pages 140–153. Springer, 2018.
- [15] M. Feldman, N. Gravin, and B. Lucier. Combinatorial auctions via posted prices. In *Proceedings of the twenty-sixth annual ACM-SIAM symposium on Discrete algorithms*, pages 123–135. SIAM, 2014.
- [16] M. Feldman, N. Gravin, and B. Lucier. Combinatorial walrasian equilibrium. *SIAM Journal on Computing*, 45(1):29–48, 2016.
- [17] A. Frank. A weighted matroid intersection algorithm. *Journal of Algorithms*, 2(4):328–336, 1981.
- [18] A. Frank. Generalized polymatroids. In *Finite and Infinite Sets (Colloquia Mathematica Societatis János Bolyai 37)*, pages 285–294, 1984.
- [19] A. Frank. *Connections in Combinatorial Optimization*. Oxford University Press, 2011.
- [20] S. Fujishige and Z. Yang. A note on Kelso and Crawford’s gross substitutes condition. *Mathematics of Operations Research*, 28(3):463–469, 2003.
- [21] T. Groves. Incentives in teams. *Econometrica: Journal of the Econometric Society*, pages 617–631, 1973.

- [22] F. Gul, E. Stacchetti, et al. Walrasian equilibrium with gross substitutes. *Journal of Economic theory*, 87(1):95–124, 1999.
- [23] J. Hsu, J. Morgenstern, R. Rogers, A. Roth, and R. Vohra. Do prices coordinate markets? In *Proceedings of the forty-eighth annual ACM symposium on Theory of Computing*, pages 440–453, 2016.
- [24] R. Huang and G.-C. Rota. On the relations of various conjectures on latin squares and straightening coefficients. *Discrete Mathematics*, 128(1-3):225–236, 1994.
- [25] A. S. Kelso Jr and V. P. Crawford. Job matching, coalition formation, and gross substitutes. *Econometrica: Journal of the Econometric Society*, pages 1483–1504, 1982.
- [26] S. Krogdahl. A combinatorial base for some optimal matroid intersection algorithms. Technical Report STAN-CS-74-468, Computer Science Department, Stanford University, Stanford, CA, U.S., 1974.
- [27] S. Krogdahl. A combinatorial proof for a weighted matroid intersection algorithm. Technical Report Computer Science Report 17, Institute of Mathematical and Physical Sciences, University of Tromso, Tromso, Norway, 1976.
- [28] S. Krogdahl. The dependence graph for bases in matroids. *Discrete Mathematics*, 19(1):47–59, 1977.
- [29] K. Murota. *Discrete Convex Analysis*. Society for Industrial and Applied Mathematics, 2003.
- [30] K. Murota. Discrete convex analysis: A tool for economics and game theory. *The Journal of Mechanism and Institution Design*, 1(1):151–273, 2016.
- [31] K. Murota and A. Shioura. M-convex function on generalized polymatroid. *Mathematics of Operations Research*, 24(1):95–105, 1999.
- [32] N. Nisan and I. Segal. The communication requirements of efficient allocations and supporting prices. *Journal of Economic Theory*, 129(1):192–224, 2006.
- [33] H. Nishimura and S. Kuroda. *A Lost Mathematician, Takeo Nakasawa: The Forgotten Father of Matroid Theory*. Springer Science & Business Media, 2009.
- [34] J. Oxley. *Matroid Theory*. Oxford University Press, 2011.
- [35] A. Schrijver. *Combinatorial optimization: polyhedra and efficiency*, volume 24. Springer Science & Business Media, 2003.
- [36] A. Shioura and A. Tamura. Gross substitutes condition and discrete concavity for multi-unit valuations: a survey. *Journal of the Operations Research Society of Japan*, 58(1):61–103, 2015.
- [37] W. Vickrey. Counterspeculation, auctions, and competitive sealed tenders. *The Journal of finance*, 16(1):8–37, 1961.
- [38] L. Walras. *Éléments d'économie politique pure, ou, Théorie de la richesse sociale*. F. Rouge, 1896.

- [39] H. Whitney. On the abstract properties of linear dependence. In *Hassler Whitney Collected Papers*, pages 147–171. Springer, 1992.