

**SMOOTHNESS IN THE L_p MINKOWSKI PROBLEM
FOR $p < 1$**

GABRIELE BIANCHI, KÁROLY J. BÖRÖCZKY, AND ANDREA COLESANTI

ABSTRACT. We discuss the smoothness and strict convexity of the solution of the L_p -Minkowski problem when $p < 1$ and the given measure has a positive density function.

1. INTRODUCTION

Given K in the class \mathcal{K}_0^n of compact convex sets in \mathbb{R}^n that have non-empty interior and contain the origin o , we write h_K and S_K to denote its support function and its surface area measure, respectively, and for $p \in \mathbb{R}$, $S_{K,p}$ to denote its L_p -area measure, where $dS_{K,p} = h_K^{1-p} dS_K$. The L_p -area measure defined by Lutwak [35] is a central notion in convexity, see say Barthe, Guédon, Mendelson and Naor [2], Böröczky, Lutwak, Yang and Zhang [5], Campi and Gronchi [10], Chou [15], Cianchi, Lutwak, Yang and Zhang [17], Gage and Hamilton [19], Haberl and Parapatits [23], Haberl and Schuster [24, 25], Haberl, Schuster and Xiao [26], He, Leng and Li [27], Henk and Linke [28], Ludwig [34], Lutwak, Yang and Zhang [37, 38], Naor [41], Naor and Romik [42], Paouris [44], Paouris and Werner [45] and Stancu [50].

The L_p Minkowski problem asks for the existence of a convex body $K \in \mathcal{K}_0^n$ whose L_p area measure is a given finite Borel measure ν on S^{n-1} . When $p = 1$ this is the classical Minkowski problem solved by Minkowski [40] for polytopes, and by Alexandrov [1] and Fenchel and Jessen [18] in general. The smoothness of the solution was clarified in a series of papers by Nirenberg [43], Cheng and Yau [14], Pogorelov [46] and Caffarelli [7, 8]. For $p > 1$ and $p \neq n$, the L_p Minkowski problem has a unique solution according to Chou and Wang [16], Guan and Lin [22] and Hug, Lutwak, Yang and Zhang [30]. The smoothness of the solution is discussed in Chou and Wang [16], Huang and Lu [29] and Lutwak and Oliker [36]. In addition, the case $p < 1$ has been intensively investigated by Böröczky, Lutwak, Yang and Zhang [4], Böröczky and Hai T. Trinh [6], Chen [13], Chen, Li and Zhu [11, 12], Ivaki [31], Jiang [32], Lu and Wang [33], Lutwak, Yang and Zhang [39], Stancu [48, 49] and Zhu [52–55].

The solution of the L_p -Minkowski problem may not be unique for $p < 1$ according to Chen, Li and Zhu [12] if $0 < p < 1$, according to Stancu [49] if $p = 0$, and according to Chou and Wang [16] if $p < 0$ small.

In this paper we are interested in this problem when $p < 1$ and ν is a measure with density with respect to the Hausdorff measure \mathcal{H}^{n-1} on S^{n-1} , i.e. in the problem

$$(1.1) \quad dS_{K,p} = f d\mathcal{H}^{n-1} \quad \text{on } S^{n-1},$$

2010 *Mathematics Subject Classification.* Primary: 52A40, secondary: 35J96.

Key words and phrases. L_p Minkowski problem, Monge-Ampère equation.

First and third authors are supported in part by the Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM). Second author is supported in part by NKFIH grants 116451 and 109789.

where f is a non-negative Borel function in S^{n-1} .

According to Chou and Wang [16], if $-n < p < 1$ and the Borel function f is bounded from above and below by positive constants, then (1.1) has a solution. More general existence results are provided by the recent works Chen, Li and Zhu [11] if $p = 0$, Chen, Li and Zhu [12] if $0 < p < 1$, and Bianchi, Böröczky and Colesanti [3] if $-n < p < 0$. In particular, it is known that (1.1) has a solution if $0 \leq p < 1$ and f is any non-negative function in $L_1(S^{n-1})$ with $\int_{S^{n-1}} f d\mathcal{H}^{n-1} > 0$, and if $-n < p < 0$ and f is any non-negative function in $L_{\frac{n}{n+p}}(S^{n-1})$ with $\int_{S^{n-1}} f d\mathcal{H}^{n-1} > 0$.

We observe that h is a non-negative positively 1-homogeneous convex function in \mathbb{R}^n which solves the Monge-Ampère equation

$$(1.2) \quad h^{1-p} \det(\nabla^2 h + hI) = f \quad \text{on } S^{n-1}$$

in the sense of measure if and only if h is the support function of a convex body $K \in \mathcal{K}_0^n$ which is the solution of (1.1) (see Section 2). Here h is the unknown non-negative (support) function on S^{n-1} to be found, $\nabla^2 h$ denote the (covariant) Hessian matrix of h with respect to an orthonormal frame on S^{n-1} , and I is the identity matrix. The function h may vanish somewhere even in the case when f is positive and continuous, and when this happens and $p < 1$ the equation (1.2) is singular at the zero set of h . Naturally, if h is C^2 , then (1.2) is a proper Monge-Ampère equation.

In this paper we study the smoothness and strict convexity of a solution $K \in \mathcal{K}_0^n$ of (1.1) assuming $\tau_2 > f > \tau_1$ for some constants $\tau_2 > \tau_1 > 0$. Concerning these aspects for $p < 1$, we summarize the known results in Theorem 1.1, and the new results in Theorem 1.2.

We say that $x \in \partial K$ is a C^1 -smooth point if there is a unique tangent hyperplane to K at x , and observe that ∂K is C^1 if and only if each $x \in \partial K$ is C^1 -smooth (see Section 2 for all definitions). In addition, we note that h_K is C^1 on S^{n-1} if and only if K is strictly convex, and h_K is strictly convex on any hyperplane avoiding the origin if and only if ∂K is C^1 . For $z \in \partial K$, the exterior normal cone at z is denoted by $N(K, z)$, and for $z \in \text{int } K$, we set $N(K, z) = \{o\}$. Theorem 1.1 (i) and (ii) are essentially due to Caffarelli [7] (see Theorem 3.6), and Theorem 1.1 (iii) is due to Chou and Wang [16]. If the function f in (1.1) is C^α for $\alpha > 0$, then Caffarelli [8] proves (iv).

Theorem 1.1 (Caffarelli, Chou, Wang). *If $K \in \mathcal{K}_0^n$ is a solution of (1.1) for $n \geq 2$ and $p < 1$, and f is bounded from above and below by positive constants, then the following assertions hold:*

- (i) *The set X_0 of the points $x \in \partial K$ with $N(K, x) \subset N(K, o)$ is closed, each point of $X = \partial K \setminus X_0$ is C^1 -smooth and X contains no segment.*
- (ii) *If $o \in \partial K$ is a C^1 -smooth point, then ∂K is C^1 .*
- (iii) *If $p \leq 2 - n$, then $o \in \text{int } K$, and hence K is strictly convex and ∂K is C^1 .*
- (iv) *If $o \in \text{int } K$ and the function f in (1.1) is positive and C^α , for some $\alpha > 0$, then ∂K is $C^{2,\alpha}$.*

Concerning strict convexity, Assertion (iii) here is optimal because Example 4.2 shows that if $2 - n < p < 1$, then it is possible that o belongs to the relative interior of an $(n - 1)$ -dimensional face of a solution K of (1.1) where f is a positive continuous function. Therefore, the only question left open is the C^1 smoothness of the boundary of the solution if $2 - n < p < 1$.

We note that if $p < 1$ and K is a solution of (1.2) with f positive and $o \in \partial K$, then

$$(1.3) \quad \dim N(K, o) \leq n - 1.$$

Therefore, Theorem 1.1 (ii) yields that ∂K is C^1 for the solution K if $n = 2$. In general, we have the following partial results.

Theorem 1.2. *If $K \in \mathcal{K}_0^n$ is a solution of (1.1) for $n \geq 2$ and $p < 1$, and f is bounded from above and below by positive constants, then the following assertions hold:*

- (i) *If $n = 2$, $n = 3$ or $n > 3$ and $p < 4 - n$, then ∂K is C^1 .*
- (ii) *If $\mathcal{H}^{n-1}(X_0) = 0$ for the X_0 in Theorem 1.1 (i), then ∂K is C^1 .*

Our results differ in some cases from the ones in Chou and Wang [16], possibly because [16] considers the equation

$$(1.4) \quad \det(\nabla^2 h + hI) = fh^{p-1} \quad \text{on } S^{n-1}$$

instead of (1.2). In the context of non-negative convex functions, being a solution of this last equation is a priori more restrictive than being a solution of (1.2), even if obviously the two notions coincide when h is positive (see Section 2 for more on this point). Chou and Wang [16] proves, under our same assumptions on f , the strict convexity of the solution h of (1.4) on hyperplanes avoiding the origin, and uses this to prove that ∂K is C^1 for the convex body K . We note that if $K \in \mathcal{K}_0^n$ is a solution of (1.4) for $p < 1$ and f is bounded from below and above by positive constants, then combining Theorem 1.2 (ii) with the simple observation (2.11) in Section 2 shows that ∂K is C^1 , as it was verified by Chou and Wang [16]. In our opinion (1.2) is the right equation to consider and using it we obtain weaker results.

To give an example of how the two equations differ, the support function h of the body K in Example 4.2 (where o belongs to the relative interior of an $(n - 1)$ -dimensional face) is a solution of (1.2) but not a solution of (1.4).

According to Chou and Wang [16] (see also Lemma 3.1 below), the Monge-Ampère equation (1.2) can be transferred to a Monge-Ampère equation

$$(1.5) \quad v^{1-p} \det(D^2 v) = g$$

for a convex function v on \mathbb{R}^{n-1} where g is a given non-negative function and D^2 stands for the Hessian in \mathbb{R}^{n-1} .

The proofs of Claims (i) and (ii) in Theorem 1.1 use as an essential tool a result proved by Caffarelli in [7] regarding smoothness and strict convexity of convex solutions of certain Monge-Ampère equation of type (1.5) (see Theorem 3.6). Proving that ∂K is C^1 is equivalent to prove that h_K is strictly convex, and [7] is the key to prove this property in $\{y \in S^{n-1} : h_K(y) > 0\}$.

The proof of Claim (i) in Theorem 1.2 is based on the following result for the singular inequality $v^{1-p} \det D^2 v \geq g$.

Proposition 1.3. *Let $\Omega \subset \mathbb{R}^n$ be an open convex set, and let v be a non-negative convex function in Ω with $S = \{x \in \Omega : v(x) = 0\}$. If for $p < 1$ and $\tau > 0$, v is the solution of*

$$(1.6) \quad v^{1-p} \det D^2 v \geq \tau \quad \text{in } \Omega \setminus S$$

in the sense of measure, and S is r -dimensional, for $r \geq 1$, then $p \geq -n + 1 + 2r$.

We mention that in Caffarelli [9] a corresponding result for $p = 1$ is established.

The underlying idea behind the proof of this result is the following: On the one hand, the graph of v near S is close to being ruled. Hence, the total variation of the derivative is “small”. On the other hand, the total variation of the derivative is “large” because of the Monge-Ampère inequality (1.6).

The inequality $p \geq -n + 1 + 2r$ in this result is close to being optimal, at least when $r = 1$. Indeed, Example 3.2 shows that, for any $p > -n + 3$, there exists a non-negative convex solution of (1.6) in Ω which vanishes on the intersection of

Ω with a line. For the version $p = 1$ of Proposition 1.3, Caffarelli [9] proves that $\dim S < n/2$ and that this inequality is optimal.

Proposition 1.3 yields actually somewhat more than Claim (i) in Theorem 1.2; namely, if $r \geq 2$ is an integer, $p < \min\{1, 2r - n\}$ and $K \in \mathcal{K}_0^n$ is a solution of (1.1) with $o \in \partial K$, then $\dim N(K, o) < r$. As a consequence, we have the following technical statements about K , where we also use Theorem 1.2 (ii) for Claim (ii).

Corollary 1.4. *If $p < 1$ and $K \in \mathcal{K}_0^n$, $n \geq 4$, is a solution of (1.1) with $o \in \partial K$, then*

- (i) $\dim N(K, o) < \frac{n+1}{2}$;
- (ii) *if in addition $n = 4, 5$ and ∂K is not C^1 , then $\dim N(K, o) = 2$ and $\dim F(K, u) = n - 1$ for some $u \in N(K, o)$.*

In Section 2 we review the notation used in this paper. Section 3 contains results and examples regarding Monge-Ampère equations in \mathbb{R}^n , namely Proposition 1.3, Example 3.2 and Proposition 3.4. This last result is the key to prove Theorem 1.2 (ii). In Section 4 we show, for the sake of completeness, how to prove Theorem 1.1 using ideas due to Caffarelli [7, 8] and Chou and Wang [16]. Theorem 1.2 and Corollary 1.4 are proved in Section 5.

2. NOTATION AND PRELIMINARIES

As usual, S^{n-1} denotes the unit sphere and o the origin in the Euclidean n -space \mathbb{R}^n . The symbol B^n denotes the unit ball in \mathbb{R}^n centred at o and ω_n denotes its volume. If $x, y \in \mathbb{R}^n$, then $\langle x, y \rangle$ is the scalar product of x and y , while $\|x\|$ is the euclidean norm of x . By $[x, y]$ we denote the segment with endpoint x and y .

We write \mathcal{H}^k for k -dimensional Hausdorff measure in \mathbb{R}^n .

We denote by ∂E , $\text{int}E$, $\text{cl}E$, and 1_E the *boundary*, *interior*, *closure*, and *characteristic function* of a set E in \mathbb{R}^n , respectively. The symbols $\text{aff}E$ and $\text{lin}E$ denote respectively the *affine hull* and the *linear hull* of E . The *dimension* $\dim E$ is the dimension of $\text{aff}E$. With the symbol $E \mid L$ we denote the orthogonal projection of E on the linear space L .

Given a function v defined on a subset of \mathbb{R}^n , Dv and D^2v denote its gradient and its Hessian, respectively.

Our next goal is to recall a standard notion of generalised solution of Monge-Ampère equations, usually referred to as *solution in the sense of measure*. Our general reference for notions and facts about Monge-Ampère equations is the survey by Trudinger and Wang [51]. Let v be a convex function defined in an open convex set Ω ; the subgradient $\partial v(x)$ of v at $x \in \Omega$ is defined as

$$\partial v(x) = \{z \in \mathbb{R}^n : v(y) \geq v(x) + \langle z, y - x \rangle \text{ for each } y \in \Omega\},$$

which is a non-empty compact convex set. Note that v is differentiable at $x \in \Omega$ if and only if $\partial v(x)$ consists of exactly one vector, which is the gradient of v at x . If $\omega \subset \Omega$ is a Borel set, then we denote by $N_v(\omega)$ the image of ω through the gradient map of v , i.e.

$$N_v(\omega) = \bigcup_{x \in \omega} \partial v(x).$$

Note that as ω is a Borel set, then $N_v(\omega)$ is measurable. Hence, we may define the Monge-Ampère measure associated to v as follows

$$(2.1) \quad \mu_v(\omega) = \mathcal{H}^n(N_v(\omega)).$$

For $p < 1$ and non-negative g on \mathbb{R}^n , we say that the non-negative convex function v satisfies the Monge-Ampère equation

$$v^{1-p} \det(D^2v) = g$$

in the sense of measure (or in the Alexandrov sense) if

$$v^{1-p} d\mu_v = g d\mathcal{H}^n.$$

Equivalently

$$\int_{\omega} v^{1-p}(x) d\mu_v(x) = \int_{\omega} g(x) dx$$

for every Borel subset ω of Ω .

A *convex body* in \mathbb{R}^n is a compact convex set with nonempty interior. The treatises Gardner [20], Gruber [21] and Schneider [47] are excellent general references for convex geometry. The function

$$h_K(u) = \max\{\langle u, y \rangle : y \in K\},$$

for $u \in \mathbb{R}^n$, is the *support function* of K . When it is clear the convex body to which we refer we will drop the subscript K from h_K and write simply h . Any convex body K is uniquely determined by its support function. A set $C \subset \mathbb{R}^n$ is a *convex cone* if $\alpha_1 u_1 + \alpha_2 u_2 \in C$ for any $u_1, u_2 \in C$ and $\alpha_1, \alpha_2 \geq 0$.

If S is a convex set in \mathbb{R}^n , then $z \in S$ is an extremal point if $z = \alpha x_1 + (1 - \alpha)x_2$ for $x_1, x_2 \in S$ and $\alpha \in (0, 1)$ imply $x_1 = x_2 = z$. We note that if S is compact and convex, then S is the convex hull of its extremal points. If C is a convex cone and $u \in C \setminus \{o\}$, we say that $\sigma = \{\lambda u : \lambda \geq 0\}$ is an extremal ray if $\alpha_1 x_1 + \alpha_2 x_2 \in \sigma$ for $x_1, x_2 \in C$ and $\alpha_1, \alpha_2 > 0$ imply $x_1, x_2 \in \sigma$. Now if $C \neq \{o\}$ is a closed convex cone such that the origin is an extremal point of C , then C is the convex hull of its extremal rays.

The *normal cone* of a convex body K at $z \in K$ is defined as

$$N(K, z) = \{u \in \mathbb{R}^n : \langle u, y \rangle \leq \langle u, z \rangle \text{ for all } y \in K\}$$

where $N(K, z) = \{o\}$ if $z \in \text{int}K$ and $\dim N(K, z) \geq 1$ if $z \in \partial K$. This definition can be written also as

$$(2.2) \quad N(K, z) = \{u \in \mathbb{R}^n : h_K(u) = \langle z, u \rangle\}.$$

In particular, $N(K, z)$ is a closed convex cone such that the origin is an extremal point, and

$$(2.3) \quad h_K(\alpha_1 u_1 + \alpha_2 u_2) = \alpha_1 h_K(u_1) + \alpha_2 h_K(u_2) \text{ for } u_1, u_2 \in N(K, z) \text{ and } \alpha_1, \alpha_2 > 0.$$

A convex body K is C^1 -smooth at $p \in \partial K$ if $N(K, p)$ is a ray, and ∂K is C^1 if each $p \in \partial K$ is a C^1 -smooth point. Therefore, ∂K is C^1 if and only if the restriction of h_K to any hyperplane not containing o is strictly convex, by (2.3).

We say that a convex body K is *strictly convex* if ∂K contains no segment. The *face* of K with outer normal $u \in \mathbb{R}^n$ is defined as

$$F(K, u) = \{z \in K : h_K(u) = \langle z, u \rangle\},$$

which lies in ∂K if $u \neq o$. Schneider [47, Thm. 1.7.4] proves that

$$(2.4) \quad \partial h_K(u) = F(K, u).$$

Therefore, K is strictly convex if and only if h_K is C^1 on $\mathbb{R}^n \setminus \{o\}$.

A crucial notion for this paper is the one of *surface area measure* S_K of a convex body K , which is a Borel measure on S^{n-1} , defined as follows. For any Borel set $\omega \subset S^{n-1}$:

$$S_K(\omega) = \mathcal{H}^{n-1}(\cup_{u \in \omega} F(K, u)) = \mathcal{H}^{n-1}(\cup_{u \in \omega} \partial h_K(u)),$$

Hence, S_K is the analogue of the Monge-Ampère measure for the restriction of h_K to S^{n-1} .

Given a convex body K containing o and $p < 1$, let $S_{K,p}$ denote the L_p area measure of K ; namely,

$$(2.5) \quad dS_{K,p} = h_K^{1-p} dS_K.$$

Let f be a positive and measurable function on S^{n-1} ; we say that h_K is a solution of (1.2) in the sense of measure if

$$(2.6) \quad \int_{\omega} h_K(y)^{1-p} dS_K(y) = \int_{\omega} f(y) d\mathcal{H}^{n-1}(y)$$

for every Borel subset ω of S^{n-1} .

In what follows we will always assume that f is bounded between two positive constants. Our first remark is that the previous definition is equivalent to the following conditions (a) and (b):

(a) $\dim N(K, o) < n$; or equivalently,

$$(2.7) \quad \mathcal{H}^{n-1}(\{y \in S^{n-1} : h_K(y) = 0\}) = \mathcal{H}^{n-1}(N(K, o) \cap S^{n-1}) = 0,$$

(b) for each Borel set $\omega \subset \{y \in S^{n-1} : h_K(y) > 0\}$, we have

$$(2.8) \quad \int_{\omega} h_K^{1-p}(y) dS_K(y) = \int_{\omega} f(y) d\mathcal{H}^{n-1}(y).$$

Moreover, condition (b) is in turn equivalent to

(b') for each Borel set $\omega \subset \{y \in S^{n-1} : h_K(y) > 0\}$, we have

$$(2.9) \quad S_K(\omega) = \int_{\omega} f(y) h_K(y)^{p-1} d\mathcal{H}^{n-1}(y).$$

To prove that (b) and (b') are equivalent is a simple exercise (in which one has to take into account the fact that h_K is continuous). Indeed, both claims are in turn equivalent to the following fact: the measure S_K is absolutely continuous with respect to \mathcal{H}^{n-1} on $S^{n-1} \setminus \{y \in S^{n-1} : h_K(y) = 0\}$, and the Radon-Nikodym derivative of S_K with respect to \mathcal{H}^{n-1} is fh_K^{p-1} .

Let us prove the equivalence between (2.6) and (a)-(b). To this end, it will be useful the following observation: the set

$$\{x \in \mathbb{R}^n : h_K(x) = 0\}$$

is a closed convex cone. Indeed, it is the set where the non-negative, convex and 1-homogeneous function h_K attains its minimum. For convenience, we set $\omega_0 = \{y \in S^{n-1} : h_K(y) = 0\}$. Assume that (2.6) holds; then (b) follows immediately. If, by contradiction, (a) is false, then ω_0 has non-empty interior so that

$$0 = \int_{\omega_0} h_K(y)^{1-p} dS_K(y) = \int_{\omega_0} f(y) d\mathcal{H}^{n-1}(y) > 0,$$

i.e. a contradiction (in the last inequality we have used the fact that f is bounded from below by a positive constant). Vice versa, assume that (a) and (b) hold. Given a Borel subset ω of S^{n-1} we may write it as the disjoint union of $\omega' = \omega \cap \omega_0$ and $\omega'' = \omega \setminus \omega'$. By (a), $\mathcal{H}^{n-1}(\omega') = 0$, moreover $h_K = 0$ on ω' ; hence,

$$\begin{aligned} \int_{\omega} h_K(y)^{1-p} dS_K(y) &= \int_{\omega''} h_K(y)^{1-p} dS_K(y) \\ &= \int_{\omega''} f(y) d\mathcal{H}^{n-1}(y) \\ &= \int_{\omega} f(y) d\mathcal{H}^{n-1}(y), \end{aligned}$$

i.e. (2.6).

Our next step is to compare the solutions considered by Chou and Wang [16] with the ones introduced here. In particular, we will show that if h_K is a solution of (1.4), then it verifies conditions (a) and (b) as well (and consequently (2.6)). Note that being a solution of (1.4) in the sense of measures means that

$$(2.10) \quad S_K(\omega) = \int_{\omega} f(y) h_K(y)^{p-1} d\mathcal{H}^{n-1}(y).$$

has to hold for every Borel subset of S^{n-1} . In particular (2.9) follows (and then (b)). Moreover, as S_K is finite, $h_K \geq 0$ and f is bounded between two positive constants, the previous relation implies that

$$\int_{S^{n-1}} h_K(y)^{p-1} d\mathcal{H}^{n-1}(y) < +\infty.$$

As $p - 1 < 0$, this yields that the set ω_0 where h_K vanishes on S^{n-1} has zero $(n - 1)$ -dimensional measure. On the other hand this is the intersection of S^{n-1} with a convex cone. Hence we get condition (a).

In addition, if we now apply (2.10) to ω_0 , we get that when h_K is a solution of (1.4) then

$$(2.11) \quad S_K(N(K, o) \cap S^{n-1}) = 0.$$

Note that (2.11) implies that $\mathcal{H}^{n-1}(X_0) = 0$, in the notation of Theorem 1.2, because $X_0 \subset \cup\{F(K, u) : u \in N(K, o) \cap S^{n-1}\}$ and (2.11) means, by definition,

$$\mathcal{H}^{n-1}\left(\cup_{u \in N(K, o) \cap S^{n-1}} F(K, u)\right) = 0.$$

Hence, applying Theorem 1.2 (ii) we deduce that if $K \in \mathcal{K}_0^n$ is a solution of (1.4) for $p < 1$ and f is bounded from below and above by positive constants, then ∂K is C^1 , as it was verified by Chou and Wang [16].

3. SOME RESULTS ON MONGE-AMPÈRE EQUATIONS IN EUCLIDEAN SPACE

Lemma 3.1 is the tool to transfer the Monge-Ampère equation (1.2) on S^{n-1} to a Euclidean Monge-Ampère equation on \mathbb{R}^{n-1} . For $e \in S^{n-1}$, we consider the restriction of a solution h of (1.2) to the hyperplane tangent to S^{n-1} at e .

Lemma 3.1. *If $e \in S^{n-1}$, h is a convex positively 1-homogeneous non-negative function on \mathbb{R}^n that is a solution of (1.2) for $p < 1$ and positive f , and $v(y) = h(y + e)$ holds for $v : e^\perp \rightarrow \mathbb{R}$, then v satisfies*

$$(3.1) \quad v^{1-p} \det(D^2 v) = g \quad \text{on } e^\perp$$

where, for $y \in e^\perp$, we have

$$g(y) = (1 + \|y\|^2)^{-\frac{n+p}{2}} f\left(\frac{e + y}{\sqrt{1 + \|y\|^2}}\right).$$

Proof. Let $h = h_K$ for $K \in \mathcal{K}_0^n$, and let

$$\tilde{S} = \{u \in S^{n-1} : h_K(u) = 0\},$$

which is a possibly empty spherically convex compact set whose spherical dimension is at most $n - 2$, by (2.7). According to (2.9), the Monge-Ampère equation for h_K can be written in the form

$$(3.2) \quad dS_K = h_K^{p-1} f d\mathcal{H}^{n-1} \quad \text{on } S^{n-1} \setminus \tilde{S}.$$

We consider $\pi : e^\perp \rightarrow S^{n-1}$ defined by

$$\pi(x) = (1 + \|x\|^2)^{-\frac{1}{2}}(x + e),$$

which is induced by the radial projection from the tangent hyperplane $e + e^\perp$ to S^{n-1} . Since $\langle \pi(x), e \rangle = (1 + \|x\|^2)^{-\frac{1}{2}}$, the Jacobian of π is

$$(3.3) \quad \det D\pi(x) = (1 + \|x\|^2)^{-\frac{n}{2}}.$$

For $x \in e^\perp$, (2.4) and writing h_K in terms of an orthonormal basis of \mathbb{R}^n containing e , yield that v satisfies

$$\partial v(x) = \partial h_K(x + e)|_{e^\perp} = F(K, x + e)|_{e^\perp} = F(K, \pi(x))|_{e^\perp}.$$

Let $S = \pi^{-1}(\tilde{S})$. For a Borel set $\omega \subset e^\perp \setminus S$, we have

$$\begin{aligned} \mathcal{H}^{n-1}(N_v(\omega)) &= \mathcal{H}^{n-1}(\cup_{x \in \omega} \partial v(x)) \\ &= \mathcal{H}^{n-1}(\cup_{u \in \pi(\omega)} (F(K, u)|_{e^\perp})) = \int_{\pi(\omega)} \langle u, e \rangle dS_K(u) \\ &= \int_{\pi(\omega)} \langle u, e \rangle h_K^{p-1}(u) f(u) d\mathcal{H}^{n-1}(u) \\ &= \int_{\omega} (1 + \|x\|^2)^{-\frac{n-p}{2}} f(\pi(x)) v(x)^{p-1} d\mathcal{H}^{n-1}(x) \end{aligned}$$

where we used at the last step that

$$v(x) = h_K(x + e) = (1 + \|x\|^2)^{\frac{1}{2}} h_K(\pi(x)).$$

In particular, v satisfies the Monge-Ampère type differential equation

$$\det D^2 v(x) = (1 + \|x\|^2)^{-\frac{n-p}{2}} f(\pi(x)) v(x)^{p-1} \quad \text{on } e^\perp \setminus S.$$

Since $\dim S \leq n - 2$ by (1.3), v satisfies (3.1) on e^\perp . \square

Having Lemma 3.1 at hand showing the need to understand related Monge-Ampère equations in Euclidean spaces, we prove Propositions 1.3 and 3.4, and quote Caffarelli's Theorem 3.6.

Proof of Proposition 1.3. Up to changing coordinate system, we may assume, without loss of generality, that $S \subset \{(x_1, x_2) \in \mathbb{R}^r \times \mathbb{R}^{n-r} : x_2 = 0\}$ and the origin is contained in the relative interior of S . Therefore, up to restricting Ω , we may also assume that v is continuous on $\text{cl } \Omega$, that $\Omega = \{(x_1, x_2) \in \mathbb{R}^r \times \mathbb{R}^{n-r} : \|x_1\| < s_1, \|x_2\| < s_2\}$ for some constants $s_1, s_2 > 0$ and that $S = \{(x_1, x_2) \in \Omega : x_2 = 0\}$.

Let $\alpha = \max_{\text{cl } \Omega} v$ and let us consider the convex body

$$M = \{(x_1, x_2, y) \in \mathbb{R}^r \times \mathbb{R}^{n-r} \times \mathbb{R} : \|x_1\| \leq s_1, \|x_2\| \leq s_2, v(x_1, x_2) \leq y \leq \alpha\}.$$

For $t \in (0, s_2/2]$, let

$$\Omega_t = \{(x_1, x_2) \in \mathbb{R}^r \times \mathbb{R}^{n-r} : \|x_1\| \leq s_1/2, \|x_2\| \leq t\}.$$

We estimate $\mathcal{H}^n(N_v(\Omega_t \setminus S))$. Let $(x_1, x_2) \in \Omega_t \setminus S$ and let $(z_1, z_2) \in \mathbb{R}^r \times \mathbb{R}^{n-r}$ belong to $\partial v(x_1, x_2)$. We prove that

$$(3.4) \quad \|z_2\| \leq \frac{2\alpha}{s_2} \quad \text{and} \quad \|z_1\| \leq \frac{4\alpha}{s_1 s_2} t.$$

If $z_2 = 0$ the first inequality in (3.4) holds true. Assume $z_2 \neq 0$. The vector $(z_1, z_2, -1)$ is an exterior normal to M at $p = (x_1, x_2, v(x_1, x_2))$. Since

$$q_1 = \left(x_1, x_2 + \frac{s_2 z_2}{2\|z_2\|}, \alpha \right) \in M$$

(because $\|x_2 + s_2 z_2 / (2\|z_2\|)\| \leq \|x_2\| + s_2/2 \leq s_2$) then $\langle q_1 - p, (z_1, z_2, -1) \rangle \leq 0$. This implies

$$\|z_2\| \leq \frac{2}{s_2} (\alpha - v(x_1, x_2))$$

and the first inequality in (3.4). Again, if $z_1 = 0$ then the second inequality (3.4) holds true. Assume $z_1 \neq 0$. We have

$$q_2 = \left(x_1 + \frac{s_1 z_1}{2\|z_1\|}, 0, v(x_1, x_2) \right) \in M,$$

because $\|x_1 + s_1 z_1 / (2\|z_1\|)\| \leq s_1$, $(x_1 + s_1 z_1 / (2\|z_1\|), 0) \in S$ and therefore $v(x_1, x_2) \geq 0 = v(x_1 + s_1 z_1 / (2\|z_1\|), 0)$. The inequality $\langle q_2 - p, (z_1, z_2, -1) \rangle \leq 0$ implies the second inequality (3.4).

The inequalities in (3.4) imply

$$(3.5) \quad \mathcal{H}^n(N_v(\Omega_t \setminus S)) \leq c t^r,$$

for a suitable constant c independent of t .

Now we estimate $\int_{\Omega_t \setminus S} v(x)^{p-1} dx$. The inclusion of the convex hull of $S \times \{0\}$ and $\{\|x_1\| \leq s_1, \|x_2\| \leq s_2, y = \alpha\}$ in M implies that $v(x_1, x_2) \leq \frac{\alpha}{s_2} \|x_2\|$ for each $(x_1, x_2) \in \Omega_t$ by the convexity of v . Using this estimate it is straightforward to compute that

$$(3.6) \quad \int_{\Omega_t \setminus S} v(x)^{p-1} dx \geq d t^{n+p-r-1},$$

for a suitable constant d independent on t . The inequalities (3.5) and (3.6) and the differential inequality satisfied by v imply, as $t \rightarrow 0^+$,

$$c t^r \geq \mathcal{H}^n(N_v(\Omega_t \setminus S)) \geq \int_{\Omega_t \setminus S} \tau v(x)^{p-1} dx \geq \tau d t^{n+p-r-1}.$$

This inequality implies $p \geq -n + 1 + 2r$. \square

Example 3.2. *Let us show that for any $p > -n + 3$ there exists a non-negative convex solution of (1.6) in $\Omega = \{(x_1, x_2) \in \mathbb{R} \times \mathbb{R}^{n-1} : x_1 \in [-1, 1], \|x_2\| \leq 1\}$ which vanish on the 1-dimensional space $S = \{(x_1, x_2) \in \mathbb{R} \times \mathbb{R}^{n-1} : x_2 = 0\}$.*

To prove this let

$$v(x_1, x_2) = \|x_2\| + f(\|x_2\|)g(x_1)$$

where $f(r) = r^\alpha$, with $\alpha = (p + n - 1)/2$, and $g(x_1) = (1 + \beta x_1^2)$, with $\beta > 0$ sufficiently small. Note that $\alpha > 1$ exactly when $p > -n + 3$.

The function v is invariant with respect to rotations around the line containing S . To compute $\det D^2 v$ at an arbitrary point, it suffices to compute it at $(x_1, 0, \dots, 0, r)$, $r \geq 0$. We get

$$\begin{aligned} v_{x_1 x_1} &= f(r)g''(x_1), \\ v_{x_1 x_i} &= 0 && \text{when } 1 < i < n, \\ v_{x_1 x_n} &= f'(r)g'(x_1), \\ v_{x_i x_i} &= \frac{1}{r} + \frac{f'(r)}{r}g(x_1) && \text{when } 1 < i < n, \\ v_{x_i x_j} &= 0 && \text{when } i \neq j, (i, j) \neq (1, n), (i, j) \neq (n, 1), \\ v_{x_n x_n} &= f''(r)g(x_1). \end{aligned}$$

The function v is convex if β is sufficiently small. Indeed, the eigenvalues of $D^2 v$ are $\frac{1}{r} + \frac{f'(r)}{r}g(x_1)$, with multiplicity $n - 2$, and those of the matrix

$$\begin{pmatrix} f g'' & f' g' \\ f' g' & f'' g \end{pmatrix}.$$

The determinant of the latter matrix is

$$2\alpha\beta r^{2(\alpha-1)} \left(\alpha - 1 - (1 + \alpha)\beta x_1^2 \right),$$

which is positive if $\beta > 0$ is sufficiently small. Thus, all eigenvalues of D^2v are positive.

We get

$$\det D^2v = \left(f''gf g'' - (f'g')^2 \right) \left(\frac{1}{r} + \frac{f'}{r}g \right)^{n-2}$$

which has the same order as $r^{2\alpha-n}$ as $r \rightarrow 0^+$. Clearly v has order r , and $v^{1-p} \det D^2v$ has order $r^{2\alpha-n+1-p}$, which is uniformly bounded from above and below for our choice of α .

The next statement is a slight modification of Lemmas 3.2 and 3.3 from Trudinger and Wang [51]. Its proof closely follows that in [51] and is given here for completeness.

Lemma 3.3. *Let v be a convex function defined on the closure of an open bounded convex set $\Omega \subset \mathbb{R}^n$ satisfying the Monge-Ampère equation*

$$\det D^2v = \nu$$

for a finite non-negative measure ν on Ω , let $v \equiv 0$ on $\partial\Omega$ and let $tE \subset \Omega \subset E$ for $t > 0$ and an origin centred ellipsoid E .

(i) *If $z \in \Omega$ satisfies $(z + sE) \cap \partial\Omega \neq \emptyset$ for $s > 0$, then*

$$|v(z)| \leq s^{1/n} c_0 \mathcal{H}^n(\Omega)^{1/n} \nu(\Omega)^{1/n}$$

for some $c_0 > 0$ depending on n, t .

(ii) *If $\nu(t\Omega) \geq b\nu(\Omega)$ for $b > 0$, then*

$$(3.7) \quad |v(0)| \geq c_1 \mathcal{H}^n(\Omega)^{1/n} \nu(\Omega)^{1/n}$$

for some $c_1 > 0$ depending on n, t and b .

(iii) *If $(z + sE) \cap \partial\Omega \neq \emptyset$ and $\nu(t\Omega) \geq b\nu(\Omega)$ then*

$$(3.8) \quad \frac{|v(z)|}{|v(o)|} \leq \frac{c_1}{c_0} s^{1/n}.$$

When $E = B^n$ the number s can be chosen as the distance of z from $\partial\Omega$. In the general case s has the same meaning in the metric induced by the norm whose unitary ball is E .

Proof. Let A be a linear transformation such that $B^n = A^{-1}E$, let $\tilde{v}(x) = v(Ax)|\det A|^{-2/n}$, $\tilde{\Omega} = A^{-1}\Omega$ and let $\tilde{\nu}$ be the measure defined for each Borel set $\omega \subset \tilde{\Omega}$ as $\tilde{\nu}(\omega) = \nu(A\omega)/|\det A|$. It is known that \tilde{v} solves

$$(3.9) \quad \det D^2\tilde{v} = \tilde{\nu} \quad \text{in } \tilde{\Omega}.$$

Moreover, $tB^n \subset \tilde{\Omega} \subset B^n$. Since $\mathcal{H}^n(\Omega) = |\det A| \mathcal{H}^n(\tilde{\Omega})$, we have

$$(3.10) \quad \frac{\mathcal{H}^n(\Omega)}{\omega_n} \leq |\det A| \leq \frac{\mathcal{H}^n(\Omega)}{\omega_n t^n}.$$

Let us prove Claim (i). Let $\tilde{z} = A^{-1}z$. Then $(\tilde{z} + sB^n) \cap \partial\tilde{\Omega} \neq \emptyset$ and if d denotes the distance of \tilde{z} from $\partial\tilde{\Omega}$ we have $d \leq s$. By choosing proper coordinates we may assume that $\tilde{z} = (0, \dots, 0, d)$, and that $\tilde{\Omega} \subset \{(x_1, \dots, x_n) \in \mathbb{R}^n : x_n > 0\}$. Then

$$\tilde{\Omega} \subset \hat{\Omega} = \{(x_1, \dots, x_n) \in \mathbb{R}^n : \|(x_1, \dots, x_{n-1})\| < 2, 0 < x_n < 4\}.$$

Let u and w be convex functions such that their graphs are convex cones with vertex at $(\tilde{z}, \tilde{v}(\tilde{z}))$ and bases $\partial\tilde{\Omega}$ and $\partial\hat{\Omega}$, respectively. Then

$$(3.11) \quad N_{\tilde{v}}(\tilde{\Omega}) \supset N_u(\tilde{\Omega}) = \partial u(\tilde{z}) \supset \partial w(\tilde{z}).$$

Since w is a convex cone over the cylinder $\widehat{\Omega}$, one can easily compute that $\mathcal{H}^n(\partial w(\tilde{z})) \geq c_2 |\tilde{v}(\tilde{z})|^n/d$, for a suitable constant $c_2 > 0$. This inequality, (3.9) and (3.11) imply

$$|\tilde{v}(\tilde{z})| \leq \left(\frac{d}{c_2}\right)^{1/n} \mathcal{H}^n(N_{\tilde{v}}(\widehat{\Omega}))^{1/n} = \left(\frac{d}{c_2}\right)^{1/n} \tilde{v}(\widehat{\Omega})^{1/n}.$$

Expressing this inequality in terms of v , Ω and ν and using $d \leq s$ and (3.10) concludes the proof of Claim (i).

Let us prove Claim (ii). There exists a unique solution w of $\det D^2 w = \widehat{v}$ in $\widetilde{\Omega}$, $w = 0$ in $\partial\widetilde{\Omega}$, where $\widehat{v} = \tilde{v}$ in $t\widetilde{\Omega}$ and $\widehat{v} = 0$ elsewhere (see Theorem 2.1 in [51]). The comparison principle for Monge-Ampère equations (see Lemma 2.4 in [51]) implies $w \geq \tilde{v}$ in $\widetilde{\Omega}$.

Let $z \in t\widetilde{\Omega}$. The distance d of z from $\partial\widetilde{\Omega}$ is larger than or equal to $(1-t)t$ (here we have used the inclusion $tB^n \subset \widetilde{\Omega}$). If $y \in \partial w(z)$ and $l(x) = \langle x, y \rangle + w(z)$ then $l(x) \leq w(x)$ for each $x \in \widetilde{\Omega}$, by definition of subgradient. In particular, we have $l(x) \leq 0$ for each $x \in \partial\widetilde{\Omega}$. This implies

$$|y| \leq \frac{|w(z)|}{d} \leq \frac{\sup_{\widetilde{\Omega}} |w|}{t(1-t)}.$$

Therefore,

$$\mathcal{H}^n(N_w(t\widetilde{\Omega})) \leq \omega_n \left(\frac{\sup_{\widetilde{\Omega}} |w|}{t(1-t)} \right)^n.$$

This inequality, the equation satisfied by w and the condition $\nu(t\Omega) \geq b\nu(\Omega)$ imply

$$\begin{aligned} \sup_{\widetilde{\Omega}} |\tilde{v}| &\geq \frac{t(1-t)}{\omega_n^{1/n}} \mathcal{H}^n(N_w(t\widetilde{\Omega}))^{1/n} = \frac{t(1-t)}{\omega_n^{1/n}} \tilde{v}(t\widetilde{\Omega})^{1/n} \\ (3.12) \quad &\geq \frac{bt(1-t)}{\omega_n^{1/n}} \tilde{v}(\widetilde{\Omega})^{1/n}. \end{aligned}$$

We claim that

$$(3.13) \quad |\tilde{v}(o)| \geq \frac{t}{1+t} \sup_{\widetilde{\Omega}} |\tilde{v}|.$$

Indeed, let $z \in \widetilde{\Omega}$ be such that $\tilde{v}(z) = \inf_{\widetilde{\Omega}} \tilde{v}$. We may clearly assume $z \neq 0$, since otherwise there is nothing to prove. By choosing proper coordinates we may assume $z = (z_1, 0, \dots, 0)$ for some $z_1 > 0$. Let l be the linear function defined on the line through o and z and such that $l(o) = \tilde{v}(o)$ and $l(z) = \tilde{v}(z)$. It is $l(s, 0, \dots, 0) = \tilde{v}(o) + s(\inf_{\widetilde{\Omega}} \tilde{v} - \tilde{v}(o))/z_1$. Since \tilde{v} is convex,

$$l(s, 0, \dots, 0) \leq \tilde{v}(s, 0, \dots, 0)$$

for each $s \notin [0, z_1]$ such that $(s, 0, \dots, 0) \in \widetilde{\Omega}$. When $s = -t$ we obtain $l(-t, 0, \dots, 0) \leq \tilde{v}(-t, 0, \dots, 0) \leq 0$. The inequality $l(-t, 0, \dots, 0) \leq 0$ and the inclusion $\widetilde{\Omega} \subset B^n$ imply (3.13).

The proof of Claim (ii) is concluded by combining (3.12) and (3.13) and expressing the obtained inequality in terms of v , Ω and ν .

Claim (iii) is a consequence of the first two claims. \square

The proof of Claim (ii) in Theorem 1.2 is based on the following proposition, which is related to a step in the proof of Theorem E (a) in [16], however our proof is substantially different from that in [16].

Proposition 3.4. *Let v be a non-negative convex function defined on the closure of an open convex set $\Omega \subset \mathbb{R}^n$, $n \geq 2$, such that $S = \{x \in \Omega : v(x) = 0\}$ is non-empty*

and compact, and v is locally strictly convex on $\Omega \setminus S$. Let $\psi : (0, \infty) \rightarrow [0, \infty)$ be monotone decreasing and not identically zero; assume that $\tau_2 > \tau_1 > 0$ and v satisfy

$$(3.14) \quad \tau_1 \psi(v) \leq \det D^2 v \leq \tau_2 \psi(v)$$

in the sense of measure on $\Omega \setminus S$. If $\dim S \leq n - 1$ and $\mu_v(S) = 0$ for the associated Monge-Ampère measure μ_v , then S is a point.

Note that (3.14) means that for each Borel set $\omega \subset \Omega \setminus S$ we have

$$\tau_1 \int_{\omega} \psi(v(x)) dx \leq \mu_v(\omega) \leq \tau_2 \int_{\omega} \psi(v(x)) dx,$$

where μ_v has been defined in (2.1).

Proof. We assume, arguing by contradiction that S is not a point. Choose coordinates so that o is the centre of mass of S . Let $L = \text{lin } S$. By assumption

$$(3.15) \quad 1 \leq \dim L \leq n - 1.$$

Let $e = (o, 1) \in \mathbb{R}^n \times \mathbb{R}$. We may assume that Ω is bounded, after possibly substituting it with a bounded open neighbourhood of S . We start by illustrating the idea of the proof.

Sketch of the proof. For any small $\varepsilon > 0$, we construct an affine function l_ε such that $l_\varepsilon(x) = \varepsilon$ for $x \in L$, and the convex set $\Omega_\varepsilon = \{v < l_\varepsilon\}$ is well-balanced; namely, there exists an ellipsoid E_ε centred at the origin such that $(1/(8n^3))E_\varepsilon \subset \Omega_\varepsilon \subset E_\varepsilon$ (see (3.19)). This is the longest part of the argument, and the main idea to construct l_ε is that the graph of l_ε cuts off the smallest volume cap from the graph of v among the hyperplanes in \mathbb{R}^{n+1} containing $L + \varepsilon e$. Subsequently, we apply Lemma 3.3 to Ω_ε and to the function $v - l_\varepsilon$ in the standard way to reach a contradiction. We show that one can choose $z \in S$ so that the corresponding parameter s , as defined in Lemma 3.3, tends to 0 as ε tends to 0. (Equivalently, S contains points whose distance from $\partial\Omega_\varepsilon$, the one induced by the norm whose unit ball is E_ε , tends to 0 as ε tends to 0.) This contradicts (3.8), since $|v(z) - l_\varepsilon(z)|/|v(o) - l_\varepsilon(o)| = \varepsilon/\varepsilon = 1$.

We divide the proof into four steps.

Step 1. Definition of l_ε and of Ω_ε .

Let $\varepsilon_0 = \min_{\partial\Omega} v > 0$ and let us consider the $(n + 1)$ -dimensional convex body

$$M = \{(x, y) \in \mathbb{R}^n \times \mathbb{R} : v(x) \leq y \leq \varepsilon_0\}.$$

For $\varepsilon \in (0, \varepsilon_0)$ define H_ε to be a hyperplane in \mathbb{R}^{n+1}

- (i) containing $L + \varepsilon e = \{(x, \varepsilon) \in \mathbb{R}^n \times \mathbb{R} : x \in L\}$ and
- (ii) cutting off the minimal volume from M (on the side containing the origin) under condition (i).

Let $r > 0$. We claim that there exists $\varepsilon_1 = \varepsilon_1(r)$ so that H_ε is the graph of an affine function l_ε for each $\varepsilon \in (0, \varepsilon_1)$, and, setting

$$\Omega_\varepsilon = \{x \in \mathbb{R}^n : v(x) < l_\varepsilon(x)\},$$

we have

$$(3.16) \quad \text{cl}\Omega_\varepsilon \subset \Omega, \quad S \subset \Omega_\varepsilon \quad \text{and} \quad \Omega_\varepsilon \cap L \subset (1 + r)S.$$

Let $F = \{(x, y) \in M : y = \varepsilon_0\}$ be the upper face of M and let \mathcal{H} be the collection of hyperplanes in \mathbb{R}^{n+1} which intersect both F and $\{(x, y) \in M : y \leq \varepsilon_0/2\}$. Since Ω is bounded and v is locally strictly convex on $\Omega \setminus S$, every hyperplane in \mathcal{H} is not a supporting hyperplane to M . Therefore, by compactness, there exists a constant $\varrho_0 > 0$ such that for every $H \in \mathcal{H}$ both components of $M \setminus H$ are of volume at least ϱ_0 . We choose $\varepsilon_1 \in (0, \varepsilon_0/2)$ such that the volume of the cap $\{(x, y) \in M : y \leq \varepsilon_1\}$ is less than ϱ_0 . This choice implies that the minimum value of the problem which

defines H_ε is less than ϱ_0 . Therefore, a minimizer H_ε does not belong to \mathcal{H} . Since $H_\varepsilon \cap \{(x, y) \in M : y \leq \varepsilon_0/2\} \neq \emptyset$, we have $H_\varepsilon \cap F = \emptyset$. In particular, H_ε is the graph of an affine function defined on \mathbb{R}^n and $\text{cl}\Omega_\varepsilon \subset \Omega$.

The inclusion $S \subset \Omega_\varepsilon$ holds because $v(x) = 0$ and $l_\varepsilon(x) = \varepsilon$ for any $x \in S$.

The origin o , being the centre of mass of S , belongs to the relative interior of S . Since $\dim S > 0$, the relative boundary of $(1+r)S$ does not intersect S . This implies $\inf_{\text{relbd}(1+r)S} v > 0$. Thus, if ε_1 satisfies

$$\varepsilon_1 < \inf_{\text{relbd}(1+r)S} v$$

in addition to the inequalities specified above then $v(x) > \varepsilon$ and $l_\varepsilon(x) = \varepsilon$ for any $x \in \text{relbd}(1+r)S$ ($l_\varepsilon(x) = \varepsilon$ is a consequence of $(1+r)S \subset L$). This implies $\Omega_\varepsilon \cap L \subset (1+r)S$.

In the rest of the proof we may assume $\varepsilon_1 < \varepsilon_1(1)$ so that

$$(3.17) \quad \Omega_\varepsilon \cap L \subset 2S.$$

Step 2. The centre of mass of Ω_ε is contained in L .

To prove this claim we have to prove that for each $w \in L^\perp \cap \mathbb{R}^n$ we have

$$(3.18) \quad \int_{\Omega_\varepsilon} \langle x, w \rangle dx = 0.$$

Indeed, for $t \in \mathbb{R}$ with $|t|$ small, let

$$F(t) = \int_{\{x \in \Omega : l_\varepsilon(x) + t\langle x, w \rangle - v(x) > 0\}} (l_\varepsilon(x) + t\langle x, w \rangle - v(x)) dx$$

be the volume cut off by the hyperplane in \mathbb{R}^{n+1} that is the graph of $x \mapsto l_\varepsilon(x) + t\langle x, w \rangle$ from M . By definition of H_ε and l_ε , F has a local minimum at $t = 0$. We have

$$\begin{aligned} \frac{F(t) - F(0)}{t} &= \int_{\{x \in \Omega : l_\varepsilon(x) - v(x) > 0\}} \langle x, w \rangle dx \\ &+ \int_{\Omega} \left(\frac{l_\varepsilon(x) - v(x)}{t} + \langle x, w \rangle \right) \left(1_{\{x : l_\varepsilon(x) + t\langle x, w \rangle - v(x) > 0\}} - 1_{\{x : l_\varepsilon(x) - v(x) > 0\}} \right) dx. \end{aligned}$$

The set where $1_{\{x : l_\varepsilon(x) + t\langle x, w \rangle - v(x) > 0\}} - 1_{\{x : l_\varepsilon(x) - v(x) > 0\}}$ differs from 0 is contained in

$$A_t = \{x \in \Omega : |l_\varepsilon(x) - v(x)| < |t\langle x, w \rangle|\}$$

and there exists c independent on t such that $\mathcal{H}^n(A_t) < ct$ and $\sup_{A_t} |l_\varepsilon(x) - v(x)| < ct$. As F has a local minimum at $t = 0$, we have

$$0 = \frac{dF}{dt}(0) = \int_{\Omega_\varepsilon} \langle x, w \rangle dx,$$

which proves (3.18).

Step 3. For any $\varepsilon \in (0, \varepsilon_1)$ there exists an ellipsoid E_ε centred at the origin such that

$$(3.19) \quad \frac{1}{8n^3} E_\varepsilon \subset \Omega_\varepsilon \subset E_\varepsilon.$$

Lemma 2.3.3 in [47] proves that any k -dimensional convex body contains its reflection, with respect to its centre of mass, scaled, with respect to the same centre of mass, by $1/k$. From the fact that the centre of mass of Ω_ε belongs to L we deduce that

$$(3.20) \quad -(\Omega_\varepsilon|L^\perp) \subset n(\Omega_\varepsilon|L^\perp).$$

According to Loewner's or John's theorems, there exists an ellipsoid \tilde{E} centred at the origin and $z_1 \in \Omega_\varepsilon$ such that

$$z_1 + \frac{1}{n} \tilde{E} \subset \Omega_\varepsilon \subset z_1 + \tilde{E}.$$

It follows from (3.20) that there exists $z_2 \in \Omega_\varepsilon$ such that $z_2|_{L^\perp} = \frac{-1}{n} z_1|_{L^\perp}$. In particular, $y_1 = \frac{1}{n+1} z_1 + \frac{n}{n+1} z_2 \in \Omega_\varepsilon$ verifies $y_1|_{L^\perp} = o$, or in other words, $y_1 \in L \cap \Omega_\varepsilon$. In addition,

$$y_1 + \frac{1}{2n^2} \tilde{E} \subset \frac{1}{n+1} \left(z_1 + \frac{1}{n} \tilde{E} \right) + \frac{n}{n+1} z_2 \subset \Omega_\varepsilon.$$

Let $m = \dim L \leq n-1$. Since $y_1 \in L \cap \Omega_\varepsilon$ and (3.17) imply $\frac{1}{2} y_1 \in S$, and since the origin is the centroid of S , we deduce that $y_2 = \frac{-1}{2m} y_1 \in S$. As $2m+1 < 2n$, we have

$$\frac{1}{4n^3} \tilde{E} \subset \frac{1}{2m+1} \left(y_1 + \frac{1}{2n^2} \tilde{E} \right) + \frac{2m}{2m+1} y_2 \subset \Omega_\varepsilon.$$

As $\Omega_\varepsilon \subset 2\tilde{E}$ follows from $o \in z_1 + \tilde{E}$, we may choose $E_\varepsilon = 2\tilde{E}$, proving (3.19).

Step 4. Application of Lemma 3.3 to $v - l_\varepsilon$ and Ω_ε and contradiction.

We observe that

$$(3.21) \quad v(x) - l_\varepsilon(x) = \begin{cases} 0 & \text{if } x \in \partial\Omega_\varepsilon \\ -\varepsilon & \text{if } x \in S. \end{cases}$$

Let ν denote the Monge-Ampère measure $\mu_{(v-l_\varepsilon)}$ restricted to Ω_ε . If Ω_0 is an open set such that $\Omega_\varepsilon \subset \Omega_0 \subset \text{cl}\Omega_0 \subset \Omega$, then the set $N_\nu(\Omega_0)$ is bounded and this implies

$$\nu(\Omega_\varepsilon) = \mathcal{H}^n(N_{(v-l_\varepsilon)}(\Omega_\varepsilon)) \leq \mathcal{H}^n(N_\nu(\Omega_0)) < \infty.$$

Let $t = 1/(8n^3)$. Formula (3.19) yields $tE_\varepsilon \subset \Omega_\varepsilon \subset E_\varepsilon$. Let us prove that

$$(3.22) \quad \nu(t\Omega_\varepsilon) \geq b\nu(\Omega_\varepsilon) \quad \text{for } b = \tau_1 t^n / \tau_2.$$

The function v is convex and attains its minimum at o , thus $v(x) \geq v(tx)$ for any $x \in \Omega_\varepsilon$. By this fact, the monotonicity of ψ , (3.14) and the assumptions on S , we deduce that

$$\begin{aligned} \nu(t\Omega_\varepsilon) &= \nu(t(\Omega_\varepsilon \setminus S)) \geq \tau_1 \int_{t(\Omega_\varepsilon \setminus S)} \psi(v(x)) dx \\ &= \tau_1 t^n \int_{\Omega_\varepsilon \setminus S} \psi(v(tz)) dz \\ &\geq \tau_1 t^n \int_{\Omega_\varepsilon \setminus S} \psi(v(z)) dz \\ &\geq \frac{\tau_1 t^n}{\tau_2} \nu(\Omega_\varepsilon \setminus S) = \frac{\tau_1 t^n}{\tau_2} \nu(\Omega_\varepsilon) \end{aligned}$$

proving (3.22).

Let $z \in \text{relbd}S$. We claim that when $\varepsilon \in (0, \varepsilon_1(r))$ then $(z + rE_\varepsilon) \cap \partial\Omega_\varepsilon \neq \emptyset$. This is a consequence of the second and third inclusion in (3.16). Indeed, since $o \in S \subset \Omega_\varepsilon \subset E_\varepsilon$, there exists $q_\varepsilon > 0$ such that $(1 + q_\varepsilon)z \in \partial E_\varepsilon$. The set $z + rE_\varepsilon$ contains the segment $[z, z + r(1 + q_\varepsilon)z]$. Since $q_\varepsilon > 0$, that segment contains the segment $[z, (1 + r)z]$. The second and third inclusion in (3.16) imply $[z, (1 + r)z] \cap \partial\Omega_\varepsilon \neq \emptyset$. This proves the claim.

Lemma 3.3 applies to this situation with $s = r$. Since $v(z) - l_\varepsilon(z) = v(o) - l_\varepsilon(o) = -\varepsilon$ (see (3.21)), (3.8) yields

$$1 = \frac{|v(z) - l_\varepsilon(z)|}{|v(o) - l_\varepsilon(o)|} \leq \frac{c_1}{c_0} r^{1/n}.$$

Since r can be any positive number, we have reached a contradiction. \square

We will actually use the following consequence of Proposition 3.4.

Corollary 3.5. *Let $\tau_2 > \tau_1 > 0$, and let g be a function defined on an open convex set $\Omega \subset \mathbb{R}^n$, $n \geq 2$, such that $\tau_2 > g(x) > \tau_1$ for $x \in \Omega$. For $p < 1$, let v be a non-negative convex solution of*

$$v^{1-p} \det D^2 v = g \quad \text{in } \Omega.$$

If $S = \{x \in \Omega : v(x) = 0\}$ is non-empty, compact and $\mu_v(S) = 0$, and v is locally strictly convex on $\Omega \setminus S$, then S is a point.

Proof. All we have to check that $\dim S \leq n - 1$. It follows from the fact that the left-hand side of the differential equation is zero on S , while the right-hand side is positive. \square

The following result by L. Caffarelli (see Theorem 1 and Corollary 1 in [7]) is the key in handling the regularity and strict convexity of the part of the boundary of a convex body K where the support function at some normal vector is positive.

Theorem 3.6 (Caffarelli). *Let $\lambda_2 > \lambda_1 > 0$, and let v be a convex function on an open convex set $\Omega \subset \mathbb{R}^n$ such that*

$$(3.23) \quad \lambda_1 \leq \det D^2 v \leq \lambda_2$$

in the sense of measure.

- (i) *If v is non-negative and $S = \{x \in \Omega : v(x) = 0\}$ is not a point, then S has no extremal point in Ω .*
- (ii) *If v is strictly convex, then v is C^1 .*

We recall that (3.23) is equivalent to saying that for each Borel set $\omega \subset \Omega$ we have

$$\lambda_1 \mathcal{H}^n(\omega) \leq \mu_v(\omega) \leq \lambda_2 \mathcal{H}^n(\omega),$$

where μ_v has been defined in (2.1).

4. PROOF OF THEOREM 1.1

The next lemma provides a tool for the proof of Theorem 1.1 (iii). The same result is also proved in Chou and Wang [16]; we present a short argument for the sake of completeness.

Lemma 4.1. *For $n \geq 2$ and $p \leq 2 - n$, if $K \in \mathcal{K}_0^n$ and there exists $c > 0$ such that $S_{K,p}(\omega) \geq c \mathcal{H}^{n-1}(\omega)$ for any Borel set $\omega \subset S^{n-1}$, then $o \in \text{int } K$.*

Proof. We suppose that $o \in \partial K$ and seek a contradiction. We choose $e \in N(K, o) \cap S^{n-1}$ such that $\{\lambda e : \lambda \geq 0\}$ is an extremal ray of $N(K, o)$. Let H^+ be a closed half space containing $\mathbb{R}e$ on the boundary such that $N(K, o) \cap \text{int } H^+ = \emptyset$. Let

$$V_0 = S^{n-1} \cap (e + B^n) \cap \text{int } H^+.$$

It follows by the condition on $S_{K,p}$ that

$$(4.1) \quad c \int_{V_0} h_K(u)^{p-1} d\mathcal{H}^{n-1} \leq \int_{V_0} h_K(u)^{p-1} dS_{K,p} = S_K(V_0) < \infty.$$

However, since h_K is convex and $h_K(e) = 0$, there exists $c_0 > 0$ such that

$$h_K(x) \leq c_0 \|x - e\| \quad \text{for } x \in e + B^n.$$

We observe that the radial projection of V_0 onto the tangent hyperplane $e + e^\perp$ to S^{n-1} at e is $e + V'_0$ for

$$V'_0 = e^\perp \cap (\sqrt{3} B^n) \cap \text{int } H^+.$$

If $y \in V'_0$, then $u = (e + y)/\|e + y\|$ verifies $\|u - e\| \geq \|y\|/2$. It follows that

$$\begin{aligned} \int_{V'_0} h_K(u)^{p-1} d\mathcal{H}^{n-1} &\geq c_0^{p-1} \int_{V'_0} \|u - e\|^{p-1} d\mathcal{H}^{n-1}(u) \\ &\geq \frac{c_0^{p-1}}{2} \int_{V'_0} \frac{\|y\|^{p-1}}{(1 + \|y\|^2)^{n/2}} d\mathcal{H}^{n-1}(y) \\ &\geq \frac{c_0^{p-1}}{2^{n+1}} \int_{V'_0} \|y\|^{p-1} d\mathcal{H}^{n-1}(y) = \infty \end{aligned}$$

as $p \leq 2 - n$. This contradicts (4.1), and hence verifies the lemma. \square

Proof of Theorem 1.1. Claim (i). For $u_0 \in S^{n-1} \setminus N(K, o)$, we choose a spherically convex open neighbourhood Ω_0 of u_0 on S^{n-1} such that for any $u \in \text{cl}\Omega_0$, we have $\langle u, u_0 \rangle > 0$ and $u \notin N(K, o)$. Let $\Omega \subset u_0^\perp$ be defined in a way such that $u_0 + \Omega$ is the radial image of Ω_0 into $u_0 + u_0^\perp$, and let v be the function on Ω defined as in Lemma 3.1 with $h = h_K$. Since h_K is positive and continuous on $\text{cl}\Omega$, we deduce from Lemma 3.1 that there exist $\lambda_2 > \lambda_1 > 0$ depending on K , u_0 and Ω_0 such that

$$(4.2) \quad \lambda_1 \leq \det D^2v \leq \lambda_2$$

on Ω .

First we claim that

$$(4.3) \quad \text{if } z \in \partial K \text{ and } N(K, z) \not\subset N(K, o), \text{ then } z \text{ is a } C^1\text{-smooth point.}$$

We suppose that $\dim N(K, z) \geq 2$, and seek a contradiction. Since $N(K, z)$ is a closed convex cone such that o is an extremal point, the property $N(K, z) \not\subset N(K, o)$ yields an $e \in (N(K, z) \cap S^{n-1}) \setminus N(K, o)$ generating an extremal ray of $N(K, z)$. We apply the construction above for $u_0 = e$. The convexity of h_K and (2.2) imply $h_K(x) \geq \langle z, x \rangle$ for $x \in \mathbb{R}^n$, with equality if and only if $x \in N(K, z)$. We define $S \subset \Omega$ by $S + e = N(K, z) \cap (\Omega + e)$ and hence o is an extremal point of S . It follows that the function \tilde{v} defined by $\tilde{v}(y) = v(y) - \langle z, y + e \rangle$ is non-negative on Ω , satisfies (4.2), and

$$S = \{y \in \Omega : \tilde{v}(y) = 0\}.$$

These properties contradict Caffarelli's Theorem 3.6 (i) as o is an extremal point of S , and in turn we conclude (4.3).

Next we show that

$$(4.4) \quad h_K \text{ is differentiable at any } u_0 \in S^{n-1} \setminus N(K, o).$$

We apply again the construction above for u_0 . If $u \in \Omega_0$ and $z \in F(K, u)$ clearly K is C^1 -smooth at z (i.e. $N(K, z)$ is a ray) by (4.3). Therefore, by (2.3), v is strictly convex on Ω and Caffarelli's Theorem 3.6 (ii) yields that v is C^1 on Ω . In turn, we conclude (4.4).

In addition, $F(K, u)$ is a unique C^1 -smooth point for $u \in \Omega_0$ (see (2.4)), yielding that $\Omega_* = \cup\{F(K, u) : u \in \Omega_0\}$ is an open subset of ∂K . Therefore $\Omega_* \subset X$, any point of Ω_* is C^1 -smooth (by (2.3)) and Ω_* contains no segment (by (2.4)), completing the proof of Claim (i).

Claim (ii). We suppose that $o \in \partial K$ is C^1 -smooth, and there exists $z \in \partial K$ such that K is not C^1 -smooth at z . Claim (i) yields that $z \in X_0$, and hence $N(K, z) \subset N(K, o)$, which is a contradiction, verifying Claim (ii).

Claim (iii). This is a consequence of Lemma 4.1 and Claim (i).

Claim (iv). This is a consequence of Lemma 3.1, Claim (i) and Caffarelli [8]. \square

Example 4.2. If $n \geq 2$ and $p \in (-n+2, 1)$, then there exists $K \in \mathcal{K}_0^n$ with C^1 boundary such that o lies in the relative interior of a facet of ∂K and $dS_{K,p} = f d\mathcal{H}^{n-1}$ for a strictly positive continuous $f : S^{n-1} \rightarrow \mathbb{R}$.

Let $q = (p+n-1)/(p+n-2)$. We have $q > 1$. Let

$$g(r) = \begin{cases} (r-1)^q & \text{when } r \geq 1; \\ 0 & \text{when } r \in [0, 1); \end{cases}$$

and $\bar{g}(x_1, \dots, x_{n-1}) = g(\|(x_1, \dots, x_{n-1})\|)$. Let $K \in \mathcal{K}_0^n$ be such that $K \cap \{x : x_n \leq 1\} = \{x : 1 \geq x_n \geq \bar{g}(x_1, \dots, x_{n-1})\}$ and $\partial K \cap \{x : x_n > 0\}$ is a C^2 surface with Gauss curvature positive at every point. Clearly $K \cap \{x : x_n = 0\}$ is a $(n-1)$ -dimensional face of K which contains o in its relative interior and has unit outer normal $(0, \dots, 0, -1)$.

To prove that $dS_{K,p} = f d\mathcal{H}^{n-1}$ for a positive continuous $f : S^{n-1} \rightarrow \mathbb{R}$, it suffices to prove that there is a neighbourhood of the South pole where $dS_{K,p}/d\mathcal{H}^{n-1}$ is continuous and bounded from above and below by positive constants. Let h be the support function of K and, for $y \in \mathbb{R}^{n-1}$, let $v(y) = h(y, -1)$ be the restriction of h to the hyperplane tangent to S^{n-1} at the South pole. It suffices to prove that in a neighbourhood U of o , v satisfies the equation $v^{1-p} \det D^2 v = G$ with a function G which is bounded from above and below by positive constants.

If $y \in U \setminus \{o\}$ we have

$$(4.5) \quad v(y) = h(y, -1) = \langle (x', \bar{g}(x')), (y, -1) \rangle \quad \text{where} \quad D\bar{g}(x') = y.$$

If U is sufficiently small then $v(y)$ depends only on $\|y\|$. Let $y = (z, 0, \dots, 0)$, with $z > 0$ small and let $r = 1 + (z/q)^{1/(q-1)}$. We have

$$D\bar{g}(r, 0, \dots, 0) = (z, 0, \dots, 0)$$

and (4.5) gives

$$\begin{aligned} v(z, 0, \dots, 0) &= rq(r-1)^{q-1} - (r-1)^q \\ &= z + \frac{q-1}{q^{n-1+p}} z^{n-1+p}. \end{aligned}$$

(Note that $n-1+p > 1$.) Clearly $v(0, \dots, 0) = h(0, \dots, 0, -1) = 0$. When $z > 0$ we have

$$\begin{aligned} v_{y_1 y_1} &= \frac{q-1}{q^{n-1+p}} (n-1-p)(n-2-p) z^{n-3+p} \\ v_{y_i y_i} &= \frac{1}{z} + \frac{q-1}{q^{n-1+p}} (n-1-p) z^{n-3+p} \quad \text{when } i \neq 1 \\ v_{y_i y_j} &= 0 \quad \text{when } i \neq j, \end{aligned}$$

and, as $z \rightarrow 0^+$

$$v(z, 0, \dots, 0)^{1-p} \det D^2 v(z, 0, \dots, 0) = c + o(1),$$

for a suitable constant $c > 0$. This implies the existence of a function G positive and continuous on U such that

$$\mathcal{H}^{n-1}(N_v(\omega \cap \{v > 0\})) = \int_{\omega \cap \{v > 0\}} nG(y)v(y)^{p-1} dy.$$

for any Borel set $\omega \subset U$. To conclude the proof that v is a solution in the sense of Alexandrov of $v^{1-p} \det D^2 v = G$ in U it remains to prove that $\mathcal{H}^{n-1}(\{y \in U : v(y) = 0\}) = 0$, but this is obvious since $\{y \in U : v(y) = 0\} = \{o\}$.

We remark that h is not a solution of (1.4) because (2.11) fails.

5. PROOFS OF THEOREM 1.2 AND COROLLARY 1.4

Proof of Theorem 1.2. We may assume that $o \in \partial K$ since otherwise ∂K is C^1 by Theorem 1.1. Let $e \in N(K, o) \cap S^{n-1}$ be such that $\langle u, e \rangle > 0$ for any $u \in N(K, o) \cap S^{n-1}$. Let v be defined on $\Omega = e^\perp$ as in Lemma 3.1 with $h = h_K$ and let $S = \{x \in e^\perp : v(x) = 0\}$. We have

$$(5.1) \quad S + e = N(K, o) \cap (e^\perp + e),$$

by (2.2). If K is not C^1 -smooth at o then $\dim S \geq 1$ and, by Proposition 1.3, $p \geq n - 4$ (note that here the dimension of the ambient space is $n - 1$). This proves Theorem 1.2 (i).

To prove Theorem 1.2 (ii) we observe that

$$N_{h_K}(e + S) = \bigcup_{u \in N(K, o)} F(K, u) = X_0,$$

where X_0 is defined as in Theorem 1.1 (i). The equality on the left in this formula follows by (2.4) and the equality on the right follows by Theorem 1.1 (i). Thus,

$$N_v(S) = X_0|e^\perp,$$

and if $\mathcal{H}^{n-1}(X_0) = 0$ then $\mu_v(S) = 0$. We observe that S is compact, by (5.1), that v is locally strictly convex, by Theorem 1.1 (i), and that $\dim S \leq n - 2$, by (1.3). Hence, Theorem 1.2 (ii) follows by Corollary 3.5 and (5.1). \square

Proof of Corollary 1.4. Claim (i) is an immediate consequence of (2.2), Proposition 1.3 and Lemma 3.1. This claim implies that when $n = 4$ or $n = 5$ and ∂K is not C^1 then $\dim N(K, o) = 2$. In this case $N(K, o) \cap S^{n-1}$ is a closed arc: let e_1 and e_2 be its endpoints. If $u \in N(K, o) \cap S^{n-1}$, $u \neq e_1$, $u \neq e_2$, then $F(K, u)$ is contained in the intersection of the two supporting hyperplanes $\{x \in \mathbb{R}^n : \langle x, e_i \rangle = h_K(e_i)\}$, $i = 1, 2$. Thus,

$$\mathcal{H}^{n-1}\left(\bigcup\{F(K, u) : u \in N(K, o) \cap S^{n-1}, u \neq e_1, u \neq e_2\}\right) = 0.$$

Therefore $\dim F(K, e_1) = n - 1$ or $\dim F(K, e_2) = n - 1$, because otherwise

$$\bigcup\{F(K, u) : u \in N(K, o) \cap S^{n-1}\},$$

which coincides with X_0 by Theorem 1.1 (i), has $(n - 1)$ -dimensional Hausdorff measure equal to zero and ∂K is C^1 by Theorem 1.2 (ii). \square

Acknowledgement We are grateful to the referees. Their observations substantially improved the paper.

REFERENCES

- [1] A. D. Alexandrov, Zur Theorie der gemischten Volumina von konvexen Körpern, III: Die Erweiterung zweier Lehrsätze Minkowskis über die konvexen Polyeder auf beliebige konvexe Flächen (in Russian). Mat. Sbornik N. S. 3 (1938), 27-46.
- [2] F. Barthe, O. Guédon, S. Mendelson, and A. Naor, A probabilistic approach to the geometry of the l_p^n -ball. Ann. of Probability 33, 480-513 (2005).
- [3] G. Bianchi, K. J. Böröczky, A. Colesanti, and D. Yang, The L_p -Minkowski problem for $-n < p < 1$, preprint
- [4] J. Böröczky, E. Lutwak, D. Yang, and G. Zhang, The logarithmic Minkowski problem. J. Amer. Math. Soc. 26, 831-852 (2013).
- [5] J. Böröczky, E. Lutwak, D. Yang, and G. Zhang, The log-Brunn-Minkowski inequality. Adv. Math. 231, 1974-1997 (2012).
- [6] K. J. Böröczky and H. T. Trinh, The planar L_p -Minkowski problem for $0 < p < 1$. Adv. Appl. Math., 87 (2017), 58-81.
- [7] L. Caffarelli, A localization property of viscosity solutions to Monge-Ampère equation and their strict convexity, Ann. Math. 131 (1990), 129-134.

- [8] L. Caffarelli, Interior $W^{2,p}$ -estimates for solutions of the Monge-Ampère equation. *Ann. Math. (2)* **131**, 135-150 (1990).
- [9] L. Caffarelli, A note on the degeneracy of convex solutions to Monge Ampère equation. *Comm. Partial Differential Equations* **18** (1993), 1213-1217.
- [10] S. Campi and P. Gronchi, The L^p -Busemann-Petty centroid inequality. *Adv. Math.* **167**, 128-141 (2002).
- [11] S. Chen, Q.-R. Li, and G. Zhu, The Logarithmic Minkowski Problem for non-symmetric measures. Submitted.
- [12] S. Chen, Q.-R. Li, and G. Zhu, The L_p Minkowski Problem for non-symmetric measures. Submitted.
- [13] W. Chen, L_p Minkowski problem with not necessarily positive data. *Adv. Math.* **201**, 77-89 (2006).
- [14] S.-Y. Cheng and S.-T. Yau, On the regularity of the solution of the n -dimensional Minkowski problem. *Comm. Pure Appl. Math.* **29**, 495-561 (1976).
- [15] K.-S. Chou, Deforming a hypersurface by its Gauss-Kronecker curvature. *Comm. Pure Appl. Math.* **38**, 867-882 (1985).
- [16] K.-S. Chou and X.-J. Wang, The L_p -Minkowski problem and the Minkowski problem in centroaffine geometry. *Adv. Math.* **205**, 33-83 (2006).
- [17] A. Cianchi, E. Lutwak, D. Yang, and G. Zhang, Affine Moser-Trudinger and Morrey-Sobolev inequalities. *Calc. Var. Partial Differential Equations* **36**, 419-436 (2009).
- [18] W. Fenchel and B. Jessen, Mengenfunktionen und konvexe Körper. *Danske Vid. Selskab. Mat.-fys. Medd.* **16**, 3 (1938), 31 pp.
- [19] M. Gage and R. Hamilton, The heat equation shrinking convex plane curves. *J. Differential Geom.* **23**, 69-96 (1986).
- [20] R. J. Gardner, *Geometric Tomography*, 2nd edition, *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 2006.
- [21] P. M. Gruber, *Convex and discrete geometry*, *Grundlehren der Mathematischen Wissenschaften*, 336. Springer, Berlin, 2007.
- [22] P. Guan and C.-S. Lin, On equation $\det(u_{ij} + \delta_{ij}u) = u^p f$ on S^n . Preprint.
- [23] C. Haberl and L. Parapatits, Centro-affine tensor valuations. arXiv:1509.03831. Submitted
- [24] C. Haberl and F. Schuster, General L_p affine isoperimetric inequalities. *J. Differential Geom.*, **83**, 1-26 (2009).
- [25] C. Haberl and F. Schuster, Asymmetric affine L_p Sobolev inequalities. *J. Funct. Anal.*, **257**, 641-658 (2009).
- [26] C. Haberl, F. Schuster, and J. Xiao, An asymmetric affine Pólya-Szegő principle. *Math. Ann.*, **352**, 517-542 (2012).
- [27] B. He, G. Leng, and K. Li, Projection problems for symmetric polytopes. *Adv. Math.*, **207**, 73-90 (2006).
- [28] M. Henk and E. Linke, Cone-volume measures of polytopes. *Adv. Math.*, **253**, 50-62 (2014).
- [29] Y. Huang and Q. Lu, On the regularity of the L_p -Minkowski problem. *Adv. Appl. Math.* **50**, 268-280 (2013).
- [30] D. Hug, E. Lutwak, D. Yang, and G. Zhang, On the L_p Minkowski problem for polytopes. *Discrete Comput. Geom.*, **33**, 699-715 (2005).
- [31] M. N. Ivaki, A flow approach to the L_{-2} Minkowski problem. *Adv. App. Math.*, **50** (2013), 445-464.
- [32] M.-Y. Jiang, Remarks on the 2-dimensional L_p -Minkowski problem. *Adv. Nonlinear Stud.*, **10**, 297-313 (2010).
- [33] J. Lu and X.-J. Wang, Rotationally symmetric solution to the L_p -Minkowski problem. *J. Differential Equations*, **254**, 983-1005 (2013).
- [34] M. Ludwig, General affine surface areas. *Adv. Math.*, **224**, 2346-2360 (2010).
- [35] E. Lutwak, The Brunn-Minkowski-Firey theory I: mixed volumes and the Minkowski problem. *J. Differential Geom.*, **38** (1993), 131-150.
- [36] E. Lutwak and V. Oliker, On the regularity of solutions to a generalization of the Minkowski problem. *J. Differential Geom.* **41**, 227-246 (1995).
- [37] E. Lutwak, D. Yang, and G. Zhang, L_p affine isoperimetric inequalities. *J. Differential Geom.* **56**, 111-132 (2000).
- [38] E. Lutwak, D. Yang, and G. Zhang, Sharp affine L_p Sobolev inequalities. *J. Differential Geom.* **62**, 17-38 (2002).
- [39] E. Lutwak, D. Yang, and G. Zhang, On the L_p -Minkowski problem. *Trans. Amer. Math. Soc.* **356**, 4359-4370 (2004).
- [40] H. Minkowski, Allgemeine Lehrsätze über die konvexen Polyeder. *Gött. Nachr.* **1897**, 198-219 (1897).

- [41] A. Naor, The surface measure and cone measure on the sphere of l_p^n . Trans. Amer. Math. Soc. **359**, 1045-1079 (2007).
- [42] A. Naor and D. Romik, Projecting the surface measure of the sphere of l_p^n . Ann. Inst. H. Poincaré Probab. Statist. **39**, 241-261 (2003).
- [43] L. Nirenberg, The Weyl and Minkowski problems in differential geometry in the large. Comm. Pure and Appl. Math. **6**, 337-394 (1953).
- [44] G. Paouris, Concentration of mass on convex bodies. Geom. Funct. Anal. **16**, 1021-1049 (2006).
- [45] G. Paouris and E. Werner, Relative entropy of cone measures and L_p centroid bodies. Proc. London Math. Soc. **104**, 253-286 (2012).
- [46] A. V. Pogorelov, The Minkowski multidimensional problem. V.H. Winston & Sons, Washington, D.C, 1978.
- [47] R. Schneider, Convex bodies: the Brunn-Minkowski theory (second edition), Encyclopedia of Mathematics and its Applications. Cambridge University Press, Cambridge, 2014.
- [48] A. Stancu, The discrete planar L_0 -Minkowski problem. Adv. Math. **167**, 160-174 (2002).
- [49] A. Stancu, On the number of solutions to the discrete two-dimensional L_0 -Minkowski problem. Adv. Math. **180**, 290-323 (2003).
- [50] A. Stancu, Centro-affine invariants for smooth convex bodies. Int. Math. Res. Not. **2012**, 2289-2320 (2012).
- [51] N. S. Trudinger and X.-J. Wang, The Monge-Ampère equation and its geometric applications, in: Handbook of geometric analysis. No. 1, 467-524, Adv. Lect. Math. (ALM), 7, Int. Press, Somerville, MA, 2008, available at <http://maths-people.anu.edu.au/~wang/publications/MA.pdf>.
- [52] G. Zhu, The logarithmic Minkowski problem for polytopes. Adv. Math. **262**, 909-931 (2014).
- [53] G. Zhu, The centro-affine Minkowski problem for polytopes. J. Differential Geom. (in press).
- [54] G. Zhu, The L_p Minkowski problem for polytopes for $0 < p < 1$. J. Func Analysis, 269 (2015), 1070-1094.
- [55] G. Zhu, The L_p Minkowski problem for polytopes for negative p . Indiana Univ. Math. J. (accepted). arXiv:1602.07774

DIPARTIMENTO DI MATEMATICA E INFORMATICA "U. DINI", UNIVERSITÀ DI FIRENZE, VIALE MORGAGNI 67/A, FIRENZE, ITALY I-50134

E-mail address: gabriele.bianchi@unifi.it

ALFRÉD RÉNYI INSTITUTE OF MATHEMATICS, HUNGARIAN ACADEMY OF SCIENCES, RELTANODA U. 13-15, H-1053 BUDAPEST, HUNGARY, AND DEPARTMENT OF MATHEMATICS, CENTRAL EUROPEAN UNIVERSITY, NADOR U 9, H-1051, BUDAPEST, HUNGARY

E-mail address: boroczky.karoly.j@renyi.mta.hu

DIPARTIMENTO DI MATEMATICA E INFORMATICA "U. DINI", UNIVERSITÀ DI FIRENZE, VIALE MORGAGNI 67/A, FIRENZE, ITALY I-50134

E-mail address: andrea.colesanti@unifi.it