

Journal of Physics: Complexity

OPEN ACCESS



CrossMark

LETTER

Probing criticality in quantum spin chains with neural networks

RECEIVED

5 May 2020

REVISED

24 June 2020

ACCEPTED FOR PUBLICATION

28 July 2020

PUBLISHED

7 August 2020

A Berezutskii^{1,2,3}, M Beketov³, D Yudin³, Z Zimborás^{4,5,6} and J D Biamonte³¹ IQB Information Technologies (IQBit), Waterloo, ON, N2J 1S7, Canada² Moscow Institute of Physics and Technology, Dolgoprudny, Moscow Region 141700, Russia³ Deep Quantum Laboratory, Skolkovo Institute of Science and Technology, Moscow 121205, Russia⁴ Wigner Research Centre for Physics, Theoretical Physics Department, Budapest, Hungary⁵ MTA-BME Lendület Quantum Information Theory Research Group, Budapest, Hungary⁶ Mathematical Institute, Budapest University of Technology and Economics, Budapest, HungaryE-mail: berezutskii@phystech.edu**Keywords:** Ising model, machine learning, phase transitions, neural networks

Original content from this work may be used under the terms of the [Creative Commons Attribution 4.0 licence](#). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



Abstract

The numerical emulation of quantum systems often requires an exponential number of degrees of freedom which translates to a computational bottleneck. Methods of machine learning have been used in adjacent fields for effective feature extraction and dimensionality reduction of high-dimensional datasets. Recent studies have revealed that neural networks are further suitable for the determination of macroscopic phases of matter and associated phase transitions as well as efficient quantum state representation. In this work, we address quantum phase transitions in quantum spin chains, namely the transverse field Ising chain and the anisotropic XY chain, and show that even neural networks with no hidden layers can be effectively trained to distinguish between magnetically ordered and disordered phases. Our neural network acts to predict the corresponding crossovers finite-size systems undergo. Our results extend to a wide class of interacting quantum many-body systems and illustrate the wide applicability of neural networks to many-body quantum physics.

1. Introduction

The concept of deep learning [1] has attracted dramatic interest over the last decade. First applied in the domain of image and natural speech recognition, algorithms for machine learning have recently shown their utility in statistical mechanics of interacting classical and quantum systems [2–17].

Solving a quantum many-body problem often implies a coarse-graining procedure to remove redundant degrees of freedom from the short-range, or the high-energy, sector of the theory. In this case, a proper elucidation of low energy properties of the system or the type of its long-range ordering encodes the macroscopic behavior. In its turn, the methodology of machine learning in multidimensional and typically non-structured datasets is inevitably linked to the effective approaches to dimensionality reduction, thereby yielding a powerful technique for the detailed analysis of classical and quantum models in many-body physics [18, 19]. Practical application of neural networks in the context of both supervised and unsupervised machine learning has now become commonplace for testing thermal, quantum, and topological phase transitions [2–11, 20] as well as for formulating effective variational wave function ansätze states [12–17, 21]. The application of machine learning to quantum-information problems has also received significant interest recently, promising to directly probe the entanglement entropy [22–24] as well as other properties. The utility of machine learning methods for quantum information purposes is driven by its great success in condensed matter physics [5, 25–38] and computational many-body methods [39–45]. In this study, we employ a specific machine learning technique to create a low-dimensional representation of microscopic states, relevant for macroscopic phase identification and probing phase transitions. More specifically, we explore phase transitions in the transverse field Ising- and the anisotropic XY chains and demonstrate that even the simplest possible neural network architecture—a binary classifier as a perceptron with no hidden neurons present—is capable of keeping track of its macroscopic

phases depending on the, e.g., external magnetic field or anisotropy parameter, without any prior knowledge. It is worth mentioning that an approach to spin models in higher dimensions has recently appeared and based on exact calculations of entanglement [46].

2. Model systems

2.1. Transverse field Ising model

One-dimensional spin models represent strongly correlated quantum systems that can be rigorously approached at equilibrium [47]. Certain non-equilibrium properties can also be extracted [48]. In the following, we focus on the one-dimensional ferromagnetic transverse field Ising model (TFIM). The TFIM naturally appears upon solving a classical two-dimensional Ising model with ferromagnetic-type nearest-neighbor exchange coupling and its exact solution dates back to the original works [49–51]. Generally, the TFIM of L spins on a chain with open boundary conditions is specified by the following Hamiltonian:

$$H = -J \sum_{i=1}^{L-1} \sigma_i^z \sigma_{i+1}^z - \tau \sum_{i=1}^L \sigma_i^x, \quad (1)$$

which represents a $2^L \times 2^L$ matrix with σ_i^α ($\alpha = x, y, z$) being a Pauli matrix acting on site i , and J and τ stand for the strength of exchange coupling and external magnetic field respectively. Interestingly, despite its relative simplicity, this model was used to describe intricate physics, e.g., the order-disorder transitions in ferroelectric crystals of KH_2PO_4 . At zero temperature, quantum fluctuations may lead to a restructuring of the ground state which is manifested by a certain non-analyticity in the ground state energy of the quantum Hamiltonian. For the case of the Hamiltonian (1), when there is no magnetic field present ($\tau = 0$) the ground state configuration is purely determined by the exchange interaction, the first term in equation (1), which favors collinear magnetic ordering. For $J > 0$, the ferromagnetic state is energetically preferable, meaning that all magnetic moments point in the same direction $\langle \sigma_i^z \rangle = +1$ (or -1), signaling the double degeneracy of the ground state. Increasing the transverse field beyond the critical value $\tau = \tau_c$ makes the system susceptible to spin flip and all the spins aligned in x direction in the limit $\tau \rightarrow \infty$, i.e., disordered in σ_z basis.

The one-dimensional TFIM can be worked out analytically by virtue of the Jordan–Wigner transformation that maps an interacting spin model onto that of free spin-polarized fermions [51, 52]. The exact solution unambiguously demonstrates a continuous quantum phase transition (QPT) upon passing through the critical field $\tau_c = 1$ (in the units of J), separating magnetically ordered ferromagnet ($\tau < \tau_c$) and disordered paramagnetic states ($\tau > \tau_c$). Although there is no exact analytical solution in higher dimensional systems, a QPT can be clearly detected [52]. It is worth noting that the phase diagram of a one-dimensional TFIM is very similar to that of a two-dimensional classical Ising model at finite temperature with a temperature-driven phase transition. Interestingly, this dualism has a strict mathematical form corresponding to the so-called Suzuki–Trotter decomposition and which maps a d -dimensional quantum model to a $d + 1$ dimensional classical one [53].

2.2. Anisotropic XY model

The XY model is yet another well-known quantum spin lattice model of magnetism. One can arrive to the isotropic version of this model by switching off the ZZ couplings in the Heisenberg Hamiltonian. In its turn, the anisotropic XY model is a generalization of it in the sense that the interaction strength in the XY plane is not isotropic anymore. In this study, we limit ourselves to the case when there is no field transverse to the interaction plane. The Hamiltonian of the model is thus given by

$$H = -J \sum_{i=1}^{L-1} \left(\frac{1+\gamma}{2} \sigma_i^x \sigma_{i+1}^x + \frac{1-\gamma}{2} \sigma_i^y \sigma_{i+1}^y \right), \quad (2)$$

where γ is the anisotropy parameter that is usually restricted to $-1 \leq \gamma \leq 1$ and J is the coupling strength which we set to 1 hereafter. If one sets $\gamma = 0$ the fully isotropic case, which possesses an additional symmetry $[H, \sigma_i^z] = 0$, is restored. On the other hand, it is also well-known that in the opposite case, i.e. $\gamma = 1$, the ground state possesses a long-range Neel order which yields

$$\sigma_i^x |\sigma\rangle = (-1)^i |\sigma\rangle \quad (3)$$

and

$$\sigma_i^y |\sigma\rangle = (-1)^i |\sigma\rangle \quad (4)$$

for $\gamma = -1$ accordingly, as is described in detail in reference [54]. It is clear that as γ decreases from 1 to -1 , the x - and y -components begin to compete. Its phase diagram is thus given by an x - and y -ferromagnetic states for $\gamma = 1$ and -1 accordingly. The model is fully isotropic at $\gamma = 0$ and undergoes a second-order phase transition at this point while the gap continuously vanishes [54, 55].

3. Methodology

3.1. General overview

The complexity of a generic quantum many-body problem grows exponentially with the size of a system (using the best known methods), making the available numerical routines computationally demanding. While machine learning has been specifically designed to coarse-grain certain information while maintaining relevant and unique features corresponding to the dataset (reminiscent to the formalism of renormalization group in statistical and high-energy physics [56]) it appears to be perfectly suited for identification of classical and quantum phases [27, 57, 58]. Indeed, sampled spin- $\frac{1}{2}$ configurations can be mapped to either binary numbers or black and white pixels which can be further classified in the form of macroscopic configurations, representing the class of problems which machine learning has been routinely used for. However, typically for quantum many-body systems we do not have predefined labels, so the use of unsupervised learning is favored. Within this paradigm we search for clusterization or associative rules that govern the behavior of a system. Unsupervised learning can also take measurement data and essentially reconstruct the wave function from individual images or snapshots. These reconstruction techniques based on machine learning are now being studied and compared to traditional techniques based on quantum state and quantum process tomography [8, 28, 59–61].

The advantage of using machine learning algorithms for exploration of both classical and QPTs is associated with finding certain features related to symmetry breaking in microscopic configurations. Particularly, phase transitions in magnetically ordered systems result in spin directions being randomized by the temperature—while the corresponding temperature can be detected as a point where the magnetization drops. When considering QPTs one typically investigates a finite region of sudden change that shrinks in the thermodynamic limit to a single point of non-analyticity [62]. Alternatively, in the vicinity of a phase transition point one can examine the behavior of the order parameter, which is known to collapse, or the correlation length that diverges [52, 63]. Passing through the phase transition point results in the ground state of a system being restructured, which is manifested by a certain non-analyticity in the ground state energy of a quantum Hamiltonian. It is therefore not surprising that there exists a final overlap between two different ground states of the system, which is regarded as a meaningful source of information on the quantum phases of a system and can be rigorously worked out within the fidelity approach [64, 65].

3.2. Sampling spin configurations

In this section, we briefly describe the sampling routine we used for the interacting spin models, described by the Hamiltonians (1) and (2). Note that the Hamiltonians (1) and (2) are sparse in the standard basis matrices with most of the elements being zero, as schematically shown in figure 1 for a system of $L = 7$ spins.



For small systems the exact diagonalization of the Hamiltonians of equations (1) and (2) is possible. Let a 2^L -dimensional vector $|g\rangle$ be the ground state of this system. In the computational basis the vector

$$|g\rangle = \sum_{i_1, i_2, \dots, i_L = \uparrow, \downarrow} \alpha_{i_1 i_2 \dots i_L} |i_1\rangle |i_2\rangle \dots |i_L\rangle, \quad (5)$$

is purely determined by 2^L complex-valued decomposition components $\alpha_{i_1 i_2 \dots i_L}$ in the basis $|i_k\rangle = \{|\uparrow\rangle, |\downarrow\rangle\}$, with $k = 1, \dots, L$, which are known to give the probability distribution $p_{i_1 i_2 \dots i_L} = |\alpha_{i_1 i_2 \dots i_L}|^2$ of a particular spin configuration $|i_1\rangle |i_2\rangle \dots |i_L\rangle$, which we refer to as a *bitstring* and later represent explicitly as strings of 0's and 1's. Thus, sampling the physical system specified by the Hamiltonian (1) might be approached by sampling each bitstring with the corresponding probabilities $p_{i_1 i_2 \dots i_L}$.

3.3. Neural network architecture

We use a neural network architecture that consists of an input layer and one output neuron, corresponding to a binary classifier. The sampled bitstrings serve as input data. Noteworthy, any hidden layers are absent. The output is prescribed to take value 0 when an input spin configuration is drawn from the ground state prescribed by $\tau_1 = 0.01$ ($\gamma_1 = -1$), whereas if the configuration is taken from τ_i (γ_i), the neuron is prescribed to take the value 1. We also discuss results of numerical simulations with other starting points, τ and γ —see section 4. The neural network architecture used is shown in figure 2.




The linear combination of the spins' z-projections $\{\sigma_i\}$ is fed into the neural network via the input layer, followed by a nonlinear activation of the output neuron

$$p(\{\sigma_i\}) = \text{sigm} \left(\sum_{i=1}^L W_i \sigma_i + b_i \right), \quad (6)$$

with $\text{sigm}(x) := (1 + e^x)^{-1}$ being the sigmoid function and the binary cross-entropy

$$H(p) = - \sum_{i=1}^{N_{\text{train}}} \{y_i \cdot \log p(y_i) + (1 - y_i) \cdot \log [1 - p(y_i)]\}, \quad (7)$$

serving as the loss-function. Such a simple form of the neural-network architecture results in high computational speed. The neural network outcome is the probability that the input state should be classified as belonging to the respective ground state specified by the control parameter value. Here, for a set of training data points $\{\sigma_i\}$ with $1 \leq i \leq N_{\text{train}}$ the neural network predicts the probability $p(y_i)$ for labels $y_i \in \{0, 1\}$. We make use of two labels, 'phase 1' and 'phase 2', namely magnetically ordered and disordered phases for



TFIM as well as X- and Y-ordered phases for the anisotropic XY model depending on parameters τ and γ respectively. While the parameters of the neural network, the weights and the biases, are updated using the *RMSProp* algorithm [66].

3.4. The algorithm

In our numerical simulations, for chains of $L = 20$ spins we explore the model described by equation (1) throughout the region $0.01J \leq \tau \leq 2J$ with $D = 40$ steps, $\tau = \{\tau_i\}_{i=1}^D$ and $N = 10^4$ spin configurations to be sampled for each value of τ_i . Afterward, a feed-forward neural network \mathbb{N}_i is trained to classify the bit-strings sampled for $\tau_1 = 0.01$ from those for τ_i with $i > 1$. Finally, we end up with $D - 1$ pairs of (P_i, τ_i) with $P_i \in (0, 1)$ being the mean output of the neural network evaluated on the samples drawn from the probability distribution given by the ground state of $H(\tau_i)$. In what follows, we show that the value of P with respect to τ dramatically changes signaling a phase transition. We apply a similar procedure to the anisotropic XY model with the anisotropy parameter $-1 \leq \gamma \leq 1$ starting with $\gamma_1 = -0.99$. The result is then averaged over 40 runs to rid possible effects caused by random initialization of the neural networks' parameters (displayed as shadows in the plots). The results described in the paper are obtained under the following conditions: we divide our initial dataset of 10^4 configurations per control parameter value into two subsets, namely training and testing set, which is in line with commonly accepted ratio of 80% for training, 20% for testing. As for the performance of the algorithm, generally, for the binary classification problem, the performance measure would be the accuracy of classification, which reached at 95% for the training set and 84% for the test set within all numerical experiments conducted.

4. Results

Below, we present and discuss the results of our numerical simulations, demonstrating how the neural network architecture and the corresponding algorithm described in section 3 are capable of probing the phase crossover point for the described models. In figure 3, we show how our setup performs for TFIM on an open chain of $L = 20$ spins together with the transverse magnetization defined as

$$m_x = \frac{1}{L} \sum_{i=1}^L \langle \sigma_i^x \rangle, \quad (8)$$

where averaging is done over the ground state. As expected, the neural network learns the order parameter due to the linearity of the latter as a function of spin projections. Note however, that while the resulting curve is remarkably close to the transverse magnetization curve, there was no information about the x -projections of the spin measurements in our setup, but only the measurements in the z -basis.

Unlike in previous studies, for example [67], the simplicity of a neural network used for the simulations makes direct visualization of the weights straightforward owing to their vectorial nature. Figure 4 clearly displays the crossover in the neighborhood of criticality, making these results intuitively clear and interpretable in contrast to usual deep learning routines [68, 69]. Each vertical row in figure 4 corresponds to a set of coefficients z -components of spins are multiplied by before transferring the whole sum to the activation function of the output neuron. Thus, the model actually mimics z -projections of spin configurations given the transverse magnetic field value τ . The latter explains why the rows in the heatmap are uniform in the ferromagnetic limit and take random values in the disordered phase. Note that the boundary coefficients are different because of the open boundary conditions.

In figure 5, we show the result for an anisotropic XY chain of $L = 20$ spins. In this plot, one can clearly see the phase crossover induced by the change of γ which is a sign of a well-studied anisotropy-induced phase




Figure 4. Heatmap of the weights W_i of the neural networks for a TFIM chain of $L = 20$ spins with open boundary conditions depending of the magnetic field strength.




Figure 5. The output of trained neural networks as a function of the anisotropy parameter γ for $L = 20$ spins on an anisotropic XY chain with open boundary conditions.




Figure 6. The output of trained neural networks as a function of the transverse magnetic field τ , for $L = 20$ spins on a TFIM chain with open boundary conditions, on condition that $\tau_0 = 1.0$ and $\tau_D = 2.0$.






Figure 7. The output of trained neural networks as a function of the anisotropy parameter γ , for $L = 20$ spins on an anisotropic XY chain with open boundary conditions, provided that $\gamma_0 = 0.5$ and $\gamma_D = 1.0$.

transition in an infinite system [70], similarly to the phase transition induced by the critical value of the magnetic field. Again, while our algorithm is given information about the z -components of spins, it is capable of exposing a phase crossover induced by the anisotropy in the x - y plain. In this case, there is no direct correspondence to any observable.



To further validate the proposed algorithm we provide outcomes of the neural network(s) for different starting reference points—that is, other than $\tau_0 = 0.01$ or $\gamma_0 = -1$ as discussed earlier. We hence perform the scan as described in section 3 while staying potentially in the same phase. The results of our numerical findings for $\tau_0 = 1$ and $\gamma_0 = 0.6$ are shown in figures 6 and 7.

To address finite-size scaling on the performance of our algorithm we provide the results of numerical simulations for phase classification depending on the number of spins L in figures 8 and 9. These findings suggest that classification robustness increases with the number of spins, while starting from $L = 20$ allowed us to correctly separate phases.

5. Conclusion

In this paper, we have considered the simplest neural network architecture with no hidden layers present and applied it to study the finite-size phase crossovers in the quantum TFIM and the quantum anisotropic XY model on a one-dimensional chain. We were able to distinguish the regions of different phases using neural networks *without* prior knowledge of the phase diagram by observing the corresponding phase boundary crossover in a finite-size system. Relative simplicity of the machine learning setup allowed us to visualize the weights of the corresponding neural network and unambiguously relate this plot to configuration of different spin orderings.

Data availability

The data that support the findings of this study are available upon reasonable request from the authors.

Acknowledgments

The authors are thankful to Anastasiia Pervishko and Sebastian Wetzel for fruitful discussions. ZZ acknowledges support from the János Bolyai Research Scholarship, the UKNP Bolyai+ Grant, and the NKFIH Grants No. K124152, K124176 KH129601, K120569, and from the Hungarian Quantum Technology National Excellence Program, Project No. 2017-1.2.1-NKP-2017-00001. The work of DY was supported by the Russian Foundation for Basic Research Project No. 19-31-90159. JB acknowledges support from the research project, Leading Research Center on Quantum Computing (Agreement No. 014/20).

References

- [1] LeCun Y, Bengio Y and Hinton G 2015 Deep learning *Nature* **521** 436–44
- [2] Zhang W, Liu J and Wei T-C 2019 Machine learning of phase transitions in the percolation and xy models *Phys. Rev. E* **99** 032142
- [3] Schindler F, Regnault N and Neupert T 2017 Probing many-body localization with neural networks *Phys. Rev. B* **95** 245134
- [4] Deng D-L, Li X and Das Sarma S 2017 Machine learning topological states *Phys. Rev. B* **96** 195145
- [5] Zhang P, Shen H and Zhai H 2018 Machine learning topological invariants with neural networks *Phys. Rev. Lett.* **120** 066401
- [6] Venderley J, Khemani V and Kim E-A 2018 Machine learning out-of-equilibrium phases of matter *Phys. Rev. Lett.* **120** 257204
- [7] Liu Y-H and Van Nieuwenburg E P L 2018 Discriminative cooperative networks for detecting phase transitions *Phys. Rev. Lett.* **120** 176401
- [8] Wang L 2016 Discovering phase transitions with unsupervised learning *Phys. Rev. B* **94** 195105
- [9] Hu W, Singh R R P and Scalettar R T 2017 Discovering phases, phase transitions, and crossovers through unsupervised machine learning: a critical examination *Phys. Rev. E* **95** 062122
- [10] Ch'ng K, Carrasquilla J, Melko R G and Khatami E 2017 Machine learning phases of strongly correlated fermions *Phys. Rev. X* **7** 031038
- [11] Ch'ng K, Vazquez N and Khatami E 2018 Unsupervised machine learning account of magnetic transitions in the Hubbard model *Phys. Rev. E* **97** 013306
- [12] Carleo G and Troyer M 2017 Solving the quantum many-body problem with artificial neural networks *Science* **355** 602–6
- [13] Glasser I, Pancotti N, August M, Rodriguez I D and Cirac J I 2018 Neural-network quantum states, string-bond states, and chiral topological states *Phys. Rev. X* **8** 011006
- [14] Cai Z and Liu J 2018 Approximating quantum many-body wave functions using artificial neural networks *Phys. Rev. B* **97** 035116
- [15] Carrasquilla J, Torlai G, Melko R G and Aolita L 2019 Reconstructing quantum states with generative models *Nat. Mach. Intell.* **1** 155–61
- [16] Hibat-Allah M, Ganahl M, Hayward L E, Melko R G and Carrasquilla J 2020 Recurrent neural network wavefunctions (arXiv:2002.02973)
- [17] Beach M J S, De Vlucht I, Golubeva A, Huembeli P, Kulchitsky B, Luo X, Melko R G, Merali E and Torlai G 2019 Qucumber: wavefunction reconstruction with neural networks *SciPost Phys.* **7** 009
- [18] Carleo G, Cirac I, Cranmer K, Daudet L, Schuld M, Tishby N, Vogt-Maranto L and Zdeborová L 2019 Machine learning and the physical sciences *Rev. Mod. Phys.* **91** 045002
- [19] Carrasquilla J 2020 Machine learning for quantum matter (arXiv:2003.11040)
- [20] Kharkov Y A, Sotskov V E, Karazeev A A, Kiktenko E O and Fedorov A K 2020 Revealing quantum chaos with machine learning *Phys. Rev. B* **101** 064406
- [21] Szabó A and Castelnovo C 2020 Neural network wave functions and the sign problem (arXiv:2002.04613)
- [22] Torlai G, Mazzola G, Carrasquilla J, Troyer M, Melko R and Carleo G 2018 Neural-network quantum state tomography *Nat. Phys.* **14** 447–50
- [23] Koch-Janusz M and Ringel Z 2018 Mutual information, neural networks and the renormalization group *Nat. Phys.* **14** 578–82
- [24] Rocchetto A, Grant E, Strelchuk S, Carleo G and Severini S 2018 Learning hard quantum distributions with variational autoencoders *Npj Quantum Inf.* **4** 28
- [25] Arsenault L-F, Lopez-Bezanilla A, von Lilienfeld O A and Millis A J 2014 Machine learning for many-body physics: the case of the Anderson impurity model *Phys. Rev. B* **90** 155136
- [26] Torlai G and Melko R G 2016 Learning thermodynamics with Boltzmann machines *Phys. Rev. B* **94** 165134
- [27] Carrasquilla J and Melko R G 2017 Machine learning phases of matter *Nat. Phys.* **13** 431
- [28] van Nieuwenburg E P L, Liu Y-H and Huber S D 2017 Learning phase transitions by confusion *Nat. Phys.* **13** 435
- [29] Wetzel S J 2017 Unsupervised learning of phase transitions: from principal component analysis to variational autoencoders *Phys. Rev. E* **96** 022140
- [30] Saito H 2017 Solving the Bose–Hubbard model with machine learning *J. Phys. Soc. Japan* **86** 093001
- [31] Mills K and Tamblyn I 2018 Deep neural networks for direct, featureless learning through observation: the case of two-dimensional spin models *Phys. Rev. E* **97** 032119
- [32] Choo K, Carleo G, Regnault N and Neupert T 2018 Symmetries and many-body excitations with neural-network quantum states *Phys. Rev. Lett.* **121** 167204
- [33] Bukov M, Day A G R, Sels D, Weinberg P, Polkovnikov A and Mehta P 2018 Reinforcement learning in different phases of quantum control *Phys. Rev. X* **8** 031086
- [34] Liu Y-H and van Nieuwenburg E P L 2018 Discriminative cooperative networks for detecting phase transitions *Phys. Rev. Lett.* **120** 176401
- [35] Shirinyan A A, Kozin V K, Hellsvik J, Pereiro M, Eriksson O and Yudin D 2019 Self-organizing maps as a method for detecting phase transitions and phase identification *Phys. Rev. B* **99** 041108
- [36] Burzawa L, Liu S and Carlson E W 2019 Classifying surface probe images in strongly correlated electronic systems via machine learning *Phys. Rev. Mater.* **3** 033805
- [37] Westerhout T, Astrakhantsev N, Tikhonov K S, Katsnelson M I and Bagrov A A 2020 Generalization properties of neural network approximations to frustrated magnet ground states *Nat. Commun.* **11** 1–8
- [38] Uvarov A, Biamonte J and Yudin D 2020 Variational quantum eigensolver for frustrated quantum systems (arXiv:2005.00544)
- [39] Huang L and Wang L 2017 Accelerated Monte Carlo simulations with restricted Boltzmann machines *Phys. Rev. B* **95** 035105
- [40] Nagai Y, Shen H, Qi Y, Liu J and Fu L 2017 Self-learning Monte Carlo method: continuous-time algorithm *Phys. Rev. B* **96** 161102
- [41] Xia R and Kais S 2018 Quantum machine learning for electronic structure calculations *Nat. Commun.* **9** 4195
- [42] Suwa H, Smith J S, Lubbers N, Batista C D, Chern G-W and Barros K 2019 Machine learning for molecular dynamics with strongly correlated electrons *Phys. Rev. B* **99** 161107
- [43] De Vlucht I J S, Iouchchenko D, Merali E, Roy P-N and Melko R G 2020 Reconstructing quantum molecular rotor ground states (arXiv:2003.14273)
- [44] Inack E M, Santoro G E, Dell'Anna L and Pilati S 2018 Projective quantum monte carlo simulations guided by unrestricted neural network states *Phys. Rev. B* **98** 235145
- [45] McNaughton B, Milošević M V, Perali A and Pilati S 2020 Boosting Monte Carlo simulations of spin glasses using autoregressive neural networks (arXiv:2002.04292)

- [46] Xu Q, Kais S, Naumov M and Sameh A Feb 2010 Exact calculation of entanglement in a 19-site two-dimensional spin system *Phys. Rev. A* **81** 022324
- [47] Pikin S A and Tsukernik V M 1966 The thermodynamics of linear spin chains in a transverse magnetic field *J. Exp. Theor. Phys.* **23** 914–6
- [48] Brandt U and Jacoby K 1977 The transverse correlation function of anisotropic $X - Y$ -chains: exact results at $T = \infty$ *Z. Phys. B* **26** 245–52
- [49] Katsura S 1962 Statistical mechanics of the anisotropic linear Heisenberg model *Phys. Rev.* **127** 1508
- [50] Schultz T D, Mattis D C and Lieb E H 1964 Two-dimensional Ising model as a soluble problem of many fermions *Rev. Mod. Phys.* **36** 856
- [51] Pfeuty P 1970 The one-dimensional Ising model with a transverse field *Ann. Phys.* **57** 79–90
- [52] Sachdev S 2011 *Quantum Phase Transitions* (Cambridge: Cambridge University Press)
- [53] Suzuki M 1976 Generalized Trotter's formula and systematic approximants of exponential operators and inner derivations with applications to many-body problems *Commun. Math. Phys.* **51** 183–90
- [54] Lieb E, Schultz T and Mattis D 1961 Two soluble models of an antiferromagnetic chain *Ann. Phys.* **16** 407–66
- [55] Luo Q, Zhao J and Wang X 2018 Fidelity susceptibility of the anisotropic xy model: the exact solution *Phys. Rev. E* **98** 022106
- [56] Mehta P and Schwab D J 2014 An exact mapping between the variational renormalization group and deep learning (arXiv:1410.3831)
- [57] Tanaka A and Tomiya A 2017 Detection of phase transition via convolutional neural networks *J. Phys. Soc. Japan* **86** 063001
- [58] Morningstar A and Melko R G 2018 Deep learning the Ising model near criticality *J. Mach. Learn. Res.* **18** 5975
- [59] Broecker P, Assaad F F and Trebst S 2017 Quantum phase recognition via unsupervised machine learning (arXiv:1707.00663)
- [60] Huembeli P, Dauphin A and Wittek P 2018 Identifying quantum phase transitions with adversarial neural networks *Phys. Rev. B* **97** 134109
- [61] Macarone Palmieri A, Kovlakov E, Bianchi F, Yudin D, Straupe S, Biamonte J D and Kulik S 2020 Experimental neural network enhanced quantum tomography *Npj Quantum Inf.* **6** 20
- [62] Vojta M 2003 Quantum phase transitions *Rep. Prog. Phys.* **66** 2069
- [63] Tsuda J, Yamanaka Y and Nishimori H 2013 Energy gap at first-order quantum phase transitions: an anomalous case *J. Phys. Soc. Japan* **82** 114004
- [64] Venuti L C and Zanardi P 2007 Quantum critical scaling of the geometric tensors *Phys. Rev. Lett.* **99** 095701
- [65] Damski B 2016 Fidelity approach to quantum phase transitions in quantum Ising model *Quantum Criticality in Condensed Matter: Phenomena, Materials and Ideas in Theory and Experiment* (Singapore: World Scientific) pp 159–82
- [66] Hinton G, Srivastava N and Swersky K 2012 Neural networks for machine learning *Lecture 6a Overview of Mini-Batch Gradient Descent (Coursera Lecture Slides)*
- [67] Arai S, Ohzeki M and Tanaka K 2018 Deep neural network detects quantum phase transition *J. Phys. Soc. Japan* **87** 033001
- [68] Montavon G, Samek W and Müller K-R 2018 Methods for interpreting and understanding deep neural networks *Digit. Signal Process.* **73** 1–15
- [69] Liu K, Greitemann J and Pollet L 2019 Learning multiple order parameters with interpretable machines *Phys. Rev. B* **99** 104410
- [70] Quan H T 2009 Finite-temperature scaling of magnetic susceptibility and the geometric phase in the XY spin chain *J. Phys. A* **42** 395002