

SHORT REPORT

Usp14 is required for spermatogenesis and ubiquitin stress responses in *Drosophila melanogaster*

Levente Kovács^{1,2,*}, Ágota Nagy^{1,2}, Margit Pál^{1,2} and Peter Deák^{1,2,‡}

ABSTRACT

Deubiquitylating (DUB) enzymes free covalently linked ubiquitin moieties from ubiquitin–ubiquitin and ubiquitin–protein conjugates, and thereby maintain the equilibrium between free and conjugated ubiquitin moieties and regulate ubiquitin-mediated cellular processes. Here, we performed genetic analyses of mutant phenotypes in *Drosophila melanogaster* and demonstrate that loss of Usp14 function results in male sterility, with defects in spermatid individualization and reduced testicular free monoubiquitin levels. These phenotypes were rescued by germline-specific overexpression of wild-type Usp14. Synergistic genetic interactions with *Ubi-p63E* and cycloheximide sensitivity suggest that ubiquitin shortage is a primary cause of male sterility. In addition, *Usp14* is predominantly expressed in testes in *Drosophila*, indicating a higher demand for this DUB in testes that is also reflected by testis-specific loss-of-function *Usp14* phenotypes. Collectively, these results suggest a major role of Usp14 in maintaining normal steady state free monoubiquitin levels during the later stages of *Drosophila* spermatogenesis.

This article has an associated First Person interview with the first author of the paper.

KEY WORDS: Deubiquitination, Deubiquitylation, Usp14, *Drosophila* spermatogenesis, Ubiquitin equilibrium

INTRODUCTION

Deubiquitylating enzymes (DUBs) are ubiquitin-specific proteases that hydrolyze isopeptide bonds between ubiquitin–ubiquitin and ubiquitin–protein conjugates, and thereby act as key regulators of ubiquitin-dependent processes (Amerik and Hochstrasser, 2004). Opposing ubiquitylation and deubiquitylation activities maintain a dynamic intracellular equilibrium between free and conjugated ubiquitin forms, in which monomeric ubiquitins are indispensable for normal cell physiology and development. About 100 DUB genes have been identified in humans, and these are classified into five or six conserved families of DUBs according to their catalytic domains (Nijman et al., 2005; Suresh et al., 2016). Basic DUB activities include processing of ubiquitin precursors and ubiquitin

recycling following deubiquitylation and/or polyubiquitin chain editing (Kim et al., 2003; Reyes-Turcu et al., 2009).

The evolutionarily conserved proteasome-associated DUBs Rpn11, Uch-L5 and Usp14 have prominent roles in protein recycling, and release ubiquitin from proteasome-bound substrate proteins prior to translocation and proteasomal degradation (Guterman and Glickman, 2004). Binding to the proteasome was previously associated with disassembling activities of Ubp6, the yeast ortholog of Usp14, and deletion of this protein led to reduced monoubiquitin levels (Leggett et al., 2002). Several pleiotropic effects of Usp14/Ubp6 loss have been identified, including synaptic transmission defects, ataxia and premature death in mice (Wilson et al., 2002). Dual roles of Ubp6 in regulating ubiquitin levels and proteasome function have been described in yeast (Hanna et al., 2007), and loss of this protein reduced the yeast prion [PSI⁺] expression and heightened sensitivity to a broad range of toxic drugs (Chernova et al., 2003). *Usp14* inactivation delays cell proliferation in mouse embryonic fibroblasts and the developing *Drosophila* eye (Lee et al., 2018). Moreover, gross perturbations of ubiquitin equilibria and reduced free monoubiquitin pools are common to all of these Usp14/Ubp6-related abnormalities.

In our previous studies of Usp5 in *Drosophila*, we found that deficiencies of free monoubiquitins triggered ubiquitin stress responses that were correlated with increased Usp14 deubiquitinase expression (Kovács et al., 2015). To investigate the biological function of this response, loss-of-function *Usp14* mutants were isolated and subjected to phenotypic characterization. We found that male *Usp14*-null flies were sterile and had sperm individualization defects and low free monoubiquitin levels in the testes. Reduced ubiquitin levels in these mutants was confirmed by high sensitivity to the translational inhibitor cycloheximide and an additive phenotype in *Usp14 Ubi-p63E* double-mutants. These phenotypes are consistent with the predominant Usp14 expression in testes.

RESULTS AND DISCUSSION

Usp14 mutant males are sterile

The *Drosophila* ortholog of Usp14 is encoded by the *CG5384* gene and has 49% overall similarity with yeast Ubp6 and 71% similarity with both mouse and human USP14 ubiquitin proteases. Catalytic ubiquitin-specific protease and proteasome binding ubiquitin-like (UBL) domains of these proteins are highly conserved (Tsou et al., 2012; Kovács et al., 2015). Our previous study showed increased expression of *Usp14* in *Usp5* mutants, suggesting roles of Usp14 in ubiquitin stress responses to ubiquitin shortages (Kovács et al., 2015), as described in yeast (Hanna et al., 2007).

To investigate the physiological consequences of Usp14 depletion, we analyzed phenotypes following PiggyBac-mediated insertional mutation of *Usp14* (to give the allele *Usp14^f*). The PiggyBac element was inserted at the start of the second exon (Fig. 1A) and resulted in substantial reductions in *Usp14* mRNA

¹Department of Genetics, University of Szeged, H-6726 Szeged, Hungary.

²Institute of Biochemistry, Biological Research Centre, H-6726 Szeged, Hungary.

*Present Address: Department of Genetics, University of Cambridge, CB2 3EH, UK

‡Author for correspondence (deakp@bio.u-szeged.hu)

© L.K., 0000-0002-3226-3740; M.P., 0000-0001-5587-1637; P.D., 0000-0002-8849-0352




Fig. 1. Male-sterile phenotype of *Usp14* mutants. (A) Diagram of the *Usp14* gene; arrowhead shows the position of the PiggyBac insertion (denoted as *Usp14^f* in text). The insertion of a single cytosine nucleotide in *Usp14^{Δ32}* mutants is highlighted in red. The protospacer adjacent motif (PAM) is in green. (B) *Usp14* expression levels in mutant and knockdown flies were determined using semi-quantitative RT-PCR analyses of WT, mutant and RNAi-transfected flies. *Usp14^{RNAi}* was controlled by the ubiquitous *Act5C-GAL4* driver and *rpL17A* was used as a loading control. (C) Quantification of the fertility of the males with indicated genotypes ($n=10$ individual males/genotype). Results are mean \pm s.d., $n=10$. * $P<0.01$ compared with WT (two-tailed Student's *t*-test). (D) Micrographs of internal male genitalia in WT and *Usp14^f* mutant flies (upper panels) and protamineB-GFP-labeled nuclei of elongated spermatids (lower panels). Arrowheads indicate seminal vesicles. Scale bars: 250 μ m.

expression levels, indicating that *Usp14^f* is a relatively strong hypomorphic allele (Fig. 1B). Although *Usp14^f* homozygotes were viable, all eclosed males were sterile (Fig. 1C, column 3) and despite normal mating behaviors, their wild-type (WT) female mating partners laid mostly unfertilized eggs. In phase-contrast and fluorescent microscopy analyses of *Usp14^f* homozygote testes, all stages through spermatid elongation were indistinguishable from those in WT flies. Although sperm bundles were formed in *Usp14^f* testes, seminal vesicles were almost empty and contained few individualized sperms (Fig. 1D). In addition, most sperm bundles were accumulated before the entrance of seminal vesicles, suggesting defects in sperm individualization. Male sterility was also observed when *Usp14^f* was situated *in trans* to a deficiency uncovering the *Usp14* locus (Fig. 1C, column 5). However, male fertility was completely restored in revertants of *Usp14^f* (Fig. 1C, column 2). Viability and fertility of females were not affected by *Usp14* mutations (data not shown).

For more accurate analyses of *Usp14* gene functions, we generated null alleles using CRISPR/Cas9-mediated mutagenesis

(Port et al., 2014). Among several indel mutants, we selected the point mutant allele *Usp14^{Δ32}* for detailed analysis. In this mutant, insertion of a cytosine after the second codon generates a frameshift and a premature stop codon at 16 base pairs downstream of the start site (Fig. 1A). Despite the resulting absence of functional Usp14, *Usp14^{Δ32}* null homozygotes were viable but males were sterile, as observed with the hypomorphic alleles (Fig. 1C, column 4). *Usp14^{Δ32}* mutant testes imitated individualization defects in *Usp14^f* males, indicating no clear differences between the phenotypes of null and the strong hypomorphic alleles. These results suggest that Usp14 is involved in the coordination of late spermatogenesis.

***Usp14* mutant male sterility is germline dependent**

The male sterile phenotype observed in *Usp14* mutants may exclusively reflect requirements of Usp14 in germline cell lineages and/or testis-forming somatic cells. To identify affected cell types following Usp14 depletion, we used the GAL4/UAS system in *Drosophila* (Duffy, 2002). Several GAL4 cell lines have specific spatio-temporal expression patterns in fly testis (White-Cooper,

2012). Herein, we used the *bam-Gal4* driver, which showed a germline-restricted expression pattern in previous studies (Chen and McKearin, 2003). The *Act5c-Gal4* driver was used exclusively to drive expression in somatic cells (White-Cooper, 2012).

Usp14 specific transgenic RNA interference was induced by mean of the germline-specific *bam-Gal4* driver and resulted in male sterility (Fig. 1C, column 8) and resulted in individualization defects that were similar to those seen in the *Usp14* hypomorphic and null mutants. However, induction of the RNAi with the somatic-specific *Act5C-Gal4* driver did not lead to male sterility or phenotypic abnormalities (Fig. 1C, column 7). Conversely, the male sterility of *Usp14^{A32}* homozygotes was rescued by transgenic *Usp14* expression under the control of the *bam-Gal4* driver (Fig. 1C, column 10), but not with the somatic *Act5C-Gal4* driver (Fig. 1C, column 9). These results suggest that germline-specific expression of *Usp14* is required for proper sperm development.

Usp14 mutation disrupts individualization complexes

The germline-dependent male sterility and deficient processing of sperm bundles to mature sperms in *Usp14* mutant testes suggest that *Usp14* facilitates sperm individualization. Cytoskeletal-membrane individualization complexes (ICs) mediate sperm individualization in elongated spermatid bundles. ICs contain a cluster of 64 actin-rich structures, known as actin cones. During sperm individualization, these move down the bundles and push excess cytoplasm and organelles into so called ‘cystic bulges’ that serve as waste bags. Concomitantly, spermatids are sheathed in plasma membranes and become individualized and move into seminal vesicles for storage until mating (Noguchi et al., 2006). Severe disturbances of this process lead to individualization defects and stagnation of sperm bundles in front of seminal vesicles.

To determine whether structures of the IC are affected in *Usp14* mutants, actin cones were visualized in WT and *Usp14^{A32}* testes after staining with fluorescently labeled phalloidin. In WT testes, triangular shaped actin cones move synchronously (Fig. 2A, upper panels) and actin cones behind the moving IC can only be observed occasionally. However, in *Usp14^{A32}* testes the majority of ICs lacked synchronous movements of actin cones, and a high

proportion of individual actin cones were spread along the sperm bundles (Fig. 2A, lower panels). Hence, *Usp14* is likely required for synchronous migration of actin cones, and disruption of ICs leads to male sterility. Strikingly similar individualization defects were observed following mutation of the testis-specific proteasome subunit *Prosa6T* (Zhong and Belote, 2007). Because the proteasome interactions of *Usp14* are well established in *Drosophila* (Lundgren et al., 2005; Lee et al., 2011), the resemblance of *Prosa6T* and *Usp14* mutant phenotypes may suggest that both subunits are required for normal testis-specific proteasome function.

Usp14 is predominantly expressed in testis

Previous studies have shown that many proteasome subunits and associated proteins have testis-specific orthologs in *Drosophila* (Belote and Zhong, 2009; Ma et al., 2002; Zhong and Belote, 2007), suggesting important roles of the proteasome in *Drosophila* testes. Because *Usp14* is a transient, but conserved, subunit of the regulatory *Drosophila* proteasome particles (Lundgren et al., 2005; Lee et al., 2011), and its mutants show testis-specific phenotypes, we investigated the possibility of differential expression of *Usp14* in testis. To investigate the expression levels of *Usp14*, testis and gonadectomized bodies of WT *Drosophila* were analyzed through semiquantitative RT-PCR experiments. *Usp14* transcripts were more abundant in testis than in other tissues (Fig. 2B). Because accessory glands and other organs were removed from testes before mRNA extraction, somatic cell representation was very low in this sample, including only epithelial cells of the testis surface. With observations of *Usp14* rescue by germline drivers (Fig. 1C), these data indicate that *Usp14* is expressed predominantly in the male germline.

Loss of *Usp14* and reduced free ubiquitin levels in testes

Loss of *Usp14* function in yeast and mouse models previously resulted in considerable reductions in monoubiquitin levels (Leggett et al., 2002; Anderson et al., 2005). *Usp14* recycles monoubiquitins from protein substrates that are targeted to the proteasome, and prevents their degradation together with the proteins targeted to the proteasome (Hanna et al., 2003;




Fig. 2. Individualization defects and reductions in monoubiquitin pools in the absence of *Usp14*. (A) Confocal micrographs of sperm bundles; dj-GFP (green) labels sperm tails and phalloidin (red) individualization complexes. Scale bars: 5 μ m. (B) Expression levels of *Usp14* in testis. Semiquantitative RT-PCR analyses of WT gonadectomized body, testis and ovary samples; *rpL17A* and tubulin were used as loading controls. (C) Monoubiquitin levels in *Usp14* mutant somatic (body) and testis tissues (also shown on Fig. 3B); western blots show a monoubiquitin band (α -Ub) at 8.5 kDa. Lanes were loaded with 4 μ g aliquots of total protein, as determined using Bradford assays.

Anderson et al., 2005). We determined monoubiquitin protein expression levels in WT and *Usp14* mutant testes and gonadectomized body samples by western blotting. We primarily observed significant differences between body and testis monoubiquitin levels (Fig. 2C, compare lane 1 and 3). Pixel density analysis of western blot images demonstrated that the monoubiquitin levels in testis are more than 3-fold higher than in the rest of the body. Greater than 80% of the monoubiquitin pool is present in testis, suggesting that ubiquitylation is prevalent during spermatogenesis. Pixel density analyses of immunoblots (Fig. 2C) revealed up to a 30% decrease in monoubiquitin concentrations in *Usp14* mutant testes, but no differences in other tissues from WT and *Usp14* mutant flies (see Fig. S1). These data show that monoubiquitin pools in testes are especially sensitive to the loss of *Usp14* function, likely contributing to male sterility and individualization defects.

The effects of monoubiquitin deficiencies on spermatogenesis were exacerbated when the *Usp14* mutation was accompanied by a hypomorphic allele of the testis-specific ubiquitin gene *Ubi-p63E*. In *Drosophila*, *Ubi-p63E* is the polyubiquitin gene that provides most of the newly synthesized ubiquitins in testis (Lu et al., 2013). *Ubi-p63E* null mutant spermatocytes show meiotic arrest with loss of monoubiquitins. The *Ubi-p63E^{EY}* hypomorph P element insertion allele, however, has moderate effects on monoubiquitin pools and permits progress of spermatogenesis to the elongated spermatid stage (Lu et al., 2013). After combining *Usp14^{Δ32}* null and *Ubi-p63E^{EY}* hypomorph mutations, we observed an increased severity of the testis phenotype compared with that in single mutants (Fig. 3A). In particular, spermatocytes from the *Usp14^{Δ32} Ubi-p63E^{EY}* double mutants had arrested meiosis and no postmeiotic and elongated spermatocytes were observed (Fig. 3A). This phenotype is consistent with the *Ubi-p63E* null phenotype (Lu et al., 2013),




Fig. 3. Lack of *Usp14* leads to monoubiquitin deficiency. (A) Phase-contrast micrographs of testes from males with indicated genotypes; arrowheads show seminal vesicles. Asterisks indicate the presence of elongated spermatids that are absent in *Usp14*; *Ubi-p63E* double mutants. Scale bars: 100 μm. (B) Monoubiquitin levels in testis of *Usp14* and *Ubi-p63E* mutants were measured using western blots with an anti-ubiquitin antibody (α-Ub). After determining protein concentrations using Bradford assays, 4 μg aliquots of total protein were loaded onto gels. Monoubiquitin contents in mutants were calculated using data from densitometric analyses and are presented as percentages of that in WT controls (mean±s.d., *n*=4) is shown in the graph. **P*<0.01 compared to WT (two-tailed Student's *t*-test). (C) Cycloheximide sensitivity of *Usp14* mutants. Data are expressed as means±s.d. deviations of three independent replicates. **P*<0.01 compared to WT data at a given concentration (two-tailed Student's *t*-test).

with considerable reductions in monoubiquitin levels (Fig. 3B). Perhaps *Usp14^{A32}* and *Ubi-63E^{EY}* synergistically maintain adequate free monoubiquitin pools in the testis. Most likely, loss of *Ubi-p63E* caused severe free ubiquitin shortage leading to early arrest of meiosis, whereas a less severe shortage of ubiquitin in *Usp14* null mutants manifests in a later stage of spermatogenesis.

Usp14 mutants are cycloheximide sensitive

The antibiotic cycloheximide is widely used as a eukaryotic protein synthesis inhibitor. It has toxic side effects that are related to rapid depletion of ubiquitin pools in treated cells. Accordingly, the expression levels of ubiquitin genes are exceptionally sensitive to this drug (Hanna et al., 2003). Previous studies have also identified mutations that reduce intracellular ubiquitin levels and lead to cycloheximide sensitivity in yeast and *Drosophila* (Leggett et al., 2002; Chernova et al., 2003; Kovács et al., 2015).

While previously studying the physiological roles of the *Drosophila* DUB Usp5, we observed a ubiquitin stress response as had been described in yeast (Hanna et al., 2007). Another study also indicates that Usp14 has an essential role (Kovács et al., 2015) that is unrelated to its testis-specific function.

Because loss of Usp5 leads to reduced monoubiquitin levels and triggers the expression of Usp14 to counteract ubiquitin degradation and boost ubiquitin recycling, it was possible that loss-of-function *Usp14* mutants are cycloheximide sensitive. To test this possibility, we compared the effects of cycloheximide in WT and hypomorph *Usp14^f* mutants.

Cycloheximide treatments of *Usp14^f* homozygote larvae (Fig. 3C) led to death in a dose-dependent manner, with an LD₅₀ of 32 μM. Viability of WT animals was not affected at this concentration. *Usp14^{f/+}* heterozygotes also showed high sensitivity to cycloheximide (LD₅₀=42 μM), indicating that one copy of *Usp14* does not compensate for ubiquitin shortages sufficiently after cycloheximide treatments. These data suggest that Usp14 functions in somatic tissues in situations when the ubiquitin equilibrium is disturbed by reduced ubiquitin expression due to cycloheximide treatment, or by abolished Ub recycling following elimination of Usp5 (Kovács et al., 2015). Demand for Usp14 activities in somatic tissues under stressed conditions indicate that Usp14 is not a testis-specific subunit, because it also has roles in somatic tissues. It is more likely that normal spermatogenic processes have high ubiquitin demands and, hence, resemble quasi-ubiquitin stress conditions in which proper functioning of Usp14 and other proteasome subunits is heavily required. Usp14 dependence of spermatogenesis may also be evolutionarily conserved because *Usp14* mutant *ataxia^f* mice are also sterile and had defects in spermatogenesis (Crimmins et al., 2009). However, loss of Usp14 function also severely affects the neuromuscular system in mice (Wilson et al., 2002; Anderson et al., 2005), suggesting that, unlike in flies, neuronal tissues of more-complex organisms are hypersensitive to ubiquitin homeostasis.

Taken together, our data demonstrate pleiotropic roles of the *Drosophila* Usp14 deubiquitylase in spermatogenesis and in the ubiquitin stress responses. Detailed phenotypic analyses of loss-of-function alleles show that *Usp14* is essential for normal spermatid differentiation. Accordingly, homozygous *Usp14* mutant males are sterile and have defective sperm individualization processes. In agreement, the phenotype associated with loss of Usp14 function is consistent with its expression pattern in testes.

Finally, we show that individualization defects were strikingly similar to the mutation of the testis-specific proteasome subunit Prosa6T (Zhong and Belote, 2007). Given that interaction of Usp14 with the proteasome is well established in *Drosophila* (Lundgren

et al., 2005; Lee et al., 2011), the resemblance of *Prosa6T* and *Usp14* mutant phenotypes may suggest that both subunits are required for normal testis-specific proteasome function.

MATERIALS AND METHODS

Drosophila stocks and methods

Drosophila lines were maintained on standard yeast and cornmeal medium at 25°C. The PiggyBac element insertion line *Usp14⁰⁰⁷⁷⁹* was obtained from Bloomington *Drosophila* Stock Center (BDSC stock number: 18368). The transgenic RNA interference line *Usp14^{KK102888}* was obtained from Vienna *Drosophila* Resource Center (VDRC stock number: v110227). All genetic markers are described in FlyBase at <http://flybase.org>. The *dj-GFP* line was obtained from Bloomington *Drosophila* Stock Center (BDSC stock number: 5417). The *protamineB-GFP* stock was kindly provided by Helen White-Cooper (School of Biosciences, Cardiff University, Cardiff, UK). The WT strain *w¹¹¹⁸* was used for comparisons in all experiments.

To reverse the *Usp14⁰⁰⁷⁷⁹* PiggyBac insertion, we crossed stock *en mass* with a PBac transposase source (BDSC 8285) and established two candidate revertant stocks. Restoration of male fertility was observed in both revertants, but only *Usp14^{f^v}* were maintained and used in experiments. CRISPR/Cas9-mediated mutagenesis was used to generate *Usp14^{A32}* null mutants using previously published tools (Port et al., 2014) with the targeting guide RNA sequence 5'-GCTACCTTTAAATGCGGGCA-3'. In detail, the targeting guide RNA sequence was cloned into pCFD3 vector containing the U6:3 ubiquitin promoter and the guide RNA backbone. This construct was injected into the *y[1] w[*] P{y[+7.7]}=nos-phiC31 \int.NLS>X; P{y[+7.7]}=CaryIP;su(Hw)attP1* (BDSC 35567) line and, through integrase-mediated site-specific transformation, we obtained a fly line constitutively expressing *Usp14*-specific guide RNAs. The guide RNA expressing line was crossed to a *nos-Cas9* expressing line (BDSC 54591) and the resulted 'cut starter' G0 progeny was further crossed to a *w; CyO/Sco* balancer line. From the subsequent generation, 100 males were individually crossed to a *w; CyO/Sco* balancer line and balanced candidate stocks were established. Candidates showing male sterility in homozygotes or over a deficiency were sequenced. In line Δ32 (Δ refers to the absence of gene function; 32 is the serial number of the candidate) we identified a single base pair insertion 3 bp downstream of the start codon resulting in a frameshift and premature stop codon 16 bp after the transcription start.

Male fertility was tested by placing individual 1-day-old males in vials with two 4-day-old WT virgin females. After 6 days, adults were discarded and the numbers of the eclosed progeny were scored.

Semiquantitative RT-PCR

Testes of 0–1-day-old *Drosophila* males were dissected in ice-cold PBS and tubes containing either 50 testes or 10 gonadectomized males were frozen at –80°C. Total RNA was isolated using Tri Reagent extraction kits (Sigma-Aldrich). RNA samples were treated with RQ1 RNase-Free DNase (Promega) and were reverse transcribed using Fermentas cDNA synthesis kits. Samples of cDNA were normalized to the rpL17A level using PCR with rpL17A forward primer, 5'-GTGATGAAGTGTGCCGACAA-3' and rpL17A reverse primer, 5'-CCTTCATTTCGCCCTTGTG-3'. *Usp14* mRNA expression levels were then determined in 20–25-cycle PCRs with the exon-specific *Usp14* RT forward primer 5'-ACGGTGGTGCCCTTCTCC-3', and *Usp14* RT reverse primer 5'-GGCGTGTGGTCTGTG-3'.

Fluorescence microscopy

Prior to native observations of GFP fluorescence, testes were dissected and mounted without extensive squashing in PBS and were then directly observed and imaged using an Olympus BX51 upright microscope with phase contrast and UV GFP filters.

Confocal microscopy was performed with testes of 0–1-day-old males expressing the sperm tail marker *dj-GFP*. In these experiments, WT or *Usp14* mutant flies were dissected at room temperature in PBS and were immediately fixed for 20 min in 4% formaldehyde. Testes were stained with Alexa Fluor 647-conjugated phalloidin (1:400; ThermoFisher Scientific, # A22287) and DAPI and were then examined under an Olympus FV 1000 confocal microscope.

SDS-PAGE and western blot analyses

Testes were dissected in ice-cold PBS buffer and were frozen in liquid nitrogen and stored at -80°C until use. Protein samples were then prepared from homogenates of 25 testes in Buffer I F, which contained 100 mM Tris-HCl pH 7.6, 150 mM NaCl, 1 mM EDTA, 100 mM N-ethylmaleimide (NEM, Sigma-Aldrich), 20 μM MG132 (Calbiochem) and 1 \times EDTA-free Complete protease inhibitor cocktail (Roche). Homogenates were centrifuged (21,000 g for 10 min at 4°C) and boiled in 1 \times Laemmli buffer. Samples containing 4 μg of total protein were loaded onto 14% SDS-acrylamide gels, were electrophoresed and subjected to western blotting. Protein concentrations were determined using Bradford assays (Bradford, 1976). Monoubiquitin forms of 8.5 kDa were detected using a mouse monoclonal anti-ubiquitin primary antibody (Sigma-Aldrich, U0508, 1:3000 dilution) and a peroxidase-conjugated AffiniPure goat anti-mouse-IgG (Jackson Immuno Research, 115-035-003, 1:30,000 dilution) secondary antibody. Western blots were developed on X-ray films, and were scanned and processed using ImageJ software.

Cycloheximide treatment

Cycloheximide (Sigma-Aldrich) was dissolved in ethanol to obtain stock solution (17.6 mM). First-instar larvae were collected and maintained in vials containing 3.5 ml of standard *Drosophila* medium supplemented with 60 μl aliquots of cycloheximide solution in ethanol. Final treatment concentrations were 0–100 μM in the final volume of 3.5 ml. Eclosing adults were scored and statistically analyzed using a Microsoft Office Excel software.

Acknowledgements

Drosophila stocks obtained from the Bloomington *Drosophila* Stock Center (NIH P40OD018537) and the Vienna *Drosophila* Resource Center (VDRRC, www.vdrcc.at) were used in this study. We also thank Flybase for providing database information (McQuilton et al., 2012).

Competing interests

The authors declare no competing or financial interests.

Author contributions

Conceptualization: P.D.; Methodology: L.K., A.N., P.D.; Validation: A.N., P.D.; Formal analysis: L.K., P.M.; Investigation: L.K., A.N., P.M.; Resources: P.D.; Data curation: A.N., P.D.; Writing - original draft: P.D.; Writing - review & editing: L.K., P.D.; Visualization: L.K., A.N.; Supervision: P.M., P.D.; Project administration: P.M., P.D.; Funding acquisition: P.D.

Funding

This work was funded by grants from the Hungarian Scientific Research Fund (OTKA-K116372), Ministry for National Economy of Hungary (GINOP-2.3.2-15-2016-00032).

Supplementary information

Supplementary information available online at <http://jcs.biologists.org/lookup/doi/10.1242/jcs.237511.supplemental>

Peer review history

The peer review history is available online at <https://jcs.biologists.org/lookup/doi/10.1242/jcs.237511.viewer-comments.pdf>

References

- Amerik, A. Y. and Hochstrasser, M. (2004). Mechanism and function of deubiquitinating enzymes. *Biochim. Biophys. Acta* **1695**, 189–207. doi:10.1016/j.bbamer.2004.10.003
- Anderson, C., Crimmins, S., Wilson, J. A., Korbel, G. A., Ploegh, H. L. and Wilson, S. M. (2005). Loss of Usp14 results in reduced levels of ubiquitin in ataxia mice. *J. Neurochem.* **95**, 724–731. doi:10.1111/j.1471-4159.2005.03409.x
- Belote, J. M. and Zhong, L. (2009). Duplicated proteasome subunit genes in *Drosophila* and their roles in spermatogenesis. *Heredity (Edinb)* **103**, 23–31. doi:10.1038/hdy.2009.23
- Bradford, M. M. (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.* **72**, 248–254. doi:10.1016/0003-2697(76)90527-3
- Chen, D. and McKearin, D. M. (2003). A discrete transcriptional silencer in the bam gene determines asymmetric division of the *Drosophila* germline stem cell. *Development* **130**, 1159–1170. doi:10.1242/dev.00325
- Chernova, T. A., Allen, K. D., Wesoloski, L. M., Shanks, J. R., Chernoff, Y. O. and Wilkinson, K. D. (2003). Pleiotropic effects of Ubp6 loss on drug sensitivities and yeast prion are due to depletion of the free ubiquitin pool. *J. Biol. Chem.* **278**, 52102–52115. doi:10.1074/jbc.M310283200
- Crimmins, S., Sutovsky, M., Chen, P.-C., Huffman, A., Wheeler, C., Swing, D. A., Roth, K., Wilson, J., Sutovsky, P. and Wilson, S. (2009). Transgenic rescue of ataxia mice reveals a male-specific sterility defect. *Dev. Biol.* **325**, 33–42. doi:10.1016/j.ydbio.2008.09.021
- Duffy, J. B. (2002). GAL4 system in *Drosophila*: a fly geneticist's swiss army knife. *Genesis* **34**, 1–15. doi:10.1002/gene.10150
- Guterman, A. and Glickman, M. H. (2004). Deubiquitinating enzymes are IN(trinsic to proteasome function). *Curr. Protein Pept. Sci.* **5**, 201–211. doi:10.2174/1389203043379756
- Hanna, J., Leggett, D. S. and Finley, D. (2003). Ubiquitin depletion as a key mediator of toxicity by translational inhibitors. *Mol. Cell. Biol.* **23**, 9251–9261. doi:10.1128/MCB.23.24.9251-9261.2003
- Hanna, J., Meides, A., Zhang, D. P. and Finley, D. (2007). A ubiquitin stress response induces altered proteasome composition. *Cell* **129**, 747–759. doi:10.1016/j.cell.2007.03.042
- Kim, J. H., Park, K. C., Chung, S. S., Bang, O. and Chung, C. H. (2003). Deubiquitinating enzymes as cellular regulators. *J. Biochem.* **134**, 9–18. doi:10.1093/jb/mvg107
- Kovács, L., Nagy, O., Pál, M., Udvardy, A., Popescu, O. and Deák, P. (2015). Role of the deubiquitylating enzyme DmUsp5 in coupling ubiquitin equilibrium to development and apoptosis in *Drosophila melanogaster*. *PLoS ONE* **10**, e0120875. doi:10.1371/journal.pone.0120875
- Lee, M. J., Lee, B. H., Hanna, J., King, R. W. and Finley, D. (2011). Trimming of ubiquitin chains by proteasome-associated deubiquitinating enzymes. *Mol. Cell. Proteomics* **10**, R110.003871. doi:10.1074/mcp.R110.003871
- Lee, J. H., Park, S., Yun, Y., Choi, W. H., Kang, M.-J. and Lee, M. J. (2018). Inactivation of Usp14 perturbs ubiquitin homeostasis and delays the cell cycle in mouse embryonic fibroblasts and in fruit fly *Drosophila*. *Cell. Physiol. Biochem.* **47**, 67–82. doi:10.1159/000489750
- Leggett, D. S., Hanna, J., Borodovsky, A., Crosas, B., Schmidt, M., Baker, R. T., Walz, T., Ploegh, H. and Finley, D. (2002). Multiple associated proteins regulate proteasome structure and function. *Mol. Cell* **10**, 495–507. doi:10.1016/S1097-2765(02)00638-X
- Lu, C., Kim, J. and Fuller, M. T. (2013). The polyubiquitin gene Ubi-p63E is essential for male meiotic cell cycle progression and germ cell differentiation in *Drosophila*. *Development* **140**, 3522–3531. doi:10.1242/dev.098947
- Lundgren, J., Masson, P., Mirzaei, Z. and Young, P. (2005). Identification and characterization of a *Drosophila* proteasome regulatory network. *Mol. Cell. Biol.* **25**, 4662–4675. doi:10.1128/MCB.25.11.4662-4675.2005
- Ma, J., Katz, E. and Belote, J. M. (2002). Expression of proteasome subunit isoforms during spermatogenesis in *Drosophila melanogaster*. *Insect Mol. Biol.* **11**, 627–639. doi:10.1046/j.1365-2583.2002.00374.x
- McQuilton, P., St Pierre, S. E. and Thurmond, J. and FlyBase Consortium (2012). FlyBase 101 – the basics of navigating FlyBase. *Nucleic Acids Res.* **40**, D706–D714. doi:10.1093/nar/gkr1030
- Nijman, S. M., Luna-Vargas, M. P. A., Velds, A., Brummelkamp, T. R., Dirac, A. M. G., Sixma, T. K. and Bernards, R. (2005). A genomic and functional inventory of deubiquitinating enzymes. *Cell* **123**, 773–786. doi:10.1016/j.cell.2005.11.007
- Noguchi, T., Lenartowska, M. and Miller, K. G. (2006). Myosin VI stabilizes an actin network during *Drosophila* spermatid individualization. *Mol. Biol. Cell* **17**, 2559–2571. doi:10.1091/mbc.e06-01-0031
- Port, F., Chen, H.-M., Lee, T. and Bullock, S. L. (2014). Optimized CRISPR/Cas tools for efficient germline and somatic genome engineering in *Drosophila*. *Proc. Natl. Acad. Sci. USA* **111**, E2967–E2976. doi:10.1073/pnas.1405500111
- Reyes-Turcu, F. E., Ventii, K. H. and Wilkinson, K. D. (2009). Regulation and cellular roles of ubiquitin-specific deubiquitinating enzymes. *Annu. Rev. Biochem.* **78**, 363–397. doi:10.1146/annurev.biochem.78.082307.091526
- Suresh, B., Lee, J., Kim, K.-S. and Ramakrishna, S. (2016). The importance of ubiquitination and deubiquitination in cellular reprogramming. *Stem Cells Int.* **2016**, 6705927. doi:10.1155/2016/6705927
- Tsou, W.-L., Sheedlo, M. J., Morrow, M. E., Blount, J. R., McGregor, K. M., Das, C. and Todi, S. V. (2012). Systematic analysis of the physiological importance of deubiquitinating enzymes. *PLoS ONE* **7**, e43112. doi:10.1371/journal.pone.0043112
- White-Cooper, H. (2012). Tissue, cell type and stage-specific ectopic gene expression and RNAi induction in the *Drosophila* testis. *Spermatogenesis* **2**, 11–22. doi:10.4161/spmg.19088
- Wilson, S. M., Bhattacharyya, B., Rachel, R. A., Coppola, V., Tessarollo, L., Householder, D. B., Fletcher, C. F., Miller, R. J., Copeland, N. G. and Jenkins, N. A. (2002). Synaptic defects in ataxia mice result from a mutation in Usp14, encoding a ubiquitin-specific protease. *Nat. Genet.* **32**, 420–425. doi:10.1038/ng1006
- Zhong, L. and Belote, J. M. (2007). The testis-specific proteasome subunit Prosalphat6T of *D. melanogaster* is required for individualization and nuclear maturation during spermatogenesis. *Development* **134**, 3517–3525. doi:10.1242/dev.004770