

Rank gradient, cost of groups and the rank versus Heegaard genus problem

Miklós Abért and Nikolay Nikolov

May 2, 2008

Abstract

We study the growth of the rank of subgroups of finite index in residually finite groups, by relating it to the notion of cost.

As a by-product, we show that the ‘Rank vs. Heegaard genus’ conjecture on hyperbolic 3-manifolds is incompatible with the ‘Fixed Price problem’ in topological dynamics.

1 Introduction

Let Γ be a finitely generated group. A *chain* in Γ is a sequence $\Gamma = \Gamma_0 \geq \Gamma_1 \geq \dots$ of subgroups of finite index in Γ . Let $T = T(\Gamma, (\Gamma_n))$ denote the *coset tree* of the chain, a rooted tree on the set of right cosets of the subgroups Γ_n with edges $(\Gamma_n g, \Gamma_{n+1} g)$ for all $g \in \Gamma$ and $n \in \mathbb{N}$. The *boundary* ∂T of T is the set of infinite rays starting from the root; it is naturally endowed with the product topology and product measure coming from the tree. The group Γ acts by automorphisms on T ; this action extends to measure preserving homeomorphisms of the boundary.

We say that a chain (Γ_n) is *Farber*, if the action of Γ on the boundary of its coset tree $T = T(\Gamma, (\Gamma_n))$ is essentially free, that is, if almost every element of ∂T has trivial stabilizer in Γ . This is the case for example when the chain consists of normal subgroups of Γ and their intersection is trivial. Note that then ∂T is simply the profinite completion of Γ with respect to (Γ_n) endowed with the normalized Haar measure.

For a group G let $d(G)$ denote the minimal number of generators (or rank) of G . Let the *rank gradient* of Γ with respect to (Γ_n) be defined as

$$\text{RG}(\Gamma, (\Gamma_n)) = \lim_{n \rightarrow \infty} \frac{d(\Gamma_n) - 1}{|\Gamma : \Gamma_n|}$$

This notion has been introduced by Lackenby [18].

Our first theorem relates the rank gradient of a Farber chain to the cost of the action of the group on the boundary of the coset tree. The analytic notion of cost was introduced by Levitt [19] and used by Gaboriau [14] to show that

free groups of different rank do not admit orbit equivalent measurable actions (see also the book of Kechris and Miller [16]).

Theorem 1 *Let (Γ_n) be a Farber chain in Γ . Then*

$$\text{RG}(\Gamma, (\Gamma_n)) = \text{cost}(E) - 1$$

where E denotes the orbit relation given by the action of Γ on the boundary of the coset tree $T(\Gamma, (\Gamma_n))$.

Theorem 1 allows us to clash two well-known problems, one in 3-manifold theory and the other in topological dynamics.

Rank vs Heegaard genus conjecture. *Let M be a compact, orientable, hyperbolic 3-manifold. Then the Heegaard genus of M equals the rank of the fundamental group of M .*

It is easy to see that the Heegaard genus is always greater or equal to the rank. The problem dates back to Waldhausen [33], who asked it for arbitrary 3-manifolds. This was proved false for Seifert manifolds by Boileau and Zieschang in [9] (see also [28]), but it remained open for hyperbolic 3-manifolds. For the above formulation, see [27, Conjecture 1.1]. Also, it is not known, whether the ratio of the two quantities can become arbitrarily large even for arbitrary 3-manifolds; the best known lower bound comes from the Boileau-Zieschang result.

A countable group Γ has *fixed price*, if every essentially free measure-preserving Borel action of Γ has the same cost. Gaboriau [14] established fixed price for a large class of groups, including free groups, higher rank non-uniform real irreducible lattices and groups containing an infinite amenable normal subgroup and asked whether the following holds.

Fixed Price problem. *Does every countable group have fixed price?*

Theorem 2 *Either the Rank vs Heegaard genus conjecture is false or the Fixed price problem has a negative solution.*

Moreover, if the Fixed Price problem has an affirmative answer, then the Rank vs Heegaard genus conjecture fails in the following strong senses. First, the ratio of the Heegaard genus and the rank of the fundamental group of a compact, orientable hyperbolic 3-manifold can get arbitrarily large. Second, the counterexamples are not exotic, rather this seems to be the general asymptotic behaviour of arithmetic hyperbolic 3-manifolds.

The contradiction between the two problems is established via the (unknown) answer to the following question: Does the rank gradient $\text{RG}(\Gamma, (\Gamma_n))$ depend on the choice of the Farber chain (Γ_n) in Γ ?

If it does, then Theorem 1 trivially provides a negative answer to the Fixed Price problem. In the other direction, there are specific (uniform or non-uniform) arithmetic lattices in $\text{SL}_2(\mathbb{C})$ (e.g. $\text{SL}_2(\mathbb{Z}[i])$), that viewed as abstract

groups, possess a chain of subgroups with vanishing rank gradient. On the other hand, being arithmetic groups, they have property (τ) with respect to congruence subgroups. Using work of Lackenby [18], this allows one to construct a covering tower of the 3-manifold corresponding to the lattice, where the Heegaard genus grows linearly. Now, if the rank gradient is independent of the chain, the rank must grow sublinearly on this tower, making the ratio of the Heegaard genus and the rank arbitrarily large.

We shall derive the independence of the rank gradient from the chain from a hypothesis that is much weaker than fixed price (see [16, Problem 25.5]).

Multiplicativity of cost-1 problem. *Let Γ be a measurable, essentially free action on (X, μ) and let H be a subgroup of Γ of finite index. Is it true that*

$$\text{cost}(H, X) - 1 = (\text{cost}(\Gamma, X) - 1) |\Gamma : H| ?$$

That is, does the cost of an action behave the same way as the rank for free groups?

The whole theory developed in this paper has a close connection to L^2 Betti numbers. The Lück approximation result [21] implies that for finitely presented groups and normal chains with trivial intersection, if we replace $d(\Gamma_n)$ with the first Betti number $\beta_1(\Gamma_n)$ in the definition of rank gradient, then the limit equals the first L^2 Betti number $\beta_1^2(\Gamma)$. This has been generalized by Farber [12] to chains satisfying his condition (see also [7] for examples showing the necessity of this condition). On the other hand, Gaboriau [13] has introduced L^2 Betti numbers of a measurable equivalence relation E and asked whether $\beta_1^2(E) = \text{cost}(E) - 1$ in general. An affirmative answer to Gaboriau's question would imply the surprising result that the asymptotic growth of $\beta_1(\Gamma_n)$ and $d(\Gamma_n)$ are equal for arbitrary groups.

Theorem 1 immediately allows us to compute the rank gradient for a class of groups where elementary methods seem to fail working.

Theorem 3 *Let Γ be a residually finite group with an infinite amenable normal subgroup and let Γ_n be a Farber chain in Γ . Then $\text{RG}(\Gamma, (\Gamma_n)) = 0$.*

This generalizes a result of Lackenby [18] who proved the result for finitely presented amenable groups.

We also answer a question of Kechris and Miller [16, Problem 35.7]. They asked whether for a countable infinite Euclidean domain D the group $\text{SL}(2, D)$ has cost 1 if and only if D has infinitely many units. The answer is negative: as we shall see, $\text{SL}(2, \mathbb{Z}[i])$ has cost 1 and $\mathbb{Z}[i]$ has finitely many units. A well-known conjecture by Thurston asserts that every hyperbolic 3-manifold virtually fibers over the circle. This would imply that any lattice in $\text{SL}(2, \mathbb{C})$ has cost 1. On the other hand, as we show in Section 5, lattices in $\text{SL}(2, \mathbb{R})$ have positive rank gradient with respect to any Farber chain.

The paper is organized as follows. In Section 2 we define measurable actions, graphings and the cost. We also introduce a new type of graphing and an

invariant, the product cost. We relate the product cost to the cost and prove a general approximation result on graphings. In Section 3 we apply this result to chains and express the product cost in terms of the rank gradient, thus proving Theorem 1. In Section 4 we use the theory developed to prove Corollary 3 and compute the rank gradient for some important classes of groups. We also discuss what happens if we relax the Farber condition on the chain. Finally, in Section 5 we consider the rank gradient of lattices in Lie groups, prove Theorem 2 and answer the question of Kechris and Miller.

Acknowledgements. The authors thank Ian Agol, Gabor Elek, Marc Lackenby and Alan Reid for helpful comments.

2 Graphings and cost

In this section we introduce Borel actions and the standard notion of cost as well as a new version of cost that will be useful in proving Theorem 1.

Let X be a standard Borel space. Let the countable group Γ act on X by Borel-automorphisms. Let μ be a Γ -invariant probability measure on X . We assume that the Γ -action has finitely many ergodic components.

Let us define the relation E on X by

$$xEy \text{ if there exists } \gamma \in \Gamma \text{ with } y = x \cdot \gamma$$

Then E is a Borel equivalence relation and every equivalence class is countable.

Since E is a subset of $X \times X$, it can also be considered as a graph on X . A *Borel subgraph* of E is a directed graph on X such that the edge set is a Borel subset of E .

Let $S \subseteq X \times X$ be an arbitrary graph on X . A path from x to y in S of length $k \geq 1$ is a sequence $x_0, x_1, \dots, x_k \in X$ such that:

- $x_0 = x, x_k = y$;
- $(x_i, x_{i+1}) \in S$ or $(x_{i+1}, x_i) \in S$ ($0 \leq i \leq k-1$).

Note that we consider undirected paths above. For $k \geq 1$ let us define the graph S^k by

$$(x, y) \in S^k \text{ if } x = y \text{ or there is a path from } x \text{ to } y \text{ in } S \text{ of length at most } k$$

We say that a subgraph S of E *spans* E , if for any $(x, y) \in E$ with $x \neq y$ there exists a path from x to y in S . Trivially, this holds if and only if

$$\bigcup_n S^n = E$$

The *edge-measure* of a Borel subgraph S of E is defined as

$$e(S) = \int_{x \in X} \deg_S(x) d\mu$$

where $\deg(x)$ is the number of edges in S with initial vertex x :

$$\deg_S x = |\{y \in X \mid (x, y) \in S\}|$$

Note that $e(S)$ may be infinite. The *cost* of E is defined as

$$\text{cost}(E) = \text{cost}(\Gamma, X) = \inf e(S)$$

where the infimum is taken over all Borel subgraphs S of E that span E . The cost of Γ is defined as

$$\text{cost}(\Gamma) = \inf \text{cost}(\Gamma, X)$$

where the infimum is taken over all ergodic, essentially free actions of Γ on a standard Borel space X .

We say that Γ has *fixed price* c if all ergodic, essentially free actions of Γ on a standard Borel space X have cost c . It is not known whether every countable group has fixed price [14].

If Γ is generated by g_1, \dots, g_d then it is easy to see that the set

$$\bigcup_{i=1}^d \bigcup_{x \in X} \{(x, x \cdot g_i)\}$$

is a spanning Borel subrelation of E of edge measure d . This implies the following.

Lemma 4 $\text{cost}(\Gamma, X) \leq d(\Gamma)$.

Now we will look at cost from another point of view, using graphings. The notion basically comes from [19] but we will need to use it somewhat differently. The advantage is that this notion will work for non-essentially-free (even finite) actions as well.

Let us consider the product space $X \times \Gamma$ where Γ is endowed with the discrete topology and the counting measure. Denote the product measure by e . A *graphing* is a Borel subset of $X \times \Gamma$. For a graphing M and $\gamma \in \Gamma$ let

$$M_\gamma = \{x \in X \mid (x, \gamma) \in M\}$$

be the γ -fiber of M . For $x \in X$ let

$$N(M, x) = \{\gamma \in \Gamma \mid x \in M_\gamma\} \text{ and } \deg_M x = |N(M, x)|$$

be the set of neighbours and the degree of x in M . Using this notation, we have

$$e(M) = \sum_{\gamma \in \Gamma} \mu(M_\gamma) = \int_{x \in X} \deg_M(x) d\mu$$

We will need a definition of powering of graphings. Let I be the graphing defined by

$$I_\gamma = \begin{cases} X & \gamma = 1 \\ \emptyset & \text{otherwise} \end{cases}$$

For a graphing M let the graphing M^\top be defined by

$$M_\gamma^\top = M_{\gamma^{-1}} \cdot \gamma^{-1} \quad (\gamma \in \Gamma)$$

and let \overline{M} be defined by

$$\overline{M} = M \cup M^\top \cup I$$

For graphings M and N let us define the graphing $M \cdot N$ by

$$(M \cdot N)_\gamma = \bigcup_{\delta \in \Gamma} (M_\delta \cap (N_{\delta^{-1}\gamma} \cdot \delta^{-1})) \quad (\gamma \in \Gamma)$$

that is,

$$(x, \gamma) \in M \cdot N \iff \exists \gamma_1, \gamma_2 \in \Gamma \text{ with } (x, \gamma_1) \in M, (x\gamma_1, \gamma_2) \in N \text{ and } \gamma_1\gamma_2 = \gamma$$

Let $M^1 = \overline{M}$ and for $k > 1$ let

$$M^k = M^{k-1} \cup (M^{k-1} \cdot \overline{M})$$

that is, $(x, \gamma) \in M^k$, if and only if there exists $l \leq k$ and $\gamma_1, \gamma_2, \dots, \gamma_l \in \Gamma$ such that $\gamma_1\gamma_2 \cdots \gamma_l = \gamma$ and

$$(x\gamma_1\gamma_2 \cdots \gamma_i, \gamma_{i+1}) \in M \text{ or } (x\gamma_1\gamma_2 \cdots \gamma_i\gamma_{i+1}, \gamma_{i+1}^{-1}) \in M \quad (i < l)$$

A graphing M is an *L-graphing* if

$$\bigcup_k M^k = X \times \Gamma$$

The *product cost* of (Γ, X) is defined as

$$\text{pcost}(\Gamma, X) = \inf e(M)$$

where the infimum is taken over all *L-graphings* M .

A graphing M is *finitely supported* if M_γ is empty for all but finitely many $\gamma \in \Gamma$. The *distance* of two graphings M, N is defined as

$$d(M, N) = e(M \Delta N) = \sum_{\gamma \in \Gamma} \mu(M_\gamma \Delta N_\gamma)$$

where $A \Delta B$ denotes the symmetric difference of A and B .

Let us fix a base O of the topology of X . We call a subset of X *cylindric* with respect to O if it is a finite union of elements of O . A graphing M is *cylindric* if for all $\gamma \in \Gamma$ the set M_γ is *cylindric*.

The following lemma says that under some assumptions on X and Γ , every *L-graphing* can be approximated with open finitely supported *cylindric L-graphings*. Note that we do not assume that X is a standard Borel space: it can also be a finite set.

Lemma 5 *Assume that X is compact, Γ is finitely generated and it acts by homeomorphisms on X . Let M be an L -graphing of finite measure and let O be a base of the topology on X . Then for all $\varepsilon > 0$ there exists an open finitely supported L -graphing N that is cylindric with respect to O and such that $d(M, N) < \varepsilon$.*

Proof. Fix a generating set g_1, g_2, \dots, g_d of Γ . Let B denote the graphing defined by

$$B_\gamma = \begin{cases} X & \gamma = g_i \text{ for some } i \leq d \\ \emptyset & \text{otherwise} \end{cases}$$

Clearly, B is an L -graphing of measure d .

List the elements of Γ as $\gamma_1, \gamma_2, \dots$ and the elements of the base O as O_1, O_2, \dots .

Let $K \supseteq M$ be an open graphing satisfying $e(K \setminus M) < \varepsilon/3$. For $n \geq 0$ let the finitely supported graphing $K(n)$ be defined by

$$K(n)_\gamma = \begin{cases} K_\gamma & \gamma = \gamma_i \text{ for some } i \leq n \\ \emptyset & \text{otherwise} \end{cases}$$

Then K has finite edge-measure, implying

$$\lim_{n \rightarrow \infty} d(K(n), K) = 0$$

For $d \geq 2$, we have

$$K^d = \bigcup_n K(n)^d$$

which yields

$$\bigcup_n K(n)^n = \bigcup_n K^n = X \times \Gamma \supseteq B.$$

It is easy to see that $K(n)^n$ is open and $K(n)^n \subseteq K(n+1)^{n+1}$ ($n \geq 2$). Since B is compact, there exists k such that, setting $L = K(k)$, we have $d(L, K) < \varepsilon/3$ and $L^k \supseteq B$. This implies that

$$\bigcup_n K^n \supseteq \bigcup_n B^n = X \times \Gamma$$

that is, L is an L -graphing.

For $n \geq 1$ let the graphing $L(n)$ be defined by

$$L(n)_\gamma = \bigcup_{\substack{1 \leq i \leq n \\ O_i \subseteq L_\gamma}} O_i$$

Then $L(n)$ is a finitely supported cylindric graphing ($n \geq 1$) and $\bigcup_n L(n) = L$. So, using the same argument as above we have

$$\bigcup_n L(n)^n = \bigcup_n L^n = X \times \Gamma \supseteq B$$

and using the compactness of B again, there exists k such that, setting $N = L(k)$, we have both $N^k \supseteq B$ and $d(N, L) < \varepsilon/3$. So N is an L -graphing of distance at most ε from M and the lemma is proved. \square

Now we will analyze the connection between Borel subgraphs and graphings. Every graphing M defines a Borel subgraph $\Phi(M)$ of E as follows:

$$\Phi(M) = \{(x, x \cdot \gamma) \mid (x, \gamma) \in M\}$$

As we will see, the map Φ is surjective. Note that it is bijective if and only if the action of Γ on X is free.

Lemma 6 *We have*

$$\text{pcost}(\Gamma, X) \geq \text{cost}(\Gamma, X).$$

If Γ acts essentially freely on X , then we have equality.

Proof. Let M be a graphing. It is easy to see that

$$\Phi(M^k) = \Phi(M)^k \quad (k \geq 2).$$

This implies that if M is an L -graphing then $\Phi(M)$ is a Borel subgraph of E that spans E .

Now for $x \in X$ the degree

$$\begin{aligned} \deg_S x &= |\{y \in X \mid \exists \gamma \in \Gamma \text{ with } x \in M_\gamma \text{ and } y = x \cdot \gamma\}| \leq \\ &\leq |\{\gamma \in \Gamma \mid x \in M_\gamma\}| = \deg_M(x) \end{aligned} \quad (1)$$

which implies

$$e(\Phi(M)) \leq e(M)$$

It follows that

$$\text{pcost}(\Gamma, X) \geq \text{cost}(\Gamma, X)$$

Assume that Γ acts essentially freely on X . Let us list the elements of Γ as $\gamma_1, \gamma_2, \dots$

Let S be a Borel subgraph of E spanning E . For each $(x, y) \in S$ let us define $f(x, y)$ to be the first element of Γ such that

$$y = x \cdot f(x, y)$$

Let us define the subset M by

$$M = \{(x, \gamma) \in X \times \Gamma \mid f(x, x \cdot \gamma) = \gamma\}$$

Then M is a graphing that satisfies $\Phi(M) = S$. Also, for almost all $x \in X$ there is equality in 1) which yields

$$e(M) = e(S)$$

However, the graphing M may not be an L -graphing. Let

$$N = \bigcup_n M^n$$

Then

$$\Phi(N) = \bigcup_n \Phi(M^n) = \bigcup_n \Phi(M)^n = \bigcup_n S^n = E$$

Let $(x, \gamma) \in X \times \Gamma \backslash N$. Then $(x, x \cdot \gamma) \in S$ so for $\delta = f(x, x \cdot \gamma) \neq \gamma$ we have $x \cdot \gamma = x \cdot \delta$, implying $\gamma\delta^{-1} \in \text{Stab}_\Gamma(x)$. Since the action of Γ is essentially free, we obtain $e(X \times \Gamma \backslash N) = 0$. But then

$$M' = M \cup (X \times \Gamma \backslash N)$$

is an L -graphing of measure $e(M') = e(S)$. This implies

$$\text{pcost}(\Gamma, X) \leq \text{cost}(\Gamma, X)$$

so equality holds as claimed. \square

3 Boundary action and rank gradient

In this section we first introduce coset trees and boundary representations. Then we describe the product cost of a boundary representation in terms of the rank gradient of the chain. This allows us to prove Theorem 1.

Let (Γ_n) be a chain in Γ . Then the *coset tree* $T = T(\Gamma, (\Gamma_n))$ of Γ with respect to (Γ_n) is defined as follows. The vertex set of T equals

$$T = \{\Gamma_n g \mid n \geq 0, g \in \Gamma\}$$

and the edge set is defined by inclusion, that is,

$$(\Gamma_n g, \Gamma_m h) \text{ is an edge in } T \text{ if } m = n + 1 \text{ and } \Gamma_n g \supseteq \Gamma_m h$$

Then T is a tree rooted at Γ and every vertex of level n has the same number of children, equal to the index $|\Gamma_n : \Gamma_{n+1}|$. The right actions of Γ on the coset spaces Γ/Γ_n respect the tree structure and so Γ acts on T by automorphisms. This action is called the *tree representation* of Γ with respect to (Γ_n) .

The boundary ∂T of T is defined as the set of infinite rays starting from the root. The boundary is naturally endowed with the product topology and product measure coming from the tree. More precisely, for $t = \Gamma_n g \in T$ let us define $\text{Sh}(t) \subseteq \partial T$, the *shadow* of t as

$$\text{Sh}(t) = \{x \in \partial T \mid t \in x\}$$

the set of rays going through t . Set the base of topology on ∂T to be the set of shadows and set the measure of a shadow to be

$$\mu(\text{Sh}(t)) = 1/|\Gamma : \Gamma_n|.$$

This turns ∂T into a totally disconnected compact space with a Borel probability measure μ . The group Γ acts on ∂T by measure-preserving homeomorphisms; we call this action the *boundary representation of Γ* with respect to (Γ_n) .

Lemma 7 *The action of Γ on ∂T is ergodic and minimal (that is, every orbit is dense).*

Proof. Let $A \subseteq \partial T$ be a measurable Γ -invariant subset such that $\mu(A) > 0$. Then using the Lebesgue density theorem, for all $\varepsilon > 0$ there exists $t \in T$ of level n with

$$\mu(\text{Sh}(t) \cap A) \geq (1 - \varepsilon)\mu(\text{Sh}(t))$$

Since Γ acts transitively on the n -th level of T , invariance implies the same inequality for all $u \in T$ of level n . Adding up, we get

$$\mu(A) = \sum_{u \in T \text{ of level } n} \mu(\text{Sh}(u) \cap A) \geq 1 - \varepsilon$$

which implies $\mu(A) = 1$.

Now let $x \in \partial T$ and let $t \in T$. Let $t' \in T$ be the vertex of the same level as t contained in x and let $g \in \Gamma$ with $t'g = t$. Then $xg \in \text{Sh}(t)$. We proved that the orbit of x is dense in ∂T . \square

There are various levels of faithfulness of a boundary representation. Let

$$\partial T_{free} = \{x \in \partial T \mid \text{Stab}_\Gamma(x) = 1\}.$$

We say that the action is *essentially free* (or that the chain is Farber), if $\mu(\partial T \setminus \partial T_{free}) = 0$. The action is *topologically free* if $\partial T \setminus \partial T_{free}$ is meagre, i.e., a countable union of nowhere dense closed sets. The action is *free*, if $\partial T_{free} = \partial T$. Note that the Farber condition has been introduced by Farber in [12] in another equivalent formulation (see also [7] for a relevant result).

It is easy to see that the following implications hold for the action of Γ on ∂T :

$$\begin{aligned} (\Gamma_n) \text{ is normal and } (\cap \Gamma_n = 1) &\implies \text{free} \implies \text{essentially free} \implies \\ &\implies \text{topologically free} \iff X_{free} \neq \emptyset \implies \text{faithful} \end{aligned}$$

For all but the third arrow it is easy to find examples showing that the reverse implications do not hold. We shall discuss these classes more in Section 4.

Note that a deep result of Stuck and Zimmer [30] tells us that every faithful ergodic measure preserving action of a higher rank semisimple real lattice on a probability space is essentially free. In particular, every faithful boundary representation of such a lattice is essentially free.

Let Γ be a group and let X be a set. A directed Γ -labeled graph is a triple (V, E, f) where (V, E) is a directed graph with vertex set V and edge list E and

f is a function from E to Γ . Note that we allow multiple edges and we make no restriction on the labeling f . For a directed Γ -labeled graph

$$G = (V, E = (e_1, \dots, e_n), f)$$

let $U(G) = (V, U(E))$ denote the undirected graph with vertex set V and edge list

$$U(E) = (\bar{e}_1, \dots, \bar{e}_n)$$

where \bar{e} denotes the unordered pair obtained from the ordered pair e .

Let G be a directed Γ -labeled graph and let $v \in V$. Then we can define a natural map

$$\Phi_v : \pi_1(U(G), v) \rightarrow \Gamma$$

from the fundamental group of $U(G)$ based at v to Γ as follows. For a loop $l = (e_1, \dots, e_k)$ in $U(G)$ starting at v let

$$\Phi_v(l) = \prod_{i=0}^{n-1} f^{\pm 1}(e_i)$$

where the sign depends on whether we travel along e_i preserving its original orientation in G or not. The following lemma is straightforward.

Lemma 8 *The map Φ_v is a group homomorphism.*

Now we express the rank gradient of a chain in terms of the product cost of the action on the boundary of the coset tree. Note that we do not make any assumptions on the boundary representation. In fact, we do not even assume that the chain is infinite!

Theorem 9 *Let (Γ_n) be any chain in Γ . Then*

$$\text{RG}(\Gamma, (\Gamma_n)) = \text{pcost}(E) - 1$$

where $E = E(\partial T(\Gamma, (\Gamma_n)))$ denotes the orbit relation on $\partial T(\Gamma, (\Gamma_n))$ defined by the action of Γ .

Proof. Let $s = \text{RG}(\Gamma, (\Gamma_n))$ and let $c = \text{pcost}(E)$.

First we show $s + 1 \geq c$. Let $\varepsilon > 0$. Then there exists n such that

$$\frac{d(\Gamma_n) - 1}{|\Gamma : \Gamma_n|} < s + \varepsilon$$

that is, Γ_n can be generated by at most

$$d = \lfloor (s + \varepsilon) |\Gamma : \Gamma_n| \rfloor + 1$$

elements where $\lfloor x \rfloor$ is the floor of x . Let h_1, h_2, \dots, h_d be such a generating set and let

$$\gamma_1, \gamma_2, \dots, \gamma_{|\Gamma : \Gamma_n|}$$

be a coset representative system for Γ_n in Γ . We can assume that $\gamma_1 = 1$. Let

$$Y = \{h_i \mid i \leq d\} \cup \{h_i^{-1} \mid i \leq d\} \cup \{1\}$$

Let us define the graphing M as follows:

$$M_\gamma = \begin{cases} \text{Sh}(\Gamma_n) & \text{if } \gamma = h_i \ (i \geq 1) \text{ or } \gamma = \gamma_i \ (i > 1) \\ \emptyset & \text{otherwise} \end{cases}$$

We claim that M is an L -graphing. First, we have

$$\overline{M}_\gamma \supseteq \begin{cases} \text{Sh}(\Gamma_n) & \text{if } \gamma \in Y \\ \text{Sh}(\Gamma_n) & \text{if } \gamma = \gamma_i \ (i \leq |\Gamma : \Gamma_n|) \\ \text{Sh}(\Gamma_n \gamma_i) & \text{if } \gamma = \gamma_i^{-1} \ (i \leq |\Gamma : \Gamma_n|) \end{cases}$$

Let $(x, \gamma) \in \partial T \times \Gamma$. Then there exists $a, b \leq |\Gamma : \Gamma_n|$ such that $x \in \text{Sh}(\Gamma_n \gamma_a)$ and $\gamma_a \gamma_b^{-1} \in \Gamma_n$. This implies that there are elements $y_1, y_2, \dots, y_k \in Y$ such that $\gamma_a \gamma_b^{-1} = y_1 y_2 \cdots y_k$. Using that $Y \subseteq \Gamma_n$ we get

$$(x, \gamma) = (x, \gamma_a^{-1} y_1 y_2 \cdots y_k \gamma_b) \in \overline{M}^{k+2}.$$

Now the edge measure of M equals

$$e(M) = \frac{1}{|\Gamma : \Gamma_n|} (d + |\Gamma : \Gamma_n| - 1) \leq 1 + s + \varepsilon$$

which implies $c \leq 1 + s$ as claimed.

Now we show that $s \leq c - 1$ holds. Let $\varepsilon > 0$. Then there exists an L -graphing M of E of edge-measure at most $c + \varepsilon/2$. Using Lemma 5 (setting the set of shadows as base of the topology) there exists a finitely supported cylindric L -graphing N of E with $d(M, N) < \varepsilon/2$. This implies that the edge-measure $e(N) < c + \varepsilon$.

Let n be a natural number such that for every $\gamma \in \Gamma$ the set N_γ is a union of shadows of some cosets of Γ_n . Let

$$V = \{\Gamma_n \gamma \mid \gamma \in \Gamma\} \text{ and } v = \Gamma_n$$

Let us define the undirected Γ -labeled graph $G = (V, E, f)$ as follows. For each $\gamma \in \Gamma$ and $w \in V$ where $\text{Sh}(w) \subseteq N_\gamma$ let us add the edge $(w, w\gamma)$ to the list E with label $f(w, w\gamma) = \gamma$.

Let us consider the map $\Phi_v : \pi_1(U(G), v) \rightarrow \Gamma$. We claim that the image of Φ_v is

$$\Phi_v(\pi_1(U(G), v)) = \Gamma_n.$$

First, let $l = (e_1, \dots, e_k)$ be a loop in $U(G)$ starting at v . Then $v\Phi_v(l) = v$ so $\Phi_v(l) \in \Gamma_n$.

Second, let $h \in \Gamma_n$. Let $x \in \text{Sh}(v)$ be an arbitrary element. Then since N is an L -graphing, we have $(x, h) \in N^k$ for some k . Thus there exist $\gamma_1, \gamma_2, \dots, \gamma_k \in \Gamma$ such that $\gamma_1 \gamma_2 \cdots \gamma_k = h$ and

$$(x \gamma_1 \gamma_2 \cdots \gamma_i, \gamma_{i+1}) \in N \text{ or } (x \gamma_1 \gamma_2 \cdots \gamma_i \gamma_{i+1}, \gamma_{i+1}^{-1}) \in N \ (i < k)$$

Let $x_i = x\gamma_1\gamma_2\cdots\gamma_i$ ($0 \leq i \leq k$). Then $vh = v$ so $x_0 = v = x_k$. Also, for all $0 \leq i < k$ there is an edge in G either from x_i to x_{i+1} labeled by γ_{i+1} or from x_{i+1} to x_i labeled by γ_{i+1}^{-1} . Thus these edges form a loop l in $U(G)$ with $\Phi_v(l) = h$. The claim follows.

The number of vertices of G equals $|\Gamma : \Gamma_n|$ while the number of edges of G equals $e(N) |\Gamma : \Gamma_n|$. Hence, the same holds for $U(G)$. Using Lemma 8 and the formula for the rank of the fundamental group of a graph we get

$$d(\Gamma_n) \leq d(\pi_1(U(G), v)) = e(N) |\Gamma : \Gamma_n| - |\Gamma : \Gamma_n| + 1$$

which yields

$$\frac{d(\Gamma_n) - 1}{|\Gamma : \Gamma_n|} \leq e(N) - 1 < c - 1 + \varepsilon.$$

This shows that $s \leq c - 1$.

The theorem is proved. \square

Now Theorem 1 follows immediately.

Proof of Theorem 1. Using Theorem 9 and Lemma 6 we get

$$\text{RG}(\Gamma, (\Gamma_n)) = \text{pcost}(E) - 1 = \text{cost}(E) - 1$$

as claimed. \square

4 Applications and examples

In this section we introduce the absolute rank gradient of a group and compute it for some important classes of groups. We will later use these results in Section 5. Then we discuss what happens if we relax the Farber condition.

Let us define the *absolute rank gradient* of Γ as

$$\text{RG}(\Gamma) = \inf_H \frac{d(H) - 1}{|\Gamma : H|}$$

where H runs through all subgroups of Γ of finite index.

Let $H, K \leq \Gamma$ be subgroups of finite index with $H \leq K$. Using the Nielsen-Schreier theorem on H and K , we get $d(H) - 1 \leq |K : H| (d(K) - 1)$ which yields

$$\frac{d(H) - 1}{|\Gamma : H|} \leq \frac{d(K) - 1}{|\Gamma : K|}$$

So for any chain (Γ_n) in Γ , the sequence $(d(\Gamma_n) - 1) / |\Gamma : \Gamma_n|$ is non-increasing and the definition of $\text{RG}(\Gamma, (\Gamma_n))$ makes sense. In fact, as the authors show in [1], if the sequence stabilizes then Γ is virtually free.

We shall make use of the following easy lemma.

Lemma 10 *Let Γ be a finitely generated, residually finite group and let $H \leq \Gamma$ be a subgroup of finite index. Then there exists a chain (Γ_n) in Γ such that $\Gamma_1 = H$ and the boundary representation of Γ with respect to (Γ_n) is free.*

Proof. Let K be the core of H , that is, $K = \bigcap_{g \in \Gamma} H^g$. Then K is a normal subgroup of Γ of finite index. For $n \geq 1$ let

$$\Delta_n = \bigcap_{L \leq \Gamma \text{ of index } n} L$$

Since Γ has only finitely many subgroups of a given index, each Δ_n is a normal subgroup of finite index in Γ . Also, since Γ is residually finite, $\bigcap_n \Delta_n = 1$. Now let $\Gamma_0 = \Gamma$, let $\Gamma_1 = H$ and for $n \geq 2$ let $\Gamma_n = K \cap \Delta_n$. Then $\bigcap_n \Gamma_n = 1$ and the chain (Γ_n) consists of normal subgroups of Γ (except for $n = 1$). Let $g \in \Gamma$ with $g \neq 1$. Then there exists $n > 1$ such that $g \notin \Gamma_n$. This implies that g acts fixed point freely on Γ/Γ_n and thus also on the boundary $\partial T(\Gamma, (\Gamma_n))$. In other words, the action of Γ on ∂T is free. \square

Theorem 1 now gives us the following on the absolute rank gradient.

Corollary 11 *Let Γ be a finitely generated, residually finite group. Then*

$$\text{RG}(\Gamma) = \text{cost}(\Gamma, \widehat{\Gamma}) - 1$$

where $\widehat{\Gamma}$ denotes the profinite completion of Γ .

Proof. Let Δ_n be as in the proof of Lemma 10. Then for all $H \leq \Gamma$ of index n , we have $\Delta_n \leq H$. Hence $\widehat{\Gamma}_{(\Delta_n)}$, the profinite completion of Γ with respect to (Δ_n) equals $\widehat{\Gamma}$ and by Theorem 1 we get

$$\text{RG}(\Gamma) = \text{RG}(\Gamma, \Delta_n) = \text{cost}(\Gamma, \widehat{\Gamma}_{(\Delta_n)}) - 1 = \text{cost}(\Gamma, \widehat{\Gamma}) - 1$$

as claimed. \square

Proof of Corollary 3. Let Γ be a finitely generated residually finite group with an infinite amenable normal subgroup. Let (Γ_n) be a Farber chain in Γ – such chain exists by Lemma 10. Let $E = E(\partial T(\Gamma, (\Gamma_n)))$. Then by [14] Γ has fixed price 1 and so, using Theorem 1 we have

$$\text{RG}(\Gamma) = \text{RG}(\Gamma, (\Gamma_n)) = \text{cost}(E) - 1 = 0.$$

\square

Note that in [1] we present an interesting alternative combinatorial proof of the corollary in the case when Γ itself is infinite amenable. However, we have no proof for the general case that does not use analysis.

Ascending HNN extensions. Let A be a finitely generated group and $f : A \rightarrow A$ a homomorphism. Then the ascending HNN extension with base A is the group Γ with presentation

$$\langle A, t \mid a^t = f(a) \forall a \in A \rangle$$

For example every free by cyclic group has such a presentation.

For each $n \in \mathbb{N}$ let $\Gamma_n = \langle A, t^n \rangle \leq \Gamma$. It is easy to see that Γ_n is a normal subgroup of index n in Γ . As $d(\Gamma_n) \leq d(A) + 1$ we get the following result, first observed by Lackenby in [18].

Proposition 12 *If the group Γ is a residually finite ascending HNN extension, then $\text{RG}(\Gamma) = 0$.*

For instance, free by cyclic groups are residually finite and so they have rank gradient 0. In fact, more generally we have the following.

Proposition 13 *Let Γ be finitely generated residually finite group which has a finitely generated normal subgroup N such that Γ/N has subgroups of arbitrarily large index (i.e. the profinite completion $\widehat{\Gamma/N}$ of Γ/N is infinite). Then $\text{RG}(\Gamma) = 0$.*

Proof. Using the assumptions one can find a sequence of subgroups $H_i < \Gamma$ such that both

$$a_i = [\Gamma : NH_i] \text{ and } b_i = [N : N \cap H_i]$$

tend to infinity as i tends to infinity. Now $[\Gamma : H_i] = a_i b_i$, $d(N \cap H_i) \leq b_i d(N)$ and $d(NH_i/N) \leq a_i d(\Gamma)$. Hence from $d(H_i) \leq d(N \cap H_i) + d((NH_i)/N)$ we obtain

$$\frac{d(H_i) - 1}{[\Gamma : H_i]} < \frac{d(\Gamma)}{b_i} + \frac{d(N)}{a_i}$$

which tends to 0 as i tends to infinity. This implies $\text{RG}(\Gamma) = 0$. \square

The result that the groups in Proposition 13 have a measurable action with cost 1, has been proved by Gaboriau [14], even without assuming residual finiteness of Γ and $|\widehat{\Gamma/N}| = \infty$, but that does not give anything on the rank gradient. We note that the condition that $\widehat{\Gamma/N}$ is infinite seems quite mild.

Now we discuss rank gradient of chains that are not Farber. As we shall see, the situation can be quite different.

Our first example is the so-called *lamplighter group*, the wreath product

$$\Gamma = \mathbb{Z}/2\mathbb{Z} \text{ wr } \mathbb{Z}.$$

This group is metabelian, in particular, it is amenable, so Corollary 3 implies that the rank gradient of any Farber chain is 0. However, consider the canonical surjections

$$\phi_n : \mathbb{Z}/2\mathbb{Z} \text{ wr } \mathbb{Z} \rightarrow \mathbb{Z}/2\mathbb{Z} \text{ wr } \mathbb{Z}/2^n\mathbb{Z} \text{ and } \pi_n : \mathbb{Z}/2\mathbb{Z} \text{ wr } \mathbb{Z}/2^n\mathbb{Z} \rightarrow \mathbb{Z}/2^n\mathbb{Z}$$

and let $\Gamma_n = \text{Ker}(\phi_n \pi_n)$. Then (Γ_n) is a normal chain, $\Gamma/\Gamma_n \cong \mathbb{Z}/2^n\mathbb{Z}$ and $\phi_n(\Gamma_n) \cong (\mathbb{Z}/2\mathbb{Z})^{2^n}$ which implies $d(\Gamma_n) \geq 2^n$. This yields

$$\text{RG}(\Gamma, (\Gamma_n)) \geq 1.$$

In the above example the chain is normal but with nontrivial intersection, and so the boundary representation is not faithful. We do not know whether there exists an amenable group Γ and a chain (Γ_n) such that the boundary representation is faithful, but the chain has positive rank gradient.

Our last example is a slight variation of one discussed by Bergeron and Gaboriau in [7]. The difference is that they estimate the first Betti number where we estimate the rank.

Proposition 14 *There exist a virtually free group Γ and an interval $[x, y) \subset \mathbb{R}$ such that for every $\alpha \in [x, y)$ there exists a subnormal chains of subgroups $\Gamma = H_0 > H_1 > H_2 > \dots$ with trivial intersection, such that $\text{RG}(\Gamma, (H_i)) = \alpha$.*

Moreover in this situation we can have the strict inequalities

$$\text{RG}(\Gamma, (H_i)) > \lim_{i \rightarrow \infty} \frac{b_{1,p}(H_i)}{|\Gamma : H_i|} > \lim_{i \rightarrow \infty} \frac{b_1(H_i)}{|\Gamma : H_i|},$$

where for a prime p the integer $b_{1,p}(H) = \dim_{\mathbb{F}_p} H/[H, H]H^p$ is the p -homology of H .

In particular, these chains are not Farber.

Proof of Proposition 14. Let $\Gamma = A * \mathbb{Z}$, where A is a finite group. Viewing Γ as a trivial HNN extension we see that Γ has a right transitive action on both the vertices and the edges of a regular $2|A|$ -valent tree T (so that the quotient I_0 is a single vertex with a looped edge). We can direct the edges of T so that Γ acts preserving this orientation. There are exactly $a = |A|$ in-edges and out-edges from every vertex of T . The action of Γ is regular on the edges of T . We fix an edge e_0 of T and a vertex v_0 at one end of e_0 so that the stabilizer of v_0 is A .

Suppose that H is a subgroup of Γ of index n . Then the quotient $I = T/H$ is a finite graph with a covering map $p : T \rightarrow I = T/H$ and H can be recovered from p as $\{g \in \Gamma \mid p(e_0) = p(e_0g)\}$.

The vertices of I are in 1-1 correspondence with the double cosets $A \backslash \Gamma / H$. Given a vertex $v \in I$ define $S_v = A^g \cap H$, where g is a fixed representative of $A \backslash \Gamma / H$ such that $p(v_0g) = v$. There are exactly n edges in I and we have $n = \sum_{v \in I} [A : S_v]$.

The Bass-Serre theory determines the structure of H as follows:

$$H = F * (*_{v \in I} S_v)$$

where the free group F is the fundamental group of I . The group F has rank $n - p + 1$, where p is the number of vertices in I .

Let X be the set of vertices of I with out-valency 1 and let $Y = I \setminus X$ be the set of vertices with out-valency $a = |A|$. Let $|X| = \mu p$. Then we have

$$n = p(\mu + (1 - \mu)|A|), \text{ hence } p = \frac{n}{\mu + (1 - \mu)|A|}.$$

We get that exactly μp of the vertex stabilizers S_v are equal to A and the rest are trivial. From the Grushko–Neumann theorem (see [15] and [24]) it now follows that the rank of H is

$$d(H) = n - p + \mu p d(A) + 1 = n \left(1 + \frac{\mu d(A) - 1}{\mu + (1 - \mu)|A|} \right) + 1$$

while $\beta_1(H) = n - np + 1$.

Now suppose that we have a sequence of finite oriented graphs with covering maps

$$I_0 \leftarrow I_1 \leftarrow I_2 \leftarrow \cdots \leftarrow T$$

with the following two properties:

1. Each vertex of each I_j has valency 2 or $2a$. The proportions $\mu_j = |X_j|/|I_j|$ of the vertices of I_j with out- and in-valency 1 form a decreasing sequence which tends to some prescribed limit $\mu = \mu_\infty \in [0, 1)$.

2. Given any integer k let B_k be the ball of radius k in the $2a$ -regular oriented tree T centered at the vertex v_0 . Then there is an integer m such that the covering map $T \rightarrow I_m$ is injective on B_k .

Then the groups H_j which correspond to the graphs I_j form a chain in Γ with trivial intersection and such that

$$\lim_{i \rightarrow \infty} \frac{d(H_i) - 1}{|\Gamma : H_i|} = 1 + \frac{\mu d(A) - 1}{\mu + (1 - \mu)|A|}$$

The first part of Proposition 14 now follows by setting $x = 1 - \frac{1}{|A|}$ (when $\mu_\infty = 0$) and $y = d(A)$ (when $\mu_\infty \rightarrow 1$).

The existence of the graphs I_j with properties 1 and 2 above is essentially proved by Bergeron and Gaboriau in [7, Section 4] in the case of a free product of two residually finite groups.

On the other hand, we have

$$\lim_{i \rightarrow \infty} \frac{\beta_1(H_i)}{|\Gamma : H_i|} = 1 - \frac{1}{\mu + (1 - \mu)|A|}$$

and similarly we have

$$b_{1,p}(H) = n \left(1 + \frac{\mu b_{1,p}(A) - 1}{\mu + (1 - \mu)|A|} \right) + 1.$$

Therefore

$$\lim_{i \rightarrow \infty} \frac{\beta_{1,p}(H_i)}{|\Gamma : H_i|} = 1 + \frac{\mu \beta_{1,p}(A) - 1}{\mu + (1 - \mu)|A|}.$$

So, if we choose the finite group A so that $d(A) > b_{1,p}(A) > 0$ and choose the chain (H_i) with limiting ratio $\mu_\infty > 0$ this gives

$$\text{RG}(\Gamma, (H_i)) > \lim_{i \rightarrow \infty} \frac{b_{1,p}(H_i)}{|\Gamma : H_i|} > \lim_{i \rightarrow \infty} \frac{b_1(H_i)}{|\Gamma : H_i|}$$

as promised. \square

5 Lattices and 3-manifolds

In this section we further discuss the rank gradient, with a special emphasis on lattices in Lie groups. We also define the Heegaard genus and derive the contradiction between the Rank vs. Heegaard genus conjecture and Fixed Price.

The following lemma follows immediately from [14, Theorem 3] saying that if $H \leq \Gamma$ of finite index, then

$$\text{cost}(H) - 1 = |\Gamma : H| (\text{cost}(\Gamma) - 1)$$

where $\text{cost}(\Gamma)$ is the infimum of costs of all measurable essentially free actions of Γ .

Lemma 15 *An affirmative answer to the Fixed Price problem implies an affirmative answer to the Multiplicativity of cost-1 problem.*

Proof. Let Γ be a finitely generated group acting on (X, μ) by measure preserving maps and let H be a subgroup of Γ of finite index. Then by our assumption we have

$$\text{cost}(H, X) - 1 = \text{cost}(H) - 1 = |\Gamma : H| (\text{cost}(\Gamma) - 1) = |\Gamma : H| (\text{cost}(\Gamma, X) - 1)$$

\square

Now we proceed to the independence of the rank gradient.

Theorem 16 *Assume that the Multiplicativity of cost-1 problem has an affirmative solution. Let (Γ_n) be a Farber chain in Γ . Then*

$$\text{RG}(\Gamma, (\Gamma_n)) = \text{RG}(\Gamma)$$

In particular, any two Farber chains have the same rank gradient in Γ .

Proof. Let Γ act on a standard Borel probability space X essentially freely. Then using the multiplicativity assumption, we have

$$\text{cost}(\Gamma, X) - 1 = \frac{\text{cost}(\Gamma_n, X) - 1}{|\Gamma : \Gamma_n|} \leq \frac{d(\Gamma_n) - 1}{|\Gamma : \Gamma_n|}$$

which implies

$$\text{cost}(\Gamma, X) - 1 \leq \text{RG}(\Gamma, (\Gamma_n)).$$

Now if (Λ_n) is another Farber chain in Γ , then using Theorem 1 and the above inequality for both chains and actions, we get

$$\begin{aligned} \text{RG}(\Gamma, (\Gamma_n)) &= \text{cost}(\Gamma, E(\Gamma, (\Gamma_n))) - 1 \leq \\ &\leq \text{RG}(\Gamma, (\Lambda_n)) = \text{cost}(\Gamma, E(\Gamma, (\Lambda_n))) - 1 \leq \text{RG}(\Gamma, (\Gamma_n)) \end{aligned}$$

so equality holds everywhere.

Let $\varepsilon > 0$ and let $H \leq \Gamma$ be a subgroup of finite index such that

$$\frac{d(H) - 1}{|\Gamma : H|} < \text{RG}(\Gamma) + \varepsilon$$

Using Lemma 10 there exists a Farber chain (Λ_n) in Γ with $\Lambda_1 = H$. This implies

$$\text{RG}(\Gamma, (\Gamma_n)) = \text{RG}(\Gamma, (\Lambda_n)) \leq \frac{d(H) - 1}{|\Gamma : H|} < \text{RG}(\Gamma) + \varepsilon$$

and so $\text{RG}(\Gamma, (\Gamma_n)) \leq \text{RG}(\Gamma)$.

On the other hand, $\text{RG}(\Gamma, (\Gamma_n)) \geq \text{RG}(\Gamma)$ by definition, so the theorem holds. \square

Actually, the above argument shows that assuming the multiplicativity of cost-1, a finitely generated residually finite group has fixed price 1 if and only if its absolute rank gradient is 0.

This could be relevant to decide whether uniform lattices have fixed price 1. Let G be a semisimple Lie group of \mathbb{R} -rank at least 2 and let Γ be a lattice in G . Gaboriau [14] shows that if Γ is non-uniform, i.e., G/Γ is not compact, then Γ has fixed price 1. Since, by a theorem of Borel, for every uniform higher rank lattice, there is a non-uniform lattice in the same ambient group, every uniform lattice in G has a measurable action of cost 1. But fixed price is not known for these lattices.

Conjecture 17 *All lattices in higher rank Lie groups have absolute rank gradient 0.*

Note that from a result of Sharma and Venkataramana [32], every non-uniform lattice Γ in G contains a subgroup of finite index generated by just 3 elements. This trivially implies $\text{RG}(\Gamma) = 0$.

Now we discuss rank 1 lattices. First, by [14], every lattice Γ in $\text{SL}(2, \mathbb{R})$ (or $\text{PSL}(2, \mathbb{R})$) has fixed price greater than 1. Using Theorem 1 we get that $\text{RG}(\Gamma) > 0$. Next we consider three examples, two of which have been studied extensively in the literature.

Example A. Let Γ be the Picard group $\text{PSL}(2, \mathbb{Z}[i])$. It is known that Γ is virtually a free by cyclic group. There is a subgroup T of index 24 in Γ which is the fundamental group of the complement of the Borromean links and

so (cf. [23]) T is an extension of the free group on 4 generators F_4 by \mathbb{Z} . From Proposition 13 it follows that $\text{RG}(\Gamma) = 0$.

Example B. Let Γ be the Bianchi group $\text{PSL}(2, \mathbb{Z}[w])$ where $w = \exp(2\pi i/3)$. Here there is a subgroup U of index 12 of Γ which is the fundamental group of the figure eight knot complement. Again, it is well known (see [10]) that U is the free by cyclic group with a presentation

$$\langle a, b, t \mid a^t = b^{-1}, b^t = b^2 ab \rangle.$$

So in this case again Γ is virtually a free by cyclic group and so $\text{RG}(\Gamma) = 0$.

Example C. Both Examples A and B are non-uniform. It is much harder to find uniform lattices of $\text{PSL}(2, \mathbb{C})$ with rank gradient zero. One such group has been found by Reid [25, Section 4, Theorem 1]. His example is a uniform arithmetic lattice Λ (that is, Λ is commensurable with the group of norm 1 elements of a suitable quaternion algebra over a number field K with just one complex embedding). Reid proves that the manifold \mathbb{H}^3/Λ has a finite cover which fibres over the circle. Group theoretically this means that Λ has a subgroup of finite index which is an ascending HNN extension with a finitely generated base group A . Proposition 12 now gives that the rank gradient of Λ is 0.

On a question of Kechris and Miller. Examples A and B present a negative answer to a question of Kechris and Miller [16, Problem 35.7]. They asked whether for a countable infinite Euclidean domain D the group $\text{SL}(2, D)$ has cost 1 if and only if D has infinitely many units. In fact, we get that these groups have rank gradient 0 and hence cost 1. We suspect that this is the general behaviour. Indeed, a well-known conjecture by Thurston asserts that every hyperbolic 3-manifold virtually fibers over the circle. This would imply that any lattice in $\text{SL}(2, \mathbb{C})$ has cost 1.

Let M be a finite closed orientable 3-manifold. A Heegaard decomposition (or splitting) for M is an expression of M as a union of two isomorphic handlebodies h_1, h_2 of genus $g \geq 0$ with boundary surfaces ∂h_i identified via a homeomorphism $f : \partial h_1 \rightarrow \partial h_2$. Such decomposition exists by [29], Section 8.3.

The Heegaard genus $g(M)$ of M is the minimal genus g of the surfaces ∂h_i in some Heegaard decomposition for M . Let $r(M) = d(\pi_1(M))$ be the rank of M .

It is easy to see that the fundamental group $\pi_1(h_i)$ surjects onto $\pi_1(M)$ and so $g(M) \geq r(M)$. In [33] Waldhausen asked if there is equality. This was proved false for Seifert manifolds by Boileau and Zieschang in [9]. Further work has been done by Schultens and Weidman who construct manifolds M with $g(M) = 4n$ and $r(M) \leq 3n$ ($n \geq 1$). However, the question of Waldhausen remained open for hyperbolic 3-manifolds. Also, until now, for all the known counterexamples the ratio $g(M)/r(M)$ was at most $3/2$. Theorem 5 addresses both these problems by proving that the ratio can get arbitrarily large even for arithmetic hyperbolic 3-manifolds.

We are ready to show that the Rank vs Heegaard genus conjecture and Fixed price problem conflict each other.

Proof of Theorem 2. Assume that the Fixed Price problem has an affirmative solution. Hence, by Lemma 15 and Theorem 16, any Farber chain has rank gradient equal to the absolute rank gradient of Γ .

Let us take the group Λ in Example C above. Take a chain $\Lambda = N_0 > N_1 > N_2 \cdots$ of *congruence* normal subgroups of Γ with trivial intersection. Define the manifolds $M_i = \mathbb{H}^3/N_i$. So we have

$$\pi_1(M_i) = N_i.$$

From results of Sarnak and Xue [26] (see also [22, p. 445, Example (f)]) it follows that if L is an arithmetic lattice of $\mathrm{PSL}(2, \mathbb{C})$, then L has property (τ) with respect to its congruence subgroups. In particular, Λ has property (τ) with respect to the chain $\{N_i\}$ and hence by a recent result of Marc Lackenby, [17, Theorem 1.5 and Corollary 1.6] we have

$$\lim_i \frac{g(M_i)}{[\Lambda : N_i]} > 0.$$

On the other hand, we have shown that $\mathrm{RG}(\Gamma) = 0$, so by Theorem 16 we have

$$\mathrm{RG}(\Gamma, (N_i)) = \lim_i \frac{r(M_i)}{[\Lambda : N_i]} = 0.$$

This implies that the ratio $g(M_i)/r(M_i)$ tends to infinity with i , thus proving the theorem. \square

Remark 1. If the manifold M is noncompact then its canonical compactification M^c is a finite 3-manifold with boundary. In this case a Heegaard decomposition of M is its expression as a union of two *compression bodies* B_1, B_2 with their positive boundaries ∂B_{i+} (both orientable surfaces of genus g) identified. The boundary of M is the negative boundary $\partial B_{1-} \cup \partial B_{2-}$. See [17], Section 3 for more details. The Heegaard genus $g(M)$ of M is defined to be the minimal genus of the surfaces identified in some decomposition for M^c as above. Again it is easy to see that in this case we have the inequality $2g(M) \geq r(M)$. Assuming the independence of the rank gradient on the chain, for Examples A or B, just as for compact manifolds, the ratio $g(M)/r(M)$ can be arbitrarily large.

Remark 2. Note that one does not need to obtain the multiplicativity of cost-1 for general measurable actions to obtain the independence of rank gradient from the chain. Trivially, it would be enough to settle this for profinite actions. Less trivially, by the recent work of the first author and Elek [3], it would be enough to settle the multiplicativity for the standard Bernoulli action $\{0, 1\}^\Gamma$.

References

- [1] M. ABÉRT AND N. NIKOLOV, The rank gradient from a combinatorial viewpoint, preprint
- [2] M. ABÉRT AND B. VIRÁG, Dimension and randomness in groups acting on rooted trees, *J. Amer. Math. Soc.* 18 (2005), 157-192.
- [3] M. ABÉRT AND G. ELEK, Dynamical properties of profinite actions, preprint
- [4] M. ABÉRT, On chains of subgroups in residually finite groups, preprint
- [5] L. BARTHOLDI AND R.I. GRIGORCHUK, On parabolic subgroups and Hecke algebras of some fractal groups, *Serdica Math. J.* 28 (2002), 47–90.
- [6] M. BEKKA, A. VALETTE, Group cohomology, harmonic functions and the first L^2 Betti number, *Potential Analysis* 6 (1997), 313-326.
- [7] N. BERGERON AND D. GABORIAU, Asymptotique des nombres de Betti, L^2 -invariants et laminations, (French. English summary) [Asymptotics of Betti numbers, L^2 -invariants and laminations], *Comment. Math. Helv.* 79 (2004), no. 2, 362–395.
- [8] A. BRUNNER, M. FRAME, Y. LEE AND N. WIELENBERG, Classifying Torsion-Free Subgroups of the Picard Group, *Transactions of the American Mathematical Society*, 282, No. 1. (1984), pp. 205-235.
- [9] M. BOILEAU AND H. ZIESCHANG, Heegaard genus of closed orientable Seifert 3-manifolds, *Invent. Math.* 76, no. 3, (1984), 455-468.
- [10] J. COSSEY, I.M.S. DEY AND S. MESKIN, Subgroups of knot groups, *Math. Z.* 121 (1971), 99-103.
- [11] A.H. DOOLEY AND V. YA. GOLODETS, The cost of an equivalence relation is determined by the cost of a finite index subrelation, preprint
- [12] M. FARBER, Geometry of growth: approximation theorems for L^2 invariants, *Math. Ann.* 311 (1998), no. 2, 335–375.
- [13] D. GABORIAU, Invariants L^2 de relations d'équivalence et de groupes, *Publ. math. Inst. Hautes Étud. Sci.*, 95 no. 1 (2002), 93-150.
- [14] D. GABORIAU, Coût des relations d'équivalence et des groupes. (French) [Cost of equivalence relations and of groups] *Invent. Math.* 139 (2000), no. 1, 41–98.
- [15] I. GRUSCHKO, Über die Basen eines freien Produktes von Gruppen. (Russian. German summary), *Rec. Math. [Mat. Sbornik] N.S.* 8 (50), (1940). 169–182.

- [16] A. KECHRIS AND B. MILLER, Topics in orbit equivalence, Lecture Notes in Mathematics, 1852. Springer-Verlag, Berlin, 2004.
- [17] M. LACKENBY, Heegaard splittings, the virtually Haken conjecture and property τ , *Invent. Math.* 164 (2006), no. 2, 317–359.
- [18] M. LACKENBY, Expanders, rank and graphs of groups, *Israel J. Math.*, 146 (2005), 357–370.
- [19] G. LEVITT, On the cost of generating an equivalence relation, *Ergodic Theory Dynam. Systems* 15 (1995), no. 6, 1173–1181.
- [20] D. D. LONG, A. LUBOTZKY AND A. REID, Heegaard genus and Property τ for hyperbolic 3-manifolds, preprint
- [21] W. LÜCK, L^2 -invariants: theory and applications to geometry and K -theory, *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*, 44. Springer-Verlag, Berlin, 2002
- [22] A. LUBOTZKY, Eigenvalues of the Laplacian, the first Betti number and the congruence subgroup problem. *Ann. of Math. (2)* 144 (1996), no. 2, 441–452.
- [23] C. MENZEL, Whitehead link complement and Borromean link complement are torus bundles, preprint
- [24] B. H. NEUMANN, On the number of generators of a free product. *J. London Math. Soc.* 18 (1943), 12–20.
- [25] A. REID, A non-Haken hyperbolic 3-manifold covered by a surface bundle, *Pacific J. Math.* 167, No. 1, (1995), 163-182.
- [26] P. SARNAK AND X. X. XUE, Bounds for multiplicities of automorphic representations. *Duke Math. J.* 64 (1991), no. 1, 207–227.
- [27] P. SHALEN, Hyperbolic volume and ranks of groups, Proceedings of the 2005 workshop on Heegaard splittings, Technion-IIT, Haifa, Israel, to appear
- [28] J. SCHULTENS AND R. WEIDMAN, On the geometric and the algebraic rank of graph manifolds, preprint
- [29] J. STILLWELL, Classical topology and combinatorial group theory, 2-nd ed. Springer Verlag, 1993.
- [30] G. STUCK AND R. J. ZIMMER, Stabilizers for ergodic actions of higher rank semisimple groups, *Ann. of Math. (2)* 139 (1994), no. 3, 723–747.

- [31] W. THURSTON, Geometry and topology of 3-manifolds, Lecture notes, Princeton University, 1978.
- [32] R. SARMA AND T. N. VENKATARAMANA, Generators of Arithmetic Groups, *Geometriae Dedicata*, 114:1 (2005), pp. 103-146.
- [33] F. WALDHAUSEN, Some problems on 3-manifolds, *Proc. Symposia in Pure Math.* 32, part 2,(1978), 313-322.