

Hydrides in the ISM. Electron induced reactivity of molecular cations

J. Zs. Mezei^{1,2}, E. Djuissi², A. Abdoulanziz², J. Boffelli², G. Farkas³, F. Iacob⁴, N. Pop⁵, D. Talbi⁶, M. Ayouz⁷, V. Kokoouline⁸ and I. F. Schneider^{2,9}

¹Institute for Nuclear Research (ATOMKI), H-4001 Debrecen, Hungary

²LOMC, CNRS, Normandie Université, le Havre, 76056 Le Havre, France

³University of Debrecen, 4032 Debrecen, Hungary

⁴West University of Timisoara, 300223 Timisoara, Romania

⁵Politechnica University of Timisoara, 300223 Timisoara, Romania

⁶LUPM, Université Montpellier, CNRS, 34095 Montpellier, France

⁷LGPM, CNRS, CentralSupélec Univ. Paris Saclay, 91190 Gif sur Yvette, France

⁸University of Central Florida, 32816 Orlando, Florida, USA

⁹LAC, CNRS, Univ. Paris-Saclay, 91405 Orsay, France

Electron-impact dissociative recombination, rovibrational (de)excitation and dissociative excitation of molecular cations are at the heart of molecular reactivity in the interstellar media and early Universe [1], being a major charge destruction path, and producing often atomic species in metastable states unaccessible through optical excitations.

These processes involve super-excited molecular states undergoing predissociation and autoionization, having thus strong resonant character. We use methods based on the Multichannel Quantum Defect Theory [2] and R-matrix theory [3], capable to account for the strong mixing between ionization and dissociative channels, open - direct mechanism - and closed - indirect mechanism, via capture into prominent Rydberg resonances correlating to the ground and excited ionic states, and for rotational effects. These features will be illustrated, comparisons with other existing theoretical and experimental results will be given and extensive data will be shown for several cations of high astrophysical and planetary relevance such as CH⁺ [2,4], N₂⁺ [5], and polyatomic systems like C₂H⁺ [6] and N₂H⁺ [7]. will be given, and perspectives on the advancement in the theoretical treatment - addressing polyatomic systems, predicting branching ratios - will be outlined.

J.Zs.M is grateful for the support from NKFIH through OTKA FK19-132989 and NKFIH-2019-2.1.11-TÉT-2020-00100 projects.

References

- [1] I. F. Schneider, O. Dulieu, and J. Robert (editors), *Eur. Phys. J. Web of Conf.* 84. (2015)
- [2] J. Zs. Mezei et al, *ACS Earth. Space. Chem.* **3**, 2376 (2019).
- [3] J. Tennyson, *Phys. Rep.* **491**, 29 (2010).
- [4] K. Chakrabarty et al, *J. Phys. B.* **51**, 104002 (2018).
- [5] A. Abdoulanziz et al, *J. Appl. Phys.* **129**, 053303 (2021).
- [6] J. Zs. Mezei et al, In preparation for *PSSST* (2021).
- [7] J. Zs. Mezei et al, In preparation for *MNRAS* (2021).