

COMMUNICATION

This document is the accepted manuscript version of the following article:

S. Giese, K. Klimov, A. Mikehazi, Z. Kelemen, D. S. Frost, S. Steinhauer, P. Müller, L. Nyulászi, C. Müller 2-(Dimethylamino)phosphinine: A Phosphorus-Containing Aniline Derivative *Ang. Chem. Int. Ed.* 2021, 60, 3581–3586. <https://doi/10.1002/anie.202014423>

2-(Dimethylamino)phosphinine: A Phosphorus Containing Aniline Derivative


Steven Giese,^[a] Katrin Klimov,^[a] Antal Mikeházi,^[b] Zsolt Kelemen,^[b] Daniel S. Frost,^[a] Simon Steinhauer,^[a] Peter Müller,^[c] László Nyulászi,^{*[b]} and Christian Müller^{*[a]}

In memory of Prof. Dr. Klaus Koch, Stellenbosch University, South Africa

Abstract: The yet unknown 2-amino-substituted λ^3, σ^2 -phosphinines are phosphorus containing aniline derivatives. Theoretical calculations show that the strong interaction of the π -donating NR₂-group with the aromatic system results in a high π -density at the phosphorus atom. We could now synthesize 2-N(CH₃)₂-functionalized phosphinines, starting from a 3-N(CH₃)₂-substituted 2-pyrone and (CH₃)₃Si-C≡P. Their reaction with CuBr·S(CH₃)₂ affords Cu(I)-complexes with the first example of a neutral phosphinine acting as a rare bridging μ_2 -P-4e donor-ligand between two Cu(I)-centers. Our experimental and theoretical investigations show, that 2-aminophosphinines are missing links in the series of known 2-donor-functionalized phosphinines.

After the successful synthesis of 2,4,6-triphenylphosphinine by Märkl in 1966 and the parent phosphinine C₅H₅P by Ashe III in 1971, several synthetic routes to access such intriguing aromatic phosphorus heterocycles have been developed.^[1] For the use of λ^3, σ^2 -phosphinines in more applied research fields, e.g. as ligands in catalytically active metal complexes or in photoluminescent coordination compounds, their specific functionalization is particularly important in order to modify their stereo-electronic properties and coordination abilities. While several polydentate phosphinines have been reported in literature, additional donor-substitution directly in 2-position of the heterocycle is relatively rare. These compounds can be considered as phosphorus containing derivatives of triarylphosphine, phenol and thiophenol (Figure 1 A, B).^[2] Heterocycles of type A have been used successfully as chelating,

small-bite angle diphosphorus ligands in several catalytic reactions.^[3] Grützmacher and co-workers have recently accessed sodium salts of phosphinin-2-ols (B, E = O) (Figure 1, C).^[4] While neutral phosphinin-2-ols show both the classical terminal 2e-donation *via* the lone-pair and the less common μ_2 -bridging 2e-lone-pair-donation (μ_2 -P-2e) to Cu(I)-centers (Figure 1, D), the anionic phosphinin-2-olate in E acts as a 4e-donor and bridges a cationic [Au(PPh₃)]⁺ and a neutral [AuCl] fragment (μ_2 -P-4e).^[5-7,8] This unprecedented bonding mode of the anionic ligand was concluded from the strong coordination shift in the ³¹P NMR spectrum to much lower frequencies compared to the uncomplexed ligand (strong negative Δ^{coord}),^[8] and the significantly elongated PC distance in the complex as compared to the ligand itself.



[a] Dr. Steven Giese, B. Sc. K. Klimov, M. Sc. D. S. Frost, Dr. S. Steinhauer, Prof. Dr. C. Müller
Freie Universität Berlin
Institut für Chemie und Biochemie

Fabeckstr. 34/36, 14195 Berlin, Germany
E-mail: c.mueller@fu-berlin.de

[b] A. Mikeházi, Dr. Z. Kelemen, Prof. Dr. L. Nyulászi,
Department of Inorganic and Analytical Chemistry
Budapest University of Technology and Economics
and MTA-BME Computation Driven Chemistry Research Group
Szt. Gellért tér 4
H-1111 Budapest, Hungary
E-mail: nyulaszi@mail.bme.hu

[c] Dr. P. Müller,
Department of Chemistry
Massachusetts Institute of Technology
77 Massachusetts Avenue, 2-325
Cambridge, MA 02139-4307, USA

Figure 1. 2-Donor-substituted phosphinines **A-C** and selected coordination compounds **D/E**, illustrating the μ_2 -P (2e) and μ_2 -P (4e) coordination mode.

Taking into account the literature known donor-functionalized phosphinines **A-C**, it seems surprising that the corresponding phosphorus containing aniline-derivatives still remain elusive. Because nitrogen is less electronegative than oxygen, the lone-pair at the nitrogen-atom in a hypothetical 2-amino-substituted phosphinine should also exert a significant interaction with the π -accepting aromatic system of the heterocycle, with pronounced consequences to the coordination chemistry of the resulting P,N-ligand. It therefore appeared intriguing to us to start an investigation on the theoretical properties, the synthesis and coordination chemistry of hitherto unknown 2-aminophosphinines. We first considered an evaluation of the electronic structure of amino-substituted P=C fragments. The π -orbital of a P=C building block is involved in conjugative interactions in a comparable manner to a C=C bond, as shown by the overall matching ionization energies of related P=C and C=C based compounds.^[9] Accordingly, the aromaticity of phosphinine is similar to that of benzene.^[10] While the P=C occupied orbitals show great similarity to the C=C orbitals, the unoccupied orbitals exhibit significant stabilization, rendering P=C units to highly efficient π -acceptors.^[11] Consequently, the usual P=C bond polarity (phosphorus is positive) can be tuned significantly by additional π -electron pair donors, such as NR₂-substituents, that are directly linked to the unsaturated carbon atom. This results in a 3c-4e interaction with a characteristic high energy orbital and a significant contribution at the phosphorus atom (Figure S1). In this way, an inversely polarized π -system with excess electrons at the phosphorus atom is formed.^[12] The most prominent examples have two amino groups at carbon, that are involved in a 6 π -electron delocalization. Accordingly, in case of phosphinidene-imidazole-ylidene, it was shown that the zwitterionic (**F'**) and the P=C double bonded (**F**) resonance structures have comparable contributions (Figure 2, S2).^[13]





Figure 2. Phosphinidene-imidazole-ylidene system **F/F'** with zwitterionic resonance structure, benzazaphosphole **G** and 2-amino-substituted phosphinine **H**, all having comparable >NC=P bonding motifs.

As a result, the phosphorus atom in **F/F'** is able to complex even two Lewis acids (e.g. boranes).^[14] Heinicke *et al.* observed for the cyclic 1,3-benzazaphosphole **G** that the strong electron donation of a single amino-group is sufficient to raise the energy of the polarized P=C bond to such an extent that a π -complex is formed with mercury, rather than the η^1 -coordination to the metal center via the in-plane lone-pair.^[15] Moreover, the formation of polymeric or oligomeric μ_2 -complexes with Cu(I) or Ag(I) were found as well.^[16] In case of the 1,3-azaphosphole, the $-\text{P}=\text{C}(\text{R})-\text{N}-$ unit is incorporated into the aromatic π -system, while in 2-aminophosphinines of type **H** with the same PCN bonding motif, the electron pair of the nitrogen substituent acts externally and, consequently, the 3c-4e system is not entirely involved in the cyclic delocalization.^[17] Thus, we were interested to evaluate the degree of interplay between the aromaticity and the polarizing effect of an external and strongly π -donating amino group. We focused primarily on the dimethylamino-substituent, as a 2-(dimethylamino)phosphinine (2-(CH₃)₂N-C₅H₄P, **1**) should be synthetically feasible (*vide infra*).

The B3LYP-D3/cc-pVTZ optimized geometry shows that the dimethylamino group in **1** is fully planar, in contrast to the pyramidal nitrogen atom in *N,N*-dimethylaniline, indicating a strong interaction between the nitrogen lone pair and the π -system. We found that the 6.8 kcal/mol rotational barrier for the N-C-bond in **1** is with 2.4 kcal/mol larger than in *N,N*-dimethylaniline. This can be attributed to the increased π -accepting property of phosphinine compared to benzene. Accordingly, the basicity of **1** is also reduced with respect to *N,N*-dimethylaniline, as shown by a decrease of the computed gas phase basicities (220 vs. 225 kcal/mol). The interaction is also clearly visualized by the shape of the intrinsic bonding orbital in **1** (Figure S3e).^[18] Accordingly, the PC(N) bond in the ring is by 0.051 Å shorter with respect to C₅H₅P, while the other PC bond distance remains virtually unchanged (Figure S4). Altogether, these geometric changes result in a small reduction of the Bird index ($I_{6,\text{C}_5\text{H}_5\text{P}} = 96.5$ vs. $I_{6,2-(\text{CH}_3)_2\text{N-C}_5\text{H}_4\text{P}} = 79.0$). Likewise, the corresponding NICS(1) value changes from -10.2 ppm to -8.9 ppm (Figure S5).^[19,20] Thus, while the amino substituent acts polarizing on the neighbouring PC bond, the aromaticity in the ring is nearly retained.

We further carried out Natural Resonance Theory (NRT) calculations on **1** (Figure 3, S6). Interestingly, it turned out that zwitterionic structures with C=N⁺ bonds have a significant contribution of about 16.0% in total, which exceeds the C=O⁺ contribution in the neutral phosphinin-2-ol (8.7%), while it is smaller than the C=O contribution in the anionic phosphinin-2-olate (40.3%).^[8] The non ionic aromatic resonance structures contribute with 47.6%. Again, this value is in-between the ones found for phosphinin-2-ol and phosphinin-2-olate (69.6 and 33.2%, respectively).^[8] Moreover, two resonance structures show two lone-pairs at the phosphorus atom contributing altogether with




12.9% in total to the electronic ground state, exceeding again the 12.5% contribution in phosphinin-2-ol.^[5,8]


COMMUNICATION

Figure 3. NRT analysis of 2-(CH₃)₂N-C₅H₄P (**1**).

The Kohn-Sham π -orbitals (Figure 4) show the n_N contribution of the (CH₃)₂N-substituent, increasing the phosphinine HOMO energy by more than 1 eV, while the energy of the LUMO (with a nodal surface at the substituting position) increases by 0.3 eV only.

**Figure 4.** Kohn-Sham-orbitals of C₅H₅P, 2-(CH₃)₂N-C₅H₄P (**1**) and 1-Me-1,3-azaphosphole.

Accordingly, **1** is a much stronger π -donor and somewhat weaker π -acceptor than the parent phosphinine C₅H₅P. The π -orbitals are delocalised throughout the ring, in accordance with the still significant aromaticity of the P-heterocycle (*vide supra*).^[20] While the in-plane phosphorus lone-pair energy for 1-Me-1,3-azaphosphole is at about the same energy as for C₅H₅P (likewise the corresponding ionization energies^[21]), the π -type HOMO (again in agreement with the ionization energies) and also the LUMO of the five-membered heterocycle are at much higher energy than in the six-membered ring, due to the presence of the donor nitrogen lone-pair, which is integrated into the π -system. In case of **1**, the HOMO energy is even higher than in 1-Me-1,3-azaphosphole, while the LUMO is at much lower energy. This renders **1** into a much “softer” ligand than 1-Me-1,3-azaphosphole. A comparison of the electrostatic potential maps of C₆H₆, C₅H₅P, (CH₃)₂N-C₆H₅, and **1** reveals additional interesting information (Figure 5). From the top view it is apparent that C₅H₅P is more positive (blue) than benzene, in accordance with its π -electron accepting property. Amino substitution induces negative charge (red) accumulation in the rings π -system, nevertheless, in *N,N*-




dimethylaniline ((CH₃)₂N-C₆H₅) nitrogen (lone pair) is the most negative. In case of **1**, however, the ring is more negative than the substituent nitrogen, in full accordance with the above discussed basicity differences of the two compounds.

Figure 5. Electrostatic potential maps for C₆H₆, C₅H₅P, (CH₃)₂N-C₆H₅ and 2-(CH₃)₂N-C₅H₄P (**1**). Parameters: r(red): -0.0250, y: -0.0125, g: 0.0000, l: 0.0125, b (blue) 0.0500. The electrostatic potential (in a.u.) is mapped on electron density isosurfaces of 0.02 e/au³.

From the side view, it is obvious that the red (electron rich) area at the phosphorus atom has a small ellipticity in the front part of the parent phosphinine C₅H₅P, in accordance with the high „s”-character of the in-plane lone-pair. In case of **1** not only the electron distribution becomes more asymmetric (elliptic), but also the negative charge (red color) increases compared to the parent phosphinine, due to the increased π -electron accumulation at the P-atom.

Having established that the amino group has an even stronger effect on the phosphinine π -system than the OH substituent, and that the perturbation approaches the anionic phosphinin-2-olate derivative, we turned our attention to the synthesis of **1**. We could already demonstrate that 3-bromo-2-pyrone reacts selectively with (CH₃)₃Si-C \equiv P to the corresponding 2-bromo-6-Si(CH₃)₃-functionalized phosphinine.^[22] We thus anticipated that a [4+2] cycloaddition reaction of phosphalkynes and 3-amino-substituted 2-pyrones might be a suitable synthetic route to access 2-aminophosphinines.^[23] Fan and Ma only recently reported on the synthesis of a series of 3-amino-functionalized 2-pyrones.^[24] After modification of their experimental procedure, we were able to access the hitherto unknown 3-(CH₃)₂N-2-pyrone **2** in reasonable yields (Scheme 1).

**Scheme 1.** Synthesis of dimethylamino-functionalized 2-pyrone **2**, TMS-substituted 2-(dimethylamino)phosphinine **3** and 2-(CH₃)₂N-C₅H₄P **1**. i): a) CuBr (0.2 eq.), 3Å MS, THF, 18 h, b) silica, toluene, 18 h; ii): a) HCl-Et₂O (2M, 4.7 eq.), r.t., 12 h; b) triethylamine (exc.), r.t. 1 h.

2-pyrone **2** indeed undergoes a [4+2] cycloaddition with (CH₃)₃Si-C \equiv P under formation of a single product, according to ³¹P{¹H} NMR spectroscopy (Scheme 1). As we could not find a ²J_(P-H) coupling in the corresponding proton-coupled ³¹P NMR spectrum, which could indicate the formation of a regioisomer with the amino-group in 3-position of the phosphorus heterocycle, we were confident that 2-(CH₃)₂N-6-TMS-phosphinine **3** (TMS = Si(CH₃)₃) had been formed selectively. The preferred formation of **3** over the possible regioisomer 2-TMS-3-(CH₃)₂N-phosphinine **3'** could

COMMUNICATION

be confirmed by means of DFT calculations. Mainly electronic but also steric effects are responsible for the regioselectivity of this cycloaddition reaction (for details see Figures S7-S9).

Phosphinine **3** shows a resonance in the $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum at $\delta(\text{ppm}) = 145.4$. This value is very similar to the one found for 2-hydroxy-phosphinine **B** (Figure 1, $\delta(\text{ppm}) = 149.1$), while 2,4,6-triaryl-phosphinines show a chemical shift at $\delta(\text{ppm}) = \sim 190$ in the $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum.^[1a] This observation is indicative for a pronounced interaction of the strongly π -donating amino group with the aromatic system of the phosphorus heterocycle, in accordance with the theoretical calculations. In analogy to our previous findings that 2-TMS-phosphinine undergoes protodesilylation under formation of the parent phosphinine $\text{C}_5\text{H}_5\text{P}$, compound **3** can easily be converted to 2-(CH_3)₂N- $\text{C}_5\text{H}_4\text{P}$ **1** in the presence of $\text{HCl}\cdot\text{Et}_2\text{O}$ and subsequent treatment with triethylamine (Scheme 1).^[7] Phosphinine **1** represents the first phosphorus containing *N,N*-dimethylaniline derivative and shows a single resonance at $\delta(\text{ppm}) = 126.5$ in the $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum. This value is close to the calculated ^{31}P NMR shift of 2-phospha-aniline (2- $\text{H}_2\text{N}-\text{C}_5\text{H}_4\text{P}$), reported by Alkorta and Elguero ($\delta(\text{ppm}) = 130.0$)^[25] and reflects again a significant interaction of the strong π -donating (CH_3)₂N-group with the π -system of the P-heterocycle. In contrast, the parent phosphinine $\text{C}_5\text{H}_5\text{P}$ shows a chemical shift at $\delta(\text{ppm}) = 206.6$ in the $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum.^[7] Theoretical calculations on Cu(I) complexes of **1** further reveal that the phosphorus atom is the preferred coordination site, and with more than one metal, also the μ_2 -coordination mode should be possible (for details see Figure S10). Therefore, **3** and **1** were reacted with $\text{CuBr}\cdot\text{S}(\text{CH}_3)_2$ in dichloromethane to afford single species, which show a resonance at $\delta(\text{ppm}) = 120.6$, respectively at $\delta(\text{ppm}) = 87.1$ in the $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum. Crystals of **4**, suitable for X-ray diffraction, could be obtained by slow crystallization from dichloromethane and, in case of **5**, by diffusion of pentane into a saturated solution of $[\text{CuBr}(\mathbf{1})]$ in acetonitrile. The crystallographic characterization of **4** reveals the presence of an infinite $(\text{CuBr})_x$ -band with the two edges decorated with phosphinine-ligands in an alternating manner (Figure 6 and S25).




Figure 6. Cu(I)-coordination polymers **4** and **5**.

The most striking feature of the structure of **4** is, that the phosphorus atom of each (CH_3)₂N-substituted phosphinine bridges two Cu(I)-centers, in analogy to phosphinin-2-ol and phosphinin-2-olate (*vide supra*). In contrast, 2-TMS-phosphinine itself forms Br-bridged dinuclear Cu(I)-complexes with the

classical η^1 -P coordination mode of the phosphorus heterocycle upon reaction with $\text{CuBr}\cdot\text{S}(\text{CH}_3)_2$.^[7] The nitrogen atom of each phosphinine in **4** is perfectly planar ($\Sigma_{\text{C-N-C}} = 360^\circ$) and the rather short N-C α bond distance of 1.331(5) Å indicates a significant N=C double bond character. The P-C α (N) bond length is with 1.80 Å considerably longer than the average P-C bond distance in free phosphinines (~ 1.70 - 1.73 Å) and in **D** (Figure 1, μ_2 -P-2e), while it is approaching the 1.82 Å found in **E** (μ_2 -P-4e).^[5,8] Furthermore, the alternation of the C-C bond lengths (between 1.358 and 1.422 Å, standard deviation 0.024 Å) is bracketed by the values found for **D** (μ_2 -P-2e, between 1.384 and 1.390 Å, standard deviation 0.002 Å) and **E** (μ_2 -P-4e, between 1.336 and 1.483 Å, standard deviation 0.064 Å).^[5,8] These structural features indicate a considerable reduction of the aromaticity of the heterocycle **4**, in accordance with the rare bridging μ_2 -P-4e coordination mode of the ligand, which is unprecedented for neutral phosphinines. However, the situation might be different in solution, as the coordination chemical shift difference of $\Delta^{\text{coord}} = -24.8$ ppm in the $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum is more in line with the μ_2 -P-2e mode, as reported for phosphinin-2-ol.^[8] It is noteworthy that for the CuCl and CuBr complexes of 1,3-benzazaphosphole (with the $-\text{P}=\text{C}-\text{N}<$ structural motif) a dynamic equilibrium has been detected between μ_1 -P bound complexes in solution, and μ_2 -P-2e bound complexes in the solid state.^[16b] Thus, it can be expected that also an average ^{31}P NMR signal appears in solutions of **4**.

The solid-state structure of **5**, along with selected bond lengths and distances is depicted in Figure 7. The crystallographic characterization of **5** reveals again the presence of a Cu(I)-coordination polymer (Figure 6). Since **5** was generated from $[\text{CuBr}(\mathbf{1})]$ in the presence of acetonitrile during crystallization, the two additional CuBr-moieties in the structure of **5** indicate the dissociation of the initially formed complex $[\text{CuBr}(\mathbf{1})]$. However, the NMR-spectroscopic investigation of the crystallization solution of **5** showed only one single resonance in the $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum with a chemical shift of $\delta = 105.7$ ppm. This can be attributed to **5**, which might again be in a dynamic equilibrium with other complexes in solution as the resulting coordination chemical shift difference of $\Delta^{\text{coord}} = -21$ ppm is again more in line with a μ_2 -P-2e coordination mode of the phosphinine.^[8]




Figure 7. Molecular structure of **5** in the crystal. Displacement ellipsoids are shown at the 50% probability level. a) Asymmetric unit; b) Representation of the infinite coordination polymer. Selected bond lengths (Å) and angles (°): P(1)-Cu(1): 2.2778(17); P(1)-Cu(2): 2.3028(17); C(1)-C(2): 1.372(8); C(2)-C(3): 1.407(9); C(3)-C(4): 1.388(9); C(4)-C(5): 1.411(8); P(1)-C(5): 1.775(6); P(1)-C(1): 1.729(6); N(1)-C(5): 1.359(8); Cu(1)-Cu(2): 2.6077(10); Cu(2)-Cu(3): 2.9710(11). C(1)-P(1)-C(5): 104.2(3); Cu(1)-P(1)-Cu(2): 69.40(5).

It should be mentioned here that the parent phosphinine C_5H_5P forms infinite coordination polymers with $CuBr \cdot S(CH_3)_2$ with the classical terminal $2e$ -donation *via* the phosphorus lone-pair.^[7] Similar to **4**, the phosphorus atom of each phosphinine ligand in **5** bridges two Cu(I)-centers of an unusual neutral $[Cu_3Br_3(CH_3CN)_3]$ unit, while the nitrogen atom is again perfectly planar ($\Sigma_{C-N-C} = 360^\circ$). The 1.775 Å P-C_α(N) bond length is longer than the average P-C bond distance in free phosphinines (~1.70–1.73 Å) but shorter than in **4**. Also the standard deviation of 0.016 Å for the ring CC distances indicates somewhat less bond length alternation than in case of **4**, which is still considerably larger than observed for **D** (μ_2 -P-2e). Thus, the π -system of ligand **1** is involved in the μ_2 -P-coordination, although this effect is larger in **3**, as shown by the comparison of the P-C_α(N) distances and the CC bond length alternation in **4** and **5**. Clearly, the π -accepting TMS-group facilitates the donation from the nitrogen lone pair of the amino group and increases the π -donor ability of the phosphorus atom. Accordingly, the electrostatic potential map of **3** (Figure S11) reveal a small depletion of the negative charge at the nitrogen atom in **3**, with some increase of negative charge at the phosphorus atom. Altogether, the μ_2 -P-4e mode seems to be a general coordination motif of 2-aminophosphinines in the solid state, at least for Cu(I).

Interestingly, the Cu(1)-Cu(2) distance of 2.6077(10) Å suggests that the phosphinine ligand **1** enables cuprophilic interactions.^[26] Similar Cu-Cu distances were reported for a Cu(I) complex of a μ_2 -P-2e coordinating 3-benzazaphosphole.^[16c] Apparently, the μ_2 -P complexation facilitates the close proximity of the Cu atoms.

In summary, we have synthesized the first phosphorus containing 2-aniline derivatives, 2-(CH₃)₂N-6-TMS-phosphinine **3** and 2-(CH₃)₂N-phosphinine **1**, starting from 3-(dimethylamino)-2-pyrone **2** and TMS-C≡P. The amino substitution increases significantly the π -electron density at the phosphorus atom in the ring, as shown by the calculated electrostatic potential map of **1**, compared to the parent phosphinine C_5H_5P . Natural Resonance Theory calculations reveal resonance structures with two lone-pairs at the phosphorus atom, contributing substantially to the electronic ground state of both **1** and **3**. Accordingly, both **1** and **3** form coordination polymers with $CuBr \cdot S(CH_3)_2$, in which the phosphorus atom of the phosphinine heterocycle bridges two Cu(I) centers in a rare μ_2 -P-4e coordination mode in the solid state, giving the first example for this bonding motif for a neutral substituted phosphinine. Our here presented results show, that 2-aminophosphinines are the most powerful neutral π -donating P,N-hybrid-ligands, which can form interesting polynuclear coordination compounds. Furthermore, the strength of the π -donation can be further modified by additional substitution. Moreover, these novel phosphorus heterocycles provide a missing link in the series of known 2-donor-functionalized phosphinines. Further studies on the complexes of **1** and **3** with other transition metals are currently under investigation.

Acknowledgements

Funding by Freie Universität Berlin and the Deutsche Forschungsgemeinschaft DFG (CM) is gratefully acknowledged. The authors thank the Scientific Computing Service of Freie Universität Berlin for the use of high-performance computing resources. LN is grateful to the Alexander von Humboldt Foundation for a reinvention, and for a BME Nanotechnology and Materials Science TKP2020 IE grant of NKFIH Hungary (BME IE-NAT TKP2020). ZK is grateful for the general support of Hungarian Academy of Science under the Premium Postdoctoral Research Program 2019.

Keywords: phosphinine • heterocycles • P,N-ligands • DFT calculations • crystallography


- [1] a) G. Märkl, *Angew. Chem.* **1966**, *78*, 907-908; b) A.J. Ashe, III, *J. Am. Chem. Soc.* **1971**, *93*, 3293-3295.
- [2] a) P. Le Floch, D. Charmichael, L. Ricard, F. Mathey, *J. Am. Chem. Soc.* **1993**, *115*, 10665-10670; b) N. Avarvari, P. Le Floch, F. Mathey, *J. Am. Chem. Soc.* **1996**, *118*, 11978-11979; c) Y. Mao, F. Mathey, *Chem. Eur. J.* **2011**, *17*, 10745-10751; d) P. Rosa, P. Le Floch, L. Ricard, F. Mathey, *J. Am. Chem. Soc.* **1997**, *119*, 9417-9423.
- [3] a) R. J. Newland, M. F. Wyatt, R. L. Wingad, S. M. Mansell, *Dalton Trans.* **2017**, *46*, 6172-6176; b) R. J. Newland, A. Smith, D. M. Smith, N. Fey, M. J. Hanton, S. M. Mansell, *Organometallics* **2018**, *37*, 1062-1073; c) R. J. Newland, M. P. Dolve, R. L. Wingad, S. M. Mansell, *New J. Chem.* **2018**, *42*, 19625-19636; d) R. J. Newland, J. M. Lynam, S. M. Mansell, *Chem. Commun.* **2018**, *54*, 5482-5485.
- [4] X. Chen, S. Alidori, F. F. Puschmann, G. Santiso-Quinones, Z. Benkő, Z. Li, G. Becker, H.-F. Grützmacher, H. Grützmacher, *Angew. Chem. Int. Ed.* **2014**, *53*, 1641-1645.
- [5] Y. Mao, K. Meng H. Lim, Y. Li, R. Ganguly, F. Mathey, *Organometallics* **2013**, *32*, 3562-3565.

- [6] a) B. Schmid, L.M. Venanzi, T. Gerfin, V. Gramlich, F. Mathey, *Inorg. Chem.* **1992**, *31*, 5117-5122; b) M. T. Reetz, E. Bohres, R. Goddard, M. C. Holthausen, W. Thiel, *Chem. Eur. J.* **1999**, *5*, 2101-2108; c) P. Rosa, L. Ricard, F. Mathey, P. Le Floch, *Organometallics*, **2000**, *19*, 5247-5250; d) M. J. Bakker, F. W. Vergeer, F. Hartl, K. Goubitz, J. Fraanje, P. Rosa, P. Le Floch, *Eur. J. Inorg. Chem.* **2000**, 843-845, e) M. J. Bakker, F. W. Vergeer, F. Hartl, P. Rosa, L. Ricard, P. Le Floch, M. J. Calhorda, *Chem. Eur. J.* **2002**, *8*, 1741-1752. f) X. Chen, Z. Li, F. Yanan, H. Grützmacher, *Eur. J. Inorg. Chem.* **2016**, 2016, 633-638.
- [7] M. H. Habicht, F. Wossidlo, T. Bens, E. A. Pidko, C. Müller, *Chem. Eur. J.* **2018**, *24*, 944-952.
- [8] Y. Hou, Z. Li, Y. Li, P. Liu, C.-Y. Su, F. Puschmann, H. Grützmacher, *Chem. Sci.* **2019**, *10*, 3168-3180.
- [9] L. Nyulászi, T. Veszprémi, J. Réffy, *J. Phys. Chem.* **1993**, *97*, 4011-4015.
- [10] a) L. Nyulászi, *Chem. Rev.* **2001**, *101*, 1229-1246. b) Z. Benkő, L. Nyulászi, *Top. Het. Chem.* **2009**, *19*, 27-83.
- [11] a) P. D. Burrow, A. J. Ashe III, D. J. Bellville, K. D. Jordan, *J. Am. Chem. Soc.* **1982**, *104*, 425-429; b) A. Modelli, J. F. Nixon, G. K. B. Clentsmith, *Chem. Phys. Lett.* **2002**, *366*, 122-127; c) A. Modelli, B. Hajgató, J. F. Nixon, L. Nyulászi *J. Phys. Chem. A* **2004**, *108*, 7440-7447.
- [12] L. Weber, *Eur. J. Inorg. Chem.* **2000**, 2425-2441.
- [13] a) G. Frison, A. Sevin, *J. Chem. Soc., Perkin Trans. 2*, **2002**, 1692-1697; b) Z. Kelemen, R. Streubel, L. Nyulászi, *RSC Adv.*, **2015**, *5*, 41795-41802.
- [14] a) A. J. Arduengo III, C. J. Carmalt, J. A. C. Clyburne, A. H. Cowley, R. Pyati, *Chem. Commun.* **1997**, 981-982; b) A. Doddi, D. Bockfeld, A. Nasr, T. Bannenberg, P. G. Jones, M. Tamm, *Chem. Eur. J.* **2015**, *21*, 16178-16189.
- [15] M. Ghalib, L. Könczöl, L. Nyulászi, P. G. Jones, G. J. Palm, J. W. Heinicke, *Dalton Trans.* **2014**, *43*, 51-54.
- [16] a) M. Ghalib, P. G. Jones, G. J. Palm, J. W. Heinicke, *RSC Advances* **2013**, *3*, 17726-17731; b) M. Ghalib, P. G. Jones, C. Schulzke, D. Sziebert, L. Nyulászi, J. W. Heinicke, *Inorg. Chem.* **2015**, *54*, 2117-2127; c) M. Ghalib, L. Könczöl, L. Nyulászi, G. J. Palm, C. Schulzke, J. W. Heinicke, *Dalton Trans.* **2015**, *44*, 1769-1774.
- [17] a) L. Nyulászi, G. Csonka, J. Réffy, T. Veszprémi, J. Heinicke, *J. Organomet. Chem.* **1989**, *373*, 49-55; b) T. Veszprémi, L. Nyulászi, J. Réffy, J. Heinicke, *J. Phys. Chem.* **1992**, *96*, 623-626; c) L. Nyulászi, T. Veszprémi, *J. Mol. Struct.* **1995**, *347*, 57-72.
- [18] G. Knizia, *J. Chem. Theory Comput.* **2013**, *9*, 4834-4843.
- [19] C.W. Bird, *Tetrahedron* **1985**, *41*, 1409-1414.
- [20] It is noteworthy that the position of the nodal surfaces differs from the parent phosphinine, due to the interacting n_N orbital.
- [21] a) T. Veszprémi, L. Nyulászi, J. Réffy, J. Heinicke, *J. Phys. Chem.* **1992**, *96*, 623-626; b) C. Batich, E. Heilbronner, V. Hornung, A. J. Ashe III, D. T. Clark, U. T. Cobley, D. Kilcast, I. Scanlan, *J. Am. Chem. Soc.* **1973**, *95*, 928-929.
- [22] M. H. Habicht, F. Wossidlo, M. Weber, C. Müller, *Chem. Eur. J.* **2016**, *22*, 12877-12883.
- [23] W. Rösch, M. Regitz, *Z. Naturforsch., B: Anorg. Chem., Org. Chem.* **1986**, *41*, 931.
- [24] W. Fan, S. Ma, *Angew. Chem. Int. Ed.* **2014**, *53*, 14542-14545.
- [25] I. Alkorta, J. Elguero, *Magn. Reson. Chem.* **2010**, *48*, S32-S37.
- [26] N. V. S. Harisomayajula, S. Makovetskiy, Y.-C. Tsai, *Chem. Eur. J.* **2019**, *2*, 8936-8954.

Entry for the Table of Contents

COMMUNICATION

The first phosphorus-containing 2-aniline-derivatives 2-(CH₃)₂N-6-(Me₃Si)-phosphinine and 2-(CH₃)₂N-phosphinine were synthesized. In-depths theoretical calculations reveal that resonance structures with two lone-pairs at the phosphorus atom exist. Consequently, the heterocycles act as π -donor-ligands and show the rare μ_2 -P-4e bridging coordination mode in the corresponding Cu(I)-complexes.



S. Giese, K. Klimov, A. Mikeházi, Z. Kelémen, D. S. Frost, S. Steinhauer, P. Müller, L. Nyulászi*, C. Müller*

Page No. – Page No.

2-(Dimethylamino)phosphinine: A Phosphorus Containing Aniline Derivative