

A Half-Space Approach to Order Dimension

Stephan Foldes · Jenő Szigeti

Received: 19 June 2006 / Accepted: 19 July 2007
© Springer Science + Business Media B.V. 2007

Abstract The aim of the present paper is to investigate the half-spaces in the convexity structure of all quasiorders on a given set and to use them in an alternative approach to classical order dimension. The main result states that linear orders can almost always be replaced by half-space quasiorders in the definition of the dimension of a partially ordered set.

Keywords Convexity · Quasiorder · Preorder · Half-space · Dimension

Mathematics Subject Classifications (2000) 06A06 · 06A07 · 52A01

1 Introduction

Within the framework of the general theory of abstract convexity (van de Vel [9]), strict quasiorders (irreflexive and transitive relations) on a set A can be thought of as convex subsets of $\{(x, y) \in A \times A \mid x \neq y\}$:

- (1) $\{(x, y) \in A \times A \mid x \neq y\}$ is a strict quasiorder,
- (2) Any intersection of strict quasiorders is a strict quasiorder,
- (3) Any nested union of strict quasiorders is a strict quasiorder.

The work of the first named author was partially supported by the European Community's Marie Curie Program (contract MTKD-CT-2004-003006). The second named author's work was supported by OTKA of Hungary No. T043034.

S. Foldes

Institute of Mathematics, Tampere University of Technology PL 553, Tampere 33101, Finland
e-mail: stephan.foldes@tut.fi

J. Szigeti (✉)

Institute of Mathematics, University of Miskolc, Miskolc-Egyetemvaros, Miskolc 3515, Hungary
e-mail: jeno.szigeti@uni-miskolc.hu

In general, a half-space is defined as a convex subset of the base set with a convex set complement. Abstract convexity theory addresses questions such as the representation of convex sets as intersections of half-spaces. For technical reasons, instead of the strict quasiorders in $\{(x, y) \in A \times A \mid x \neq y\}$, we shall consider the ordinary (reflexive) quasiorders in $A \times A$ (there is a natural one to one correspondence between them). We can use half-space quasiorders to define the half-space dimension of a quasiordered set, in a similar way as linear orders are used to define the order dimension of a partially ordered set. The aim of the present paper is to investigate the half-space quasiorders and to study the above dimension concept for quasiorders, along the lines of the classical theory of order dimension (see e.g. [1, 2, 7, 8]). Our main result (Theorem 2.16) states that linear orders can almost always be replaced by half-space quasiorders in the definition of the order dimension. Since there are considerably more half-spaces than linear orders, establishing upper bounds on order dimension can be easier using representations of partial orders as intersections of half-spaces. In order to demonstrate this, we give a simple proof for the “difficult” part of the classical Dushnik–Miller theorem (in [2]) about the dimension of the direct product of chains.

In Section 2 we provide some simple characterizations of half-spaces and examine the relationship between half-spaces and linear orders. A standard construction together with a complete description of half-spaces is also given. In the rest of Section 2, we show the tight connection between half-space dimension and classical order dimension. It turns out, that the half-space dimension and the order dimension of a partially ordered set can be different only for half-space partial orders.

In Section 3 we deal with direct products. First we prove that the direct product of quasiorders can be a half-space only in one exceptional situation. Then we use half-spaces to obtain the exact upper bound for the dimension in the above mentioned theorem of Dushnik and Miller.

2 Half-Spaces and the Dimension of Quasiordered Sets

A *quasiorder* γ on the set A is a reflexive and transitive relation:

$$\Delta_A = \{(a, a) \mid a \in A\} \subseteq \gamma \subseteq A \times A$$

and $(x, y) \in \gamma, (y, z) \in \gamma$ imply $(x, z) \in \gamma$ for all $x, y, z \in A$. The containment relation \subseteq provides a natural complete lattice structure on the set $\text{Quord}(A)$ of all quasiorders on A : $(\text{Quord}(A), \vee, \cap)$. If γ is a partial order, then we frequently use the standard notations $x \leq_\gamma y$ and $x <_\gamma y$ for $(x, y) \in \gamma$ and for $(x, y) \in \gamma, x \neq y$. For a relation γ , the inverse of γ is $\gamma^{-1} = \{(y, x) \mid (x, y) \in \gamma\}$ and for a quasiorder the intersection $\gamma \cap \gamma^{-1}$ is an equivalence on A . The equivalence class of an element $a \in A$ is denoted by $[a]_{\gamma \cap \gamma^{-1}}$, thus

$$A/(\gamma \cap \gamma^{-1}) = \{[a]_{\gamma \cap \gamma^{-1}} \mid a \in A\}.$$

It is well known that γ induces a natural partial order r_γ (in order to avoid repeated indices, we write \leq^γ instead of \leq_{r_γ}) on the above quotient set: for $a, b \in A$

$$[a]_{\gamma \cap \gamma^{-1}} \leq^\gamma [b]_{\gamma \cap \gamma^{-1}} \text{ if and only if } (x, y) \in \gamma \text{ for some } x \in [a]_{\gamma \cap \gamma^{-1}} \text{ and } y \in [b]_{\gamma \cap \gamma^{-1}}.$$

Also $[a]_{\gamma \cap \gamma^{-1}} \leq^\gamma [b]_{\gamma \cap \gamma^{-1}}$ holds if and only if $(x, y) \in \gamma$ for all $x \in [a]_{\gamma \cap \gamma^{-1}}$ and for all $y \in [b]_{\gamma \cap \gamma^{-1}}$. 53 54

A quasiorder $\alpha \subseteq A \times A$ is said to be a *half-space on A* if it has a “strong” complement in the lattice $(\text{Quord}(A), \subseteq)$, i.e. if $\alpha \cup \beta = A \times A$ and $\alpha \cap \beta = \Delta_A$ hold for some quasiorder $\beta \subseteq A \times A$. Clearly, this complement β is also a half-space and it is uniquely determined by α : $\beta = \Delta_A \cup ((A \times A) \setminus \alpha)$. It follows, that α is a half-space if and only if $\Delta_A \cup ((A \times A) \setminus \alpha)$ is transitive. The simplest examples of half-spaces are linear orders, the identity Δ_A and the full relation $A \times A$ on any set A . Complementary half-spaces are put into a pair of the form $\alpha \updownarrow \beta$ and can be characterized in the lattice $(\text{Quord}(A), \vee, \cap)$ as follows. 55 56 57 58 59 60 61 62

Proposition 2.1 *For any quasiorders $\alpha, \beta \in \text{Quord}(A)$ the following are equivalent:* 63

- (1) $\alpha \updownarrow \beta$ is a pair of complementary half-spaces, i.e. $\alpha \cap \beta = \Delta_A$ and $\alpha \cup \beta = A \times A$. 64 65
- (2) $\alpha \cap \beta = \Delta_A$ and $(\alpha \cap \gamma) \vee (\beta \cap \gamma) = \gamma$ for all $\gamma \in \text{Quord}(A)$. 66
- (3) $\alpha \cap \beta = \Delta_A$ and $(\alpha \cap \gamma) \cup (\beta \cap \gamma) = \gamma$ for all $\gamma \in \text{Quord}(A)$. 67

Proof (1) \implies (2): 68

$$\gamma = (A \times A) \cap \gamma = (\alpha \cup \beta) \cap \gamma = (\alpha \cap \gamma) \cup (\beta \cap \gamma) \subseteq (\alpha \cap \gamma) \vee (\beta \cap \gamma) \subseteq \gamma.$$

(2) \implies (1): Suppose that $\alpha \cup \beta \neq A \times A$, then $(a, b) \notin \alpha \cup \beta$ for some $a, b \in A$. Since $\gamma(a, b) = \Delta_A \cup \{(a, b)\}$ is a quasiorder on A , we have 69 70

$$(\alpha \cap \gamma(a, b)) \vee (\beta \cap \gamma(a, b)) = \gamma(a, b)$$

in contradiction with $\alpha \cap \gamma(a, b) = \beta \cap \gamma(a, b) = \Delta_A$. 71

(1) \implies (3) and (3) \implies (2) trivially. \square 72

For a half-space α the inverse relation α^{-1} is also a half-space, if $\alpha \updownarrow \beta$ for $\alpha, \beta \in \text{Quord}(A)$, then $\alpha^{-1} \updownarrow \beta^{-1}$. If $B \subseteq A$ is a subset, then the restriction of a quasiorder to B yields a quasiorder on B and a similar statement holds for half-spaces, $\alpha \updownarrow \beta$ implies that $\alpha \cap (B \times B) \updownarrow \beta \cap (B \times B)$. This observation leads to another characterization of half-spaces, which will be repeatedly used in the sequel. 73 74 75 76 77

Proposition 2.2 *For a quasiorder $\alpha \in \text{Quord}(A)$ the following are equivalent:* 78

- (1) α is a half-space. 79
- (2) $\alpha \cap (B \times B)$ is a half-space (on B) for any three element subset $B \subseteq A$. 80
- (3) For any $x, y, z \in A$ the relations $(x, y) \notin \alpha$, $(y, x) \notin \alpha$ and $(x, z) \in \alpha$, $z \neq x$ imply that $(y, z) \in \alpha$. 81 82
- (4) For any $x, y, z \in A$ the relations $(z, y) \notin \alpha$, $(y, z) \notin \alpha$ and $(x, z) \in \alpha$, $x \neq z$ imply that $(x, y) \in \alpha$. 83 84

Proof (1) \implies (2): This is a special case of our claim preceding Proposition 2.2. 85

(2) \implies (3): Let $(x, y) \notin \alpha$, $(y, x) \notin \alpha$ and $(x, z) \in \alpha$, $z \neq x$ for the elements $x, y, z \in A$ and take the three element subset $B = \{x, y, z\}$ of A . Suppose that $(y, z) \notin \alpha$ and consider the complementary half-space $\delta \subseteq B \times B$ of $\alpha \cap (B \times B)$. 86 87 88

89 Now

$$(\alpha \cap (B \times B)) \cup \delta = B \times B$$

90 implies that $(x, y) \in \delta$ and $(y, z) \in \delta$, whence $(x, z) \in (\alpha \cap (B \times B)) \cap \delta = \Delta_B$ can be
91 derived in contradiction with $z \neq x$.

92 (3) \implies (4): Let $(z, y) \notin \alpha$, $(y, z) \notin \alpha$ and $(x, z) \in \alpha$, $x \neq z$ for the elements
93 $x, y, z \in A$ and suppose that $(x, y) \notin \alpha$. Clearly, $(y, x) \in \alpha$ would imply $(y, z) \in \alpha$,
94 a contradiction. Thus $(x, y) \notin \alpha$, $(y, x) \notin \alpha$ and $(x, z) \in \alpha$, $x \neq z$, whence we obtain
95 that $(y, z) \in \alpha$, a contradiction. It follows that $(x, y) \in \alpha$.

96 (4) \implies (1): In order to see the transitivity of $\beta = \Delta_A \cup ((A \times A) \setminus \alpha)$ let $(x, y) \in \beta$,
97 $(y, z) \in \beta$, $x \neq y$ and suppose that $(x, z) \notin \beta$. We have either $(z, y) \notin \alpha$ or $(z, y) \in \alpha$.
98 In the first case $(z, y) \notin \alpha$, $(y, z) \notin \alpha$ and $(x, z) \in \alpha$, $x \neq z$ would imply that $(x, y) \in$
99 $\alpha \cap \beta = \Delta_A$, a contradiction. In the second case $(x, z) \in \alpha$ and $(z, y) \in \alpha$ would imply
100 that $(x, y) \in \alpha \cap \beta = \Delta_A$, a contradiction again. Thus we have $(x, z) \in \beta$. \square

101 **Proposition 2.3** *If α is a half-space quasiorder on A , then the induced partial order*
102 *r_α is a half-space quasiorder on $A/(\alpha \cap \alpha^{-1})$.*

103 *Proof* We can use part (3) in Proposition 2.2. If $([x]_{\alpha \cap \alpha^{-1}}, [y]_{\alpha \cap \alpha^{-1}}) \notin r_\alpha$,
104 $([y]_{\alpha \cap \alpha^{-1}}, [x]_{\alpha \cap \alpha^{-1}}) \notin r_\alpha$ and $([x]_{\alpha \cap \alpha^{-1}}, [z]_{\alpha \cap \alpha^{-1}}) \in r_\alpha$, $[z]_{\alpha \cap \alpha^{-1}} \neq [x]_{\alpha \cap \alpha^{-1}}$, then we have
105 $(x, y) \notin \alpha$, $(y, x) \notin \alpha$ and $(x, z) \in \alpha$, $z \neq x$. Since α is a half-space, we obtain first
106 $(y, z) \in \alpha$ and then $([y]_{\alpha \cap \alpha^{-1}}, [z]_{\alpha \cap \alpha^{-1}}) \in r_\alpha$. \square

107 **Proposition 2.4** *If $\gamma \subseteq A \times A$ is a quasiorder and $\gamma \subseteq \alpha$ for some half-space α on A ,*
108 *then there exists a half-space τ on A , such that $\gamma \subseteq \tau \subseteq \alpha$ and $\tau \cap \tau^{-1} = \gamma \cap \gamma^{-1}$.*

109 *Proof* Let R be a linear extension of the induced partial order r_γ and define the
110 relation $\tau \subseteq A \times A$ as follows:

$$\tau = \alpha \setminus \{(a, b) \in \alpha \cap \alpha^{-1} \mid [b]_{\gamma \cap \gamma^{-1}} <_R [a]_{\gamma \cap \gamma^{-1}}\}.$$

111 Since $(x, y) \in \gamma$ implies that $(x, y) \in \alpha$ and $[x]_{\gamma \cap \gamma^{-1}} \leq_R [y]_{\gamma \cap \gamma^{-1}}$, we obtain that
112 $(x, y) \in \tau$. Thus $\gamma \subseteq \tau \subseteq \alpha$ and $\gamma \cap \gamma^{-1} \subseteq \tau \cap \tau^{-1}$. If $(x, y) \in \tau \cap \tau^{-1}$, then the
113 relations $[y]_{\gamma \cap \gamma^{-1}} <_R [x]_{\gamma \cap \gamma^{-1}}$ and $[x]_{\gamma \cap \gamma^{-1}} <_R [y]_{\gamma \cap \gamma^{-1}}$ are not satisfied, whence
114 $[x]_{\gamma \cap \gamma^{-1}} = [y]_{\gamma \cap \gamma^{-1}}$ and $(x, y) \in \gamma \cap \gamma^{-1}$ can be derived. It follows, that $\tau \cap \tau^{-1} \subseteq$
115 $\gamma \cap \gamma^{-1}$ and hence $\tau \cap \tau^{-1} = \gamma \cap \gamma^{-1}$.

116 In order to see the transitivity of τ take $(x, y) \in \tau$ and $(y, z) \in \tau$. Now $(x, y) \in$
117 α and $(y, z) \in \alpha$ imply that $(x, z) \in \alpha$. Suppose that $(x, z) \notin \tau$, whence $(x, z) \in \alpha \cap$
118 α^{-1} and $[z]_{\gamma \cap \gamma^{-1}} <_R [x]_{\gamma \cap \gamma^{-1}}$ follow. The relations $(y, z) \in \alpha$ and $(z, x) \in \alpha$ imply that
119 $(y, x) \in \alpha$ and hence $(x, y) \in \alpha \cap \alpha^{-1}$. Similarly, $(z, x) \in \alpha$ and $(x, y) \in \alpha$ imply that
120 $(y, z) \in \alpha \cap \alpha^{-1}$. In view of $(x, y) \in \tau$ and $(y, z) \in \tau$ we have $[x]_{\gamma \cap \gamma^{-1}} \leq_R [y]_{\gamma \cap \gamma^{-1}}$ and
121 $[y]_{\gamma \cap \gamma^{-1}} \leq_R [z]_{\gamma \cap \gamma^{-1}}$, whence we obtain that $[x]_{\gamma \cap \gamma^{-1}} \leq_R [z]_{\gamma \cap \gamma^{-1}}$, a contradiction.

122 In order to prove that τ is a half-space we can use part (3) of Proposition 2.2.
123 Take $x, y, z \in A$ such that $(x, y) \notin \tau$, $(y, x) \notin \tau$ and $(x, z) \in \tau$, $z \neq x$. Now $(x, y) \notin \tau$
124 implies that either $(x, y) \notin \alpha$ or $(x, y) \in \alpha \cap \alpha^{-1}$ with $[y]_{\gamma \cap \gamma^{-1}} <_R [x]_{\gamma \cap \gamma^{-1}}$. Similarly,
125 $(y, x) \notin \tau$ implies that either $(y, x) \notin \alpha$ or $(y, x) \in \alpha \cap \alpha^{-1}$ with $[x]_{\gamma \cap \gamma^{-1}} <_R [y]_{\gamma \cap \gamma^{-1}}$.
126 It is easy to check that the only possibility to have $(x, y) \notin \tau$ and $(y, x) \notin \tau$ at the same
127 time is the case when $(x, y) \notin \alpha$ and $(y, x) \notin \alpha$. Since α is a half-space, $(x, y) \notin \alpha$,
128 $(y, x) \notin \alpha$ and $(x, z) \in \alpha$, $z \neq x$ imply that $(y, z) \in \alpha$. Suppose that $(y, z) \in \alpha \cap \alpha^{-1}$,

then $(x, z) \in \alpha$ and the transitivity of α imply that $(x, y) \in \alpha$, a contradiction. Thus we have $(y, z) \notin \alpha \cap \alpha^{-1}$, whence $(y, z) \in \tau$ follows. \square

Proposition 2.5 *Let the partial order α on A be a half-space. If λ is a linear order on A , then*

$$\alpha[\lambda] = \alpha \cup (\lambda \setminus (\alpha \cup \alpha^{-1}))$$

is a linear extension of α on A and $\alpha = \alpha[\lambda] \cap \alpha[\lambda^{-1}]$.

Proof In order to see the transitivity of $\alpha[\lambda]$ take $(x, y) \in \alpha[\lambda]$ and $(y, z) \in \alpha[\lambda]$ with $x \neq y \neq z$. Clearly, $(x, y) \in \alpha$ and $(y, z) \in \alpha$ imply $(x, z) \in \alpha$. If $(x, y) \in \alpha$ and $(y, z) \in \lambda \setminus (\alpha \cup \alpha^{-1})$, then $(y, z) \notin \alpha$, $(z, y) \notin \alpha$ and $(x, y) \in \alpha$, $x \neq y$, whence $(x, z) \in \alpha$ can be derived by part (4) of Proposition 2.2. Similarly, $(x, y) \in \lambda \setminus (\alpha \cup \alpha^{-1})$ and $(y, z) \in \alpha$ imply $(x, z) \in \alpha$ by part (3) of Proposition 2.2. If we have $(x, y) \in \lambda \setminus (\alpha \cup \alpha^{-1})$ and $(y, z) \in \lambda \setminus (\alpha \cup \alpha^{-1})$, then $(x, y) \in \lambda$ and $(y, z) \in \lambda$ imply $(x, z) \in \lambda$. Since $(x, y) \notin \alpha \cup \alpha^{-1}$ and $(y, z) \notin \alpha \cup \alpha^{-1}$ imply that $(x, y) \in \beta \cap \beta^{-1}$ and $(y, z) \in \beta \cap \beta^{-1}$ (here β is the complementary half-space of α), the transitivity of $\beta \cap \beta^{-1}$ gives that $(x, z) \in \beta \cap \beta^{-1}$, i.e. that $(x, z) \notin \alpha \cup \alpha^{-1}$. It follows that $(x, z) \in \lambda \setminus (\alpha \cup \alpha^{-1})$.

Suppose that $(x, y) \in \alpha[\lambda]$ and $(y, x) \in \alpha[\lambda]$, then $(x, y) \in \alpha$ and $(y, x) \in \lambda \setminus (\alpha \cup \alpha^{-1})$ is impossible. Similarly, $(x, y) \in \lambda \setminus (\alpha \cup \alpha^{-1})$ and $(y, x) \in \alpha$ is also impossible. Thus we have either $(x, y) \in \alpha$, $(y, x) \in \alpha$ or $(x, y) \in \lambda \setminus (\alpha \cup \alpha^{-1})$, $(y, x) \in \lambda \setminus (\alpha \cup \alpha^{-1})$, in both cases $x = y$ follows by the antisymmetric properties of α and λ , respectively.

Suppose that $(x, y) \notin \alpha$ and $(y, x) \notin \alpha$, then $(x, y) \notin \alpha \cup \alpha^{-1}$. Now $(x, y) \in \lambda$ implies $(x, y) \in \lambda \setminus (\alpha \cup \alpha^{-1})$ and $(y, x) \in \lambda$ implies $(y, x) \in \lambda \setminus (\alpha \cup \alpha^{-1})$. We proved that $\alpha[\lambda]$ is a linear order.

Using $\alpha \cap (\lambda \setminus (\alpha \cup \alpha^{-1})) = \alpha \cap (\lambda^{-1} \setminus (\alpha \cup \alpha^{-1})) = \emptyset$ and $\lambda \cap \lambda^{-1} = \Delta_A$, it is straightforward to see that $\alpha = \alpha[\lambda] \cap \alpha[\lambda^{-1}]$. \square

Corollary 2.6 *If α is a half-space quasiorder on A , then the induced partial order is of the form $r_\alpha = R_1 \cap R_2$ for some linear orders R_1 and R_2 on $A/(\alpha \cap \alpha^{-1})$, i.e. r_α has order dimension at most 2.*

Proof The partial order r_α is a half-space on $A/(\alpha \cap \alpha^{-1})$ by Proposition 2.3. If R is an arbitrary linear order on $A/(\alpha \cap \alpha^{-1})$, then $R_1 = r_\alpha[R]$ and $R_2 = r_\alpha[R^{-1}]$ are linear orders on $A/(\alpha \cap \alpha^{-1})$ with $r_\alpha = r_\alpha[R] \cap r_\alpha[R^{-1}]$ by Proposition 2.5. \square

We remark that Corollary 2.6 does not characterize half-spaces entirely.

As already noted, any linear order λ on A is an example of a half-space: $\lambda \uparrow \lambda^{-1}$. Let $f : A \rightarrow X$ be a function, $Y \subseteq X$ a subset, and R a linear order on X . Define the following relations on A :

$$\ker_Y(f) = \Delta_A \cup \{(a, b) \in A \times A \mid f(a) = f(b) \in Y\},$$

$$f^{-1}(R) = \Delta_A \cup \{(a, b) \in A \times A \mid f(a) <_R f(b)\}.$$

Note that $\ker_X(f)$ is the ordinary kernel

$$\ker(f) = \{(a, b) \in A \times A \mid f(a) = f(b)\}.$$

164 The following is a standard construction of a half-space using a linear order.

165 **Proposition 2.7** *Let (A, γ) be a quasiordered set, (X, ρ) a partially ordered set and*
 166 *$f : A \longrightarrow X$ a (γ, ρ) quasiorder preserving function: $(x, y) \in \gamma \implies (f(x), f(y)) \in \rho$*
 167 *for all $x, y \in A$. If $Y \subseteq X$ is a subset, R is a linear extension of ρ on X and $\gamma \cap$*
 168 *$\ker(f) \subseteq \ker_Y(f)$, then*

$$\alpha = \ker_Y(f) \cup f^{-1}(R)$$

169 *is a half-space extension of γ and $\alpha \cap \alpha^{-1} = \ker_Y(f)$.*

170 *If $Y = \emptyset$, then $\ker_Y(f) = \Delta_A$ and $\ker_Y(f) \cup f^{-1}(R) = f^{-1}(R)$ is a partial order.*

171 *If $Y = X$, then $\ker_Y(f) = \ker(f)$ (now $\gamma \cap \ker(f) \subseteq \ker_Y(f)$ is automatically sat-*
 172 *isfied) and $\ker_Y(f) \cup f^{-1}(R) = \ker(f) \cup f^{-1}(R)$ is a half-space extension of γ . In*
 173 *particular, if $\kappa : A \longrightarrow A/(\gamma \cap \gamma^{-1})$ is the canonical surjection and R is a linear*
 174 *extension of the induced partial order r_γ on $A/(\gamma \cap \gamma^{-1})$, then $\ker(\kappa) \cup \kappa^{-1}(R) =$*
 175 *$(\gamma \cap \gamma^{-1}) \cup \kappa^{-1}(R)$ is a half-space extension of γ .*

176 *Proof* The containment $\gamma \subseteq \ker_Y(f) \cup f^{-1}(R)$ is a consequence of $\rho \subseteq R$, $\gamma \cap$
 177 $\ker(f) \subseteq \ker_Y(f)$ and of the quasiorder preserving property of f . It is easy to see
 178 that $\ker_Y(f) \cup f^{-1}(R)$ and $\ker_{X \setminus Y}(f) \cup f^{-1}(R^{-1})$ are quasiorders on A . We have

$$(\ker_Y(f) \cup f^{-1}(R)) \cup (\ker_{X \setminus Y}(f) \cup f^{-1}(R^{-1})) = A \times A$$

179 and

$$(\ker_Y(f) \cup f^{-1}(R)) \cap (\ker_{X \setminus Y}(f) \cup f^{-1}(R^{-1})) = \Delta_A,$$

180 thus $\ker_Y(f) \cup f^{-1}(R) \upharpoonright \ker_{X \setminus Y}(f) \cup f^{-1}(R^{-1})$. Also $\alpha \cap \alpha^{-1} = \ker_Y(f)$ is obvious.
 181 To conclude the proof, it is enough to note that κ is a (γ, r_γ) quasiorder preserving
 182 function. \square

183 **Proposition 2.8** *Let (A, γ) be a quasiordered set, (X, ρ) a partially ordered set*
 184 *and $f : A \longrightarrow X$ a completely (γ, ρ) quasiorder preserving function: $(x, y) \in \gamma \iff$*
 185 *$(f(x), f(y)) \in \rho$ for all $x, y \in A$. If $Y_i \subseteq X$, $i \in I$ is a collection of subsets, $\gamma \cap$*
 186 *$\ker(f) \subseteq \ker_{Y_i}(f)$ for all $i \in I$ and $\{R_i \mid i \in I\}$ is a set of linear extensions of ρ with*
 187 *$\bigcap_{i \in I} R_i = \rho$, then*

$$\bigcap_{i \in I} (\ker_{Y_i}(f) \cup f^{-1}(R_i)) = \gamma,$$

188 *where the half-spaces $\ker_{Y_i}(f) \cup f^{-1}(R_i)$, $i \in I$ are described in Proposition 2.7. In*
 189 *particular, if $\kappa : A \longrightarrow A/(\gamma \cap \gamma^{-1})$ is the canonical surjection and $\{R_i \mid i \in I\}$ is a*
 190 *set of linear extensions of the induced partial order r_γ on $A/(\gamma \cap \gamma^{-1})$ with $\bigcap_{i \in I} R_i = r_\gamma$,*
 191 *then*

$$\bigcap_{i \in I} (\ker(\kappa) \cup \kappa^{-1}(R_i)) = \bigcap_{i \in I} ((\gamma \cap \gamma^{-1}) \cup \kappa^{-1}(R_i)) = \gamma.$$

192 *Proof* We only have to show that

$$\bigcap_{i \in I} (\ker_{Y_i}(f) \cup f^{-1}(R_i)) \subseteq \gamma.$$

In view of the definition of $\ker_{Y_i}(f) \cup f^{-1}(R_i)$, the relation

$$(a, b) \in \bigcap_{i \in I} (\ker_{Y_i}(f) \cup f^{-1}(R_i))$$

ensures that $f(a) \leq_{R_i} f(b)$ for all $i \in I$. Now $\bigcap_{i \in I} R_i = \rho$ implies $f(a) \leq_{\rho} f(b)$, whence we obtain $(a, b) \in \gamma$. To conclude the proof, it is enough to note that κ is completely (γ, r_{γ}) quasiorder preserving. \square

The following is now a straightforward consequence.

Theorem 2.9 Any quasiorder on A can be obtained as an intersection of half-space quasiorders on A .

In terms of the classification of convexities by separation axioms (van de Vel [9]) the above theorem means that the convexity on the base set $\{(x, y) \in A \times A \mid x \neq y\}$ whose convex sets are the strict quasiorders on A is an S_3 convexity, i.e. a convex set K can be always separated from any element of the base set not in K by complementary half-spaces. This is the case in the standard convexity of a Euclidean space and, as Szpilrajn's theorem [7] shows, in the convexity on the set $\{(x, y) \in A \times A \mid x \neq y\}$ whose convex sets are the strict partial orders on A plus $\{(x, y) \in A \times A \mid x \neq y\}$ itself. However, it is not difficult to see that, unlike in Euclidean space, in quasiorder convexity or in the coarser partial order convexity, disjoint convex sets cannot always be separated by complementary half-spaces. A counterexample with respect to both the quasiorder and partial order convexities is provided, for $A = \{1, 2, 3, 4\}$, by the partial orders $\{(1, 2), (3, 4)\}$ and $\{(1, 4), (3, 2)\}$.

Theorem 2.9 enables us to define a *half-space realizer* of a quasiorder $\gamma \subseteq A \times A$ as a set $\{\alpha_i \mid i \in I\}$ of half-spaces on A with $\bigcap_{i \in I} \alpha_i = \gamma$. The *half-space dimension* $\text{hsdim}(A, \gamma)$ of a quasiordered set (A, γ) is the minimum of the cardinalities of the half-space realizers of γ . The close analogy between the half-space dimension and the usual order dimension of a partially ordered set can be seen immediately. The observation preceding Proposition 2.2 guarantees that

$$\text{hsdim}(B, \gamma \cap (B \times B)) \leq \text{hsdim}(A, \gamma)$$

for any subset $B \subseteq A$. Since any linear order is a half-space, for a partially ordered set (A, γ) we have $\text{hsdim}(A, \gamma) \leq \dim(A, \gamma)$, where \dim denotes the order dimension. In general, here we can not expect equality. The partial order of the four element Boolean lattice M_2 is a half-space, thus $\text{hsdim}(M_2, \leq) = 1$, while $\dim(M_2, \leq) = 2$. The next inequality is also a straightforward consequence of Proposition 2.8.

Corollary 2.10 For a quasiordered set (A, γ) we have

$$\text{hsdim}(A, \gamma) \leq \dim(A/(\gamma \cap \gamma^{-1}), r_{\gamma}).$$

The following theorem gives a complete description of half-space quasiorders.

Theorem 2.11 If $\alpha \subseteq A \times A$ is a relation, then the following are equivalent.

- (1) α is a half-space quasiorder on A .

- 227 (2) *There exists an equivalence relation ε on A , a linear order R on the factor set*
 228 *A/ε and a function $t : A/\varepsilon \rightarrow \{0, 1\}$ with $t([a]_\varepsilon) = 0$ where $[a]_\varepsilon = \{a\}$ such that*

$$\alpha = \Delta_A \cup \{(a, b) \in A \times A \mid [a]_\varepsilon = [b]_\varepsilon \text{ and } t([a]_\varepsilon) = 1\} \cup \{(a, b) \in A \times A \mid [a]_\varepsilon <_R [b]_\varepsilon\}.$$

 229 (3) *There exist a set X , a subset $Y \subseteq X$, a linear order R on X and a function $f : A \rightarrow X$ such that $\alpha = \ker_Y(f) \cup f^{-1}(R)$.*
 231 (4) *There exists an equivalence relation ε on A such that α is either the full or the identity relation on each ε -equivalence class, and any irredundant set of*
 232 *representatives of the ε -equivalence classes is linearly ordered by α .*
 233

234 *Proof* (1) \implies (2): Let $\alpha \upharpoonright \beta$ be complementary half-spaces and take

$$\varepsilon = (\alpha \cap \alpha^{-1}) \cup (\beta \cap \beta^{-1}).$$

235 Clearly, ε is reflexive and symmetric. Assume that $(x, y) \in \alpha \cap \alpha^{-1}$ and $(y, z) \in \beta \cap \beta^{-1}$. Since $\alpha \cup \beta = A \times A$, we have either $(x, z) \in \alpha$ or $(x, z) \in \beta$. In the first case
 236 $(y, x) \in \alpha$ implies that $(y, z) \in \alpha \cap \beta = \Delta_A$. In the second case $(z, y) \in \beta$ implies that
 237 $(x, y) \in \alpha \cap \beta = \Delta_A$. Thus $(x, y) \in \alpha \cap \alpha^{-1}$ and $(y, z) \in \beta \cap \beta^{-1}$ imply $x = y$ or $y = z$.
 238 Similarly, $(x, y) \in \beta \cap \beta^{-1}$ and $(y, z) \in \alpha \cap \alpha^{-1}$ also imply $x = y$ or $y = z$. In view
 239 of the above observations, it is easy to see that ε is transitive. We also have $[a]_\varepsilon =$
 240 $[a]_{\alpha \cap \alpha^{-1}} \cup [a]_{\beta \cap \beta^{-1}}$ and $[a]_{\alpha \cap \alpha^{-1}} = \{a\}$ or $[a]_{\beta \cap \beta^{-1}} = \{a\}$ for all $a \in A$.

241 We claim that $(a, b) \in \alpha$ and $[a]_\varepsilon \neq [b]_\varepsilon$ imply that $(x, y) \in \alpha$ for all $x \in [a]_\varepsilon$ and
 242 for all $y \in [b]_\varepsilon$. Suppose that $(x, y) \notin \alpha$, then $(x, y) \in \beta$. In view of $(x, a), (y, b) \in \varepsilon$
 243 we have the following cases. (1) $(x, a), (b, y) \in \alpha$, whence $(x, y) \in \alpha$ can be obtained,
 244 a contradiction. (2) $(x, a) \in \alpha$ and $(y, b) \in \beta$, whence $(x, b) \in \alpha \cap \beta = \Delta_A$ can be ob-
 245 tained in contradiction with $[x]_\varepsilon = [a]_\varepsilon \neq [b]_\varepsilon$. (3) $(a, x) \in \beta$ and $(b, y) \in \alpha$, whence
 246 $(a, y) \in \alpha \cap \beta = \Delta_A$ can be obtained in contradiction with $[a]_\varepsilon \neq [b]_\varepsilon = [y]_\varepsilon$. (iv)
 247 $(a, x), (y, b) \in \beta$, whence $(a, b) \in \alpha \cap \beta = \Delta_A$ can be obtained in contradiction with
 248 $[a]_\varepsilon \neq [b]_\varepsilon$. Thus the claim is proved.

249 Using our claim it is straightforward to check that

$$R = \{([a]_\varepsilon, [b]_\varepsilon) \mid (a, b) \in \alpha\}$$

251 is a linear order on A/ε . For $a \in A$ let

$$t([a]_\varepsilon) = \begin{cases} 1 & \text{if } [a]_\varepsilon = [a]_{\alpha \cap \alpha^{-1}} \neq \{a\} \\ 0 & \text{otherwise} \end{cases}.$$

252 Clearly, t is well defined, moreover $[a]_\varepsilon = \{a\}$ implies $[a]_\varepsilon = [a]_{\alpha \cap \alpha^{-1}} = \{a\}$ and
 253 $t([a]_\varepsilon) = 0$. If $t([a]_\varepsilon) = 1$, then $[a]_\varepsilon = [a]_{\alpha \cap \alpha^{-1}}$ and $[a]_\varepsilon = [b]_\varepsilon$ implies that $(a, b) \in \alpha$.
 254 It follows that

$$\Delta_A \cup \{(a, b) \in A \times A \mid [a]_\varepsilon = [b]_\varepsilon \text{ and } t([a]_\varepsilon) = 1\} \cup \{(a, b) \in A \times A \mid [a]_\varepsilon <_R [b]_\varepsilon\} \subseteq \alpha.$$

255 If $[a]_\varepsilon = [b]_\varepsilon$, then $(a, b) \in \alpha$ and $a \neq b$ implies that $[a]_\varepsilon = [a]_{\alpha \cap \alpha^{-1}} \neq \{a\}$, whence

$$\alpha \subseteq \Delta_A \cup \{(a, b) \in A \times A \mid [a]_\varepsilon = [b]_\varepsilon \text{ and } t([a]_\varepsilon) = 1\} \cup \{(a, b) \in A \times A \mid [a]_\varepsilon <_R [b]_\varepsilon\}$$

256 can be obtained.

257 (2) \implies (3): It is straightforward to see that $\alpha = \ker_Y(f) \cup f^{-1}(R)$, where $X =$
 258 A/ε , $Y = \{[a]_\varepsilon \mid a \in A \text{ and } t([a]_\varepsilon) = 1\}$ and $f : A \rightarrow X$ is the canonical surjection.
 259 Thus any half-space quasiorder can be obtained by the standard construction of
 260 Proposition 2.7.

(3) \implies (1): This implication is a part of Proposition 2.7. 261

(2) \iff (4): Condition (4) is simply a reformulation of (2). \square 262

Remark 2.12 The triple $(f, Y \subseteq X, R)$ given in the (2) \implies (3) part of the above 263
proof has the following universal property. If $g : A \longrightarrow U$ is a function, $V \subseteq U$ is 264
a subset and S is a linear order on U such that 265

$$\alpha = \ker_V(g) \cup f^{-1}(S),$$

then there exists a unique function $h : X \longrightarrow U$ with $h \circ f = g$, moreover $h(Y) \subseteq V$, 266
 $g^{-1}(\{y\})$ is a one element set for all $y \in h(X \setminus Y) \cap V$ and h is $(<_R, <_S)$ strict order 267
preserving 268

In view of the above characterization of the half-space α , an equivalence class $[a]_\varepsilon$ 269
is called a *box* of α , such a box is called *full* if $t([a]_\varepsilon) = 1$ and *empty* if $t([a]_\varepsilon) = 0$ (note 270
that a one element box is always empty). A subset $B \subseteq A$ is a box of the half-space 271
 α , if and only if there are no elements $b_1, b_2 \in B$ such that $(b_1, b_2) \in \alpha$, $(b_2, b_1) \notin \alpha$ 272
and B is maximal with respect to this property. A box is empty if $\alpha \cap (B \times B) = \Delta_B$ 273
and full if $|B| > 1$ and $B \times B \subseteq \alpha$. 274

In certain situations it is also convenient to give a half-space as 275

$$\alpha = (B_w, w \in W, \leq_w, t),$$

where the subsets $B_w \subseteq A$, $w \in W$ are the boxes of α , the linear order \leq_w is given 276
on the index set W and $t(B_w) = 1$ or $t(B_w) = 0$ shows that B_w is full or empty. If W is 277
finite, then we can write $W = \{1, 2, \dots, n\}$ and $\alpha = (B_1 < B_2 < \dots < B_n, t)$. If $\alpha \updownarrow \beta$ is 278
a complementary pair of half-spaces, then α and β have the same boxes, a full α -box 279
is an empty β -box and a full β -box is an empty α -box, moreover the linear order of 280
the boxes in α and β are opposite to each other. It is also clear that $[a]_{\alpha \cap \alpha^{-1}} = \{a\}$ if 281
 $[a]_\varepsilon$ is empty and $[a]_{\alpha \cap \alpha^{-1}} = [a]_\varepsilon$ if $[a]_\varepsilon$ is full. 282

With reference to the terminology of interval decompositions and lexicographic 283
sums of partial orders and more general relations (see e.g. [3–6]), it is clear from 284
Condition (4) of Theorem 2.11 that half-space quasiorders are precisely the lexico- 285
graphic relational sums of trivial and full binary relations over a linear order, i.e. they 286
are the binary relations decomposable into intervals such that the restriction to each 287
interval is a trivial or full relation and the quotient is a linear order. 288

Theorem 2.13 If (A, γ) is a quasiordered set and $\{\alpha_i \mid i \in I\}$ is a half-space realizer 289
of γ with $|I| \geq 2$, then there exists an I -indexed family R_i , $i \in I$ of linear extensions of 290
the induced partial order r_γ on $A/(\gamma \cap \gamma^{-1})$ such that 291

$$\bigcap_{i \in I} R_i = r_\gamma.$$

Proof By Proposition 2.4, for each $i \in I$ there exists a half-space τ_i on A such that 292
 $\gamma \subseteq \tau_i \subseteq \alpha_i$ and $\tau_i \cap \tau_i^{-1} = \gamma \cap \gamma^{-1}$. Clearly, $\bigcap_{i \in I} \alpha_i = \gamma$ implies that $\bigcap_{i \in I} \tau_i = \gamma$, whence 293

$$\bigcap_{i \in I} r_{\tau_i} = r_\gamma$$

can be derived for the induced partial orders $r_{\tau_i}, i \in I$ on $A/(\tau_i \cap \tau_i^{-1}) = A/(\gamma \cap \gamma^{-1})$. Using the notation $\pi_i = r_{\tau_i}$, Proposition 2.3 ensures that each partial order π_i is a half-space on $P = A/(\gamma \cap \gamma^{-1})$. We claim, that

$$\rho = \Delta_P \cup \left(\left(\bigcup_{i \in I} \pi_i \right)^{-1} \setminus \left(\bigcup_{i \in I} \pi_i \right) \right)$$

is partial order on P . The reflexive and antisymmetric properties of ρ can be immediately seen. In order to prove the transitivity of ρ consider the pairs $(x, y) \in \rho$ and $(y, z) \in \rho$ with $x, y, z \in P$ being different. We have $(y, x) \in \pi_j, (z, y) \in \pi_k$ for some $j, k \in I$ and $(x, y) \notin \bigcup_{i \in I} \pi_i, (y, z) \notin \bigcup_{i \in I} \pi_i$. If $(z, y) \in \pi_j$, then the transitivity of π_j implies $(z, x) \in \pi_j$. If $(z, y) \notin \pi_j$, then $(y, z) \notin \pi_j$ and the half-space property of π_j imply that $(z, x) \in \pi_j$ (see part (3) of Proposition 2.2). It follows that $(x, z) \in \left(\bigcup_{i \in I} \pi_i \right)^{-1}$. Suppose that $(x, z) \in \bigcup_{i \in I} \pi_i$, then $(x, z) \in \pi_t$ for some $t \in I$. If $(z, y) \in \pi_t$, then the transitivity of π_t gives that $(x, y) \in \pi_t$, a contradiction. If $(z, y) \notin \pi_t$, then $(y, z) \notin \pi_t$ and the half-space property of π_t gives that $(x, y) \in \pi_t$ (see part (4) of Proposition 2.2), another contradiction. Thus $(x, z) \notin \bigcup_{i \in I} \pi_i$, whence $(x, z) \in \rho$ follows. Let $\sigma_i \subseteq P \times P$ denote the complementary half-space of π_i and consider the following equivalence relation:

$$\Theta = \bigcap_{i \in I} (\sigma_i \cap \sigma_i^{-1})$$

on P . Since $\pi_i^{-1} \cap \sigma_i^{-1} = \Delta_P$ for all $i \in I$, we have $\rho \cap \Theta = \Delta_P$ and hence $\rho^{-1} \cap \Theta = \Delta_P$. Now we prove the containments $\Theta \circ \rho \subseteq \rho$ and $\rho \circ \Theta \subseteq \rho$. If $(x, y) \in \Theta$ and $(y, z) \in \rho$ for the elements $x, y, z \in P$ with x, y, z being different, then $(z, y) \in \pi_j$ for some $j \in I$ and $(y, z) \notin \bigcup_{i \in I} \pi_i$. In view of $(x, y) \in \sigma_j \cap \sigma_j^{-1}$, we have $(x, y) \notin \pi_j$ and $(y, x) \notin \pi_j$. Using part (4) in Proposition 2.2, we obtain that $(z, x) \in \pi_j$ and $(x, z) \in \left(\bigcup_{i \in I} \pi_i \right)^{-1}$. Suppose that $(x, z) \in \bigcup_{i \in I} \pi_i$, then $(x, z) \in \pi_k$ follows for some $k \in I$. Since $(x, y) \in \sigma_k \cap \sigma_k^{-1}$ implies that $(x, y) \notin \pi_k$ and $(y, x) \notin \pi_k$, the application of part (3) in Proposition 2.2 yields $(y, z) \in \pi_k$, a contradiction. Thus we have $(x, z) \notin \bigcup_{i \in I} \pi_i$, whence $(x, z) \in \rho$ follows. A similar argument shows that $\rho \circ \Theta \subseteq \rho$.

Fix a linear order μ on P , then $\mu \cap \Theta$ and $\mu^{-1} \cap \Theta$ are partial orders. Using the above properties of ρ and Θ , it is straightforward to see that $\rho \cup (\mu \cap \Theta)$ and $\rho \cup (\mu^{-1} \cap \Theta)$ are also partial orders on P .

Let $\lambda \supseteq \rho \cup (\mu \cap \Theta)$ and $\lambda^* \supseteq \rho \cup (\mu^{-1} \cap \Theta)$ be linear extensions on P and fix an index $i^* \in I$. In view of Proposition 2.5, we can consider the linear orders $R_i = \pi_i[\lambda]$, $i \in I \setminus \{i^*\}$ and $R_{i^*} = \pi_{i^*}[\lambda^*]$ on P (note that $I \setminus \{i^*\}$ is not empty). Since $\pi_i \subseteq R_i$ for all $i \in I$, the inclusion

$$r_\gamma = \bigcap_{i \in I} \pi_i \subseteq \bigcap_{i \in I} R_i$$

is obvious. In order to prove the reverse containment let $(x, y) \notin \bigcap_{i \in I} \pi_i$ for some $x, y \in P$. We have $(x, y) \notin \pi_j$ for some $j \in I$. If $(y, x) \in \bigcup_{i \in I} \pi_i$, then $(y, x) \in \pi_k \subseteq R_k$ and hence $(x, y) \notin R_k$ for some $k \in I$. If $(y, x) \notin \bigcup_{i \in I} \pi_i$, then we distinguish two cases.

First suppose that $(x, y) \in \bigcup_{i \in I} \pi_i$. Then $(y, x) \in \rho \subseteq \lambda \cap \lambda^*$ and the relations $(x, y) \notin \pi_j$, $(y, x) \notin \pi_j$ imply that $(y, x) \in \pi_j[\lambda]$ (or $(y, x) \in \pi_{i^*}[\lambda^*]$ if $j = i^*$), whence $(x, y) \notin R_j$ follows.

Next suppose that $(x, y) \notin \bigcup_{i \in I} \pi_i$. Then $(x, y) \in \Theta$ and the linearity of μ gives that we have either $(y, x) \in \mu \cap \Theta$ or $(y, x) \in \mu^{-1} \cap \Theta$. If $(y, x) \in \mu \cap \Theta \subseteq \lambda$, then $(y, x) \in \pi_i[\lambda]$ and hence $(x, y) \notin \pi_i[\lambda] = R_i$ for all $i \in I \setminus \{i^*\}$. If $(y, x) \in \mu^{-1} \cap \Theta \subseteq \lambda^*$, then $(y, x) \in \pi_{i^*}[\lambda^*]$ and hence $(x, y) \notin \pi_{i^*}[\lambda^*] = R_{i^*}$. \square

Remark 2.14 Another possibility to construct the linear orders R_i in the above proof is the following. Fix a well ordering $<$ on I and for $i \in I \setminus \{i^*\}$ let

$$R_i = \pi_i \cup ((\sigma_i \cap \sigma_i^{-1}) \cap \Lambda) \cup (\Theta \cap \mu),$$

$$R_{i^*} = \pi_{i^*} \cup ((\sigma_{i^*} \cap \sigma_{i^*}^{-1}) \cap \Lambda) \cup (\Theta \cap \mu^{-1}),$$

where $\Lambda = \{(x, y) \mid (y, x) \in \pi_k \text{ and } (x, y) \in \bigcap_{i \in I, i < k} (\sigma_i \cap \sigma_i^{-1}) \text{ for some } k \in I\}$.

In view of Corollaries 2.6 and 2.10, the above Theorem 2.13 yields the following.

Theorem 2.15 *If (A, γ) is a quasiordered set and $\text{hsdim}(A, \gamma) = 1$, then γ is a half-space and*

$\dim(A/(\gamma \cap \gamma^{-1}), r_\gamma) = 1$ if γ has no empty box with more than one element,

$\dim(A/(\gamma \cap \gamma^{-1}), r_\gamma) = 2$ if γ has an empty box with more than one element.

If $\text{hsdim}(A, \gamma) \geq 2$, then we have

$$\dim(A/(\gamma \cap \gamma^{-1}), r_\gamma) = \text{hs dim}(A, \gamma).$$

Theorem 2.16 *If (A, γ) is a partially ordered set and $\text{hsdim}(A, \gamma) = 1$, then γ is a half-space and*

$\dim(A, \gamma) = 1$ if γ is a linear order,

$\dim(A, \gamma) = 2$ if γ is not a linear order.

If $\text{hsdim}(A, \gamma) \geq 2$, then we have

$$\dim(A, \gamma) = \text{hs dim}(A, \gamma).$$

346 3 Direct Product Irreducibility of Half-Space Quasiorders

347 If $(A_i, \gamma_i), i \in I$ is a family of quasiordered sets, then

$$\prod_{i \in I} \gamma_i = \{(\mathbf{a}, \mathbf{b}) \mid \mathbf{a}, \mathbf{b} \in \prod_{i \in I} A_i \text{ and } (\mathbf{a}(i), \mathbf{b}(i)) \in \gamma_i \text{ for all } i \in I\}$$

348 is a quasiorder on the product set $\prod_{i \in I} A_i$ (here \mathbf{a} and \mathbf{b} are functions $I \longrightarrow \bigcup_{i \in I} A_i$ such
349 that $\mathbf{a}(i), \mathbf{b}(i) \in A_i$ for all $i \in I$). We call $(\prod_{i \in I} A_i, \prod_{i \in I} \gamma_i)$ the direct product of the above
350 family. The kernel of the natural surjection

$$\varphi : \prod_{i \in I} A_i \longrightarrow \prod_{i \in I} A_i / (\gamma_i \cap \gamma_i^{-1})$$

351 is $\prod_{i \in I} (\gamma_i \cap \gamma_i^{-1})$, whence we obtain a natural bijection

$$\left(\prod_{i \in I} A_i \right) / \left(\prod_{i \in I} (\gamma_i \cap \gamma_i^{-1}) \right) \longrightarrow \prod_{i \in I} A_i / (\gamma_i \cap \gamma_i^{-1}).$$

352 It is easy to see that

$$\left(\prod_{i \in I} \gamma_i \right) \cap \left(\prod_{i \in I} \gamma_i \right)^{-1} = \prod_{i \in I} (\gamma_i \cap \gamma_i^{-1}) \text{ and } r = \prod_{i \in I} r_{\gamma_i},$$

353 where r is the partial order on $\prod_{i \in I} A_i / (\gamma_i \cap \gamma_i^{-1})$ induced by the quasiorder $\prod_{i \in I} \gamma_i$.

354 The product of non-trivial partial orders is never a linear order. In contrast, the
355 product of two half-spaces can be a half-space again: the four element Boolean lattice
356 M_2 is a product of two-element chains. We show that this is the only possibility to get
357 a non-trivial half-space as a product of quasiorders.

358 **Lemma 3.1** *Let $(A_i, \gamma_i), i \in I$ be a family of quasiordered sets and let $j, k \in I, j \neq k$
359 be indices such that $a_j \neq c_j, (a_j, c_j) \in \gamma_j, (a_j, b_j) \notin \gamma_j$ for some $a_j, b_j, c_j \in A_j$ and $\gamma_k \neq$
360 $A_k \times A_k$ with $|A_k| > 1$. Then $\prod_{i \in I} \gamma_i$ is not a half-space on $\prod_{i \in I} A_i$.*

361 *Proof* Let $\mathbf{u} \in \prod_{i \in I} A_i$ be an arbitrary element and $x_k, y_k \in A_k$ such that $(x_k, y_k) \notin \gamma_k$.

362 Define $\mathbf{a}, \mathbf{b}, \mathbf{c} \in \prod_{i \in I} A_i$ as follows: for an index $i \in I$ let

$$\mathbf{a}(i) = \begin{cases} a_j & \text{if } i = j \\ y_k & \text{if } i = k \\ \mathbf{u}(i) & \text{if } i \in I \setminus \{j, k\} \end{cases}, \quad \mathbf{b}(i) = \begin{cases} b_j & \text{if } i = j \\ x_k & \text{if } i = k \\ \mathbf{u}(i) & \text{if } i \in I \setminus \{j, k\} \end{cases},$$

$$\mathbf{c}(i) = \begin{cases} c_j & \text{if } i = j \\ y_k & \text{if } i = k \\ \mathbf{u}(i) & \text{if } i \in I \setminus \{j, k\} \end{cases}.$$

363 Clearly, $(a_j, b_j) \notin \gamma_j$ implies $(\mathbf{a}, \mathbf{b}) \notin \prod_{i \in I} \gamma_i$ and $(x_k, y_k) \notin \gamma_k$ implies $(\mathbf{b}, \mathbf{a}) \notin \prod_{i \in I} \gamma_i$. Since

364 $(\mathbf{a}, \mathbf{c}) \in \prod_{i \in I} \gamma_i$ and $(x_k, y_k) \notin \gamma_k$ implies $(\mathbf{b}, \mathbf{c}) \notin \prod_{i \in I} \gamma_i$, we can use part (3) in Proposition 2.2

to see that $\prod_{i \in I} \gamma_i$ is not a half-space (we note that $\mathbf{c} \neq \mathbf{a}$ is an immediate consequence 365
of $a_j \neq c_j$). □ 366

Lemma 3.2 *If (A, γ) is a quasiordered set such that there are no elements $a, b, c \in A$ 367
with $a \neq c$, $(a, c) \in \gamma$ and $(a, b) \notin \gamma$, then $\gamma \in \{\Delta_A, A \times A\}$ or $\gamma = (B_1 < B_2)$ is a 368
half-space with a full lower box B_1 (or $|B_1| = 1$) and an empty upper box B_2 . 369*

Proof If $\gamma \notin \{\Delta_A, A \times A\}$ satisfies the above conditions, then for each $a \in A$ we 370
have either $(a, x) \in \gamma$ for all $x \in A$ or $(a, y) \notin \gamma$ for all $y \in A$. Take 371

$$B_1 = \{a \in A \mid (a, x) \in \gamma \text{ for all } x \in A\} \text{ and } B_2 = \{a \in A \mid (a, y) \notin \gamma \text{ for all } y \in A\},$$

then $B_1 \cup B_2 = A$, $B_1 \cap B_2 = \emptyset$ and $\gamma = B_1 \times A = (B_1 \times B_1) \cup (B_1 \times B_2)$ is a half- 372
space, with a full lower box B_1 (or $|B_1| = 1$) and an empty upper box B_2 . Thus we 373
can write $\gamma = (B_1 < B_2)$. □ 374

Lemma 3.3 *Let $\gamma_i = (B_{i1} < B_{i2})$, $1 \leq i \leq 2$ be half-spaces on A_i with full lower boxes 375
 B_{i1} (or $|B_{i1}| = 1$) and empty upper boxes B_{i2} . Then we have the following. 376*

- (1) $\Delta_{A_1 \times A_2} \neq \gamma_1 \times \gamma_2 \neq (A_1 \times A_2) \times (A_1 \times A_2)$ and take $\mathbf{a} = (a_{11}, a_{21})$, $\mathbf{b} =$ 377
 (a_{11}, a_{21}) , $\mathbf{c} = (a_{12}, a_{22})$, where $a_{ij} \in B_{ij}$, $i, j \in \{1, 2\}$ are arbitrary elements. Then 378
 $\mathbf{a} \neq \mathbf{c}$, $(\mathbf{a}, \mathbf{c}) \in \gamma_1 \times \gamma_2$ and $(\mathbf{a}, \mathbf{b}) \notin \gamma_1 \times \gamma_2$. 379
- (2) $\gamma_1 \times \gamma_2$ is a half-space if and only if $|B_{ij}| = 1$ for all $i, j \in \{1, 2\}$. 380

Proof 381

- (1): Obvious. 382
- (2): If $|B_{ij}| = 1$ for all $i, j \in \{1, 2\}$, then it is clear that $A_1 \times A_2$ is a four element set 383
and $\gamma_1 \times \gamma_2$ is a partial order relation on $A_1 \times A_2$ providing a lattice isomorphic 384
to M_2 , which is a half-space as we have already noted. 385

Suppose now, that $|B_{11}| > 1$ and take $a', a'' \in B_{11}$ such that $a' \neq a''$. Let $\mathbf{z} = (a', b)$, 386
 $\mathbf{x} = (a'', b)$ and $\mathbf{y} = (a, c)$, where $a \in B_{12}$, $b \in B_{22}$, $c \in B_{21}$ are arbitrary elements. 387
Since $(\mathbf{x}, \mathbf{y}) \notin \gamma_1 \times \gamma_2$, $(\mathbf{y}, \mathbf{x}) \notin \gamma_1 \times \gamma_2$ and $(\mathbf{x}, \mathbf{z}) \in \gamma_1 \times \gamma_2$, $(\mathbf{y}, \mathbf{z}) \notin \gamma_1 \times \gamma_2$, we can 388
apply part (3) in Proposition 2.2 to derive that $\gamma_1 \times \gamma_2$ is not a half-space. 389

If $|B_{12}| > 1$ then take $a', a'' \in B_{12}$ such that $a' \neq a''$. Let $\mathbf{z} = (a', b)$, $\mathbf{x} = (a', c)$ 390
and $\mathbf{y} = (a'', c)$, where $b \in B_{22}$, $c \in B_{21}$ are arbitrary elements. Since $(\mathbf{x}, \mathbf{y}) \notin \gamma_1 \times$ 391
 γ_2 , $(\mathbf{y}, \mathbf{x}) \notin \gamma_1 \times \gamma_2$ and $(\mathbf{x}, \mathbf{z}) \in \gamma_1 \times \gamma_2$, $(\mathbf{y}, \mathbf{z}) \notin \gamma_1 \times \gamma_2$, we can apply part (3) in 392
Proposition 2.2 to derive that $\gamma_1 \times \gamma_2$ is not a half-space. 393

The cases $|B_{21}| > 1$ and $|B_{22}| > 1$ can be treated analogously. □ 394

Theorem 3.4 *If (A_i, γ_i) , $i \in I$ is a family of non-trivial quasiordered sets (i.e. $\Delta_{A_i} \neq$ 395
 $\gamma_i \neq A_i \times A_i$ for all $i \in I$), then the following are equivalent. 396*

- (1) $\prod_{i \in I} \gamma_i$ is a half-space on $\prod_{i \in I} A_i$. 397
- (2) Either $I = \{1\}$ and γ_1 is a half-space or $I = \{1, 2\}$ and (A_1, γ_1) , (A_2, γ_2) are two- 398
element chains. 399

Proof 400

- (2) \implies (1): It is an immediate consequence of part (2) in Lemma 3.3. 401

(1) \implies (2): It is enough to deal with the case $|I| \geq 2$. Using Lemma 3.1, we obtain that there is no $j \in I$ such that $a_j \neq c_j$, $(a_j, c_j) \in \gamma_j$, $(a_j, b_j) \notin \gamma_j$ for some $a_j, b_j, c_j \in A_j$. In view of Lemma 3.2, each γ_j is a half-space on A_j of the form $\gamma_j = (B_{j1} < B_{j2})$ with a full lower box B_{j1} (or $|B_{j1}| = 1$) and an empty upper box B_{j2} . If $|I| \geq 3$, then we have different indices $i_1, i_2, i_3 \in I$ and

$$\prod_{i \in I} \gamma_i = (\gamma_{i_1} \times \gamma_{i_2}) \times \gamma_{i_3} \times \left(\prod_{i \in I \setminus \{i_1, i_2, i_3\}} \gamma_i \right),$$

where $\gamma_{i_1} \times \gamma_{i_2}$ has the property described in part (1) of Lemma 3.3. Since $\gamma_{i_3} \neq A_{i_3} \times A_{i_3}$ with $|A_{i_3}| > 1$, Lemma 3.1 ensures that our product is not a half-space, a contradiction. Thus $|I| = 2$ and part (2) in Lemma 3.3 gives that (A_1, γ_1) and (A_2, γ_2) are two-element chains (here we assumed $I = \{1, 2\}$). \square

Remark 3.5 If $\gamma_j = \Delta_{A_j}$ and $|A_j| > 1$ for some $j \in I$, then $\prod_{i \in I} \gamma_i$ is disconnected, hence not a non-trivial half-space (because $\prod_{i \in I} \gamma_i = \Delta$ would be the only possibility to get a half-space). If $\gamma_j = A_j \times A_j$ for some $j \in I$, then γ_j has no effect on whether the product $\prod_{i \in I} \gamma_i$ is a half-space (in other words $\prod_{i \in I} \gamma_i$ is a half-space if and only if $\prod_{i \in I \setminus \{j\}} \gamma_i$ is a half-space).

Now, as promised in the introduction, we illustrate the use of half-spaces in a short proof of the following statement.

Theorem 3.5 (Dushnik–Miller) *If (A_i, R_i) , $i \in I$ is a family of non-trivial linearly ordered sets (chains) with $|I| \geq 2$, then*

$$\dim \left(\prod_{i \in I} A_i, \prod_{i \in I} R_i \right) \leq |I|.$$

Proof For an index $j \in I$ let π_j denote the natural $\prod_{i \in I} A_i \rightarrow A_j$ projection. We have

$$\bigcap_{j \in I} \{(\mathbf{a}, \mathbf{b}) \mid \mathbf{a}, \mathbf{b} \in \prod_{i \in I} A_i \text{ and } \mathbf{a}(j) \leq_{R_j} \mathbf{b}(j)\} = \prod_{i \in I} R_i,$$

for the half-spaces

$$\{(\mathbf{a}, \mathbf{b}) \mid \mathbf{a}, \mathbf{b} \in \prod_{i \in I} A_i \text{ and } \mathbf{a}(j) \leq_{R_j} \mathbf{b}(j)\} = \ker(\pi_j) \cup \pi_j^{-1}(R_j), \quad j \in I$$

(see Proposition 2.7). If $\prod_{i \in I} R_i$ is not a half-space, then

$$\dim \left(\prod_{i \in I} A_i, \prod_{i \in I} R_i \right) = \text{hs dim} \left(\prod_{i \in I} A_i, \prod_{i \in I} R_i \right) \leq |I|$$

by Theorem 2.16. If $\prod_{i \in I} R_i$ is a half-space, then $|I| = 2$ and $(\prod_{i \in I} A_i, \prod_{i \in I} R_i)$ is the four element Boolean lattice by Theorem 3.4. \square

Acknowledgement The initial version of this paper was prepared while the first named author was at the Alfred Renyi Institute of Mathematics, Hungarian Academy of Sciences.

References

1. Bonnet, R., Pouzet, M.: Linear extensions of ordered sets. In: Rival, I. (ed.) *Ordered Sets*, Proceedings of the Nato Advanced Study Institute Conference held in Banff, pp. 125–170, August 28–September 12, 1981. Reidel Publishing, Dordrecht (1982)
2. Dushnik, B., Miller, E.W.: Partially ordered sets. *Am. J. Math.* **63**, 600–610 (1941)
3. Foldes, S.: On intervals in relational structures. *Zeitschrift Math. Log. Grund. Math.* **26**, 97–101 (1980)
4. Foldes, S., Radeleczki, S.: On interval decomposition lattices. *Discus. Math., Gen. Algebra Appl.* **24**, 95–114 (2004)
5. Hausdorff, F.: Grundzüge einer Theorie der geordneten Mengen. *Math. Ann.* **65**(4), 435–505 (1908)
6. Körtesi, P., Radeleczki, S., Szilágyi, Sz.: Congruences and isotone maps on partially ordered sets. *Math. Pannonica* **16**(1), 39–55 (2005)
7. Szpilrajn, E.: Sur l'extension de l'ordre partiel. *Fundam. Math.* **16**, 386–389 (1930)
8. Trotter, W.T.: *Combinatorics and Partially Ordered Sets, Dimension Theory*. The Johns Hopkins University Press, Baltimore (1992)
9. van de Vel, M.L.J.: *Theory of Convex Structures*, North-Holland Mathematical Library, 50. North-Holland Publishing, Amsterdam (1993)