

# A note on the Turán number of a Berge odd cycle

Dániel Gerbner

Alfréd Rényi Institute of Mathematics, Hungarian Academy of Sciences

P.O.B. 127, Budapest H-1364, Hungary.

gerbner@renyi.hu

May 25, 2021

## Abstract

In this note we obtain upper bounds on the number of hyperedges in 3-uniform hypergraphs not containing a Berge cycle of given odd length. We improve the bound given by Füredi and Özkahya in 2017. The result follows from a more general theorem. We also obtain some new results for Berge cliques.

**Keywords:** Berge, hypergraph, cycle, Turán number

## 1 Introduction

We say that a hypergraph  $\mathcal{H}$  is a Berge copy of a graph  $F$  (in short:  $\mathcal{H}$  is a Berge- $F$ ) if  $V(F) \subset V(\mathcal{H})$  and there is a bijection  $f : E(F) \rightarrow E(\mathcal{H})$  such that for any  $e \in E(F)$  we have  $e \subset f(e)$ . This definition was introduced by Gerbner and Palmer [11], extending the well-established notion of Berge cycles and paths. Note that there are several non-uniform Berge copies of  $F$ , and a hypergraph  $\mathcal{H}$  is a Berge copy of several graphs. A particular copy of  $F$  defining a Berge- $F$  is called its *core*. Note that there can be multiple cores in a Berge- $F$ .

We denote by  $ex_r(n, \text{Berge-}F)$  the largest number of hyperedges in an  $r$ -uniform Berge- $F$ -free hypergraph on  $n$  vertices. There are several papers dealing with  $ex_r(n, \text{Berge-}C_k)$  (e.g. [8, 14, 15, 16]) or  $ex_r(n, \text{Berge-}F)$  in general (e.g. [9, 10, 11, 12, 20]). For a short survey on this topic see Subsection 5.2.2 in [13].

In this note we consider  $ex_3(n, \text{Berge-}C_k)$ . In the case  $k = 5$ , this was first studied by Bollobás and Győri [2]. They showed  $ex_3(n, \text{Berge-}C_5) \leq \sqrt{2}n^{3/2} + 4.5n$ . This bound was improved to  $(0.254 + o(1))n^{3/2}$  by Ergemlidze, Győri and Methuku [5]. For cycles of any length, Győri and Lemons [15, 16] proved  $ex_r(n, \text{Berge-}C_k) = O(n^{1+1/\lfloor k/2 \rfloor})$ . The constant factors were improved by Jiang and Ma [18], and in the case  $k$  is even by Gerbner, Methuku and Vizer [10]. In the 3-uniform case, Füredi and Özkahya [8] obtained better constant factors (depending on  $k$ ). In the case  $k$  is even, further improvements were obtained by Gerbner, Methuku and Vizer [10] and by Gerbner, Methuku and Palmer [9].

A closely related area is counting triangles in  $C_k$ -free graphs. More generally, let  $ex(n, H, F)$  denote the maximum number of copies of  $H$  in an  $F$ -free graph on  $n$  vertices. After some sporadic results, the systematic study of these problems (often called *generalized Turán problems*) was initiated by Alon and Shikhelman [1]. Their connection to Berge hypergraphs was established by Gerbner and Palmer [12], who proved

$$ex(n, K_r, F) \leq ex_r(n, \text{Berge-}F) \leq ex(n, K_r, F) + ex(n, F)$$

for any  $r$ ,  $n$  and  $F$ .

Counting triangles in  $C_k$ -free graphs and counting hyperedges in Berge- $C_k$ -free 3-uniform hypergraphs was handled together already by Bollobás and Győri [2] for  $C_5$ , and by Füredi and Özkahya [8], who proved  $ex(n, K_3, C_{2k}) \leq \frac{2k-3}{3}ex(n, C_{2k})$  and  $ex_3(n, \text{Berge-}C_{2k}) \leq \frac{2k}{3}ex(n, C_{2k})$ . Their upper bound for  $ex(n, K_3, C_{2k})$  is still the best known bound, but their other upper bound was improved to  $ex_3(n, \text{Berge-}C_{2k}) \leq \frac{2k-3}{3}ex(n, C_{2k})$  by Gerbner, Methuku and Vizer [10] in the case  $k \geq 5$  and by Gerbner, Methuku and Palmer [9] in the case  $k = 3, 4$ .

In the case of forbidden cycles of any odd length, the number of triangles was first studied by Györi and Li [17], who proved<sup>1</sup>  $ex(n, K_3, C_{2k+1}) \leq \frac{(2k-2)(16k-1)}{3} ex(n, C_{2k})$ . It was improved independently by Füredi and Özkahya [8] and by Alon and Shikhelman [1]. The latter had the stronger bound  $ex(n, K_3, C_{2k+1}) \leq \frac{16(k-1)}{3} ex(\lceil n/2 \rceil, C_{2k})$ . In the case  $k = 2$ , the current best bound  $ex(n, K_3, C_5) \leq 0.231975n^{3/2}$  is due to Ergemlidze and Methuku [6].

Füredi and Özkahya [8] obtained the currently best upper bound on the Berge version by showing

$$ex_3(n, \text{Berge-}C_{2k+1}) \leq ex(n, K_3, C_{2k+1}) + 4ex(n, C_{2k}) + 12ex_3^{lin}(n, \text{Berge-}C_{2k+1}), \quad (1.1)$$

where  $ex_r^{lin}(n, \text{Berge-}F)$  denotes the largest number of hyperedges in an  $r$ -uniform Berge- $F$ -free linear hypergraph on  $n$  vertices. Recall that a linear hypergraph is one in which any two hyperedges share at most one vertex.

In this note we improve the bound (1.1). Recall that we have  $ex_3(n, \text{Berge-}C_{2k+1}) \geq ex(n, K_3, C_{2k+1})$ , thus we cannot hope for a huge improvement, especially as  $ex(n, K_3, C_{2k+1})$  might be the largest of the three terms. Indeed, the best upper bound currently known is  $O(n^{1+1/k})$  for all the three terms, but the dependence of the known upper bound in  $k$  is the largest for  $ex(n, K_3, C_{2k+1})$  (we will state these bounds after Theorem 1.2).

Recall that in case of  $C_{2k}$ , the two upper bounds obtained by Füredi and Özkahya [8] were  $ex(n, K_3, C_{2k}) \leq \frac{2k-3}{3} ex(n, C_{2k})$  and  $ex_3(n, \text{Berge-}C_{2k}) \leq \frac{2k}{3} ex(n, C_{2k})$ , and the Berge bound was improved in [10, 9] to match the generalized Turán bound. Our goal would be to do the same here and get rid of the terms  $4ex(n, C_{2k+1}) + 12ex_3^{lin}(n, \text{Berge-}C_{2k+1})$  in (1.1). We cannot achieve that, but we decrease these additional terms. Recall that the currently best bound for the generalized Turán problem is  $ex(n, K_3, C_{2k+1}) \leq \frac{16(k-1)}{3} ex(\lceil n/2 \rceil, C_{2k})$  by Alon and Shikhelman [1]. Our new upper bound on  $ex_3(n, \text{Berge-}C_{2k+1})$  is larger than that bound by  $ex_3^{lin}(n, \text{Berge-}C_{2k+1})$ . We wonder if it is an example of a more general phenomenon and similar bounds could be obtained for other graphs.

The way we use the linearity involves subdividing an edge  $uv$ , i.e. deleting it and adding  $uw$  and  $vw$  for a new vertex  $w$ . Our method uses only the following two properties of  $C_{2k+1}$ : it can be obtained from  $C_{2k}$  by subdividing an edge and deleting a vertex from  $C_{2k+1}$  we obtain a path. In the next theorem we state our result in the most general form.

**Theorem 1.1.** *Let  $F$  be a connected graph obtained from  $F_0$  by subdividing an edge and  $F'$  be obtained from  $F$  by deleting a vertex. Let  $c = c(n)$  be such that  $ex(n, K_{r-1}, F') \leq cn$  for every  $n$ . Then we have*

- (i)  $ex_r(n, \text{Berge-}F) \leq ex(n, K_r, F) + 2^{r-1} ex(n, F_0) + ex_r^{lin}(n, \text{Berge-}F)$ ,
- (ii)  $ex_r(n, \text{Berge-}F) \leq \max\{1, \frac{2c}{r}\} 2^{r-1} ex(n, F_0) + ex_r^{lin}(n, \text{Berge-}F)$ .

In the case  $F = C_{2k+1}$  we have  $F_0 = C_{2k}$  and  $F' = P_{2k}$ , the path on  $2k$  vertices. A theorem of Luo [19] shows  $ex(n, K_{r-1}, P_{2k}) \leq \frac{n}{2k-1} \binom{2k-1}{r-1}$ , but what we need for the 3-uniform case is the Erdős-Gallai theorem [4] showing  $ex(n, P_{2k}) \leq (k-1)n$ . Using this, (ii) of Theorem 1.1 gives  $ex_3(n, \text{Berge-}C_{2k+1}) \leq \frac{8k-8}{3} ex(n, C_{2k}) + ex_3^{lin}(n, \text{Berge-}C_{2k+1})$  if  $k > 2$ . We can improve this a little bit.

**Theorem 1.2.** *If  $k > 2$ , then  $ex_3(n, \text{Berge-}C_{2k+1}) \leq \frac{16k-16}{3} ex(\lceil n/2 \rceil, C_{2k}) + ex_3^{lin}(n, \text{Berge-}C_{2k+1})$*   

$$\leq \left( \frac{1280k-1280}{3} \sqrt{k} \log k \right) \lceil n/2 \rceil^{1+1/k} + 2kn^{1+1/k} + 9kn + \frac{16k-16}{3} 10k^2 \lceil n/2 \rceil.$$

The bound in Theorem 1.2 is currently stronger than the bound given by (i) of Theorem 1.1 for  $F = C_{2k+1}$  and  $r = 3$ . However, an improvement on  $ex(n, K_3, C_{2k+1})$  would immediately improve the bound in (i). Any significant improvement would make (i) stronger than Theorem 1.2 for  $F = C_{2k+1}$ .

The second inequality in Theorem 1.2 follows from known results. Füredi and Özkahya [8] proved  $ex_3^{lin}(n, \text{Berge-}C_{2k+1}) \leq 2kn^{1+1/k} + 9kn$ , and Bukh and Jiang [3] obtained the strongest bound on the Turán number of even cycles by showing  $ex(n, C_{2k}) \leq 80\sqrt{k} \log kn^{1+1/k} + 10k^2n$ . As we do not have good lower bounds on  $ex(n, C_{2k})$ , we cannot be sure that the first term is actually the larger term. However, if  $ex_3^{lin}(n, \text{Berge-}C_{2k+1})$  is the larger term, then our improvement on the upper bound of  $ex_3(n, \text{Berge-}C_{2k+1})$  is more significant, as we changed the constant factor of

<sup>1</sup>We note that the bound is incorrectly stated in their paper [17].

that term from 12 to 1. Obviously we have  $ex_3^{lin}(n, \text{Berge-}C_{2k+1}) \leq ex_3(n, \text{Berge-}C_{2k+1})$ , hence further improvement is impossible here.

We prove Theorem 1.1 by combining the ideas of [8] and [1] with the methods developed in [9, 10]. In the next section we state some lemmas needed for the proof. We give a new proof of a lemma by Gerbner, Methuku and Palmer [9], and we strengthen the lemma a little bit. This strengthens results on  $ex_r(n, \text{Berge-}K_k)$  for some values of  $r$ ,  $k$  and  $n$ . In Section 3 we prove Theorems 1.1 and 1.2.

## 2 Lemmas

We say that a graph  $G$  is red-blue if each of its edges is colored with one of the colors red and blue. For a red-blue graph  $G$ , we denote by  $G_{red}$  the subgraph spanned by the red edges and  $G_{blue}$  the subgraph spanned by the blue edges. For two graphs  $H$  and  $G$  we denote by  $N(H, G)$  the number of subgraphs of  $G$  that are isomorphic to  $H$ . Let  $g_r(G) = |E(G_{red})| + N(K_r, G_{blue})$ .

**Lemma 2.1** (Gerbner, Methuku, Palmer [9]). *For any graph  $F$  and integers  $r$  and  $n$ , there is a red-blue  $F$ -free graph  $G$  on  $n$  vertices, such that  $ex_r(n, \text{Berge-}F) \leq g_r(G)$ .*

Note that an essentially equivalent version was obtained by Füredi, Kostochka and Luo [7]. The proof of Lemma 2.1 relies on a lemma about bipartite graphs (hidden in the proof of Lemma 2 in [9]). If  $M$  is a matching and  $ab$  is an edge in  $M$ , then with a slight abuse of notation we say  $M(a) = b$  and  $M(b) = a$ .

**Lemma 2.2.** *Let  $\Gamma$  be a finite bipartite graph with parts  $A$  and  $B$  and let  $M$  be a largest matching in  $\Gamma$ . Let  $B'$  denote the set of vertices in  $B$  that are incident to  $M$ . Then we can partition  $A$  into  $A_1$  and  $A_2$  and partition  $B'$  into  $B_1$  and  $B_2$  such that for  $a \in A_1$  we have  $M(a) \in B_1$ , and every neighbor of the vertices of  $A_2$  is in  $B_2$ .*

Here we present a proof that is built on the same principle, but is somewhat simpler than the proof found in [9]. Before that, let us recall the well-known notion of alternating paths. Given a bipartite graph  $\Gamma$  and a matching  $M$  in it, a path  $P$  in  $\Gamma$  is called *alternating* if its first edge is not in  $M$ , and then it alternates between edges in  $M$  and edges not in  $M$ , finishing with an edge not in  $M$ . It is well-known and easy to see that deleting the edges of  $P$  from  $M$  and replacing them with the edges of  $P$  that were not in  $M$ , we obtain another matching, that is larger than  $M$ .

*Proof.* First we build a set  $V' \subset V(\Gamma)$  in the following way. Let  $V_0$  be the set of vertices in  $A$  that are not incident to any edges of  $M$ . Then in the first step we add to  $V_0$  the set of vertices in  $B$  that are neighbors of a vertex in  $V_0$ , to obtain  $V_1$ . In the second step we add to  $V_1$  the vertices in  $A$  that are connected to a vertex in  $V_1$  by an edge in  $M$ , to obtain  $V_2$ . Similarly, in the  $i$ th step, if  $i$  is odd we add to  $V_{i-1}$  the set of vertices in  $B$  that are neighbors of a vertex in  $V_{i-1}$ , while if  $i$  is even, we add to  $V_{i-1}$  the vertices in  $A$  that are connected to a vertex in  $V_{i-1}$  by an edge in  $M$  (i.e.  $M(b)$  for some  $b \in B \cap V_{i-1}$ ), to obtain  $V_i$ . After finitely many steps,  $V_i$  does not increase anymore, let  $V'$  be the resulting set of vertices.

We claim that no vertex from  $B \setminus B'$  can be in  $V'$ . Indeed, such a vertex could be reached by an alternating path from a vertex in  $A$  that is not incident to  $M$ , thus  $M$  is not a largest matching, a contradiction.

Then let  $A_2 = A \cap V'$ ,  $A_1 = A \setminus A_2$ ,  $B_2 = B' \cap V'$  and  $B_1 = B' \setminus B_2$ . A vertex in  $A_2$  cannot be connected to a vertex  $v$  not in  $B_2$ , as  $v$  could be added to  $V'$  then. Similarly, for a vertex  $u \in A_1$ ,  $M(u)$  has to be in  $B_1$ , otherwise  $M(u)$  is in  $B_2$  and then  $u$  can be added to  $V'$ . □

Let us briefly describe how we can apply this lemma to obtain Lemma 2.1. We take a Berge- $F$ -free  $r$ -uniform hypergraph  $\mathcal{H}$  on  $n$  vertices. Let  $A$  be the set of hyperedges in  $\mathcal{H}$  and  $B$  be the set of sub-edges of these hyperedges (by edge and sub-edge we always mean an edge of size two, i.e. a pair of vertices). We connect  $a \in A$  to  $b \in B$  if  $a \supset b$ . Let  $\Gamma$  denote this auxiliary bipartite graph. Let  $M$  be an arbitrary largest matching and  $B'$  be the vertices of  $B$  incident to the edges in  $M$ . It is easy to see that the elements of  $B'$  form an  $F$ -free graph which we call  $G$ . Indeed, otherwise  $M$  defines the bijection between a copy of  $F$  and hyperedges in  $\mathcal{H}$  to form a Berge- $F$ .

Now we apply Lemma 2.2 to  $\Gamma$  and  $M$ . We define a red-blue coloring of  $G$  by taking the edges of  $G$  in  $B_1$  to be the red edges, and the edges of  $G$  in  $B_2$  to be the blue edges. We have

$|\mathcal{H}| = |A_1| + |A_2| = |B_1| + |A_2| = |E(G_{red})| + |A_2|$ . As hyperedges in  $A_2$  have all their neighbors in  $B_2$ , they each contain a blue  $K_r$ , which is distinct from the other blue  $r$ -cliques obtained this way, showing  $|A_2| \leq N(K_r, G_{blue})$ .

Let us remark here that Lemma 2.2 also gives some information on the structure of  $G$ . If there is  $a \in A_1$  that has a neighbor  $b \in B \setminus B'$ , then we could obtain another matching  $M'$  by changing the neighbor of  $a$  to  $b$ , i.e.  $M'(a) = b$  and if  $a' \neq a$ , then  $M'(a') = M(a')$ . Then  $B'$  is replaced by  $B'' = B' \setminus \{M(a)\} \cup \{b\}$ . In this case the same partition of  $A$  into  $A_1$  and  $A_2$ , and the partition of  $B''$  into  $B_2$  and  $B'' \setminus B_2$  satisfies Lemma 2.2. This means for  $G$  that we can delete the (red) edge  $M(a)$  and replace it with the edge  $b$ , to obtain another  $F$ -free graph.

If on the other hand the vertices in  $A_1$  have all their neighbors in  $B'$ , then we could recolor the red edges to blue. Therefore, in  $G$  we can delete an edge and add another edge so that the resulting graph is still  $F$ -free. Let  $\alpha = \alpha_{F,n}$  be the largest value of  $g_r(G')$ , where  $G'$  is an  $n$ -vertex  $F$ -free blue-red graph. Assume that each  $n$ -vertex  $F$ -free blue-red graph  $G'$  with  $g_r(G') = \alpha$  is not monoblue and we cannot delete an edge and add another edge to  $G'$  so that the resulting graph is still  $F$ -free. Then by the above,  $G$  cannot be one of these graphs, thus  $ex_r(n, \text{Berge-}F) \leq g_r(G) < \alpha$ . This is usually a negligible improvement, as we often do not even know the order of magnitude.

However, if  $F = K_k$ , Gerbner, Methuku and Palmer [9] proved that  $\alpha_{K_k,n} = \max\{g_r(T_B(n, k-1)), g_r(T_R(n, k-1))\}$ , where  $T_B(n, k-1)$  is the monoblue Turán graph  $T(n, k-1)$  and  $T_R(n, k-1)$  is the monored Turán graph  $T(n, k-1)$ . We mention without going into the details that their proof also shows that for any other graphs  $G$  we have  $g_r(G) < \alpha_{K_k,n}$ . As we cannot delete an edge from  $T(n, k-1)$  and add another edge to obtain a  $K_k$ -free graphs, we do have an improvement. For example, if  $r = 4$  and  $k = 5$ , then the result in [9] determines  $ex_4(n, \text{Berge-}K_k)$  for  $n \geq 11$ . For  $n = 10$ ,  $T(10, 4)$  has 36 copies of  $K_4$  and 37 edges. Therefore, (as  $ex(n, K_r, F)$  is a lower bound on  $ex_r(n, \text{Berge-}F)$ ), we have  $36 \leq ex_4(n, \text{Berge-}K_k) \leq 37$ . With our new observation, we know  $ex_4(n, \text{Berge-}K_k) = 36$ .

### 3 Proof of Theorems 1.1 and 1.2

Let  $\mathcal{H}$  be a Berge- $F$ -free  $r$ -graph on  $n$  vertices. We say that an edge  $uv$  with  $u, v \in V(\mathcal{H})$  is  $t$ -heavy if  $u, v$  are contained together in exactly  $t$  hyperedges. First we will build a linear subhypergraph  $\mathcal{H}_1$  in a greedy way: if we can find a hyperedge  $H$  that does not share an edge with any hyperedge in  $\mathcal{H}_1$ , we add  $H$  to  $\mathcal{H}_1$ , and then repeat this procedure. By definition,  $\mathcal{H}_1$  is linear. Let  $\mathcal{H}_2$  consist of the remaining hyperedges. Note that  $|\mathcal{H}| = |\mathcal{H}_1| + |\mathcal{H}_2| \leq ex_r^{lin}(n, \text{Berge-}F) + |\mathcal{H}_2|$ , and the remainder of the proof is for proving the needed upper bound on  $|\mathcal{H}_2|$ .

We build an auxiliary bipartite graph  $\Gamma$  in the usual way: let  $A$  be the set of hyperedges in  $\mathcal{H}_2$  and  $B$  be the set of sub-edges of these hyperedges. We connect  $a \in A$  to  $b \in B$  if  $a \supset b$ . We will let  $M$  be a largest matching in  $\Gamma$ , however, we do not choose  $M$  arbitrarily. Let  $M_0$  be an arbitrary largest matching in  $\Gamma$ . Let  $B'$  be the set of vertices in  $B$  that are incident to some edge of  $M_0$  and  $A_0$  denote the set of vertices in  $A$  that are incident to some edge of  $M_0$ . Now a hyperedge  $a \in A_0$  contains a sub-edge  $M_0(a)$ , at least one sub-edge  $b_0$  shared with a hyperedge in  $\mathcal{H}_1$ , maybe some sub-edges that are matched to some other  $a' \in A$ , and maybe some other sub-edges  $b \in B \setminus B'$ . We have the option to replace in  $M_0$  the edge between  $a$  and  $M_0(a)$  with any of the edges of  $\Gamma$  between  $a$  and an unused sub-edge of  $a$ , to obtain another largest matching. We will build a largest matching  $M$ , that contains the same vertices ( $A_0$ ) from  $A$  as  $M_0$ .

For  $a \in A_0$ , we pick  $M(a)$  to be one of the sub-edges  $b \in B$  of  $a$  (potentially we let  $M(a) = M_0(a)$ ) in the following way:  $M(a)$  should share exactly one vertex with  $b_0$  (where  $b_0$  is a sub-edge that is also a sub-edge of a hyperedge in  $\mathcal{H}_1$ ) if possible. We go through the hyperedges greedily; as long as there is a hyperedge  $a \in A_0$  such that  $M_0(a)$  can be changed in this way, we execute the change (it is possible that  $M_0(a)$  cannot be changed originally, but later a sub-edge of  $a$  that is  $M_0(a')$  becomes free to use, when  $M(a')$  is chosen to be different from  $M_0(a')$ ). This process finishes after finitely many (at most  $|A_0|$ ) steps, as we change  $M_0(a)$  to  $M(a)$  at most once for every  $a \in A_0$ . After this, we rename the unchanged  $M_0(a)$  to  $M(a)$ .

The resulting matching  $M$  has the following property: for every  $a \in A_0$ ,  $a$  shares a sub-edge  $b_0$  with a hyperedge in  $\mathcal{H}_1$ , such that that either  $M(a)$  shares exactly one vertex with  $b_0$ , or all the sub-edges of  $a$  sharing exactly one vertex with  $b_0$  are  $M(a')$  for some  $a' \in A_0$ .

Now we can apply Lemma 2.2 to  $\Gamma$  and  $M$  to obtain  $A_1, A_2, B_1, B_2$ . Let us call the elements of  $B_1$  red edges and the elements of  $B_2$  blue edges. Let  $G$  be the graph consisting of all the red and blue edges. Then  $G$  is obviously  $F$ -free.

Let us now take a random partition of  $V(\mathcal{H})$  into  $V_1$  and  $V_2$ . For every  $a \in A_0$ , we look at  $b = M(a)$ . If the two vertices of  $b$  are in one part, and all the other vertices of  $a$  are in the other part, we keep  $a$ , otherwise we delete it. Let  $A^*$  denote the set of elements in  $A$  that are not deleted (note that elements in  $A \setminus A_0$  are never deleted, thus are in  $A^*$ ). Let  $G'$  be the graph consisting of the elements of  $B'$  that are connected by an edge in  $M$  to an element of  $A^*$ . Then  $G'$  is obviously  $F$ -free, as it is a subgraph of  $G$ .

**Claim 3.1.**  $G'$  is  $F_0$ -free, where  $F_0$  is any graph for which  $F$  can be obtained from  $F_0$  by subdividing an edge of  $F_0$ .

*Proof.* Let us assume we are given a copy  $Q$  of  $F_0$  in  $G'$  such that  $uv$  is the edge that needs to be subdivided to obtain  $F$ . Observe that there is no edge between  $V_1$  and  $V_2$  in  $G'$ , thus  $Q$  is in one of them, say  $V_1$ . Let  $w$  be a vertex of  $M(uv)$  with  $u \neq w \neq v$ , then  $w \in V_2$ , thus  $w$  is not in  $Q$ .

We say that a hyperedge  $H$  in  $\mathcal{H}$  is *good* if  $H$  contains  $u$  and  $w$  for some  $w \in M(uv) \setminus \{u, v\}$  and  $H$  is not  $M(e)$  for any edge  $e$  of  $Q$ . If there is a good hyperedge, then we build a Berge- $F$  with the following core: we subdivide  $uv$  with  $w$ . For each edge  $e$  of this core we assign  $M(e)$  except for  $uw$  (where we assign  $H$ ) and  $vw$  (where we assign  $M(uv)$ ). This way we obtain a Berge- $F$ , a contradiction.

$M(uv)$  shares at least one sub-edge with a hyperedge  $H \in \mathcal{H}_1$ . If the sub-edge shares exactly one vertex with  $uv$ , then  $H$  is good and we are done. Thus every sub-edge of  $M(uv)$  shared with a hyperedge in  $\mathcal{H}_1$  has to contain none or both of  $u$  and  $v$ . In both cases, when we tried to change  $M_0(M(uv))$  when constructing  $M$ , we failed, because all such edges are matched to some other hyperedges of  $\mathcal{H}_2$ . In particular,  $uw$  is  $M(a)$  for some  $a \in A_0$  and for some  $w \in M(u, v) \setminus \{u, v\}$ . Observe that  $w$  is in  $V_2$ , thus  $M(a)$  has vertices from both parts  $V_1$  and  $V_2$ , hence  $a$  cannot be in  $A^*$  by the definition of  $A^*$ . This implies  $a$  is good, finishing the proof.  $\square$

The above claim implies  $G'$  has at most  $ex(n, F_0)$  edges. For an arbitrary  $a \in A$ , the probability that  $a$  is in  $A^*$  is at least  $1/2^{r-1}$ . Let  $S$  be any subset of  $A$ , then we have that the expected value of the number of hyperedges in  $A^* \cap S$  is at least  $|S|/2^{r-1}$ , thus there is a partition with  $|A^* \cap S| \geq |S|/2^{r-1}$ .

There are  $|B_1| = |A_1|$  red edges in  $G$ , and there is a random partition where at least  $|A_1|/2^{r-1}$  elements of  $A_1$  are undeleted, hence there are at least  $|A_1|/2^{r-1}$  red edges in  $G'$ . This implies  $|A_1|/2^{r-1} \leq ex(n, F_0)$ . Hence there are at most  $2^{r-1}ex(n, F_0)$  red edges altogether. For the total number of edges in  $G$  we can use the same argument: there is a random partition where at least  $|A_0|/2^{r-1}$  hyperedges in  $A_0$  are undeleted, thus for the  $G'$  defined by that partition, we have  $|A_0| = |E(G)| \leq 2^{r-1}|E(G')| \leq 2^{r-1}ex(n, F_0)$ .

Observe that we have  $|\mathcal{H}_2| = |A_1| + |A_2| \leq |A_1| + N(K_r, G_{blue}) \leq |A_1| + ex(n, K_r, F)$ , hence we are done with the proof of (i).

Note that  $G$  is not necessarily  $F_0$ -free, but it is  $F$ -free. Let  $m$  be the number of blue edges in  $G$ , then  $G$  has at most  $2^{r-1}ex(n, F_0) - m$  red edges. An argument of Gerbner, Methuku and Vizer [10] bounds the number of  $r$ -cliques in  $F$ -free graphs with the given number of vertices and edges. For sake of completeness, we include the argument here.

Let  $d(v)$  be the degree of  $v$  in  $G_{blue}$ . Obviously the neighborhood of every vertex in  $G_{blue}$  is  $F'$ -free. An  $F'$ -free graph on  $d(v)$  vertices contains at most  $ex(d(v), K_{r-1}, F') \leq cd(v)$  copies of  $K_{r-1}$ . Thus  $v$  is contained in at most  $cd(v)$  copies of  $K_r$  in  $G_{blue}$ . If we sum, for each vertex, the number of  $K_r$ 's containing a vertex, then each  $K_r$  is counted  $r$  times. On the other hand as  $\sum_{v \in V(G_{blue})} d(v) = 2|E(G_{blue})| = 2m$ , we have  $\sum_{v \in V(G_{blue})} cd(v) = 2cm$ . This gives that the number of blue  $K_r$ 's is at most  $2cm/r$ . Thus we have

$$g_r(G) \leq 2^{r-1}ex(n, F_0) - m + 2cm/r \leq \max \left\{ 1, \frac{2c}{r} \right\} (2^{r-1}ex(n, F_0) - m + m) = \max \left\{ 1, \frac{2c}{r} \right\} 2^{r-1}ex(n, F_0).$$

The above inequality, together with Lemma 2.1 implies that  $|\mathcal{H}_2| \leq \max \left\{ 1, \frac{2c}{r} \right\} 2^{r-1}ex(n, F_0)$ , finishing the proof of (ii).

Now we show how to obtain the small improvement needed to prove Theorem 1.2. It is based on the proof of the upper bound on  $ex(n, K_3, C_{2k+1})$  in [1]. If  $n$  is odd, replace it by  $n + 1$ . As



the stated upper bound is the same in both cases, obvious monotonicity conditions show we can do this. Thus we can assume  $n$  is even. When we take the random partition into  $V_1$  and  $V_2$ , first we take a random partition into  $n/2$  sets  $U_1, \dots, U_{n/2}$  of size 2, and then randomly put one vertex into  $V_1$  and the other into  $V_2$ . The obtained graph  $G'$  will be  $C_{2k}$ -free, and it is divided into two components, hence it has at most  $ex(|V_1|, C_{2k}) + ex(|V_2|, C_{2k})$  edges. The way we chose  $V_1$  ensures the above sum is  $2ex(\lceil n/2 \rceil, C_{2k})$ . Then we can go through every step of the remaining part of the proof to obtain the result we need, if for an arbitrary  $a \in A$ , the probability that  $a$  is in  $A^*$  is still at least  $1/2^{r-1} = 1/4$ . We will separate into cases according to the intersection of  $a$  with the parts  $U_i$ . In case the three vertices of  $a$  are in three different  $U_i$ 's, the probability is  $1/4$ . In case  $a$  contains  $U_i$  for some  $i$ , there are two cases. If  $M(a) = U_i$ , then the probability is 0, otherwise it is  $1/2$ . As  $M(a) = U_i$  happens with probability  $1/3$  (having the condition that  $a$  contains  $U_i$ ), for every  $i$  we have that the probability of  $a$  being in  $A^*$  if  $a$  contains  $U_i$  is  $\frac{2}{3} \cdot \frac{1}{2} \geq 1/4$ .

This gives the first inequality of Theorem 1.2. As we have mentioned after the statement, the second inequality follows from earlier results, stated there.

**Acknowledgements:** Research supported by the National Research, Development and Innovation Office - NKFIH under the grants SNN 129364, KH 130371 and K 116769 and by the János Bolyai Research Fellowship of the Hungarian Academy of Sciences.

## References

- [1] N. Alon and C. Shikhelman, Many  $T$  copies in  $H$ -free graphs, *J. Comb. Theory, Ser. B*, **121** (2016), 146–172.
- [2] B. Bollobás and E. Győri, Pentagons vs. triangles, *Discrete Math.*, **308**(19) (2018), 4332–4336.
- [3] B. Bukh and Z. Jiang, A bound on the number of edges in graphs without an even cycle, *Comb. Probab. Comput.* **26**(1) (2017), 1–15. Erratum-ibid. **26**(6), 952–953. For the version incorporating erratum see [arXiv:1403.1601](https://arxiv.org/abs/1403.1601)
- [4] P. Erdős and T. Gallai, On maximal paths and circuits of graphs, *Acta Math. Acad. Sci. Hung.*, **10** (1959), 337–356.
- [5] B. Ergemlidze, E. Győri and A. Methuku, 3-uniform hypergraphs without a cycle of length five, *Electronic J. Comb.*, **27**(2) (2020), P2.16.
- [6] B. Ergemlidze and A. Methuku, Triangles in  $C_5$ -free graphs and Hypergraphs of Girth Six, *arXiv:1811.11873* (2018).
- [7] Z. Füredi, A. Kostochka and R. Luo, Avoiding long Berge cycles, *J. Comb. Theory, Ser. B*, **137** (2019), 55–64.
- [8] Z. Füredi and L. Özkahya, On 3-uniform hypergraphs without a cycle of a given length, *Discrete Appl. Math.*, **216** (2017), 582–588.
- [9] D. Gerbner, A. Methuku and C. Palmer, General lemmas for Berge-Turán hypergraph problems, *Eur. J. Comb.*, **86** (2020), 103082.
- [10] D. Gerbner, A. Methuku and M. Vizer, Asymptotics for the Turán number of Berge- $K_{2,t}$ , *J. Comb. Theory, Ser. B*, **137** (2019), 264–290.
- [11] D. Gerbner and C. Palmer, Extremal Results for Berge Hypergraphs, *SIAM J. Discrete Math.*, **31**(4) (2017), 2314–2327.
- [12] D. Gerbner and C. Palmer, Counting copies of a fixed subgraph in  $F$ -free graphs, *Eur. J. Comb.* **82** (2019), 103001.
- [13] D. Gerbner and B. Patkós, Extremal Finite Set Theory, 1st Edition, CRC Press, 2018.
- [14] E. Győri, Triangle-free hypergraphs, *Comb. Probab. Comput.*, **15**(1-2) (2006) 185–191.
- [15] E. Győri and N. Lemons, 3-uniform hypergraphs avoiding a given odd cycle, *Combinatorica* **32** (2012), 187–203.

- [16] E. Győri and N. Lemons, Hypergraphs with no cycle of a given length, *Comb. Probab. Comput.*, **21**(1-2) (2012), 193–201.
- [17] E. Győri and H. Li, The maximum number of triangles in  $C_{2k+1}$ -free graphs, *Comb. Probab. Comput.*, **21**(1-2) (2012), 187–191.
- [18] T. Jiang and J. Ma, Cycles of given lengths in hypergraphs, *J. Comb. Theory, Ser. B* **133** (2018), 54–77.
- [19] R. Luo, The maximum number of cliques in graphs without long cycles, *J. Comb. Theory, Ser. B*, **128** (2018), 219–226.
- [20] C. Palmer, M. Tait, C. Timmons and A.Z. Wagner, Turán numbers for Berge-hypergraphs and related extremal problems, *Discrete Math.*, **342**(6) (2019), 1553–1563.