



# An Almost Optimal Bound on the Number of Intersections of Two Simple Polygons

Eyal Ackerman<sup>1</sup> · Balázs Keszegh<sup>2,3</sup> · Günter Rote<sup>4</sup>

Received: 16 August 2020 / Revised: 31 July 2022 / Accepted: 31 July 2022

© The Author(s) 2022

## Abstract

What is the maximum number of intersections of the boundaries of a simple  $m$ -gon and a simple  $n$ -gon? This is a basic question in combinatorial geometry, and the answer is easy if at least one of  $m$  and  $n$  is even: If both  $m$  and  $n$  are even, then every pair of sides may cross and so the answer is  $mn$ . If exactly one polygon, say the  $n$ -gon, has an odd number of sides, it can intersect each side of the  $m$ -gon polygon at most  $n - 1$  times; hence there are at most  $mn - m$  intersections. It is not hard to construct examples that meet these bounds. If both  $m$  and  $n$  are odd, the best known construction has  $mn - (m + n) + 3$  intersections, and it is conjectured that this is the maximum. However, the best known upper bound is only  $mn - (m + \lceil n/6 \rceil)$ , for  $m \geq n$ . We prove a new upper bound of  $mn - (m + n) + C$  for some constant  $C$ , which is optimal apart from the value of  $C$ .

**Keywords** Simple polygon · Ramsey theory · Combinatorial geometry

**Mathematics Subject Classification** 52C45

---

Editor in Charge: Kenneth Clarkson

---

Eyal Ackerman  
ackerman@sci.haifa.ac.il

Balázs Keszegh  
keszegh@renyi.hu

Günter Rote  
rote@inf.fu-berlin.de

<sup>1</sup> Department of Mathematics, Physics, and Computer Science, University of Haifa at Oranim, 36006 Tivon, Israel

<sup>2</sup> Alfréd Rényi Institute of Mathematics, 1053 Budapest, Hungary

<sup>3</sup> MTA-ELTE Lendület Combinatorial Geometry Research Group, ELTE Eötvös Loránd University, Budapest, Hungary

<sup>4</sup> Department of Computer Science, Freie Universität Berlin, Takustr. 9, 14195 Berlin, Germany

## 1 Introduction

To determine the union of two or more geometric objects in the plane is one of the basic computational geometric problems. In strong relation to that, determining the maximum complexity of the union of two or more geometric objects is a basic extremal geometric problem. We study this problem when the two objects are simple polygons.

Let  $P$  and  $Q$  be two simple polygons with  $m$  and  $n$  sides, respectively, where  $m, n \geq 3$ . We are interested in the maximum number of intersection points of the boundaries of  $P$  and  $Q$ , hence we assume that this number is finite, that is, there are no sides of  $P$  and  $Q$  that overlap.

This problem was first studied in 1993 by Dillencourt et al. [3]. The cases when  $m$  or  $n$  is even are solved there. If  $m$  and  $n$  are both even, then every pair of sides may cross and so the answer is  $mn$ . Figure 1a shows one of many ways to achieve this number. If one polygon, say  $Q$ , has an odd number  $n$  of sides, no line segment  $s$  can be intersected  $n$  times by  $Q$ , because otherwise each side of  $Q$  would have to flip from one side of  $s$  to the other side. Thus, each side of the  $m$ -gon  $P$  is intersected at most  $n - 1$  times, for a total of at most  $mn - m$  intersections. It is easy to see that this bound is tight when  $P$  has an even number of sides, see Fig. 1b.

When both  $m$  and  $n$  are odd, the situation is more difficult; the bound that is obtained by the above argument remains at  $mn - \max\{m, n\}$ , because the set of  $m$  intersections that are necessarily “missing” due to the odd parity of  $n$  might conceivably overlap with the  $n$  intersections that are “missing” due to the odd parity of  $m$ . However, the best known family of examples gives only  $mn - (m + n) + 3 = (m - 1)(n - 1) + 2$  intersection points, see Fig. 1c.


**Conjecture 1** *Let  $P$  and  $Q$  be simple polygons with  $m$  and  $n$  sides, respectively, such that  $m, n \geq 3$  are odd numbers. If they intersect in a finite number of points, then there are at most  $mn - (m + n) + 3$  intersection points between sides of  $P$  and sides of  $Q$ .*

In [3] an unrecoverable error appears in a claimed proof of Conjecture 1. Another attempted proof [8] also turned out to have a fault. The only correct improvement over the trivial upper bound is an upper bound of  $mn - (m + \lceil n/6 \rceil)$  for  $m \geq n$ , due to Černý et al. [2]. We will briefly discuss their proof in Sect. 2. We improve the upper bound to  $mn - (m + n) + O(1)$ , which is optimal apart from an additional constant:

**Theorem 1** *There is an absolute constant  $C$  such that the following holds. Suppose that  $P$  and  $Q$  are simple polygons with  $m$  and  $n$  sides, respectively, such that  $m$  and  $n$  are odd numbers. If they intersect in a finite number of points, then there are at least  $m + n - C$  pairs of a side of  $P$  and a side of  $Q$  that do not intersect. Hence, there are at most  $mn - (m + n) + C$  intersections.*

The value of the constant  $C$  that we obtain in our proof is around  $2^{2^{67}}$ . We did not make a large effort to optimize this value, and obviously, there is ample space for improvement.

A preliminary version of this paper was presented at the *36th International Symposium on Computational Geometry* (SoCG 2020) in June 2020 [1].



**Fig. 1** **a** Optimal construction for  $m = n = 8$ , with  $8 \times 8 = 64$  intersections. **b** Optimal construction for  $m = 8, n = 7$ , with  $8 \times 6 = 48$  intersections. **c** Lower-bound construction for  $m = 9, n = 7$ . There are  $8 \times 6 + 2 = 50$  intersections

## 2 Overview of the Proof

First we establish the crucial statement that the odd parity of  $m$  and  $n$  allows us to *associate* to any two consecutive sides of one polygon a pair of consecutive sides of the other polygon with a restricted intersection pattern among the four involved sides (Lemma 5 and Fig. 5). This is the only place where we use the fact that  $P$  and  $Q$  are polygons, rather than merely two sets of pairwise disjoint segments, and that each of them has an odd number of sides.

A simple observation (Observation 2) relates the bound on  $C$  in Theorem 1 to the number of connected components of the bipartite “disjointness graph” between the polygon sides of  $P$  and  $Q$ . Our goal is therefore to show that there are few connected components.

We proceed to consider *two* pairs of associated pairs of sides (four consecutive pairs with eight sides in total). Unless they form a special structure, they cannot belong to four different connected components (Lemma 7). (Four is the maximum number of components that they could conceivably have.) The proof involves a case analysis with a moderate number of cases. This structural statement allows us to reduce the bound on the number of components by a constant factor, and thereby, we can already improve the best previous result on the number of intersections (Proposition 9 in Sect. 6).

Finally, to get a constant bound on the number of components, our strategy is to use Ramsey-theoretic arguments like the Erdős–Szekeres Theorem on caps and cups or the pigeonhole principle (see Sect. 7) in order to impose additional structure on the configurations that we have to analyze. This is the place in the argument where we give up control over the constant  $C$  in exchange for useful properties that allow us to derive a contradiction. This eventually boils down again to a moderate number of cases (Sect. 8.2).

By contrast, the proof of the bound  $mn - (m + \lceil n/6 \rceil)$  for  $m \geq n$  by Černý et al. proceeds in a more local manner. The core of their argument [2, Lemma 3], which is proved by case analysis, is that it is impossible to have six consecutive sides of one polygon together with six distinct sides of the other polygon forming a perfect matching in the disjointness graph. This statement is used to bound the number of components of the disjointness graph. (Lemma 8 below uses a similar argument.)

### 3 General Assumptions and Notations

Let  $P$  and  $Q$  be two simple polygons with sides  $p_0, p_1, \dots, p_{m-1}$  and  $q_0, q_1, \dots, q_{n-1}$ . We assume that  $m \geq 3$  and  $n \geq 3$  are odd numbers. Indices are taken modulo  $m$  or  $n$ , respectively. We consider the sides  $p_i$  and  $q_j$  as closed line segments. The condition that the polygon  $P$  is simple means that its edges are pairwise disjoint except for the unavoidable common endpoints between *consecutive* sides  $p_i$  and  $p_{i+1}$ . Throughout this paper, unless stated otherwise, we regard a polygon as a piecewise linear closed curve, and we disregard the region that it encloses. Thus, by intersections between  $P$  and  $Q$ , we mean intersection points between the polygon *boundaries*.

*General Position.* The basic assumption that is made throughout the paper is that  $P$  and  $Q$  have only finitely many intersection points, i.e., there are no overlapping edges. In addition, we will assume from now on that no vertex of one polygon lies on the other polygon. This assumption will be justified in Sect. 9 by a non-trivial perturbation. Thus, every intersection point between  $P$  and  $Q$  is an interior point of two polygon sides.

Furthermore, we assume for convenience that no three vertices of  $P$  and  $Q$  combined lie on a line, and no two sides of  $P$  and  $Q$  combined are parallel. Indeed, given that no vertex of one polygon lies on the other polygon, this can be achieved easily by another small perturbation of the sides without changing the intersection pattern.

*The Disjointness Graph.* As in [2], our basic tool of analysis is the *disjointness graph* of  $P$  and  $Q$ , which we denote by  $G^D = (V^D, E^D)$ . (Its original name in [2] is *non-intersection graph*.) It is a bipartite graph with node set  $V^D = \{p_0, p_1, \dots, p_{m-1}\} \cup \{q_0, q_1, \dots, q_{n-1}\}$  and edge set  $E^D = \{(p_i, q_j) \mid p_i \cap q_j = \emptyset\}$ . (Since we are interested in the situation where almost all pairs of edges intersect, the disjointness graph is more useful than its more commonly used complement, the intersection graph.) Our goal is to bound from above the number of connected components of  $G^D$ :

**Observation 2** *If  $G^D$  has at most  $C$  connected components, then  $G^D$  has at least  $m + n - C$  edges. Thus, there are at least  $m + n - C$  pairs of a side of  $P$  and a side of  $Q$  that do not intersect, and there are at most  $mn - (m + n) + C$  crossings between  $P$  and  $Q$ .*

For a polygon side  $s$  of  $P$  or  $Q$ ,  $CC(s)$  denotes the connected component of the disjointness graph  $G^D$  to which  $s$  belongs.

*Geometric Notions.* Let  $s$  and  $s'$  be two line segments. We denote by  $\ell(s)$  the line through  $s$  and by  $I(s, s')$  the intersection of  $\ell(s)$  and  $\ell(s')$ , see Fig. 4. We say that  $s$  and  $s'$  are *avoiding* if neither of them contains  $I(s, s')$ . (This requirement is stronger than just disjointness.) If  $s$  and  $s'$  are avoiding or share an endpoint, we denote by  $\mathbf{r}_{s'}(s)$  the ray from  $I(s, s')$  to infinity that contains  $s$ , and by  $\mathbf{r}_s(s')$  the ray from  $I(s, s')$  to infinity that contains  $s'$ . Moreover, we denote by  $\text{Cone}(s, s')$  the convex cone with apex  $I(s, s')$  between these two rays.

**Observation 3** *If a segment  $s''$  that does not go through  $I(s, s')$  has one of its endpoints in the interior of  $\text{Cone}(s, s')$ , then  $s''$  cannot intersect both  $\mathbf{r}_{s'}(s)$  and  $\mathbf{r}_s(s')$ . In particular, it cannot intersect both  $s$  and  $s'$ .*




Fig. 2 The edge-labeled multigraph  $G_0$  in Proposition 4




Fig. 3 The unfolded graph  $G'_0$

### 4 Associated Pairs of Consecutive Sides


We begin with the following seemingly unrelated claim concerning a specific small edge-labeled multigraph. Let  $G_0 = (V_0, E_0)$  be the undirected multigraph shown in Fig. 2. It has four nodes  $V_0 = \{I, II, III, IV\}$  and five edges  $E_0 = \{e_1 = \{II, IV\}, e_2 = \{I, IV\}, e_3 = \{I, II\}, e_4 = \{I, III\}, e_5 = \{I, III\}\}$ . Every edge  $e_i \in E_0$  has a label  $L(e_i) \in \{a, b, *\}$  as follows:  $L(e_1) = *$ ,  $L(e_2) = L(e_4) = a$ ,  $L(e_3) = L(e_5) = b$ .

**Proposition 4** *If  $W$  is a closed walk in  $G_0$  of odd length, then  $W$  contains two cyclically consecutive edges of labels  $a$  and  $b$ .*

**Proof** Suppose for contradiction that  $W$  does not contain two consecutive edges of labels  $a$  and  $b$ . Since  $W$  cannot switch between the  $a$ -edges and the  $b$ -edges in I or III, we can split I (resp., III) into two nodes  $I_a$  and  $I_b$  (resp.,  $III_a$  and  $III_b$ ) such that every  $a$ -labeled edge that is incident to I (resp., III) in  $G_0$  becomes incident to  $I_a$  (resp.,  $III_a$ ) and every  $b$ -labeled edge that is incident to I (resp., III) in  $G_0$  becomes incident to  $I_b$  (resp.,  $III_b$ ). In the resulting graph  $G'_0$ , which is shown in Fig. 3, we can find a closed walk  $W'$  that corresponds to  $W$  and that uses the edges with the same name as  $W$ . Since  $G'_0$  is a path, every closed walk has even length. Thus,  $W$  cannot have odd length.  $\square$

**Lemma 5** *Let  $p_a$  and  $p_b$  be two sides of  $P$  that are either consecutive or avoiding such that  $CC(p_a) \neq CC(p_b)$ . Then there are two consecutive sides  $q_i, q_{i\pm 1}$  of  $Q$  such that  $(p_a, q_i), (p_b, q_{i\pm 1}) \in E^D$  and  $(p_a, q_{i\pm 1}), (p_b, q_i) \notin E^D$ . Furthermore,  $I(p_a, p_b) \in \text{Cone}(q_i, q_{i\pm 1})$  or  $I(q_i, q_{i\pm 1}) \in \text{Cone}(p_a, p_b)$ .*

The sign ‘ $\pm$ ’ is needed since we do not know which of the consecutive sides intersects  $p_i$  and is disjoint from  $p_{i+1}$ .



**Fig. 4** How an odd polygon  $Q$  can intersect two segments. The segments  $p_a$  and  $p_b$  are avoiding, whereas  $s$  and  $s'$  are disjoint but non-avoiding. In this situation, we say that  $s$  *stabs*  $s'$

**Proof** We may assume without loss of generality that  $I(p_a, p_b)$  is the origin,  $p_a$  lies on the positive  $x$ -axis and the interior of  $p_b$  is above the  $x$ -axis. The lines  $\ell(p_a)$  and  $\ell(p_b)$  partition the plane into four convex cones (“quadrants”). Denote them in counterclockwise order by I, II, III, IV, starting with  $I = \text{Cone}(p_a, p_b)$ , see Fig. 4.


Every side of  $Q$  must intersect  $p_a$  or  $p_b$  (maybe both), since  $\text{CC}(p_a) \neq \text{CC}(p_b)$ . One can now check that traversing the sides of  $Q$  in order generates a closed walk  $W$  in the graph  $G_0$  of Fig. 2. For example, a side of  $Q$  that we traverse from its endpoint in I to its endpoint in III and that intersects  $p_a$  corresponds to traversing the edge  $e_4 = \{I, III\}$  from I to III, whose label is  $L(e_4) = a$ . We do not care which of  $p_a$  and  $p_b$  are crossed when we move between II and IV.

It follows from Proposition 4 that  $Q$  has two consecutive sides  $q_i, q_{i\pm 1}$  such that  $q_i$  intersects  $p_b$  and does not intersect  $p_a$ , while  $q_{i\pm 1}$  intersects  $p_a$  and does not intersect  $p_b$ . Hence,  $(p_a, q_i), (p_b, q_{i\pm 1}) \in E^D$  and  $(p_a, q_{i\pm 1}), (p_b, q_i) \notin E^D$ . Furthermore,  $I(q_i, q_{i\pm 1})$  must be either in I or III as these are the only nodes in  $G_0$  that are incident both to an edge labeled  $a$  and an edge labeled  $b$ . In the latter case  $I(p_a, p_b) \in \text{Cone}(q_i, q_{i\pm 1})$ , and in the former case  $I(q_i, q_{i\pm 1}) \in \text{Cone}(p_a, p_b)$ .  $\square$


Let  $p_i, p_{i+1}$  be two sides of  $P$  such that  $\text{CC}(p_i) \neq \text{CC}(p_{i+1})$ . Then by Lemma 5 there are sides  $q_j, q_{j\pm 1}$  of  $Q$  such that  $(p_i, q_j), (p_{i+1}, q_{j\pm 1}) \in E^D$ . We say that such a pair  $q_j, q_{j\pm 1}$  is *associated* to  $p_i, p_{i+1}$ . By Lemma 5 we have  $I(q_j, q_{j\pm 1}) \in \text{Cone}(p_i, p_{i+1})$  or  $I(p_i, p_{i+1}) \in \text{Cone}(q_j, q_{j\pm 1})$ . If the first condition holds we say that  $p_i, p_{i+1}$  is *hooking* and  $q_j, q_{j\pm 1}$  is *hooked*, see Fig. 5. In the second case we say that  $p_i, p_{i+1}$  is *hooked* and  $q_j, q_{j\pm 1}$  is *hooking*. Note that it is possible that a pair of consecutive sides is associated with several pairs and that it is both hooking and hooked (with respect to two different pairs from the other polygon or even with respect to a single pair, as in Fig. 5c).

**Observation 6** (Axis Property) *If the pair  $p_i, p_{i+1}$  and the pair  $q_j, q_{j\pm 1}$  are associated such that  $(p_i, q_j), (p_{i+1}, q_{j\pm 1}) \in E^D$ , then the line through  $I(p_i, p_{i+1})$  and  $I(q_j, q_{j\pm 1})$  separates  $p_i$  and  $q_{j\pm 1}$  on the one side from  $p_{i+1}$  and  $q_j$  on the other side.*

We call this line the *axis* of the associated pairs. In our figures it appears as a dotted line when it is shown.



**Fig. 5** Hooking and hooked pairs of consecutive sides. **a** The pair  $p_i, p_{i+1}$  is hooking and the associated pair  $q_j, q_{j\pm 1}$  is hooked. **b** Vice versa. **c** Both pairs are both hooking and hooked



**Fig. 6** The pair  $p_i, p_{i+1}$  is hooking with respect to the pair  $q_{i'}, q_{i'\pm 1}$ , and  $p_j, p_{j+1}$  is hooked with respect to  $q_{j'}, q_{j'\pm 1}$

### 5 The Principal Structure Lemma About Pairs of Associated Pairs

**Lemma 7** Let  $p_i, p_{i+1}, p_j, p_{j+1}$  be two pairs of consecutive sides of  $P$  that belong to four different connected components of  $G^D$ . Then it is impossible that both  $p_i, p_{i+1}$  and  $p_j, p_{j+1}$  are hooked or that both pairs are hooking.

Figure 6 shows a scenario with four different components, together with the associated pairs of  $Q$ . The combinatorial structure of such a configuration is unique up to relabeling.

**Proof** Suppose by contradiction that both pairs  $p_i, p_{i+1}$  and  $p_j, p_{j+1}$  are hooking or both of these pairs are hooked. Let  $q_{i'}, q_{i'\pm 1}$  and  $q_{j'}, q_{j'\pm 1}$  be associated (hooked or hooking) pairs of  $p_i, p_{i+1}$  and  $p_j, p_{j+1}$ , respectively, such that  $(p_i, q_{i'}), (p_{i+1}, q_{i'\pm 1}) \in E^D$  and  $(p_j, q_{j'}), (p_{j+1}, q_{j'\pm 1}) \in E^D$ . Recall that the existence of such pairs follows from Lemma 5.

For better readability, we rename  $p_i, p_{i+1}$  and  $q_{i'}, q_{i'\pm 1}$  as  $a, b$  and  $A, B$ , and we rename  $p_j, p_{j+1}$  and  $q_{j'}, q_{j'\pm 1}$  as  $a', b'$  and  $A', B'$ . The small letters denote sides of  $P$  and the capital letters denote sides of  $Q$ . In the new notation,  $a, b$  are consecutive sides of  $P$  with an associated pair  $A, B$  of consecutive sides of  $Q$ , and  $a', b'$  are two other consecutive sides of  $P$  with an associated pair  $A', B'$  of consecutive sides of  $Q$ . The disjointness graph  $G^D$  contains the edges  $(a, A), (b, B), (a', A'), (b', B')$ . Since




Fig. 7 Normalizing the position of  $a, b, A', B'$

$a, b, a', b'$  belong to different connected components of  $G^D$ , it follows that the nodes  $A, B, A', B'$ , to which they are connected, belong to the same four different connected components. There can be no more edges among these eight nodes, and they induce a matching in  $G^D$ . One can remember as a rule that every side of  $P$  intersects every side of  $Q$  among the eight involved sides, except when their names differ only in their capitalization.

Suppose first that both  $a, b$  and  $a', b'$  are hooking. Hence,  $I(A, B) \in \text{Cone}(a, b)$  and  $I(A', B') \in \text{Cone}(a', b')$ . Since each of  $A'$  and  $B'$  intersects each of  $a$  and  $b$  they must lie as in Fig. 7a. To facilitate the future discussion, we will now normalize the positions of these four sides.

We first ensure that the intersection  $I(A', b)$  is directly adjacent to the two polygon vertices  $I(a, b)$  and  $I(A', B')$  in the arrangement of the four sides, as shown in Fig. 7b. This can be achieved by swapping the labels  $a, A$  with the labels  $b, B$  if necessary, and by independently swapping the labels  $a', A'$  with  $b', B'$  if necessary. Our assumptions are invariant under these swaps.

By an affine transformation we may finally assume that  $I(A', b)$  is the origin;  $b$  lies on the  $x$ -axis and is directed to the right; and  $A'$  lies on the  $y$ -axis and is directed upwards. Then  $a$  has a positive slope and its interior is in the upper half-plane, and  $B'$  has a positive slope and its interior is to the right of the  $y$ -axis, see Fig. 7c.

The arrangement of the lines through  $a, b, A', B'$  has 11 faces, some of which are marked as  $F_1, \dots, F_6$  in Fig. 7. Our current assumption is that both  $a, b$  and  $a', b'$  are hooking: The hooking of  $a, b$  means that  $I(A, B) \in \text{Cone}(a, b) = F_1 \cup F_2 \cup F_3$ . By the Axis Property (Observation 6), the line through  $I(A', B')$  and  $I(a', b')$  must separate  $A'$  from  $B'$ . Therefore, the vertex  $I(a', b')$  can lie only in  $F_2 \cup F_4 \cup F_5 \cup F_6$ . Thus, based on the faces that contain  $I(A, B)$  and  $I(a', b')$ , there are 12 cases to consider. Some of these cases are symmetric, and all can be easily dismissed, as follows.

In the figures, the four sides  $a', b', A', B'$ , which are associated to the second associated pair are dashed. All dashed sides of one polygon must intersect all solid sides of the other polygon.

1.  $I(A, B) \in F_1$  and  $I(a', b') \in F_2$ , see Fig. 8 (symmetric to  $I(A, B) \in F_2$  and  $I(a', b') \in F_4$ ). Let  $r_a$  (resp.,  $r_b$ ) be the ray on  $\ell(a)$  (resp.,  $\ell(b)$ ) that goes from the right endpoint of  $a$  (resp.,  $b$ ) to the right. Since  $a'$  is not allowed to cross  $b$ , the






Fig. 8 Case 1:  $I(A, B) \in F_1, I(a', b') \in F_2$




Fig. 9 Case 2:  $I(A, B) \in F_1, I(a', b') \in F_4$

only way for  $a'$  to intersect  $A$  is by crossing  $r_b$ . Similarly, in order to intersect  $B$ ,  $a'$  has to cross  $r_a$ . However, it cannot intersect both  $r_a$  and  $r_b$ , by Observation 3. Since we did not use the assumption that  $A, B$  are hooked, the analysis holds for the symmetric case 6,  $I(A, B) \in F_2$  and  $I(a', b') \in F_4$ , as well.

2.  $I(A, B) \in F_1$  and  $I(a', b') \in F_4$ , see Fig. 9. Since  $a'$  is not allowed to cross  $b$ , the only way for  $a'$  to intersect  $B$  is by crossing  $r_b$ . However, in this case  $a'$  cannot intersect  $A$ .
3.  $I(A, B) \in F_1$  and  $I(a', b') \in F_5$ , see Fig. 10 (symmetric to  $I(A, B) \in F_3$  and  $I(a', b') \in F_4$ ). Both  $a'$  and  $b'$  must intersect  $A$ , and they have to go below the line  $\ell(b)$  to do so. However,  $a'$  can only cross  $\ell(b)$  to the right of  $b$ , and  $b'$  can only cross  $\ell(b)$  to the left of  $b$ , and therefore they cross  $A$  from different sides. This is impossible, because  $a'$  and  $b'$  start from the same point.
4.  $I(A, B) \in F_1$  and  $I(a', b') \in F_6$ . If one of the polygon sides  $a'$  and  $b'$  has an endpoint in  $F_4$  (see Fig. 11a), then this side cannot intersect  $B$ . So assume otherwise, see Fig. 11b. The side  $a'$  intersects  $B'$  and is disjoint from  $A'$ , while  $b'$  is disjoint from  $B'$  and intersects  $A'$ . (Due to space limitation some line segments are drawn schematically as curves.) Thus, each of  $a'$  and  $b'$  has an endpoint in





Fig. 10 Case 3:  $I(A, B) \in F_1$  and  $I(a', b') \in F_5$



(a) At least one of the sides  $a'$  and  $b'$  has an endpoint in  $F_4$ .

(b) None of the sides  $a'$  and  $b'$  has an endpoint in  $F_4$ .

Fig. 11 Case 4:  $I(A, B) \in F_1$  (or  $I(A, B) \in F_2$ , which is similar) and  $I(a', b') \in F_6$

$F_2 \cup F_5$ . But then  $I(A, B) \in \text{Cone}(a', b')$  and it follows from Observation 3 that neither  $A$  nor  $B$  can intersect both  $a'$  and  $b'$ .

5.  $I(A, B) \in F_2$  and  $I(a', b') \in F_2$ , see Fig. 12. Since  $a', b'$  is hooking,  $I(A', B') \in \text{Cone}(a', b')$ , and the line segments  $a', b', A', b, B'$  enclose a convex pentagon. The polygon side  $A$  must intersect  $b, a'$ , and  $b'$ , but it is restricted to  $F_2 \cup F_4$ . It follows that  $A$  must intersect three sides of the pentagon, which is impossible. (This is in fact the only place where we need the assumption that  $a', b'$  is hooking.)
6.  $I(A, B) \in F_2$  and  $I(a', b') \in F_4$ . This is symmetric to case 1.
7.  $I(A, B) \in F_2$  and  $I(a', b') \in F_5$ , see Fig. 13 (symmetric to  $I(A, B) \in F_3$  and  $I(a', b') \in F_2$ ). Then  $A$  is restricted to  $F_2 \cup F_4$ , while  $a'$  and  $b'$  do not intersect  $F_2$  and  $F_4$ . Therefore  $A$  can intersect neither  $a'$  nor  $b'$ .
8.  $I(A, B) \in F_2$  and  $I(a', b') \in F_6$ . This case is very similar to case 4, where  $I(A, B) \in F_1$  and  $I(a', b') \in F_6$ , see Fig. 11. If one of the polygon sides  $a'$




Fig. 12 Case 5:  $I(A, B) \in F_2, I(a', b') \in F_2$

and  $b'$  has an endpoint in  $F_4$ , then it cannot intersect  $B$ . Otherwise,  $I(A, B) \in \text{Cone}(a', b')$  and therefore, neither  $A$  nor  $B$  can intersect both  $a'$  and  $b'$ .

- 9.  $I(A, B) \in F_3$  and  $I(a', b') \in F_2$ . This is symmetric to case 7.
- 10.  $I(A, B) \in F_3$  and  $I(a', b') \in F_4$ . This is symmetric to case 3.
- 11.  $I(A, B) \in F_3$  and  $I(a', b') \in F_5$ , see Fig. 14. Then the intersection of  $b'$  and  $A$  can lie only in the lower left quadrant. It follows that the triangle whose vertices are  $I(a', b')$ ,  $I(a', A)$ , and  $I(A, b')$  contains  $a$  and does not contain  $I(A, B)$ . This in turn implies that  $B$  cannot intersect both  $b'$  and  $a$ , without intersecting  $B'$ .
- 12.  $I(A, B) \in F_3$  and  $I(a', b') \in F_6$ , see Fig. 15. As in case 4, we may assume that neither  $a'$  nor  $b'$  has an endpoint in  $F_4$ , since then this side could not intersect  $B$ . We may also assume that  $I(A, B) \notin \text{Cone}(a', b')$  for otherwise neither  $A$  nor  $B$  intersects both of  $a'$  and  $b'$ , according to Observation 3. If  $a'$  has an endpoint in  $F_2$ , then it cannot intersect  $B$  (see Fig. 15a). Otherwise, if  $a'$  has an endpoint in  $F_5$ , then  $B$  cannot intersect  $b'$  (Fig. 15b).

We have finished the case that  $a, b$  and  $a', b'$  are hooking. Suppose now that  $a, b$  and  $a', b'$  are hooked with respect to  $A, B$  and  $A', B'$ , respectively. Then  $A, B$  is hooking with respect to  $a, b$  and  $A', B'$  is hooking with respect to  $a', b'$ . Recall that  $A, B, A'$ , and  $B'$  belong to four different connected components. Hence, this case can be handled as above, after exchanging the capital letters with the small letters (i.e., exchanging  $P$  and  $Q$ ). □

### 6 A Weaker Bound

The principal structure lemma is already powerful enough to get an improvement over the previous best bound:

**Lemma 8**  $G^D$  has at most  $(n + 5)/2$  connected components.




Fig. 13 Case 7:  $I(A, B) \in F_2, I(a', b') \in F_5$




Fig. 14 Case 11:  $I(A, B) \in F_3$  and  $I(a', b') \in F_5$

**Proof** Partition the sides  $q_0, q_1, \dots, q_{n-1}$  of  $Q$  into  $(n-1)/2$  disjoint pairs  $q_{2i}, q_{2i+1}$ , discarding the last side  $q_{n-1}$ . Let  $H_+$  denote the subset of these pairs that are hooked. Suppose first that this set contains some pair  $q_{2i_0}, q_{2i_0+1}$  of sides that are in two different connected components. Combining  $q_{2i_0}, q_{2i_0+1}$  with any of the remaining pairs  $q_{2i}, q_{2i+1}$  of  $H_+$ , Lemma 7 tells us that the sides  $q_{2i}$  and  $q_{2i+1}$  must either belong to the same connected component, or one of them must belong to  $CC(q_{2i_0})$  or  $CC(q_{2i_0+1})$ . In other words, each remaining pair contributes at most one “new” connected component, and it follows that the sides in  $H_+$  belong to at most  $|H_+| + 1$  connected components. This conclusion holds also in the case that  $H_+$  contains no pair  $q_{2i_0}, q_{2i_0+1}$  of sides that are in different connected components.

The same argument works for the complementary subset  $H_-$  of pairs that are not hooked, but hooking. Along with  $CC(q_{n-1})$  there are at most  $(|H_+|+1)+(|H_-|+1)+1 = (n-1)/2 + 3 = (n+5)/2$  components.  $\square$

Together with Observation 2, this already improves the previous bound of  $mn - (m + \lceil n/6 \rceil)$  for a large range of parameters, namely when  $m \geq n \geq 11$ :




Fig. 15 Case 12:  $I(A, B) \in F_3$  and  $I(a', b') \in F_6$

**Proposition 9** Let  $P$  and  $Q$  be simple polygons with  $m$  and  $n$  sides, respectively, such that  $m$  and  $n$  are odd and  $m \geq n \geq 3$ . Then there are at most  $mn - (m + (n - 5)/2)$  intersection points between  $P$  and  $Q$ .

### 7 Ramsey-Theoretic Tools

We recall some classic results. A tournament is a directed graph that contains between every pair of nodes  $x, y$  either the arc  $(x, y)$  or the arc  $(y, x)$  but not both. A tournament is *transitive* if for every three nodes  $x, y, z$  the existence of the arcs  $(x, y)$  and  $(y, z)$  implies the existence of the arc  $(x, z)$ . Equivalently, the nodes can be ordered on a line such that all arcs are in the same direction. The following is easy to prove by induction.

**Lemma 10** (Erdős and Moser [5]) Every tournament on a node set  $V$  contains a transitive sub-tournament on  $1 + \lfloor \log_2 |V| \rfloor$  nodes.

**Proof** Choose  $v \in V$  arbitrarily, and let  $N \subseteq V - \{v\}$  with  $|N| \geq (|V| - 1)/2$  be the set of in-neighbors of  $v$  or the set of out-neighbors of  $v$ , whichever is larger. Then  $v$  together with a transitive sub-tournament of  $N$  gives a transitive sub-tournament of size one larger. □

A set of points  $p_1, p_2, \dots, p_r$  in the plane sorted by  $x$ -coordinates (and with distinct  $x$ -coordinates) forms an  $r$ -cup (resp.,  $r$ -cap) if  $p_i$  is below (resp., above) the line through  $p_{i-1}$  and  $p_{i+1}$  for every  $i$  with  $1 < i < r$ .

**Theorem 11** (Erdős–Szekeres Theorem for caps and cups in point sets [6]) For any two integers  $r \geq 2$  and  $s \geq 2$ , the value  $ES(r, s) := \binom{r+s-4}{r-2}$  fulfills the following statement:

Suppose that  $P$  is a set of  $ES(r, s) + 1$  points in the plane with distinct  $x$ -coordinates such that no three points of  $P$  lie on a line. Then  $P$  contains an  $r$ -cup or an  $s$ -cap.

Moreover,  $ES(r, s)$  is the smallest value that fulfills the statement.

A similar statement holds for lines by the standard point-line duality. A set of lines  $\ell_1, \ell_2, \dots, \ell_r$  sorted by slope forms an  $r$ -cup (resp.,  $r$ -cap) if  $\ell_{i-1}$  and  $\ell_{i+1}$  intersect below (resp., above)  $\ell_i$  for every  $1 < i < r$ .

**Theorem 12** (Erdős–Szekeres Theorem for lines) *For the numbers  $ES(r, s)$  from Theorem 11, the following statement holds for any two integers  $r \geq 2$  and  $s \geq 2$ :*

*If  $L$  is a set of  $ES(r, s) + 1$  non-vertical lines in the plane no two of which are parallel and no three of which intersect at a common point, then  $L$  contains an  $r$ -cup or an  $s$ -cap.*

**Theorem 13** (Erdős–Szekeres Theorem for monotone subsequences [6]) *For any integer  $r \geq 0$ , a sequence of  $r^2 + 1$  distinct numbers contains either an increasing subsequence of length  $r + 1$  or a decreasing subsequence of length  $r + 1$ .*

## 8 Proof of Theorem 1

### 8.1 Imposing More Structure on the Examples

Going back to the proof of Theorem 1, recall that in light of Observation 2 it is enough to prove that  $G^D$ , the disjointness graph of  $P$  and  $Q$ , has at most constantly many connected components.

We will use the following constants:  $C_6 := 6$ ;  $C_5 := (C_6)^2 + 1 = 37$ ;  $C_4 := ES(C_5, C_5) + 1 = \binom{70}{35} + 1 = 112,186,277,816,662,845,433 < 2^{67}$ ;  $C_3 := 2^{C_4 - 1}$ ;  $C_2 := C_3 + 5$ ;  $C_1 := 8C_2$ ;  $C := C_1 - 1 < 2^{2^{67}}$ .

We claim that  $G^D$  has at most  $C$  connected components. Suppose that  $G^D$  has at least  $C_1 = C + 1$  connected components, numbered as  $1, 2, \dots, C_1$ . For each connected component  $j$ , we find two consecutive sides  $q_{i_j}, q_{i_j+1}$  of  $Q$  such that  $CC(q_{i_j}) = j$  and  $CC(q_{i_j+1}) \neq j$ . We call  $q_{i_j}$  the *primary* side and  $q_{i_j+1}$  the *companion* side of the pair. We take these  $C_1$  consecutive pairs in their cyclic order along  $Q$  and remove every second pair. This ensures that the remaining  $C_1/2$  pairs are disjoint in the sense that no side of  $Q$  belongs to two different pairs.

We apply Lemma 5 to each of the remaining  $C_1/2$  pairs  $q_{i_j}, q_{i_j+1}$  and find an associated pair  $p_{k_j}, p_{k_j \pm 1}$  such that  $(q_{i_j}, p_{k_j}), (q_{i_j+1}, p_{k_j \pm 1}) \in E^D$ . Therefore,  $CC(q_{i_j}) = CC(p_{k_j})$  and  $CC(q_{i_j+1}) = CC(p_{k_j \pm 1}) \neq CC(q_{i_j})$ . Again, we call  $p_{k_j}$  the *primary* side and  $p_{k_j \pm 1}$  the *companion* side. As before, we delete half of the pairs  $p_{k_j}, p_{k_j \pm 1}$  in cyclic order along  $P$ , along with their associated pairs from  $Q$ , and thus we ensure that the remaining  $C_1/4$  pairs are disjoint also on  $P$ .

At least  $C_1/8$  of the remaining pairs  $q_{i_j}, q_{i_j+1}$  are hooking or at least  $C_1/8$  of them are hooked. We may assume that at least  $C_2 = C_1/8$  of the pairs  $q_{i_j}, q_{i_j+1}$  are hooking with respect to their associated pair,  $p_{k_j}, p_{k_j \pm 1}$ , for otherwise,  $p_{k_j}, p_{k_j \pm 1}$  is

hooking with respect to  $q_{i_j}, q_{i_{j+1}}$  and we may switch the roles of  $P$  and  $Q$ . Let us denote by  $Q_2$  the set of  $C_2$  hooking consecutive pairs  $(q_{i_j}, q_{i_{j\pm 1}})$  at which we have arrived. (Because of the potential switch, we have to denote the companion side by  $q_{i_{j\pm 1}}$  instead of  $q_{i_{j+1}}$  from now on.)

By construction, all  $C_2$  primary sides  $q_{i_j}$  of these pairs belong to distinct components. We now argue that all  $C_2$  adjacent companion sides  $q_{i_{j\pm 1}}$  with at most one exception lie in the same connected component, provided that  $C_2 \geq 4$ .

We model the problem by a graph whose nodes are the connected components of  $G^D$ . For each of the  $C_2$  pairs  $q_{i_j}, q_{i_{j\pm 1}}$ , we insert an edge between  $CC(q_{i_j})$  and  $CC(q_{i_{j\pm 1}})$ . The result is a multigraph with  $C_2$  edges and without loops. Two disjoint edges would represent two consecutive pairs of the form  $(q_{i_j}, q_{i_{j\pm 1}})$  whose four sides are in four distinct connected components, but this is a contradiction to Lemma 7. Thus, the graph has no two disjoint edges, and such graphs are easily classified: they are the triangle (cycle on three vertices) and the star graphs  $K_{1r}$ , possibly with multiple edges. Overall, the graph involves at least  $C_2 \geq 4$  distinct connected components  $CC(q_{i_j})$ , and therefore the triangle graph is excluded. Let  $v$  be the central vertex of the star. There can be at most one  $j$  with  $CC(q_{i_j}) = v$ , and we discard it. All other sides  $q_{i_j}$  have  $CC(q_{i_j}) \neq v$ , and therefore  $CC(q_{i_{j\pm 1}})$  must be the other endpoint of the edge, that is,  $v$ .

In summary, we have found  $C_2 - 1$  adjacent pairs  $q_{i_j}, q_{i_{j\pm 1}}$  with the following properties.

- The primary sides  $q_{i_j}$  belong to  $C_2 - 1$  distinct components.
- All companion sides  $q_{i_{j\pm 1}}$  belong to the same component, distinct from the other  $C_2 - 1$  components.
- All  $2C_2 - 2$  sides of the pairs  $q_{i_j}, q_{i_{j\pm 1}}$  are distinct.
- Each  $q_{i_j}, q_{i_{j\pm 1}}$  is hooking with respect to an associated pair  $p_{k_j}, p_{k_{j\pm 1}}$ .
- All  $2C_2 - 2$  sides of the pairs  $p_{k_j}, p_{k_{j\pm 1}}$  are distinct.


Let us denote by  $Q'_2$  the set of  $C_2 - 1$  sides  $q_{i_j}$ .

**Proposition 14** *There are no six distinct sides  $q_a, q_b, q_c, q_d, q_e, q_f$  among the  $C_2 - 1$  sides  $q_{i_j} \in Q'_2$  such that  $q_a, q_b$  are avoiding or consecutive,  $q_c, q_d$  are avoiding or consecutive, and  $q_e, q_f$  are avoiding or consecutive.*

**Proof** Suppose for contradiction that six such sides exist. It follows from Lemma 5 that there are two consecutive sides  $p_{a'}$  and  $p_{b'}$  of  $P$  such that  $CC(p_{a'}) = CC(q_a)$  and  $CC(p_{b'}) = CC(q_b)$ .

Similarly, we find a pair of consecutive sides  $p_{c'}$  and  $p_{d'}$  of  $P$  such that  $CC(p_{c'}) = CC(q_c)$  and  $CC(p_{d'}) = CC(q_d)$ , and the same story for  $e$  and  $f$ . By the pigeonhole principle, two of the three consecutive pairs  $(p_{a'}, p_{b'}), (p_{c'}, p_{d'}), (p_{e'}, p_{f'})$  are hooking or two of them are hooked. This contradicts Lemma 7. □

Define a complete graph whose nodes are the  $C_2 - 1$  sides  $q_{i_j} \in Q'_2$ , and color an edge  $(q_{i_j}, q_{i_k})$  red if  $q_{i_j}$  and  $q_{i_k}$  are avoiding or consecutive and blue otherwise. Proposition 14 says that this graph contains no red matching of size three. This means that we can get rid of all red edges by removing at most four nodes. To see this, pick any red edge and remove its two nodes from the graph. If any red edge remains, remove its



**Fig. 16** The seven sides  $a_0, a_1, \dots, a_6$ . The lines  $\ell_0, \dots, \ell_6$  form a 7-cap. The side  $a_0$  could be between any two consecutive intersection points of  $\ell_0$  with the rest of the lines

two nodes. Then all red edges are gone, because otherwise we would find a matching with three red edges.

We conclude that there is a blue clique of size  $C_3 = C_2 - 5$ , i.e., there is a set  $Q_3 \subset Q'_2$  of  $C_3$  polygon sides among the  $C_2 - 1$  sides  $q_{ij} \in Q'_2$  that are pairwise non-avoiding and disjoint, i.e., they do not share a common endpoint.

Our next goal is to find a subset of seven segments in  $Q_3$  that are arranged as in Fig. 16. To define this precisely, we say for two segments  $s$  and  $s'$  that  $s$  *stabs*  $s'$  if  $I(s, s') \in s'$ , see Fig. 4. Among any two non-avoiding and non-consecutive sides  $s$  and  $s'$ , either  $s$  stabs  $s'$  or  $s'$  stabs  $s$ , but not both. Define a tournament  $T$  whose nodes are the  $C_3$  sides  $q_{ij} \in Q_3$ , and the arc between each pair of nodes is oriented towards the stabbed side. It follows from Lemma 10 that  $T$  has a transitive sub-tournament of size  $1 + \lfloor \log_2 C_3 \rfloor = C_4$ .

Furthermore, since  $C_4 = ES(C_5, C_5) + 1$ , it follows from Theorem 12 that there is a subset of  $C_5$  sides such that the lines through them form a  $C_5$ -cup or a  $C_5$ -cap. By a vertical reflection if needed, we may assume that they form a  $C_5$ -cap.


We now reorder these  $C_5$  sides  $q_{ij}$  of  $Q$  in stabbing order, according to the transitive sub-tournament mentioned above. By the Erdős–Szekeres Theorem on monotone subsequences (Theorem 13), there is a subsequence of size  $C_6 + 1 = \sqrt{C_5 - 1} + 1 = 7$  such that their slopes form a monotone sequence. By a horizontal reflection if needed, we may assume that they have decreasing slopes.

We rename these seven segments to  $a_0, a_1, \dots, a_6$ , and we denote the line  $\ell(a_i)$  by  $\ell_i$ , see Fig. 16. We have achieved the following properties:

- The lines  $\ell_0, \dots, \ell_6$  form a 7-cap, with decreasing slopes in this order.
- The segments  $a_i$  are pairwise disjoint and non-avoiding.
- $a_i$  stabs  $a_j$  for every  $i < j$ .

These properties allow  $a_0$  to lie between any two consecutive intersections on  $\ell_0$ . There is no such flexibility for the other sides: Every side  $a_j$  is stabbed by every preceding side  $a_i$ . For  $1 \leq i < j$ ,  $a_i$  cannot stab  $a_j$  from the right, because then  $a_0$  would not be able to stab  $a_i$ . Hence, the arrangement of the sides  $a_1, \dots, a_6$  must be exactly as





**Fig. 17** The subgraph of  $G^D$  induced on two pairs of consecutive sides  $a_i, b_i$  and  $a_j, b_j$  of  $P$  and their associated partner pairs  $A_i, B_i$  and  $A_j, B_j$  of  $Q$ . Parts of  $P$  and  $Q$  are shown to indicate consecutive sides. The dashed edges may or may not be present

shown in Fig. 16, in the sense that the order of endpoints and intersection points along each line  $\ell_i$  is fixed. We will ignore  $a_0$  from now on.

### 8.2 Finalizing the Analysis

Recall that every  $a_i$  is the primary side of two consecutive sides  $a_i, b_i$  of  $Q$  that are hooking with respect to an associated pair  $A_i, B_i$  of consecutive sides of  $P$ . The sides  $a_i$  and  $A_i$  are the primary sides and  $b_i$  and  $B_i$  are the companion sides. All these  $4 \times 6$  sides are distinct, and they intersect as follows:  $a_i$  intersects  $B_i$  and is disjoint from  $A_i$ ;  $b_i$  intersects  $A_i$  and is disjoint from  $B_i$ ; and  $I(A_i, B_i) \in \text{Cone}(a_i, b_i)$ .

Figure 17 summarizes the intersection pattern among these sides. A side  $A_i$  must intersect every side  $a_j$  with  $j \neq i$  and every side  $b_j$  since  $\text{CC}(A_i) = \text{CC}(a_i) \neq \text{CC}(a_j)$  and  $\text{CC}(A_i) = \text{CC}(a_i) \neq \text{CC}(b_i) = \text{CC}(b_j)$ . (Recall that all companion sides  $b_i$  belong to the same component.) Similarly, every side  $B_i$  must intersect every side  $a_j$ . We have no information about the intersection between  $B_i$  and  $b_j$ , as these sides belong to the same connected component. We will now derive a contradiction through a series of case distinctions.

**Case 1:** There are three segments  $A_i$  with the property that  $A_i$  crosses  $\ell_i$  to the left of  $a_i$ . Without loss of generality, assume that these segments are  $A_1, A_2, A_3$ , see Fig. 18. The segments  $A_1, A_2, A_3$  must not cross because  $P$  is a simple polygon. Therefore  $A_1$  intersects  $a_2$  to the right of  $I(a_1, a_2)$  because otherwise  $A_1$  would cross  $A_2$  on the way between its intersections with  $\ell_2$  and with  $a_1$ .  $A_3$  must cross  $\ell_3, a_2, a_1$  in this order, as shown. But then  $A_1$  and  $A_3$  (and  $a_2$ ) block  $A_2$  from intersecting  $a_3$ .

**Case 2:** There at most two segments  $A_i$  with the property that  $A_i$  crosses  $\ell_i$  to the left of  $a_i$ . In this case, we simply discard these segments. We select four of the remaining segments and renumber them from 1 to 4. From now on, we can make the following assumption:

**General Assumption:** For every  $1 \leq i \leq 4$ , the segment  $A_i$  does not cross  $\ell_i$  at all, or it crosses  $\ell_i$  to the right of  $a_i$ .




Fig. 18 The assumed intersection points between  $A_i$  and  $\ell_i$  are marked





Fig. 19 Case 2.1,  $I(A_3, B_3) = \text{left}(A_3)$  and  $I(a_3, b_3) = \text{right}(a_3)$ . A hypothetical segment  $A_2$  is shown as a dashed curve. The side  $a_2$  and the part of  $\ell_2$  to the left of  $a_2$  is blocked for  $A_2$

This implies that  $A_3$  must intersect the sides  $a_2, a_1, a_4$  in this order, and it is determined in which cell of the arrangement of the lines  $\ell_1, \ell_2, \ell_3, \ell_4$  the left endpoint of  $A_3$  lies (see Figs. 16 and 19). For the right endpoint, we have a choice of two cells, depending on whether  $A_3$  intersects  $\ell_3$  or not.

We denote by  $\text{left}(s)$  and  $\text{right}(s)$  the left and right endpoints of a segment  $s$ . We distinguish four cases, based on whether the common endpoint of  $A_3$  and  $B_3$  lies at  $\text{left}(A_3)$  or  $\text{right}(A_3)$ , and whether the common endpoint of  $a_3$  and  $b_3$  lies at  $\text{left}(a_3)$  or  $\text{right}(a_3)$ .

**Case 2.1:**  $I(A_3, B_3) = \text{left}(A_3)$  and  $I(a_3, b_3) = \text{right}(a_3)$ , see Fig. 19.



**Fig. 20** Case 2.2.  $I(A_3, B_3) = \text{left}(A_3)$ ,  $I(a_3, b_3) = \text{left}(a_3)$ ,  $\ell(A_3)$  does not intersect  $a_3$

As indicated in the figure, we leave it open whether and where  $A_3$  intersects  $\ell_3$ . We know that  $b_3$  must lie below  $\ell_3$  because  $I(A_3, B_3) \in \text{Cone}(a_3, b_3)$ . We claim that  $A_2$  cannot have the required intersections with  $a_1, a_3$ , and  $b_3$ . Let us first consider  $a_1$ : It is cut into three pieces by  $A_3$  and  $B_3$ .

If  $A_2$  intersects the middle piece of  $a_1$  in the wedge between  $A_3$  and  $B_3$ , then  $A_2$  intersects exactly one of  $a_3$  and  $b_3$  inside the wedge, as these parts together with  $a_1$  are three sides of a convex pentagon. If  $A_2$  intersects  $a_3$ , then it has crossed  $\ell_3$  and it cannot cross  $b_3$  thereafter. If  $A_2$  intersects  $b_3$ , it must cross  $\ell_4$  before leaving the wedge, and then it cannot cross  $a_3$  thereafter.

Suppose now that  $A_2$  crosses the bottom piece of  $a_1$ . Then it cannot go around  $A_3$  and  $B_3$  to the right in order to reach  $a_3$  because it would have to intersect  $\ell_4$  twice.  $A_2$  also cannot pass to the left of  $A_3$  and  $B_3$  because it cannot cross  $\ell_2$  through  $a_2$  or, by the general assumption, to the Suppose finally that  $A_2$  crosses the top piece of  $a_1$ . Then it would have to cross  $\ell_3$  twice before reaching  $b_3$ .

**Case 2.2:**  $I(A_3, B_3) = \text{left}(A_3)$  and  $I(a_3, b_3) = \text{left}(a_3)$ .

If  $\ell(A_3)$  does not intersect  $a_3$ , we derive a contradiction as follows, see Fig. 20. We know that the sides  $a_2, a_3, a_4$  must be arranged as shown. The segment  $A_3$  crosses  $a_2$  but not  $a_3$ . Now, the parts of  $a_3$  and  $A_3$  to the left of  $\ell_2$  form two opposite sides of a quadrilateral, as shown in the figure. If this quadrilateral were not convex, then either  $\ell(A_3)$  would intersect  $a_3$ , which we have excluded by assumption, or  $\ell_3$  would intersect  $A_3$  left of  $a_3$ , contradicting the General Assumption. Thus, the sides  $a_3$  and  $A_3$  violate the Axis Property (Observation 6), which requires  $a_3$  and  $A_3$  to lie on different sides of the line through  $I(A_3, B_3)$  and  $I(a_3, b_3)$ .

Looking back at the proofs so far, we have seen that the configuration of the segments  $a_1, a_2, a_3, a_4$  according to Fig. 16 in connection with the particular case assumptions make the situation sufficiently constrained that the case can be dismissed by looking at the drawing. The treatment of the other cases will be proofs by picture in a similar way, but we will not always spell out the arguments in such detail.

If  $\ell(A_3)$  intersects  $a_3$ , the situation must be as shown in Fig. 22: the pair  $A_3, B_3$  is hooked by  $a_3$  and  $b_3$ . The analysis of Case 2.1 (Fig. 19) applies verbatim, except that the word “pentagon” must be replaced by “hexagon”.

**Case 2.3:**  $I(A_3, B_3) = \text{right}(A_3)$  and  $I(a_3, b_3) = \text{right}(a_3)$ .

If  $A_3$  lies entirely below  $\ell_3$ , then  $A_3$  together with  $a_3$  violates the Axis Property (Observation 6), see Fig. 21.




Fig. 21 Case 2.3.  $I(A_3, B_3) = \text{right}(A_3)$ , and  $I(a_3, b_3) = \text{right}(a_3)$ ,  $A_3$  lies below  $\ell_3$




Fig. 22 Case 2.2,  $I(A_3, B_3) = \text{left}(A_3)$ ,  $I(a_3, b_3) = \text{left}(a_3)$ , and  $\ell(A_3)$  intersects  $A_3$ . A hypothetical segment  $A_2$  is shown as a dashed curve




Fig. 23 Case 2.3.  $A_3$  intersects  $\ell_3$

Let us therefore assume that  $A_3$  intersects  $\ell_3$  (to the right of  $a_3$ ), and thus  $\text{right}(A_3) = I(A_3, B_3)$  lies above  $\ell_3$ , see Fig. 23a. Then  $b_3$  must also lie above  $\ell_3$ , because  $a_3, b_3$  is supposed to be hooking, that is,  $I(A_3, B_3) \in \text{Cone}(a_3, b_3)$ .

It follows that  $A_3$  cannot intersect  $\ell_3$  to the right of  $I(a_3, a_4)$  (the option shown as a dashed curve), because otherwise it would miss  $b_3$ :  $b_3$  is blocked by  $a_4$ .




Fig. 24 Case 2.4.  $A_3$  intersects  $\ell_3$




Fig. 25 Case 2.4.  $A_3$  lies below  $\ell_3$

Therefore, the situation looks as shown in Fig. 23a. Figure 23b shows the position of the relevant pieces. The segments  $a_4, B_3, a_3, b_3, A_3$  enclose a convex pentagon. Now, the segment  $A_2$  should intersect  $a_3, b_3$ , and  $a_4$  without crossing  $A_3$  and  $B_3$ , like the dashed curve in the figure. This is impossible.

**Case 2.4:**  $I(A_3, B_3) = \text{right}(A_3)$  and  $I(a_3, b_3) = \text{left}(a_3)$ .

If  $A_3$  intersects  $\ell_3$  (to the right of  $a_3$ ), then  $A_3$  together with  $a_3$  violates the Axis Property (Observation 6), see Fig. 24. We thus assume that  $A_3$  lies entirely below  $\ell_3$ .

If  $\ell(A_3)$  passes above  $I(a_3, b_3) = \text{left}(a_3)$ , the sides  $a_3$  and  $A_3$  violate the Axis Property, see Fig. 25a. On the other hand, if  $\ell(A_3)$  passes below  $I(a_3, b_3) = \text{left}(a_3)$ , as shown in Fig. 25b, then  $b_3$  must cross  $\ell_1$  to the right of  $a_1$  in order to reach  $A_2$ . Again by the Axis Property,  $B_3$  must remain above the dotted axis line through  $I(A_3, B_3) = \text{right}(A_3)$  and  $I(a_3, b_3) = \text{left}(a_3)$ . On  $\ell_1, b_3$  separates  $a_1$  from the axis line, and hence  $a_1$  lies below the axis line. Therefore  $B_3$  and  $a_1$  cannot intersect. This concludes the proof of Theorem 1.

## 9 Degenerate Cases

We will now justify our general-position assumption that we introduced in Sect. 3 stating that no vertex of one polygon lies on the boundary of the other polygon. That is, we will show that one can get rid of such degenerate cases without decreasing the number of intersection points. There are two types of degenerate cases:

**vertex-vertex:** Two vertices from different polygons coincide.

**vertex-side:** A vertex of one polygon lies on a side of the other polygon.




Fig. 26 The dependence graph may contain cycles

We will first use a global perturbation governed by a parametric system of linear equations to get rid of vertex-vertex degeneracies. Vertex-side degeneracies can then be treated locally by shifting sides one by one.

### 9.1 Coincident Vertices

Finding a good perturbation is delicate because a side may be involved in multiple degeneracies. We will simultaneously perturb all vertex-vertex degeneracies while leaving vertex-side degeneracies intact.

For each vertex-side degeneracy, some point  $A$  of one polygon lies on a side  $BC$  of the other polygon: There is a ratio  $\lambda = \lambda_{ABC} \in (0, 1)$  such that

$$A = \lambda B + (1 - \lambda)C. \tag{1}$$

Our perturbation will maintain this relation, with the same value  $\lambda$ . We say that  $A$  depends on  $B$  and  $C$ . We can set up a *dependence graph* that has arcs  $(A, B)$  and  $(A, C)$ , for each such vertex-side degeneracy. This graph may contain cycles. The polygon vertices that don't depend on other vertices are called *free*.

We specify the movement of the free vertices as follows: Free vertices that are not involved in a degeneracy stay where they are. For each pair  $(P_i, Q_j)$  of coincident vertices, we leave  $Q_j$  stationary but we move  $P_i = P_i(\alpha) := P_i^0 + \alpha E_i$  linearly, with constant speed in an appropriate direction  $E_i$ . We choose each direction  $E_i$  independently so that the two polygons have at least one nondegenerate intersection in the neighborhood of the original configuration, assuming that all other points stay where they are.

Theorem 17 below shows that we can perform such a motion in a continuous way while simultaneously maintaining all vertex-side degeneracies (1).

In this theorem, we consider a general system of finitely many points  $A_1, A_2, \dots, A_N$ , which are partitioned into *dependent* points and *free* points. We denote the indices of the dependent points by  $D \subseteq [N]$  and those of the free points by  $F = [N] \setminus D$ . Each dependent point  $A_i$  is a convex combination of other points:

$$A_i = \sum_{j \in [N] \setminus \{i\}} \lambda_{ij} A_j \tag{2}$$

with

$$\sum_{j \in [N] \setminus \{i\}} \lambda_{ij} = 1,$$

whereas each free point  $A_i$  is given “externally”, depending linearly on some parameter  $\alpha$ :

$$A_i = A_i^0 + \alpha E_i, \quad i \in F. \tag{3}$$

We will use the following simple observation.

**Observation 15** *If (2) holds for a dependent point  $A_i$ , then either*

- (i) *all points  $A_j$  on which  $A_i$  depends coincide with  $A_i$ , or*
- (ii) *some point  $A_j$  on which  $A_i$  depends is strictly larger than  $A_i$ , in the lexicographic order according to the coordinates  $(x, y)$ .*

The *dependence graph* contains a node for every point  $A_i$ , and contains the arc  $(i, j)$  whenever  $\lambda_{ij} > 0$ . The next lemma and theorem hold for a general system of the form (2)–(3), under the following assumption about the dependence graph: From every dependent point, one must be able to reach a free point. In other words, there must be no component of dependent points that depend only on each other. It is easy to show that this condition is fulfilled in our setting, see Lemma 18.

**Lemma 16** *Suppose that from every dependent point, one can reach a free point in the dependence graph. Then, in every solution of the system of equations (2), all points lie in the convex hull of the free points.*

**Proof** We start with some dependent point  $A_i$ . Our first goal is to reach a free point that is lexicographically at least as large as  $A_i$ . We apply Observation 15 to  $A_i$ . In case (i), we repeatedly apply Observation 15 to all points  $A_j$  on which  $A_i$  depends. Eventually, we conclude that all points that are reachable from  $A_i$  coincide with  $A_i$ , or we get a point  $A_k$  that is lexicographically strictly larger. In the first case, there is, among the points that coincide with  $A_i$ , a free point, by assumption. In the second case, we repeat the process with  $A_k$  until we eventually arrive at a free point.

Thus, we conclude that the  $x$ -coordinate of  $A_i$  is not larger than the largest  $x$ -coordinate of the free points. Since the proof is not affected by rotations, (or in fact, arbitrary affine transformations) we obtain the claimed result. □

**Theorem 17** *Suppose that from every dependent point, one can reach a free point in the dependence graph. If the system of equations (2)–(3) has a solution for  $\alpha = 0$ , then it has a unique solution for all  $\alpha$ , and the solution depends linearly on  $\alpha$ .*

**Proof** The system of equations (2)–(3) actually decomposes into two independent systems of linear equations: one for the  $x$ -coordinates, and one for the  $y$ -coordinates. (In  $d$  dimensions, there would be  $d$  such systems.)

If we substitute the explicit values from (3) into (2), we get a  $|D| \times |D|$  system of linear equations with parametric right-hand sides where the unknowns are the  $x$ -coordinates (resp.,  $y$ -coordinates) of the  $|D|$  dependent points. We are done if we show

that the coefficient matrix of this system, whose entries are the coefficients  $\lambda_{ij}$ , has non-zero determinant. A system of linear equations with zero determinant has either no solution or infinitely many solutions that form an affine subspace. Let us look at the set of solutions for  $\alpha = 0$ . We can exclude the first possibility, because we have assumed a solution for  $\alpha = 0$ . To exclude the second possibility, assume that there are infinitely many solutions. Since they form an affine subspace, there must be solutions where some coordinate of some point takes arbitrarily high values. This contradicts Lemma 16.  $\square$

In order to apply Theorem 17, we have to show that the assumption is fulfilled in our system:

**Lemma 18** *In the dependence graph derived from the system of dependencies (1) describing the vertex-side degeneracies, one can reach a free point from every dependent point.*

**Proof** We look at our given (initial) configuration. We take some dependent point  $A_i$  and apply Observation 15 to it. We know that the points on which  $A_i$  depends don't coincide with  $A_i$ , and thus we find a lexicographically strictly larger point. Repeating the argument as long as necessary, we must eventually arrive at a free point that is reachable from  $A_i$ .  $\square$

One final point: We have chosen each movement  $P_i = P_i^0 + \alpha E_i$  of a free point that is involved in a vertex-vertex degeneracy in such a way that there are no degeneracies in the vicinity of  $P_i$ , assuming that the other points do not move. In reality, the other endpoints of the four involved sides may move on linear trajectories. However, as we start from coinciding points, the movement of the other endpoints has a lower-order effect. To see this, consider the marked distance from  $Q_j$  to the intersection between  $Q_{j-1}Q_j$  and  $P_iP_{i-1}$  in Fig. 27.  $P_i$  and  $Q_j$  are free, and  $Q_j$  remains stationary. If  $P_i$  moves linearly as  $\alpha E_i$ , the marked distance changes like  $k\alpha + O(\alpha^2)$  for some constant  $k > 0$  if all other points remain fixed, see Fig. 27b: The distance would have a linear growth of the form  $k\alpha$  if the sides would undergo a parallel shift, but in fact the sides are rotated. If one or both endpoints of an edge move by a distance  $O(\alpha)$ , the rotation angle is bounded by  $O(\alpha)$ . Since the leading side  $\alpha E_i$  of the shaded triangle is of length  $O(\alpha)$ , the effect of the rotation on the side lengths of the triangle is of order  $O(\alpha^2)$ . (This can be checked by using the sine law.) An additional linear movement of  $P_{i-1}$  (see Fig. 27c) or  $Q_{j-1}$  by a distance  $O(\alpha)$  will cause an additional change of direction by  $O(\alpha)$ . Still, the side lengths of the triangle change only by  $O(\alpha^2)$ . Thus, for small enough  $\alpha$ , the linear growth  $k\alpha$  dominates, and there will be no degenerate situations.

The morphing procedure goes back to Floater and Gotsman [7]. The idea for proving unique solvability in Theorem 17 was used, in a different context, in [4].

## 9.2 Vertex-Side Degeneracies

Once there are no more coincident vertices, it is easy to get rid of the remaining degeneracies by looking at the sides one by one, see Fig. 28.






Fig. 27 Lower-order effects due to movement of endpoints




Fig. 28 Perturbing a blue polygon side to get rid of collinear vertices without decreasing the number of intersection points: **a** before, **b** after

If we perform a small parallel shift of a side, an intersection point may disappear or multiply into two intersection points, depending on the direction of the shift. However, the average number of intersections over the two possible directions is always the same as before. Thus, there exists a small parallel shift that does not decrease the number of intersection points.

### 9.3 The Combinatorial Counting Model

Instead of counting intersection points, one could count pairs of sides that intersect. A vertex-side degeneracy is then counted twice, and a vertex-vertex degeneracy is counted four times. We leave this variation for future research.

In [3], a more refined counting model is used: a further distinction is made for vertex-side degeneracies, in accordance with the number of intersection points that can be achieved by perturbation: if one polygon *touches* the other (i.e., two adjacent sides lie on the same side of the side of the other polygon), the intersection is counted twice. If one polygon *crosses* the other, it is counted as one intersection. In vertex-vertex degeneracies, depending on the angles in which the sides meet, the number of intersections that one can get after perturbation can be any number between 1 and 4. With this type of accounting, our results carry over.

## 10 Discussion

We have shown that the number of intersections of a simple  $n$ -gon and a simple  $m$ -gon, for odd  $n$  and  $m$ , is at most  $mn - (m + n) + C$ , for some constant  $C$ . This bound matches the best lower bound  $mn - (m + n) + 3$  up to a constant. However, our constant  $C$  is huge, and we have not made much effort to improve it. For example, it can be improved by replacing Lemma 10 with a less general result which suffices for our purposes. Recall that we use Lemma 10 to obtain a transitive sub-tournament in a tournament whose vertices represent pairwise disjoint segments and whose edges represent stabbing relations. Instead of the logarithmic-size transitive sub-tournament obtained using Lemma 10, one can get a polynomial-size transitive sub-tournament using Dilworth's Theorem as follows (see [9] for a similar argument): Let  $s$  and  $s'$  be two disjoint segments such that  $s$  stabs  $s'$  or vice versa. Let  $s <_1 s'$  (resp.,  $s <_2 s'$ ) if  $s$  stabs  $s'$  such that the right extension of  $s$  intersects  $s'$  and the slope of  $s$  is smaller (resp., larger) than the slope of  $s'$ . Similarly, let  $s <_3 s'$  (resp.,  $s <_4 s'$ ) if  $s$  stabs  $s'$  such that the left extension of  $s$  intersects  $s'$  and the slope of  $s$  is smaller (resp., larger) than the slope of  $s'$ . Every pair of segments is comparable by exactly one of these four relations and this implies that each relation  $<_i$  is transitive. From Dilworth's Theorem it follows that the above-mentioned tournament has  $N^{1/4}$  vertices such that the relation between each pair of their corresponding segments is  $<_1$ , or  $N^{3/4}$  vertices such that the relation between each pair of their corresponding segments is not  $<_1$  (where  $N$  is the size of the tournament). In the second case we can apply Dilworth's Theorem once or twice more in a similar way, and conclude that there are always  $i \in \{1, 2, 3, 4\}$  and  $N^{1/4}$  vertices such that the relation between each pair of their corresponding segments is  $<_i$ . Hence, the tournament has a transitive sub-tournament of size  $N^{1/4}$ .<sup>1</sup>


This gives  $C < (2^{67})^4 = 2^{268}$ , which is obviously still very far from the truth, so determining the exact answer for this very basic question remains an interesting open problem. We believe that our approach to bound the number of connected components of the disjointness graph is a right approach, that is, we conjecture:

**Conjecture 2** *Suppose that  $P$  and  $Q$  are simple polygons with  $m$  and  $n$  sides, respectively, such that  $m$  and  $n$  are odd numbers. Then there are at most three connected components in the disjointness graph of the sides of  $P$  and  $Q$ .*

If both polygons are allowed to self-intersect, then a simple construction shows that the upper bound  $mn - \max\{m, n\}$  is the correct one [3, 8]. However, if only one of the polygons is allowed to self-intersect then we do not know of such a construction.

Our results should carry over to *pseudolinear* polygons, that is, polygons whose sides can be extended to an arrangement of pseudolines, but we did not check the details of this extension. In fact, we already used the freedom to draw curved edges in our schematic drawings, like in Figs. 11b, 14, and 15. Moreover, the computer verifications mentioned below in Sect. 10.2 use none of the properties of straight

<sup>1</sup> It may be interesting to determine the maximum size of such a sub-tournament that one can guarantee for every set of  $N$  disjoint segments where for each pair of them one segment stabs the other. A recursive construction starting with three segments whose corresponding tournament is a directed cycle, then replacing each segment with the same "flattened" construction shows that this number is at most  $O(N^{\log_3 2}) \approx O(N^{0.631})$ .



**Fig. 29** A best lower bound construction for  $m = n = 7$ . It is symmetric with respect to the horizontal green axis. In the left drawing of the disjointness graph, the nodes are arranged in the natural order as the edges appear along the polygon boundaries. The right drawing makes the structure as a forest clearer

lines, as opposed to pseudoline arrangements. Obviously, Conjectures 1 and 2 extend to pseudolinear polygons as well.

### 10.1 Alternative Lower-Bound Constructions

Studying the best lower bound constructions (conjectured to be optimal) may provide insights that may help to improve the upper bound. For example, the disjointness graph of the construction on Fig. 1c has three components: one edge and two stars. If we replace a side that corresponds to a leaf in the disjointness graph by a narrow zigzag of three sides, the number of intersections increases by  $2(m - 1)$  or  $2(n - 1)$ , depending of which polygon side is replaced. This way we can get a best lower bound construction for  $m, n + 2$  or  $m + 2, n$  from a best lower bound construction for odd  $m$  and  $n$ . Therefore, there are many ways to create best lower bound constructions. Note that each step as above replaces one leaf by three leaves in the disjointness graph, and thus repeating it would yield the existence of large-degree vertices and constantly many vertices that dominate the disjointness graph. Thus, it is tempting to conjecture that an optimal construction must possess these properties apart from having three connected components (which are necessarily trees). However, this does not seem to be true. We have a construction for  $m = n = 7$  which is significantly different from the one on Fig. 1c and whose disjointness graph has maximum degree 3, see Fig. 29. We believe that this construction can be extended for greater values of  $m = n$  while maintaining maximum degree 3, but the figure becomes too complicated to draw, so we do not go into further details.

## 10.2 Computer-Supported Proofs

We have independently confirmed Lemma 7 concerning the intersection structure of pairs of associated pairs of sides, which is our principal workhorse, by a computer enumeration.

Lemma 7 involves eight polygon sides. Thus, we enumerated all 158830 arrangements of eight lines. Actually, we enumerated pseudoline arrangements, but it is known that all pseudoline arrangements with less than nine lines are stretchable to line arrangements. For each arrangement, we tried all possible ways of assigning the sides  $p_i, p_{i+1}, p_j, p_{j+1}$  and  $q_i', q_{i'\pm 1}, q_{j'}, q_{j'\pm 1}$  (in the notation of Lemma 7) to these lines. Since there are symmetries in the configuration of the eight sides, we did not have to try all  $8! = 40320$  assignments, but only 840. Once this assignment is given, we know where the vertices between consecutive sides, like  $p_i, p_{i+1}$ , are in the arrangement. This specifies one endpoint of each side. It is then easy to check whether it is possible to realize the required intersection pattern of a perfect matching among the eight involved sides. We first took into account the *required* intersections among the sides, for example, between  $p_i$  and  $q_{j'}$ . On each line, these required intersection points must lie on one side of the vertex that we already know. If this is the case, we get a minimum range on the line that the side must cover. We then check if all *forbidden* intersections (for example, between  $p_i$  and  $q_{i'}$ , or between  $p_i$  and  $p_j$ ) are avoided.

After a few days of computing (and several weeks of programming), the result was that there were, up to isomorphism, 14 configurations with the required intersection pattern. In 13 of them, some associated pair did not conform to the hooking/hooked pattern of Lemma 5. The remaining configuration is the one in Fig. 6.

We had some hope to attack the case of *three* pairs of associated pairs of sides and obtain a more powerful statement than Proposition 14. However, since there are too many (pseudo-)line arrangements with 12 lines, this is out of reach for the simple approach outlined above.

## 10.3 The Union of the Polygon Areas

Instead of polygon boundaries, we can consider polygons as regions and ask about the complexity of the union of the polygon areas. The bound on boundary intersections gives an upper bound on the boundary of this union: Every boundary intersection contributes to the complexity of the union, and in the worst case all the  $m + n$  vertices of the two polygons contribute to the complexity of the boundary in addition.

In the examples of Fig. 1, this worst case arises: all polygon vertices contribute to the boundary of the union of the polygon areas. This gives a lower bound of  $mn - 3$  for the maximum complexity of the union of an  $m$ -gon and an  $n$ -gon and it is conjectured that  $mn - 3$  is in fact the exact value. Theorem 1 implies the upper bound  $mn + C$  for some absolute constant  $C$ .

One can also ask about the *intersection* of the polygon areas. Here, the relation to our problem is not clear.

**Acknowledgements** We thank the reviewers for helpful suggestions.

**Funding** Eyal Ackerman: The main part of this work was performed during a visit to Freie Universität Berlin which was supported by the Freie Universität Alumni Program. Balázs Keszegh: Research supported by the Lendület program of the Hungarian Academy of Sciences (MTA), under the grant LP2017-19/2017, by the János Bolyai Research Scholarship of the Hungarian Academy of Sciences, by the National Research, Development and Innovation Office—NKFIH under the grant K 116769, K 132696, and FK 132060, and by the ÚNKP-21-5 New National Excellence Program of the Ministry for Innovation and Technology from the source of the National Research, Development and Innovation Fund.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References

1. Ackerman, E., Keszegh, B., Rote, G.: An almost optimal bound on the number of intersections of two simple polygons. In: 36th International Symposium on Computational Geometry. Leibniz Int. Proc. Inform., vol. 164, # 1. Leibniz-Zent. Inform., Wadern (2020)
2. Černý, J., Kára, J., Král', D., Podbrdský, P., Sotáková, M., Šámal, R.: On the number of intersections of two polygons. *Comment. Math. Univ. Carolin.* **44**(2), 217–228 (2003)
3. Dillencourt, M.B., Mount, D.M., Saalfeld, A.: On the maximum number of intersections of two polyhedra in 2 and 3 dimensions. In: 5th Canadian Conference on Computational Geometry (Waterloo 1993), pp. 49–54. University of Waterloo (1993)
4. Dujmović, V., Frati, F., Gonçalves, D., Morin, P., Rote, G.: Every collinear set in a planar graph is free. *Discrete Comput. Geom.* **65**(4), 999–1027 (2021)
5. Erdős, P., Moser, L.: On the representation of directed graphs as unions of orderings. *Magyar Tud. Akad. Mat. Kutató Int. Közl.* **9**, 125–132 (1964)
6. Erdős, P., Szekeres, G.: A combinatorial problem in geometry. *Compos. Math.* **2**, 463–470 (1935)
7. Floater, M.S., Gotsman, C.: How to morph tilings injectively. *J. Comput. Appl. Math.* **101**(1–2), 117–129 (1999)
8. Günther, F.: The maximum number of intersections of two polygons (2012). [arXiv:1207.0996](https://arxiv.org/abs/1207.0996)
9. Pach, J., Töröcsik, J.: Some geometric applications of Dilworth's theorem. *Discrete Comput. Geom.* **12**(1), 1–7 (1994)

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.