

RIGID CYLINDRICAL FRAMEWORKS WITH TWO COINCIDENT POINTS

B. JACKSON, V. E. KASZANITZKY, AND A. NIXON

ABSTRACT. We develop a rigidity theory for frameworks in \mathbb{R}^3 which have two coincident points but are otherwise generic and only infinitesimal motions which are tangential to a family of cylinders induced by the realisation are considered. We then apply our results to show that vertex splitting, under the additional assumption that the new edge is redundant, preserves the property of being generically globally rigid on families of concentric cylinders.

1. INTRODUCTION

A *framework* (G, p) in \mathbb{R}^d is the combination of a finite, simple graph $G = (V, E)$ and a map $p : V \rightarrow \mathbb{R}^d$. It is *rigid* if every edge-length preserving continuous motion of the vertices arises as a congruence of \mathbb{R}^d (see, for example, [12] for basic definitions and background). The theory of generic rigidity aims to characterise the graphs G for which (G, p) is rigid for all generic choices of p . This was accomplished by Laman [6] for $d = 2$, but is a long-standing open problem for $d \geq 3$.

We are interested in frameworks in \mathbb{R}^3 whose motions are restricted such that all vertices are realised on a fixed surface and only those continuous motions that keep (G, p) on the surface are considered. Generic rigidity in this context has been characterised for various surfaces [8, 9]. In this paper we consider frameworks on concentric cylinders in which two of the vertices are mapped to the same point, but are otherwise generic. For such frameworks we give a combinatorial characterisation of rigidity.

Frameworks in \mathbb{R}^2 with two coincident points were studied in [3] where the following deletion-contraction characterisation of rigidity was proved. A graph G is *uv-rigid* in \mathbb{R}^2 if there exists a realisation p of G in \mathbb{R}^2 such that $p(u) = p(v)$, $p|_{V-v}$ is generic and the framework (G, p) is rigid. We use $G - uv$ to denote the graph formed from G by deleting the edge uv if it exists and G/uv to denote the graph which arises from G by contracting the vertices u and v .

Theorem 1.1. [3] *Let $G = (V, E)$ be a graph and let $u, v \in V$ be distinct vertices. Then G is uv-rigid in \mathbb{R}^2 if and only if $G - uv$ and G/uv are both rigid in \mathbb{R}^2 .*

Our main result is an analogue of this theorem for frameworks on concentric cylinders. One motivation for studying frameworks on cylinders with coincident points is an ongoing work to understand global rigidity for frameworks on concentric cylinders [4, 5]. In particular we utilise our main result to prove that vertex splitting, under the additional assumption that the new edge is redundant, preserves generic global rigidity for families of concentric cylinders.

In Section 2 we provide background on frameworks on concentric cylinders. In Section 3 we define a count matroid $\mathcal{M}_{uv}(G)$ on a graph G with two distinguished vertices u and v . In Section 4 we derive an inductive construction for graphs whose edge set is independent in

$\mathcal{M}_{uv}(G)$. We then use this construction to prove our characterisation of rigidity on families of concentric cylinders for frameworks in which u and v are coincident but are otherwise generic. In Section 5 we discuss global rigidity and apply our coincident point result to prove that the vertex splitting operation preserves generic global rigidity for frameworks on families of concentric cylinders when the new edge is redundant. Finally, in Section 6 we comment on extensions to other surfaces.

In this paper we consider simple graphs only as parallel edges correspond to the same distance constraint and thus one of them is always redundant.

2. FRAMEWORKS ON CONCENTRIC CYLINDERS

Let $G = (V, E)$ where $V = \{v_1, \dots, v_n\}$. We will consider realisations of G on a family of concentric cylinders $\mathcal{Y} = \mathcal{Y}_1 \cup \mathcal{Y}_2 \cup \dots \cup \mathcal{Y}_k$ where $\mathcal{Y}_i = \{(x, y, z) \in \mathbb{R}^3 : x^2 + y^2 = r_i\}$ and $r = (r_1, \dots, r_k)$ is a vector of positive real numbers. A *framework* (G, p) on \mathcal{Y} is an ordered pair consisting of a graph G and a realisation p such that $p(v_i) \in \mathcal{Y}$ for all $v_i \in V$.

Two frameworks (G, p) and (G, q) on \mathcal{Y} are *equivalent* if $\|p(v_i) - p(v_j)\| = \|q(v_i) - q(v_j)\|$ for all edges $v_i v_j \in E$. Moreover (G, p) and (G, q) on \mathcal{Y} are *congruent* if $\|p(v_i) - p(v_j)\| = \|q(v_i) - q(v_j)\|$ for all pairs of vertices $v_i, v_j \in V$. The framework (G, p) is *rigid* on \mathcal{Y} if there exists an $\epsilon > 0$ such that every framework (G, q) on \mathcal{Y} which is equivalent to (G, p) , and has $\|p(v_i) - q(v_i)\| < \epsilon$ for all $1 \leq i \leq n$, is congruent to (G, p) . Moreover (G, p) is *minimally rigid* if (G, p) is rigid but $(G - e, p)$ is not for any $e \in E$. The framework (G, p) is *generic* on \mathcal{Y} if $\text{td}[\mathbb{Q}(r, p) : \mathbb{Q}(r)] = 2n$.

It was shown in [8] that a generic framework (G, p) on any family of concentric cylinders is rigid if and only if it is infinitesimally rigid in the following sense. An *infinitesimal flex* s of (G, p) on \mathcal{Y} is a map $s : V \rightarrow \mathbb{R}^3$ such that $s(v_i)$ is tangential to \mathcal{Y} at $p(v_i)$ for all $v_i \in V$ and $(p(v_j) - p(v_i)) \cdot (s(v_j) - s(v_i)) = 0$ for all $v_j v_i \in E$. The framework (G, p) is *infinitesimally rigid* on \mathcal{Y} if every infinitesimal flex is an infinitesimal isometry of \mathbb{R}^3 .

The *rigidity matrix* $R_{\mathcal{Y}}(G, p)$ is the $(|E| + |V|) \times 3|V|$ matrix

$$R_{\mathcal{Y}}(G, p) = \begin{pmatrix} R_3(G, p) \\ S(G, p) \end{pmatrix}$$

where: $R_3(G, p)$ has rows indexed by E and 3-tuples of columns indexed by V in which, for $e = v_i v_j \in E$, the submatrices in row e and columns v_i and v_j are $p(v_i) - p(v_j)$ and $p(v_j) - p(v_i)$, respectively, and all other entries are zero; $S(G, p)$ has rows indexed by V and 3-tuples of columns indexed by V in which, for $v_i \in V$, the submatrix in row v_i and column v_i is $\bar{p}(v_i) = (x_i, y_i, 0)$ when $p(v_i) = (x_i, y_i, z_i)$. The *rigidity matroid* $\mathcal{R}^{\mathcal{Y}}(G)$ is the row matroid of $R_{\mathcal{Y}}(G, p)$ for any generic p .

A graph $G = (V, E)$ is $(2, 2)$ -*sparse* if $|E'| \leq 2|V'| - 2$ for all subgraphs (V', E') of G . Moreover G is $(2, 2)$ -*tight* if G is $(2, 2)$ -sparse and $|E| = 2|V| - 2$.

The following characterisation of generic rigidity on \mathcal{Y} was proved in [8].

Theorem 2.1. *Let (G, p) be a generic framework on a union of concentric cylinders \mathcal{Y} . Then (G, p) is minimally rigid if and only if G is a complete graph on at most 3 vertices or G is $(2, 2)$ -tight and simple.*

2.1. Coincident realisations on concentric cylinders. Let $G = (V, E)$ be a graph and suppose $u, v \in V$. A framework (G, p) on \mathcal{Y} is *uv-coincident* if $p(u) = p(v)$. A *generic uv-coincident framework* is a *uv-coincident* framework (G, p) for which $(G - u, p|_{V-u})$ is generic. We denote the *uv-coincident cylinder rigidity matroid* by $\mathcal{R}_{uv}^{\mathcal{Y}}(G)$ (this is the row matroid of $R_{\mathcal{Y}}(G, p)$ for any generic *uv-coincident* realisation (G, p)). Note that the matroid depends on G but not on the choice of generic *uv-coincident* realisation. That is, for any

two generic uv -coincident realisations (G, p) and (G, p') on \mathcal{Y} , we get the same matroid. We also use $r_{uv}(G)$ to denote the rank of $\mathcal{R}_{uv}^{\mathcal{Y}}(G)$. We say that G is *uv-rigid* if $r_{uv}(G) = 2|V| - 2$ and that G is *minimally uv-rigid* if G is *uv-rigid* and $|E| = 2|V| - 2$.

Note that the term *uv-rigid* and the notation $r_{uv}(G)$ refer to generic realisations on a fixed family of concentric cylinders \mathcal{Y} , and hence appear to depend on \mathcal{Y} . We will see, however, that this is not the case since our characterisation of $\mathcal{R}_{uv}^{\mathcal{Y}}(G)$ depends only on the graph G .

3. A COUNT MATROID

In this section we define a count matroid $\mathcal{M}_{uv}(G)$ on the edge set of a graph G with two distinguished vertices u and v . We will show that $\mathcal{M}_{uv}(G)$ is equal to $\mathcal{R}_{uv}^{\mathcal{Y}}(G)$ in Section 4.

Let $G = (V, E)$ be a graph. For some $X \subseteq V$ let $G[X]$ denote the subgraph of G induced by X and let $E_G(X)$ be the set of edges of $G[X]$. Thus $i_G(X) = |E_G(X)|$. For a family $\mathcal{S} = \{S_1, S_2, \dots, S_k\}$, where $S_i \subseteq V$ for all $i = 1, \dots, k$, we define $E_G(\mathcal{S}) = \bigcup_{i=1}^k E_G(S_i)$ and put $i_G(\mathcal{S}) = |E_G(\mathcal{S})|$. We also define $\text{cov}(\mathcal{S}) = \{(x, y) : x, y \in V, \{x, y\} \subseteq S_i \text{ for some } 1 \leq i \leq k\}$. We say that \mathcal{S} *covers* a set $F \subseteq E$ if $F \subseteq \text{cov}(\mathcal{S})$. The degree of a vertex w is denoted by $d_G(w)$. We let $N_G(w) = \{z \in V : wz \in E\}$ denote the *neighbours* of w in G . We may omit the subscripts referring to G if the graph is clear from the context.

Let $G = (V, E)$ be a graph and $u, v \in V$ be two distinct vertices of G . Let $\mathcal{H} = \{H_1, \dots, H_k\}$ be a family with $H_i \subseteq V$, $1 \leq i \leq k$. We say that \mathcal{H} is *uv-compatible* if $u, v \in H_i$ and $|H_i| \geq 3$ hold for all $1 \leq i \leq k$. See Figure 1 for an example. We define the *value* of subsets of V and of *uv-compatible* families as follows. For a nonempty subset $H \subseteq V$, we let

$$\text{val}(H) = 2|H| - t_H,$$

where $t_H = 4$ if $H = \{u, v\}$, $t_H = 3$ if $H \neq \{u, v\}$ and $|H| \in \{2, 3\}$, and $t_H = 2$ otherwise. We will often denote t_{H_i} by t_i for short. For a *uv-compatible* family $\mathcal{H} = \{H_1, H_2, \dots, H_k\}$ we let

$$\text{val}(\mathcal{H}) = \sum_{i=1}^k \text{val}(H_i) - 2(k-1).$$

Note that if $\mathcal{H} = \{H\}$ is a *uv-compatible* family containing only one set then the two definitions are compatible, i.e. $\text{val}(\mathcal{H}) = \text{val}(H)$ holds.

We say that G is *uv-sparse* if for all $H \subseteq V$ with $|H| \geq 2$ we have $i_G(H) \leq \text{val}(H)$ and for all *uv-compatible* families \mathcal{H} we have $i_G(\mathcal{H}) \leq \text{val}(\mathcal{H})$. Note that if G is *uv-sparse* then G is simple and $uv \notin E$ must hold. A set $H \subseteq V$ of vertices with $|H| \geq 2$ (resp. a *uv-compatible* family $\mathcal{H} = \{H_1, \dots, H_k\}$) is called *tight* if $i_G(H) = \text{val}(H)$ (resp. $i_G(\mathcal{H}) = \text{val}(\mathcal{H})$) holds. We will show that the edge sets of the *uv-sparse* subgraphs of G form the independent sets of a matroid that we will denote by $\mathcal{M}_{uv}(G)$.

The next lemmas will enable us to 'uncross' tight sets and tight *uv-compatible* families in a sparse graph. The first result follows immediately from the definition of the i - and val -functions.

Lemma 3.1. *Let $X, Y \subseteq V$ be distinct vertex sets in G . Then*

- (a) $i(X) + i(Y) \leq i(X \cup Y) + i(X \cap Y)$ and
- (b) if $X \cap Y \neq \emptyset$, then $\text{val}(X) + \text{val}(Y) + t_X + t_Y = \text{val}(X \cup Y) + \text{val}(X \cap Y) + t_{X \cup Y} + t_{X \cap Y}$.

Lemma 3.2. *Let $\mathcal{H} = \{H_1, \dots, H_k\}$ be a *uv-compatible* family in G .*

- (a) *Suppose $|H_i \cap H_j| \geq 3$ for some pair $1 \leq i < j \leq k$. Then there is a *uv-compatible* family \mathcal{H}' with $\text{cov}(\mathcal{H}) \subseteq \text{cov}(\mathcal{H}')$ and $\text{val}(\mathcal{H}') < \text{val}(\mathcal{H})$.*
- (b) *Suppose G is *uv-sparse* and \mathcal{H} is tight. Then $H_i \cap H_j = \{u, v\}$ for all $1 \leq i \leq k$.*

Proof. (a) We may assume that $i = k - 1$, $j = k$. Let $\mathcal{H}' = \{H_1, \dots, H_{k-2}, H_{k-1} \cup H_k\}$. Using Lemma 3.1(b) we have $\text{val}(H_{k-1}) + \text{val}(H_k) \geq \text{val}(H_{k-1} \cup H_k) + \text{val}(H_{k-1} \cap H_k)$. Hence

$$\begin{aligned} \text{val}(\mathcal{H}) &= \sum_{l=1}^k \text{val}(H_l) - 2(k-1) = \sum_{l=1}^{k-2} \text{val}(H_l) - 2((k-1)-1) + \text{val}(H_{k-1}) + \text{val}(H_k) - 2 \\ &\geq \sum_{l=1}^{k-2} \text{val}(H_l) + \text{val}(H_{k-1} \cup H_k) - 2((k-1)-1) + \text{val}(H_{k-1} \cap H_k) - 2 > \text{val}(\mathcal{H}'). \end{aligned}$$

Clearly, we have $\text{cov}(\mathcal{H}) \subseteq \text{cov}(\mathcal{H}')$.

(b) Since \mathcal{H} is tight, if $|H_i \cap H_j| \geq 3$ for some pair $1 \leq i < j \leq k$ then, by (a), we have $\text{val}(\mathcal{H}') < \text{val}(\mathcal{H}) = i(\mathcal{H}) \leq i(\mathcal{H}')$. This contradicts the uv -sparsity of G . Hence $H_i \cap H_j = \{u, v\}$ for all $1 \leq i \leq k$. \square

Lemma 3.3. *Let $\mathcal{H} = \{H_1, \dots, H_k\}$ be a uv -compatible family with $H_i \cap H_j = \{u, v\}$ for all $1 \leq i < j \leq k$ and $|H_k| \geq 4$. Then $\mathcal{H}' = \{H_1, \dots, H_{k-2}, H_{k-1} \cup H_k\}$ is a uv -compatible family with $\text{cov}(\mathcal{H}) \subset \text{cov}(\mathcal{H}')$ and for which $\text{val}(\mathcal{H}') \leq \text{val}(\mathcal{H}) + 1$ with equality only if $|H_{k-1}| = 3$. Furthermore, if G is uv -sparse, \mathcal{H} is tight and $|H_{k-1}| \geq 4$, then \mathcal{H}' is tight.*

Proof. Using Lemma 3.1(b) and the facts that $t_k = t_{H_{k-1} \cup H_k} = 2$ and $t_{H_{k-1} \cap H_k} = 4$ we have $\text{val}(H_{k-1}) + \text{val}(H_k) = \text{val}(H_{k-1} \cup H_k) + \text{val}(H_{k-1} \cap H_k) + 4 - t_{k-1} = \text{val}(H_{k-1} \cup H_k) + 4 - t_{k-1}$. Hence

$$\begin{aligned} \text{val}(\mathcal{H}) &= \sum_{l=1}^k \text{val}(H_l) - 2(k-1) = \sum_{l=1}^{k-2} \text{val}(H_l) - 2((k-1)-1) + \text{val}(H_{k-1}) + \text{val}(H_k) - 2 \\ &= \sum_{l=1}^{k-2} \text{val}(H_l) + \text{val}(H_{k-1} \cup H_k) - 2((k-1)-1) + 2 - t_{k-1} \\ &= \text{val}(\mathcal{H}') + 2 - t_{k-1}. \end{aligned}$$

Thus $\text{val}(\mathcal{H}') \leq \text{val}(\mathcal{H}) + 1$ with equality only if $|H_{k-1}| = 3$. Clearly, we have $\text{cov}(\mathcal{H}) \subset \text{cov}(\mathcal{H}')$.

Now suppose G is uv -sparse, \mathcal{H} is tight and $|H_{k-1}| \geq 4$. Then $\text{val}(\mathcal{H}') \leq \text{val}(\mathcal{H}) = i(\mathcal{H}) = i(\mathcal{H}')$, so \mathcal{H}' is tight. \square

Lemma 3.4. *Let $G = (V, E)$ be uv -sparse and let $X, Y \subseteq V$ be tight sets in G with $X \cap Y \neq \emptyset$ and $|X|, |Y| \geq 4$. Then $|X \cap Y| \notin \{2, 3\}$ and $X \cup Y$ and $X \cap Y$ are both tight.*

Proof. We have

$$\begin{aligned} 2|X| - 2 + 2|Y| - 2 &= i(X) + i(Y) \leq i(X \cup Y) + i(X \cap Y) \\ &\leq 2|X \cup Y| - t_{X \cup Y} + 2|X \cap Y| - t_{X \cap Y} = 2|X| + 2|Y| - 2 - t_{X \cap Y}. \end{aligned}$$

This implies that $t_{X \cap Y} = 2$ and equality holds throughout. Thus $X \cup Y$ and $X \cap Y$ are both tight and either $|X \cap Y| \geq 4$ or $|X \cap Y| = 1$. \square

Lemma 3.5. *Let $\mathcal{H} = \{H_1, \dots, H_k\}$ be a uv -compatible family with $H_j \cap H_l = \{u, v\}$ for all $1 \leq j < l \leq k$, and let $Y \subseteq V$ be a set of vertices with $|Y| \geq 4$, and $|Y \cap \{u, v\}| \leq 1$. Suppose that for some $1 \leq i \leq k$ either $|Y \cap H_i| \geq 2$, or $|Y \cap H_i| = 1$ and $|H_i| \geq 4$. Then there is a uv -compatible family \mathcal{H}' with $\text{cov}(\mathcal{H}) \cup \text{cov}(Y) \subseteq \text{cov}(\mathcal{H}')$ and $\text{val}(\mathcal{H}') \leq \text{val}(\mathcal{H}) + \text{val}(Y)$. Furthermore, if G is uv -sparse and \mathcal{H} and Y are both tight then \mathcal{H}' and $Y \cap H_i$ are also tight.*

Proof. Let $S = \{H_i \in \mathcal{H} : |Y \cap H_i| \geq 2 \text{ or } |Y \cap H_i| = 1 \text{ and } |H_i| \geq 4\}$. Renumbering the sets of \mathcal{H} , if necessary, we may assume that $S = \{H_i \in \mathcal{H} : j \leq i \leq k\}$, for some $j \leq k$. Let $X = Y \cup (\cup_{i=j}^k H_i)$ and $\mathcal{H}' = \{H_1, \dots, H_{j-1}, X\}$. Then $\text{cov}(\mathcal{H}) \cup \text{cov}(Y) \subseteq \text{cov}(\mathcal{H}')$ and

$$|X| = \sum_{i=j}^k |H_i| + |Y| - 2(k-j) - \sum_{i=j}^k |H_i \cap Y| + |Y \cap \{u, v\}|(k-j).$$

This gives

$$\begin{aligned} \text{val}(\mathcal{H}) + \text{val}(Y) &= \sum_{i=1}^k \text{val}(H_i) - 2(k-1) + \text{val}(Y) \\ &= \sum_{i=1}^{j-1} \text{val}(H_i) - 2(j-1) + \sum_{i=j}^k (2|H_i| - t_i) - 2(k-j) + (2|Y| - 2) \\ &= \sum_{i=1}^{j-1} \text{val}(H_i) + (2|X| - 2) - 2(j-1) + 4(k-j) - \sum_{i=j}^k t_{H_i} \\ &\quad + 2 \sum_{i=j}^k |Y \cap H_i| - 2(k-j) - 2|Y \cap \{u, v\}|(k-j) \\ &\geq \sum_{i=1}^{j-1} \text{val}(H_i) + \text{val}(X) - 2(j-1) + \sum_{i=j}^k (2|Y \cap H_i| - t_{H_i}). \end{aligned}$$

If $|Y \cap H_i| \geq 2$ then $\text{val}(Y \cap H_i) = 2|Y \cap H_i| - t_{Y \cap H_i} \leq 2|Y \cap H_i| - t_{H_i}$. On the other hand, if $|Y \cap H_i| = 1$ and $|H_i| \geq 4$, then $t_{Y \cap H_i} = 2 = t_{H_i}$ and we have $\text{val}(Y \cap H_i) = 2|Y \cap H_i| - t_{H_i}$. Thus, in both cases,

$$\text{val}(\mathcal{H}) + \text{val}(Y) \geq \text{val}(\mathcal{H}') + \sum_{i=j}^k \text{val}(Y \cap H_i)$$

and so $\text{val}(\mathcal{H}') \leq \text{val}(\mathcal{H}) + \text{val}(Y)$.

Now, suppose that G is uv -sparse and \mathcal{H} and Y are tight. Then we have

$$\begin{aligned} i(\mathcal{H}') + \sum_{i=j}^k i(Y \cap H_i) &\geq i(\mathcal{H}) + i(Y) = \text{val}(\mathcal{H}) + \text{val}(Y) \geq \\ &\geq \text{val}(\mathcal{H}') + \sum_{i=j}^k \text{val}(Y \cap H_i) \geq i(\mathcal{H}') + \sum_{i=j}^k i(Y \cap H_i), \end{aligned}$$

where the first inequality follows from the fact that edges spanned by \mathcal{H} or Y are spanned by \mathcal{H}' and if some edge is spanned by both \mathcal{H} and Y then it is spanned by $Y \cap H_i$ for some i . The equality holds because \mathcal{H} and Y are tight, and the second inequality holds by our calculations above. The last inequality holds because G is uv -sparse. Hence equality must hold everywhere, which implies that \mathcal{H}' is tight and that $Y \cap H_i$ is also tight for all $j \leq i \leq k$. \square

Lemma 3.6. *Let $\mathcal{H} = \{H_1, \dots, H_k\}$ be a uv -compatible family with $H_i \cap H_j = \{u, v\}$ for all $1 \leq i < j \leq k$, and let $Y \subseteq V$ be a set of vertices with $|Y| \geq 4$, $Y \cap \{u, v\} = \emptyset$ and $|Y \cap H_i| \leq 1$ for all $1 \leq i \leq k$. Suppose that $|Y \cap H_i| = |Y \cap H_j| = 1$ for some pair*

$1 \leq i < j \leq k$. Then there is a uv -compatible family \mathcal{H}' with $\text{cov}(\mathcal{H}) \cup \text{cov}(Y) \subseteq \text{cov}(\mathcal{H}')$ for which $\text{val}(\mathcal{H}') \leq \text{val}(\mathcal{H}) + \text{val}(Y)$. Furthermore, if G is uv -sparse and \mathcal{H} and Y are both tight, then \mathcal{H}' is tight and $|H_i| = |H_j| = 3$.

Proof. We may assume that $i = k-1$ and $j = k$. Let $\mathcal{H}' = \{H_1, \dots, H_{k-2}, H_{k-1} \cup H_k \cup Y\}$. We have $\text{cov}(\mathcal{H}) \cup \text{cov}(Y) \subseteq \text{cov}(\mathcal{H}')$ and

$$\begin{aligned} \text{val}(\mathcal{H}) + \text{val}(Y) &= \sum_{i=1}^k \text{val}(H_i) - 2(k-1) + \text{val}(Y) \\ &= \sum_{i=1}^{k-2} \text{val}(H_i) - 2((k-1)-1) - 2 + \text{val}(H_{k-1}) + \text{val}(H_k) + \text{val}(Y). \end{aligned}$$

Using Lemma 3.1(b) twice and the fact that $|H_{k-1} \cap (H_k \cup Y)| = 3$ we obtain

$$\begin{aligned} \text{val}(H_{k-1}) + \text{val}(H_k) + \text{val}(Y) &= \text{val}(H_{k-1}) + \text{val}(H_k \cup Y) + 2 - t_{H_k} \\ &= \text{val}(H_{k-1} \cup H_k \cup Y) + 8 - t_{H_{k-1}} - t_{H_k} \\ &\geq \text{val}(H_{k-1} \cup H_k \cup Y) + 2, \end{aligned}$$

with equality only if $|H_{k-1}| = |H_k| = 3$. Thus $\text{val}(\mathcal{H}') \leq \text{val}(\mathcal{H}) + \text{val}(Y)$ as claimed.

Now suppose that G is uv -sparse. and \mathcal{H} and Y are both tight. Then we have

$$i(\mathcal{H}) + i(Y) = \text{val}(\mathcal{H}) + \text{val}(Y) \geq \text{val}(\mathcal{H}') \geq i(\mathcal{H}') \geq i(\mathcal{H}) + i(Y)$$

where the last inequality follows since $|Y \cap H_{k-1}| = |Y \cap H_k| = 1$ and $|Y \cap H_i| \leq 1$ for all $1 \leq i \leq k$. Hence equality must hold throughout. Thus \mathcal{H}' is tight and $|H_{k-1}| = |H_k| = 3$. \square

Lemma 3.7. Let $G = (V, E)$ be uv -sparse and suppose that there is a tight uv -compatible family in G . Then there is a unique tight uv -compatible family \mathcal{H}_{\max} in G for which $\text{cov}(\mathcal{H}) \subseteq \text{cov}(\mathcal{H}_{\max})$ for all tight uv -compatible families \mathcal{H} of G . In addition, if $\mathcal{H}_{\max} = \{X_1, X_2, \dots, X_k\}$ and $|X_1| \geq |X_2| \geq \dots \geq |X_k|$, then:

- (a) $X_i \cap X_j = \{u, v\}$ for all $1 \leq i < j \leq k$;
- (b) $|X_i| = 3$ for all $2 \leq i \leq k$;
- (c) $N(u, v) \subseteq V(\mathcal{H}_{\max})$.

Furthermore, if $Y \subseteq V$ is tight, $|Y| \geq 4$, $\text{cov}(Y) \not\subseteq \text{cov}(\mathcal{H}_{\max})$, and $Y \cap X_i \neq \emptyset$ for some $1 \leq i \leq k$, then $|Y \cap X_i| = 1$, $|X_i| = 3$, $Y \cap \{u, v\} = \emptyset$, and $Y \cap X_j = \emptyset$ for all $j \neq i$.

Proof. Let $\mathcal{H}_1 = \{X_1, X_2, \dots, X_k\}$ be a tight uv -compatible family in G labeled such that $|X_1| \geq |X_2| \geq \dots \geq |X_k|$ and suppose that $\text{cov}(\mathcal{H}_1)$ is maximal with respect to inclusion. Then Lemmas 3.2 and 3.3 imply that $X_i \cap X_j = \{u, v\}$ holds for all $1 \leq i < j \leq k$ and $|X_i| = 3$ for all $2 \leq i \leq k$. Suppose for a contradiction that $\mathcal{H}_2 = \{Y_1, Y_2, \dots, Y_l\}$ is another tight uv -compatible family whose cover is maximal, labeled so that $|Y_1| \geq |Y_2| \geq \dots \geq |Y_l|$. We will use notation $X_i = \{u, v, x_i\}$ for $2 \leq i \leq k$ and $Y_j = \{u, v, y_j\}$ for $2 \leq j \leq l$. Without loss of generality we can assume that if $|X_1| = |Y_1| = 3$ then $X_1 \neq Y_1$.

We define two uv -compatible families as follows: let

$$\mathcal{H}_\cap = \{Z \subseteq V : |Z| \geq 3 \text{ and } X_i \cap Y_j = Z \text{ for some } X_i \in \mathcal{H}_1, Y_j \in \mathcal{H}_2\};$$

let

$$\mathcal{H}_\cup = \{X_1 \cup Y_1\} \cup \{X_i : 2 \leq i \leq k \text{ and } x_i \notin X_1 \cup Y_1\} \cup \{Y_j : 2 \leq j \leq l \text{ and } y_j \notin X_1 \cup Y_1\}$$

if $|X_1 \cap Y_1| \geq 3$, and

$$\mathcal{H}_\cup = \{X_1\} \cup \{Y_1\} \cup \{X_i : 2 \leq i \leq k \text{ and } x_i \notin X_1 \cup Y_1\} \cup \{Y_j : 2 \leq j \leq l \text{ and } y_j \notin X_1 \cup Y_1\}$$

if $|X_1 \cap Y_1| = 2$.

It is easy to see that \mathcal{H}_\cup and \mathcal{H}_\cap are both uv -compatible. For convenience we rename the families as $\mathcal{H}_\cup = \{A_1, \dots, A_p\}$ and $\mathcal{H}_\cap = \{B_1, \dots, B_q\}$, where $A_1 = X_1 \cup Y_1$ and $B_1 = X_1 \cap Y_1$ if $|X_1 \cap Y_1| \geq 3$, and $A_1 = X_1$ and $A_2 = Y_1$ if $|X_1 \cap Y_1| = 2$. It follows from their construction that $|A_i| = 3$ for all $3 \leq i \leq p$ and $|B_j| = 3$ for all $2 \leq j \leq q$ and also at least one of $|A_2| = 3$, $|B_1| = 3$ holds. It can be seen easily that $p + q = k + l$. We also have $i(\mathcal{H}_1) + i(\mathcal{H}_2) \leq i(\mathcal{H}_\cup) + i(\mathcal{H}_\cap)$, since the family \mathcal{H}_\cup spans all the edges spanned by \mathcal{H}_1 or \mathcal{H}_2 and \mathcal{H}_\cap spans all the edges spanned by both \mathcal{H}_1 and \mathcal{H}_2 . Thus

$$\begin{aligned} \text{val}(X_1) + 3(k-1) - 2(k-1) + \text{val}(Y_1) + 3(l-1) - 2(l-1) &= \text{val}(\mathcal{H}_1) + \text{val}(\mathcal{H}_2) \\ &= i(\mathcal{H}_1) + i(\mathcal{H}_2) \leq i(\mathcal{H}_\cup) + i(\mathcal{H}_\cap) \leq \text{val}(\mathcal{H}_\cup) + \text{val}(\mathcal{H}_\cap) \\ &= \text{val}(A_1) + \max\{\text{val}(A_2), \text{val}(B_1)\} + 3(p-1) - 2(p-1) + 3(q-1) - 2(q-1). \end{aligned}$$

We will show that equality occurs at both ends of the above inequality. Since $k-1+l-1 = p-1+q-1$, it will suffice to show that $\text{val}(X_1) + \text{val}(Y_1) \geq \text{val}(A_1) + \max\{\text{val}(A_2), \text{val}(B_1)\}$. This is immediate if $|X_1 \cap Y_1| = 2$ and follows from Lemma 3.1(b) when $|X_1 \cap Y_1| \geq 3$.

Hence equality must hold throughout the displayed inequality. In particular, \mathcal{H}_\cup and \mathcal{H}_\cap are both tight. Since $\text{cov}(\mathcal{H}_1) \cup \text{cov}(\mathcal{H}_2) \subseteq \text{cov}(\mathcal{H}_\cup)$, the maximality of the covers implies that $\text{cov}(\mathcal{H}_1) = \text{cov}(\mathcal{H}_2)$ which in turn gives $\mathcal{H}_1 = \mathcal{H}_2$.

We have now shown that $\mathcal{H}_1 = \mathcal{H}_{\max}$ is unique and that properties (a) and (b) hold. To see that (c) holds choose $x \in N(u, v)$ and suppose that $x \notin V(\mathcal{H}_{\max})$. Let $\mathcal{H}' = \mathcal{H}_{\max} + \{u, v, x\}$. Then $i(\mathcal{H}') \geq i(\mathcal{H}_{\max}) + 1$ and $\text{val}(\mathcal{H}') = \text{val}(\mathcal{H}_{\max}) + 1$, so \mathcal{H}' is tight and hence contradicts the maximality of \mathcal{H}_{\max} .

To complete the proof we suppose that $Y \subseteq V$ is tight, $|Y| \geq 4$, $\text{cov}(Y) \not\subseteq \text{cov}(\mathcal{H}_{\max})$, and $Y \cap X_i \neq \emptyset$ for some $1 \leq i \leq k$. If $\{u, v\} \subseteq Y$ then $\mathcal{H} = \{Y\}$ would be a uv -compatible family with $\text{cov}(\mathcal{H}) \not\subseteq \text{cov}(\mathcal{H}_{\max})$. This would contradict the maximality of \mathcal{H}_{\max} and hence $\{u, v\} \not\subseteq Y$. If $|Y \cap X_i| \geq 2$ or $|Y \cap X_i| = 1$ and $|X_i| \geq 4$ then Lemma 3.5 would imply that there exists a uv -compatible family \mathcal{H}' with $\text{cov}(\mathcal{H}_{\max}) \cup \text{cov}(Y) \subseteq \text{cov}(\mathcal{H}')$. Hence $|Y \cap X_i| \leq 1$ and $|X_i| = 3$. This tells us that $|Y \cap X_j| \leq 1$ for all j and hence $\text{cov}(Y) \cap \text{cov}(\mathcal{H}_{\max}) = \emptyset$. If $Y \cap \{u, v\} \neq \emptyset$ then putting $\mathcal{H}' = \mathcal{H}_{\max} \cup \{Y \cup \{u, v\}\}$ we have $i(\mathcal{H}') \geq i(\mathcal{H}) + 2|Y| - 2$ and $\text{val}(\mathcal{H}') = \text{val}(\mathcal{H}) + 2|Y| - 2$, so \mathcal{H}' would contradict the maximality of \mathcal{H}_{\max} . Thus $Y \cap \{u, v\} = \emptyset$. If $Y \cap X_j \neq \emptyset$ for some $j \neq i$ then Lemma 3.6 now gives us a tight uv -compatible family \mathcal{H}' with $\text{cov}(\mathcal{H}_{\max}) \cup \text{cov}(Y) \subseteq \text{cov}(\mathcal{H}')$. Hence $Y \cap X_j = \emptyset$ for all $j \neq i$. \square

Note that Lemma 3.7 tells us in particular that if G is uv -sparse and $Y \subseteq V$ is tight with $\{u, v\} \cap Y \neq \emptyset$, then $Y \subseteq X_i$ for some $X_i \in \mathcal{H}_{\max}$.

3.1. The matroid and its rank function. We first remind the reader of the simple $(2, 2)$ -sparse matroid and its rank function. Given a graph $G = (V, E)$, a set $F \subseteq E$ is independent if and only if it is simple and induces a $(2, 2)$ -sparse subgraph. A system $\mathcal{K} = \{H_1, \dots, H_k\}$ of subsets of V is *thin* if $|H_i \cap H_j| \leq 1$ for all pairs $1 \leq i, j \leq k$ with equality only if $|H_i| = 2$ or $|H_j| = 2$. The value of the system \mathcal{K} is given by $\sum_{H_i \in \mathcal{K}} \text{val}(H_i)$.

Now we define the count matroid $\mathcal{M}_{uv}(G)$. Let $G = (V, E)$ be a graph and $u, v \in V$ be distinct vertices of G . We will prove that the family of sets

$$(3.1) \quad \mathcal{I}_G = \{F : F \subseteq E \text{ and } (V, F) \text{ is } uv\text{-sparse}\}$$

defines a matroid $\mathcal{M}_{uv}(G)$ on E and characterise the rank function of this matroid. We need the following definition.

Let $\mathcal{H} = \{X_1, \dots, X_t\}$ be a uv -compatible family and let H_1, \dots, H_k be subsets of V of size at least two. The system $\mathcal{K} = \{\mathcal{H}, H_1, \dots, H_k\}$ is a uv -cover of $F \subseteq E$ if $F \subseteq$

$\text{cov}(\mathcal{H}) \cup \text{cov}(\{H_1, \dots, H_k\})$. It is *thin* if

- (i) $\{H_1, \dots, H_k\}$ is thin,
- (ii) $X_i \cap X_j = \{u, v\}$ for all pairs $1 \leq i, j \leq t$, and
- (iii) $|H_i \cap X_j| \leq 1$ for all $1 \leq i \leq k$, $1 \leq j \leq t$.

The value of the system \mathcal{K} is given by $\text{val}(\mathcal{K}) = \text{val}(\mathcal{H}) + \sum_{i=1}^k \text{val}(H_i)$.

We will show that the rank of an arbitrary subset $F \subseteq E$ in $\mathcal{M}_{uv}(G)$ is given by

$$(3.2) \quad r(F) = \min\{\text{val}(\mathcal{K}) : \mathcal{K} \text{ is a thin cover of } F\}.$$




FIGURE 1. An example of a simple (2,2)-tight graph G which is not independent in $\mathcal{M}_{uv}(G)$.

Let $G = (V, E)$ be the graph shown in Figure 1. It is not difficult to see that G is (2,2)-sparse and simple, and hence E is independent in the simple (2,2)-sparse matroid. We will show that E is not independent in $\mathcal{M}_{uv}(G)$. Consider the following sets: $X_1 = \{u, v, v_1\}$, $X_2 = \{u, v, v_2\}$ and $X_3 = \{u, v, v_3, v_4, v_5\}$. Then $\mathcal{H} = \{X_1, X_2, X_3\}$ is a uv -compatible family of G with $\text{val}(\mathcal{H}) = \text{val}(X_1) + \text{val}(X_2) + \text{val}(X_3) - 2 \cdot 2 = (2 \cdot 3 - 3) + (2 \cdot 3 - 3) + (2 \cdot 5 - 2) - 4 = 10$ and $\text{cov}(\mathcal{H}) = E - v_1v_2$. Hence $i_G(\mathcal{H}) = 11 > \text{val}(\mathcal{H})$ so E is dependent in $\mathcal{M}_{uv}(G)$.

Theorem 3.8. *Let $G = (V, E)$ be a graph and $u, v \in V$ be distinct vertices of G . Then $\mathcal{M}_{uv}(G) = (E, \mathcal{I}_G)$ is a matroid on ground-set E , where \mathcal{I}_G is defined by (3.1). The rank of a set $E' \subseteq E$ in $\mathcal{M}_{uv}(G)$ is equal to*

$$\min\{\text{val}(\mathcal{K}) : \mathcal{K} \text{ is either a thin cover or a thin } uv\text{-cover of } E'\}.$$

Proof. Let $\mathcal{I} = \mathcal{I}_G$, let $E' \subseteq E$ and let $F \subseteq E'$ be a maximal subset of E' in \mathcal{I} . Since $F \in \mathcal{I}$ we have $|F| \leq \text{val}(\mathcal{K})$ for all (uv) -covers \mathcal{K} of E' . We shall prove that there is a thin (uv) -cover \mathcal{K} of E' with $|F| = \text{val}(\mathcal{K})$, from which the theorem will follow.

Let $J = (V, F)$ denote the subgraph defined by the edge set F . First suppose that there is no tight uv -compatible family in J and consider the following cover of F :

$$\mathcal{K}_1 = \{H_1, H_2, \dots, H_k\},$$

where H_1, H_2, \dots, H_t are the maximal tight sets with size at least four in J for some $t \leq k$ and H_{t+1}, \dots, H_k are the pairs of end vertices of edges in $J' = (V, F - \cup_{i=1}^t E(H_i))$. Clearly \mathcal{K}_1 is a cover of F . It is thin by Lemma 3.4. Thus

$$|F| = \sum_{j=1}^k |E_J(H_j)| = \sum_{j=1}^k (2|H_j| - t_j) = \text{val}(\mathcal{K}_1)$$

follows. We claim that \mathcal{K}_1 is a cover of E' . To see this consider an edge $ab = e \in E' - F$. Since F is a maximal subset of E' in \mathcal{I} we have $F + e \notin \mathcal{I}$. By our assumption there is no tight uv -compatible family in J , and hence there must be a tight set X in J with $a, b \in X$. Hence $X \subseteq H_i$ for some $1 \leq i \leq k$ which implies that \mathcal{K}_1 covers e .

Next suppose that there is a tight uv -compatible family in J and consider the following uv -cover of F :

$$\mathcal{K}_2 = \{\mathcal{H}_{\max}, H_1, H_2, \dots, H_k\},$$

where $\mathcal{H}_{\max} = \{X_1, X_2, \dots, X_l\}$ is the uv -compatible family of G for which $\text{cov}(\mathcal{H}_{\max})$ is maximal (given by Lemma 3.7) and H_1, H_2, \dots, H_t are the maximal tight sets with size at least four of $J' = (V, F - E(\mathcal{H}_{\max}))$ for some $t \leq k$ and H_{t+1}, \dots, H_k are the pairs of end vertices of edges in $J'' = (V, F - E(\mathcal{H}_{\max}) - \cup_{i=1}^t E(H_i))$. Then \mathcal{K}_2 is a uv -cover of F . By Lemmas 3.4 and 3.7, the uv -cover \mathcal{K}_2 is thin, and hence

$$|F| = \sum_{i=1}^l |E_J(X_i)| + \sum_{j=1}^k |E_J(H_j)| = \sum_{i=1}^l (2|X_i| - t_i) - 2(l-1) + \sum_{j=1}^k (2|H_j| - t_j) = \text{val}(\mathcal{K}_2).$$

We claim that \mathcal{K}_2 is a uv -cover of E' . As above, let $ab = e \in E' - F$ be an edge. By the maximality of F we have $F + e \notin \mathcal{I}$. Thus either there is a tight set $X \subseteq V$ in J with $a, b \in \text{cov}(X)$ or there is a tight uv -compatible family $\mathcal{H}' = \{Y_1, \dots, Y_t\}$ in J with $a, b \in Y_i$ for some $1 \leq i \leq t$.

In the latter case Lemma 3.7 implies that $\text{cov}(\mathcal{H}') \subseteq \text{cov}(\mathcal{H}_{\max})$ and hence e is covered by \mathcal{K}_2 . In the former case, when $a, b \in X$ for some tight set X in J , we have $|X| \geq 5$ since if $|X| = 2, 3$ or 4 then X induces a complete graph in J and $e = ab$ would be an edge of F . Lemma 3.7 now gives $|X \cap \cup_{i=1}^l X_i| \leq 1$. Then $E(X) \subseteq E(J')$ and hence $X \subseteq H_i$ for some $1 \leq i \leq k$, since every edge of J' induces a tight set and every tight set is contained in a maximal tight set. Thus e is covered by \mathcal{K}_2 , as claimed. \square

4. CHARACTERISATION OF THE uv -COINCIDENT CYLINDER RIGIDITY MATROID

Our aim is to show that the uv -coincident cylinder rigidity matroid $\mathcal{R}_{uv}^{\mathcal{Y}}(G)$ of a graph $G = (V, E)$ is equal to the count matroid $\mathcal{M}_{uv}(G)$. To simplify terminology we will say that G is *independent in $\mathcal{R}_{uv}^{\mathcal{Y}}$* , respectively *$\mathcal{M}_{uv}$* , if E is independent in $\mathcal{R}_{uv}^{\mathcal{Y}}(G)$, respectively $\mathcal{M}_{uv}(G)$.

We first show that independence in $\mathcal{R}_{uv}^{\mathcal{Y}}$ implies independence in \mathcal{M}_{uv} . Let G/uv denote the graph obtained from G by contracting the vertex pair u, v into a new vertex z_{uv} (and deleting the resulting loops and parallel copies of edges). Given a uv -coincident realisation (G, p) of G on \mathcal{Y} we obtain a realisation $(G/uv, p_{uv})$ of G/uv on \mathcal{Y} by putting $p_{uv}(z_{uv}) = p(u) = p(v)$ and $p_{uv}(x) = p(x)$ for all $x \in V - \{u, v\}$. Furthermore, each vector in the kernel of $R_{\mathcal{Y}}(G/uv, p_{uv})$ determines a vector in the kernel of $R_{\mathcal{Y}}(G, p)$ in a natural way. It follows that

$$(4.1) \quad \dim \text{Ker} R_{\mathcal{Y}}(G, p) \geq \dim \text{Ker} R_{\mathcal{Y}}(G/uv, p_{uv}).$$

We can use this fact to prove that independence in $\mathcal{R}_{uv}^{\mathcal{Y}}$ implies independence in \mathcal{M}_{uv} .

Lemma 4.1. *Let $G = (V, E)$ be a graph and let $u, v \in V$ be distinct vertices. If G is independent in $\mathcal{R}_{uv}^{\mathcal{Y}}$ then G is independent in \mathcal{M}_{uv} .*

Proof. Let (G, p) be an independent uv -coincident realisation of G . Independence implies that $i(X) \leq \text{val}(X)$ holds for all $X \subseteq V$ by Theorem 2.1. Since $p(u) = p(v)$, $uv \notin E$ follows.

Let $\mathcal{H} = \{X_1, \dots, X_k\}$ be a uv -compatible family and consider the subgraph $F = (\cup_{i=1}^k X_i, \cup_{i=1}^k E(X_i))$. By contracting the vertex pair u, v in F we obtain the graph F/uv , in which $\mathcal{H}_{uv} = \{X_1/uv, \dots, X_k/uv\}$ is a cover where X_i/uv denotes the set that we get from X_i by identifying u and v . Thus we get $r(F/uv) \leq \sum_{i=1}^k \text{val}(X_i/uv) = \sum_{i=1}^k (2(|X_i| - 1) - t_i)$ by using (3.2). This bound and (4.1) imply that $\dim \text{Ker } R_{\mathcal{Y}}(F, p) \geq \dim \text{Ker } R_{\mathcal{Y}}(F/uv, p_{uv}) \geq 2(|\cup_{i=1}^k X_i| - 1) - \sum_{i=1}^k (2|X_i| - (t_i + 2))$. Since (G, p) is an independent uv -coincident realisation of G , we have

$$\begin{aligned} i_F(\mathcal{H}) = |F| &\leq 2 \left| \bigcup_{i=1}^k X_i \right| - \left(2 \left(\left| \bigcup_{i=1}^k X_i \right| - 1 \right) - \sum_{i=1}^k (2|X_i| - (t_i + 2)) \right) = \\ &= \sum_{i=1}^k (2|X_i| - t_i) - 2(k - 1) = \text{val}(\mathcal{H}). \end{aligned}$$

Thus G is independent in \mathcal{M}_{uv} , as claimed. \square

We next define operations on uv -sparse graphs and use them to show that independence in \mathcal{M}_{uv} implies independence in $\mathcal{R}_{uv}^{\mathcal{Y}}$.

The (two-dimensional versions of) the well-known Henneberg operations are as follows. Let $G = (V, E)$ be a graph. The *0-extension* operation (on a pair of distinct vertices $a, b \in V$) adds a new vertex z and two edges za, zb to G . The *1-extension* operation (on edge $ab \in E$ and vertex $c \in V - \{a, b\}$) deletes the edge ab , adds a new vertex z and edges za, zb, zc .

We shall need the following specialized versions. Let $u, v \in V$ be two distinct vertices. The *0- uv -extension* operation is a 0-extension on a pair a, b with $\{a, b\} \neq \{u, v\}$. The *1- uv -extension* operation is a 1-extension on some edge ab and vertex c for which $\{u, v\}$ is not a subset of $\{a, b, c\}$. The inverse operations are called *0- uv -reduction* and *1- uv -reduction*, respectively.

We will also need two further moves. The *vertex-to- K_4* move deletes a vertex w and substitutes in a copy of K_4 with $V(K_4) \cap V(G) = \{w\}$ and with an arbitrary replacement of edges xw by edges xy with $y \in V(K_4)$. The inverse operation is known as a *K_4 -contraction*. A *vertex-to-4-cycle* move takes a vertex w with neighbours v_1, v_2, \dots, v_k for any $k \geq 2$, splits w into two new vertices w, w' with $w' \notin V(G)$, adds edges $wv_1, w'v_1, wv_2, w'v_2$ and then arbitrarily replaces edges xw with edges xy where $x \in \{v_3, \dots, v_k\}$ and $y \in \{w, w'\}$. The inverse move is known as a *4-cycle-contraction*. The only difference in the specialised versions of these moves are that we require $|V(K_4) \cap \{u, v\}| \leq 1$ in a uv - K_4 -contraction and similarly $|V(C_4) \cap \{u, v\}| \leq 1$ in a uv -4-cycle-contraction.

We first consider the 0-extension and 1-extension operations. It was shown in [8] that these operations preserve independence in $\mathcal{R}^{\mathcal{Y}}$. The same arguments can be used to verify analogous results for $\mathcal{R}_{uv}^{\mathcal{Y}}$.

Lemma 4.2. *Let $G = (V, E)$ be independent in $\mathcal{R}_{uv}^{\mathcal{Y}}$ and suppose that G' is obtained from G by a 0- uv -extension or a 1- uv -extension. Then G' is independent in $\mathcal{R}_{uv}^{\mathcal{Y}}$.*

In the case of 0-extensions we will also need the following result.

Lemma 4.3. *Let (G, p) be a generic realisation of a graph $G = (V, E)$ and $v \in V$. Suppose that $R_{\mathcal{Y}}(G, p)$ has linearly independent rows. Let G' be obtained by performing a 0-extension which adds a new vertex u to G . Put $p'(a) = p(a)$ for all $a \in V$, and put $p'(u) = p(v)$. Then $R_{\mathcal{Y}}(G', p')$ has linearly independent rows.*

Proof. The 0-extension adds 3 rows and 3 columns to $R_{\mathcal{Y}}(G, p)$, the 3 columns being 0 everywhere except the 3 new rows. The genericness of p and the fact that $uv \notin E$ implies the new 3×3 block is invertible. Hence $R_{\mathcal{Y}}(G', p')$ has linearly independent rows so G' is independent in $\mathcal{R}_{uv}^{\mathcal{Y}}$. \square

We next consider the vertex-to-4-cycle operation. It was shown in [9] that this operation preserves independence in $\mathcal{R}^{\mathcal{Y}}$. A similar argument would yield the analogous result for $\mathcal{R}_{uv}^{\mathcal{Y}}$ but we will need a stronger result that a vertex-to-4-cycle move which creates two coincident vertices preserves independence in $\mathcal{R}^{\mathcal{Y}}$.

Lemma 4.4. *Suppose (G, p) is a framework on \mathcal{Y} , $R_{\mathcal{Y}}(G, p)$ has linearly independent rows and $w \in V$ with neighbours v_1, v_2, \dots, v_k . Suppose further that $p(w) - p(v_1), p(w) - p(v_2)$ and $\bar{p}(w)$ are linearly independent where $\bar{p}(w)$ is the projection of $p(w)$ onto the plane $z = 0$. Let G' be obtained by performing a vertex-to-4-cycle operation at w in G such that v_1 and v_2 are both adjacent to w and w' in G' . Put $p'(a) = p(a)$ for all $a \in V - w$ and put $p'(w) = p'(w') = p(w)$. Then $R_{\mathcal{Y}}(G', p')$ has linearly independent rows.*

Proof. We will construct $R_{\mathcal{Y}}(G', p')$ from $R_{\mathcal{Y}}(G, p)$ by a series of simple matrix operations that preserve the independence of the rows.

We first add three zero columns corresponding to w' . We then add three rows corresponding to the edges $w'v_1, w'v_2$ and the vertex w' . Adding these rows increases the rank by 3 since $p(w) - p(v_1), p(w) - p(v_2)$ and $\bar{p}(w)$ are linearly independent so the 3×3 matrix formed by the entries in the columns corresponding to w' and the rows corresponding to $w'v_1, w'v_2, w'$ is non-singular and the rest of the entries in these columns are zero. The matrix M we obtain by this modification has the following form:

$$\begin{array}{cc}
 & \overbrace{\hspace{1.5cm}}^w & \overbrace{\hspace{1.5cm}}^{w'} & \\
 \begin{array}{l} (wv_1) \\ (wv_2) \\ \vdots \\ (wv_i) \\ \vdots \\ (w'v_1) \\ (w'v_2) \\ \vdots \\ w \\ w' \end{array} & \begin{array}{|ccc|} \hline p(w) - p(v_1) & \mathbf{0} & \star \\ p(w) - p(v_2) & \mathbf{0} & \star \\ \vdots & \vdots & \vdots \\ p(w) - p(v_i) & \mathbf{0} & \star \\ \vdots & \vdots & \vdots \\ \mathbf{0} & p(w) - p(v_1) & \star \\ \mathbf{0} & p(w) - p(v_2) & \star \\ \vdots & \vdots & \vdots \\ \hline \bar{p}(w) & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \bar{p}(w) & \mathbf{0} \\ \vdots & \vdots & \vdots \\ \hline \end{array} & = M
 \end{array}$$

To obtain $R_{\mathcal{Y}}(G', p')$ from M we need to modify some of the rows in M corresponding to edges (wv_i) into the form of rows corresponding to edges $(w'v_i)$, i.e. we need to move the entries in the columns of w to the columns of w' and replace them with zeros. We will do this one by one.

Since $(p(w) - p(v_1)), (p(w) - p(v_2))$ and $\bar{p}(w)$ are linearly independent, for every $3 \leq i \leq k$ there exist unique values α, β, γ such that $\alpha(p(w) - p(v_1)) + \beta(p(w) - p(v_2)) + \gamma\bar{p}(w) = (p(w) - p(v_i))$. Now subtract the row of (wv_1) multiplied by α , the row of (wv_2) multiplied by β and the row of w multiplied by γ from the row of (wv_i) in M . Then add the row of $(w'v_1)$ multiplied by α , the row of $(w'v_2)$ multiplied by β and the row of w' multiplied by γ to the same row (and change its label from (wv_i) to $(w'v_i)$) for every neighbour v_i of w'

in G' to obtain $R_{\mathcal{Y}}(G', p')$. These operations also preserve independence, thus we conclude that the rows of $R_{\mathcal{Y}}(G', p')$ are independent. \square

Corollary 4.5. *Let G be independent in $\mathcal{R}_{uv}^{\mathcal{Y}}$ and suppose that G' is obtained from G by a vertex-to-4-cycle operation. Then G' is independent in $\mathcal{R}_{uv}^{\mathcal{Y}}$.*

Proof. We choose a generic uv -coincident realisation (G, p) . Then (G, p) satisfies the hypotheses of Lemma 4.4. Hence G' has a uv -coincident realisation (G', p') such that $R_{\mathcal{Y}}(G', p')$ has linearly independent rows. It follows that every generic uv -coincident realisation is independent. \square

We next consider a generalisation of the vertex-to- K_4 operation. It was shown in [8] that this operation preserves independence in $\mathcal{R}^{\mathcal{Y}}$. We will need an analogous result for uv -coincident realisations.

Lemma 4.6. *Let $G = (V, E)$ be a graph with $|E| = 2|V| - 2$ and let $u, v \in V$ be distinct vertices. Suppose $H \subset G$ is chosen so that either:*

- (a) $u, v \in V(H)$, H is minimally uv -rigid on \mathcal{Y} and G/H is minimally rigid on \mathcal{Y} , or
 - (b) $|\{u, v\} \cap V(H)| \leq 1$, H is minimally rigid on \mathcal{Y} and G/H is minimally uv -rigid on \mathcal{Y} . (Taking z to be the vertex of G/H obtained by contracting H when $\{u, v\} \cap V(H) = z$.)
- Then G is uv -rigid on \mathcal{Y} .*

Proof. (a) Let $|V| = n$, $|V(H)| = r$ and consider $R_{\mathcal{Y}}(G, p)$ where (G, p) is a generic uv -coincident framework on \mathcal{Y} and $p = (p(v_1), p(v_2), \dots, p(v_n))$. By reordering rows and columns if necessary we can write $R_{\mathcal{Y}}(G, p)$ in the form

$$\begin{pmatrix} R_{\mathcal{Y}}(H, p|_H) & 0 \\ M_1(p) & M_2(p) \end{pmatrix}$$

where $M_2(p)$ is a square matrix with $3(n - r)$ rows.

Suppose, for a contradiction, that G is not uv -rigid. Then there exists a vector $m \in \ker R_{\mathcal{Y}}(G, p)$ which is not an infinitesimal isometry of \mathcal{Y} . Since $(H, p|_H)$ is uv -rigid we may suppose that $m = (0, \dots, 0, m_{r+1}, \dots, m_n)$. Consider the realisation (G, p') where $p' = (p(v_r), p(v_r), \dots, p(v_r), p(v_{r+1}), \dots, p(v_n))$ and define the realisation $(G/H, p^*)$ by setting $p^* = (p(v_r), p(v_{r+1}), \dots, p(v_n))$. Since p^* is generic, $(G/H, p^*)$ is infinitesimally rigid on \mathcal{Y} by assumption.

Now, $M_2(p)$ is square with the nonzero vector $(m_{r+1}, \dots, m_n) \in \ker M_2(p)$. Hence $\text{rank } M_2(p) < 3(n - r)$. Since p is generic, we also have $\text{rank } M_2(p') < 3(n - r)$ and hence there exists a nonzero vector $m' \in \ker M_2(p')$. Therefore we have

$$(R_{\mathcal{Y}}(G/H, p^*)) \begin{pmatrix} 0 \\ m' \end{pmatrix} = \begin{pmatrix} p(v_r) & 0 \\ \star & M_2(p') \end{pmatrix} \begin{pmatrix} 0 \\ m' \end{pmatrix} = 0,$$

contradicting the infinitesimal rigidity of $(G/H, p^*)$.

(b) A similar proof holds. We choose a generic uv -coincident framework (G, p) , a vector $m \in \ker R_{\mathcal{Y}}(G, p)$ which is not an infinitesimal isometry of \mathbb{R}^3 , and uv -coincident realisations (G, p') and $(G/H, p^*)$ as above. We then use the facts that H is rigid on \mathcal{Y} and G/H is uv -rigid on \mathcal{Y} to obtain a contradiction. \square

We next consider the 0- uv -reduction, 1- uv -reduction, uv - K_4 -contraction and uv -4-cycle contraction operations.

Lemma 4.7. *Let $G = (V, E)$ be a graph and let $u, v \in V$ be distinct vertices. Suppose that $|E| = 2|V| - 2$, G is independent in \mathcal{M}_{uv} , and $d(w) \geq 3$ for all $w \in V$. Then either there is a vertex $z \in V - \{u, v\}$ with $d(z) = 3$ and $|N(z) \cap \{u, v\}| \leq 1$ or there is a 4-cycle in G which contains both u and v .*

Proof. Since $|E| = 2|V| - 2$ and $d(w) \geq 3$ for all $w \in V$, there are at least 4 vertices of degree 3. Since G is independent in \mathcal{M}_{uv} , G has at most two vertices which are adjacent to both u and v . Hence, if there is no vertex $z \in V - \{u, v\}$ with $d(z) = 3$ and $|N(z) \cap \{u, v\}| \leq 1$, then the vertices of degree 3 must induce a C_4 in G which contains both u and v . \square

Lemma 4.8. *Let $G = (V, E)$ be a graph and let $u, v \in V$ be distinct vertices. Suppose that G is independent in \mathcal{M}_{uv} , and there are vertices a, b such that a, u, b, v is a cycle in G . Then the uv -4-cycle contraction which merges u and v results in a simple graph G' which is $(2, 2)$ -sparse.*

Proof. The independence of G in \mathcal{M}_{uv} implies that there is no vertex other than a, b that is adjacent with both u and v . Thus G' is simple. Suppose G is not $(2, 2)$ -sparse. Then there exists a $(2, 2)$ -tight set X in G that contains u, v and exactly one of a and b , say a . Let $\{X, \{u, v, b\}\} = \mathcal{H}$. Then $i(\mathcal{H}) = 2|X| - 2 + 2$ and $\text{val}(\mathcal{H}) = 2|X| - 2 + 3 - 2$ which contradicts the independence of G in \mathcal{M}_{uv} . \square

Lemma 4.9. *Let $G = (V, E)$ be a graph and let $u, v \in V$ be distinct vertices. Suppose that G is independent in \mathcal{M}_{uv} and let $z \in V - \{u, v\}$ with $N(z) = \{v_1, v_2, v_3\}$ and $|N(z) \cap \{u, v\}| \leq 1$. Then either:*

- (a) *there is a 1-reduction at z which leads to a graph which is independent in \mathcal{M}_{uv} , or*
- (b) *z and its neighbours induce a copy of K_4 in G , or*
- (c) *$v_i \in \{u, v\}$ and $v_j v_k \in E$ for some $\{i, j, k\} = \{1, 2, 3\}$, and there is a tight uv -compatible family $\{X_1, X_2, \dots, X_k\}$ in G such that $X_1 = N(z) \cup \{u, v, z\}$ and $i(X_1) \geq 2|X_1| - 4$.*

Proof. Suppose (a) does not occur. Then, for all $1 \leq i < j \leq 3$, either $v_i v_j \in E$, or there exists a tight uv -compatible family \mathcal{H}_{ij} in $G - z$ with $v_i v_j \in \text{cov}(\mathcal{H}_{ij})$ or there exists a tight set X_{ij} in $G - z$ with $\{v_i, v_j\} \subset X_{ij}$ and $\{u, v\} \not\subset X_{ij}$. If the second alternative occurs we may assume that \mathcal{H}_{ij} has been chosen to be the unique tight uv -compatible family in $G - z$ with maximal cover. If $G[v_1, v_2, v_3] \cong K_3$ then (b) occurs. So we may assume that $v_1 v_2 \notin E$.

We first show that $v_i v_j \notin E$ and that \mathcal{H}_{ij} exists for some $1 \leq i < j \leq 3$. Suppose \mathcal{H}_{12} does not exist. Then X_{12} exists. If $v_3 \in X_{12}$ then $X_{12} + z$ contradicts the independence of G in \mathcal{M}_{uv} . Hence $v_3 \notin X_{12}$. If $v_1 v_3, v_2 v_3 \in E$ then $X_{12} \cup \{v_3, z\}$ contradicts the independence of G in \mathcal{M}_{uv} . Hence suppose that $v_1 v_3 \notin E$. If X_{13} exists, then $X_{12} \cup X_{13} \cup \{z\}$ contradicts the independence of G in \mathcal{M}_{uv} . Hence \mathcal{H}_{13} exists. Relabeling if necessary we assume that $\mathcal{H}_{12} = \{X_1, X_2, \dots, X_k\}$ exists.

Since $v_1 v_2 \in \text{cov}(\mathcal{H}_{12})$ we have $v_1, v_2 \in X_i$ for some $1 \leq i \leq k$. If $v_3 \in X_i$ then $|X_i| \geq 4$, since $|N(z) \cap \{u, v\}| \leq 1$, and the uv -compatible family obtained from \mathcal{H}_{12} by replacing X_i by $X_i + z$ will contradict the independence of G in \mathcal{M}_{uv} . Hence $v_3 \notin X_i$.

Suppose that $\{v_1, v_2\} \cap \{u, v\} = \emptyset$. Then $|X_i| \geq 4$ and neither $v_1 v_3$ nor $v_2 v_3$ are covered by \mathcal{H}_{12} . The maximality of $\text{cov}(\mathcal{H}_{12})$ now implies that \mathcal{H}_{13} and \mathcal{H}_{23} do not exist. If $v_1 v_3, v_2 v_3 \in E$, then the uv -compatible family obtained from \mathcal{H}_{12} by replacing X_i by $X_i + v_3$ will contradict the maximality of $\text{cov}(\mathcal{H}_{12})$. Relabeling if necessary, we may suppose that $v_1 v_3 \notin E$, and hence X_{13} exists. Then $X_i \cap X_{13} \neq \emptyset$, $|X_i| \geq 4$, $|X_{13}| \geq 4$ and $v_1 v_3 \in \text{cov}(X_{13}) \setminus \text{cov}(\mathcal{H}_{12})$. This contradicts the final part of Lemma 3.7. Hence $\{v_1, v_2\} \cap \{u, v\} \neq \emptyset$ and we may assume, without loss of generality, that $u = v_1$.

If $v_3 \notin V(\mathcal{H}_{12})$, then Lemma 3.7(c) implies that $v_1v_3 \notin E$ and hence X_{13} exists. This contradicts the final part of Lemma 3.7 since $u \in X_{13} \cap X_i$. Hence $v_3 \in X_j$ for some $X_j \in \mathcal{H}_{12} - X_i$. The final part of Lemma 3.7 now implies that X_{23} does not exist and hence $v_2v_3 \in E$.

Let $X = X_i \cup X_j \cup \{z\}$ and $\mathcal{H} = (\mathcal{H}_{12} \setminus \{X_i, X_j\}) \cup \{X\}$. Then the facts that G is independent in \mathcal{M}_{uv} and \mathcal{H}_{12} is tight imply that $|X_i| = 3 = |X_j|$ (so $X = N(z) \cup \{u, v, z\}$), and that \mathcal{H} is a tight uv -compatible family in G with $i(X) \geq 2|X| - 4$. \square

Lemma 4.10. *Let $G = (V, E)$ be a graph and let $u, v \in V$ be distinct vertices. Suppose that G is independent in \mathcal{M}_{uv} , $\mathcal{H} = \{X_1, X_2, \dots, X_k\}$ is a tight uv -compatible family in G and that $\mathcal{H} - X_i$ is not tight for all $1 \leq i \leq k$. Then either:*

- (a) $k = 1$ and X_1 is tight;
- (b) $k = 2$, $|X_1| = |X_2| = 3$ and $i(X_1) = i(X_2) = 2$;
- (c) $k = 2$, $|X_1| \geq 4$, $i(X_1) = 2|X_1| - 3$, $|X_2| = 3$ and $i(X_2) = 2$; or
- (d) $k = 2$, $|X_i| \geq 4$ and $i(X_i) = 2|X_i| - 3$ for all $i \in \{1, 2\}$.

Proof. We have $i(\mathcal{H} - X_i) < \text{val}(\mathcal{H} - X_i)$, and hence $i(X_i) \geq 2|X_i| - 3$ if $|X_i| \geq 4$ and $i(X_i) = 2$ if $|X_i| = 3$. The fact that G is independent in \mathcal{M}_{uv} and \mathcal{H} is tight now imply that $k = 1$ or 2 and that the sets in \mathcal{H} satisfy the assertions in the lemma. \square

Note that if alternative (d) holds then $X_1 \cup X_2$ is tight so we can reduce to alternative (a).

Lemma 4.11. *Let $G = (V, E)$ be a graph and let $u, v \in V$ be distinct vertices. Suppose that G is independent in \mathcal{M}_{uv} and that there exists a subgraph H of G isomorphic to K_4 . Then either:*

- (a) there is a vertex $x \in V - V(H)$ such that $|N(x) \cap V(H)| = 2$,
- (b) $|V(H) \cap \{u, v\}| = 1 = |N(V(H)) \cap \{u, v\}|$,
- (c) there is a tight uv -compatible family $\{X_1, X_2, \dots, X_k\}$ in G such that $X_1 = V(H) \cup \{u, v\}$, $|X_1| = 6$ and $i(X_1) = 8$,
- (d) there is a tight uv -compatible family $\{X_1, X_2, \dots, X_k\}$ in G such that $X_1 = V(H) \cup \{u, v, a\}$ for some $a \in V - (V(H) \cup \{u, v\})$, $|X_1| = 6$ and $i(X_1) = 8$, or
- (e) the contraction of H gives a graph G' which is independent in \mathcal{M}_{uv} .

Proof. Let G' be the result of a K_4 contraction applied to H with w being the contracted vertex. Suppose that (a), (b) and (e) fail. It is easy to check that G' is $(2, 2)$ -sparse. Since (a) fails, G' is simple. Since (b) fails, $uv \notin E(G')$. Since (e), there is a uv -compatible family $\mathcal{H} = \{X_1, X_2, \dots, X_k\}$ for which $\text{val}(\mathcal{H}) < i(\mathcal{H})$ and $w \in V(\mathcal{H})$. Without loss of generality we may assume $w \in X_1$. If $|X_1| \geq 4$ then we get a contradiction as the uv -compatible family $\mathcal{H}' = \{(H_1 - w) \cup V(H), X_2, \dots, X_k\}$ of G violates independence. If $|X_1| = 3$ and $V(H) \cap \{u, v\} = \emptyset$ then \mathcal{H}' is the uv -compatible family described in (c). Finally if $|X_1| = 3$ and $V(K_4) \cap \{u, v\} = u$ then $\mathcal{H}'' = \{V(H) \cup \{u, v, a\}, X_2, \dots, X_k\}$ is the uv -compatible family described in (d). \square

Lemma 4.12. *Let $G = (V, E)$ be a graph and let $u, v \in V$ be distinct vertices. Suppose that G is independent in \mathcal{M}_{uv} , $z \in V - \{u, v\}$ is a vertex of degree 3 with $N(z) = \{v_1, v_2, v_3\}$, $|N(z) \cap \{u, v\}| \leq 1$ and $G[N(z) + z]$ is isomorphic to K_4 . Suppose further that there is a vertex $x \in V - \{z, v_1, v_2, v_3\}$ such that $N(x) \cap N(z) = \{v_2, v_3\}$ and $\{v_1, x\} \neq \{u, v\}$. Then the uv -4-cycle contraction operation which contracts x and z into a single vertex x leads to a graph G' which is independent in \mathcal{M}_{uv} .*

Proof. Suppose G' is not independent in \mathcal{M}_{uv} . Since $G' = G - z + v_1x$ and $xv_1 \notin E$, there exists either a tight uv -compatible family \mathcal{H} in $G - z$ with $xv_1 \in \text{cov}(\mathcal{H})$, or a tight set X in $G - z$ with $\{x, v_1\} \subset X$. Set $Y = \{z, v_1, v_2, v_3, x\}$. Then Y is tight in G .

Suppose X exists. Then $X \cup Y$ and $X \cap Y$ are tight by Lemma 3.4. Since $\{v_1, x\} \subseteq X \cap Y$ and no proper subset of Y containing v_1 and x is tight, we have $X \cap Y = Y$. This implies that $z \in X$ contradicting the choice of X . Hence $\mathcal{H} = \{X_1, X_2, \dots, X_k\}$ exists.

Since $xv_1 \in \text{cov}(\mathcal{H})$, we may assume, without loss of generality, that $x, v_1 \in X_1$. Then $x, v_1 \in X_1 \cap Y$. Since $|\{u, v\} \cap Y| \leq 1$ by the hypotheses of the lemma, Lemma 3.5 implies that $X_1 \cap Y$ is tight. Since no proper subset of Y containing v_1 and x is tight we have $X_1 \cap Y = Y$. This implies that $z \in X_1$ and contradicts the choice of \mathcal{H} . \square

Theorem 4.13. *Let $G = (V, E)$ be a graph and let $u, v \in V$ be distinct vertices. Then G is independent in $\mathcal{R}_{uv}^{\mathcal{Y}}$ if and only if G is independent in \mathcal{M}_{uv} .*

Proof. Necessity follows from Lemma 4.1. Now suppose that G is independent in \mathcal{M}_{uv} . We prove that G is independent in $\mathcal{R}_{uv}^{\mathcal{Y}}$ by induction on $|V|$. It is straightforward to check that G is independent in $\mathcal{R}_{uv}^{\mathcal{Y}}$ when $|V| \leq 4$. Hence we may assume that $|V| \geq 5$. By extending $|E|$ to a base of $\mathcal{M}_{uv}(K_{|V|})$ if necessary, we may also assume that $|E| = 2|V| - 2$.

Case 1. G contains a vertex of degree 2. First suppose that u has degree 2. Then $G - u$ is $(2, 2)$ -sparse. Hence, by Theorem 2.1, $R_{\mathcal{Y}}(G - u, p)$ has linearly independent rows for any generic p . We can now use Lemma 4.3 to show that G is independent in $\mathcal{R}_{uv}^{\mathcal{Y}}$.

Now, suppose that there is a vertex $w \in V - \{u, v\}$ with $d(w) = 2$. Let $N(w) = \{a, b\}$. Clearly, $a \neq b$ holds. If $\{a, b\} = \{u, v\}$ then let $\mathcal{H} = \{\{u, v, w\}, \{V - w\}\}$, where $|V - w| \geq 4$. We have

$$2|V| - 2 = |E| = i_E(\mathcal{H}) \leq \text{val}(\mathcal{H}) = 2 \cdot 3 - 3 + 2(|V| - 1) - 2 - 2 = 2|V| - 3,$$

a contradiction. Hence $\{a, b\} \neq \{u, v\}$, which implies that the 0 - uv -reduction operation can be applied at w to obtain a graph $G' = (V - w, E')$ that is independent in \mathcal{M}_{uv} and satisfies $|E'| = 2|V - w| - 2$. By induction, G' is independent in $\mathcal{R}_{uv}^{\mathcal{Y}}$. Now Lemma 4.2 implies that G is independent in $\mathcal{R}_{uv}^{\mathcal{Y}}$.

Case 2. There is a 4-cycle in G containing u and v . By Lemma 4.8, we may apply a uv -4-cycle-contraction (contracting u and v) to obtain a graph H which is simple and $(2, 2)$ -sparse. Theorem 2.1 implies that any generic realisation (H, p) on \mathcal{Y} is infinitesimally rigid. Now we can use Lemma 4.4 to show that G is independent in $\mathcal{R}_{uv}^{\mathcal{Y}}$.

Henceforth we assume that Cases 1 and 2 do not occur.

Case 3. There is a proper tight set X containing u and v . Since Case 1 does not occur, we may suppose X is a maximal proper tight set (where proper means $X \neq V$ and maximal means there is no vertex $w \in V - X$ with more than one neighbour in X). Now by the maximality of X , G/X is simple and $|V - X| \geq 3$. Hence G/X is $(2, 2)$ -tight. Theorem 2.1 implies that any generic framework $(G/X_1, p)$ on \mathcal{Y} is infinitesimally rigid. We may now apply Lemma 4.6(a) to show that G is independent in $\mathcal{R}_{uv}^{\mathcal{Y}}$.

Henceforth we may assume that Case 3 does not occur.

Case 4. There is a degree three vertex z in G which is contained in a subgraph $H \cong K_4$, and a vertex $x \in V - V(H)$ such that $|V(H) \cap N(x)| = 2$. If $\{u, v\} \not\subset V(H) \cup \{x\}$ then we may apply Lemma 4.12 to find a graph G' which is independent in \mathcal{M}_{uv} . We can now use Corollary 4.5 to show that G is independent in $\mathcal{R}_{uv}^{\mathcal{Y}}$. Thus we may suppose that $\{u, v\} \subset V(H) \cup \{x\}$. Then $H \cup \{x\}$ is tight. This contradicts the assumption that Case 1 (if $H \cup \{x\} = V$) or Case 3 (if $H \cup \{x\} \neq V$) do not occur.

A vertex z of degree 3 in G is *bad* if either

- $z \in \{u, v\}$, or
- z is adjacent to both u and v , or
- $X = N(z) \cup \{u, v, z\}$ satisfies alternative (c) of Lemma 4.9 and $i(X) \geq 2|X| - 3$, or
- z belongs to a subgraph $H \cong K_4$ satisfying alternative (b) of Lemma 4.11.

Otherwise we say that z is *good*.

Case 5. All degree three vertices are bad. We may use Lemma 4.7 and the fact that Case 2 does not occur to deduce there exists a degree three vertex $v_1 \in V \setminus \{u, v\}$ with $|N(v_1) \cap \{u, v\}| \leq 1$. Since v_1 is bad $X = N(v_1) \cup \{u, v, v_1\}$ satisfies alternative (c) of Lemma 4.9 and $i(X) \geq 2|X| - 3$, or v_1 belongs to a subgraph $H \cong K_4$ satisfying alternative (b) of Lemma 4.11. If the first alternative occurs then we may use the facts that G is independent in \mathcal{M}_{uv} and Case 2 does not occur to deduce that $i(X) = 2|X| - 3 = 7$. It follows that, in both cases, we may relabel the vertices of $H = G[N(v_1) \cup \{u, v, v_1\}]$ such that H is one of the graphs shown in Figure 2.




FIGURE 2. The two alternatives for H .

The fact that G is $(2, 2)$ -sparse implies that, in both cases, there exists a (necessarily bad) degree three vertex $v_4 \in V \setminus V(H)$. Since Case 2 does not occur, v_4 is not adjacent to both u and v . Hence v_4 also belongs to a subgraph H' which is isomorphic to one of the graphs shown in Figure 2. Since Case 2 does not occur, $v_4 \in V(H')$. Since G is $(2, 2)$ -sparse, $V(H) \cap V(H') = \{u, v, v_3\}$. Now $H \cup H'$ is one of the graphs shown in Figure 3. Since all three graphs are tight, we may use the fact that Case 3 does not occur to deduce that $G = H \cup H'$. The fact that Case 1 does not occur now tells us that G is not the graph in Figure 3(a). The graph in Figure 3(b) cannot be equal to G since $X = N(v_1) \cup \{u, v, v_1\}$ does not belong to a tight uv -compatible family (so v_1 is not bad). Hence G is as shown in Figure 3(c).




FIGURE 3. The three alternatives for G .

We will complete the discussion of this case by showing that G is minimally uv -rigid on \mathcal{Y} . Let (G, p) be a generic uv -coincident realisation of G on \mathcal{Y} and m be an infinitesimal motion of (G, p) with $m(u) = 0$. Since K_4 is rigid, $m(w) = 0$ for all $w \in V(H) - v$. In particular $m(v_3) = 0$ and hence $m(w) = 0$ for all $w \in V$.

Case 6. None of the previous cases occur. Let z_1, z_2, \dots, z_k be the good degree three vertices in G . If the edge set of some 1-reduction of G at z_i is independent in \mathcal{M}_{uv} then we may apply induction to the reduced graph and then apply Lemma 4.2 to deduce that G is independent in $\mathcal{R}_{uv}^{\mathcal{Y}}$. Hence we may assume that alternative (b) or (c) of Lemma 4.9 holds for z_i . Similarly, if alternative (b) of Lemma 4.9 holds for z_i and z_i is contained in a K_4 -subgraph whose contraction results in a graph which is independent in \mathcal{M}_{uv} , then we may apply induction to the reduced graph and then apply Lemma 4.6(b) to deduce that G is independent in $\mathcal{R}_{uv}^{\mathcal{Y}}$. Since Case 4 does not occur, it follows that, for every good degree 3 vertex z_i , there exists a tight uv -compatible family \mathcal{H}_i as described in alternative (c) or (d) of Lemma 4.11, or alternative (c) of Lemma 4.9. We may assume that the first alternatives holds for all $1 \leq i \leq k$ and that the second alternative holds for $l+1 \leq i \leq k$. Choose a subgraph $H_i \cong K_4$ that contains z_i and satisfies alternative (c) or (d) of Lemma 4.11 for each $1 \leq i \leq k$ and let X_i be the element of \mathcal{H}_i which contains $V(H_i)$. Let $X_i = \{z_i, u, v\} \cup N(z_i)$ for each $l+1 \leq i \leq k$. With these definitions we have $i(X_i) = 2|X_i| - 4$ for all $1 \leq i \leq k$.

Let $X = \bigcup_{i=1}^k X_i$. We will show by induction that $i(X) \geq 2|X| - 4$. Suppose that we have $i(X') \geq 2|X'| - 4$ for some $X' = \bigcup_{i=1}^s X_i$ and some $1 \leq s \leq k$. If $i(X' \cup X_{s+1}) \leq 2|X' \cup X_{s+1}| - 5$, then Lemma 3.1(a) implies that $i(X' \cap X_{s+1}) \geq 2|X' \cap X_{s+1}| - 3$, contradicting the fact that no subset of X_{s+1} that contains u, v (in each of the three possibilities for X_{s+1}) satisfies this inequality.

We may apply Lemma 4.10 to a minimal tight uv -compatible subfamily of \mathcal{H}_i for all $1 \leq i \leq k$, and use the facts that Cases 2 and 3 do not occur to deduce that alternative (c) of Lemma 4.10 must hold. Hence there exist sets Y_i and $\{u, v, y_i\}$ in $\mathcal{H}_i - X_i$ with $i(Y_i) = 2|Y_i| - 3$ and $i(\{u, v, y_i\}) = 2$. Lemma 3.2(b) implies that $Y_i \cap X_i = \{u, v\} = Y_i \cap \{u, v, y_i\}$ for all $1 \leq i \leq k$. The fact that we are not in Case 2 also implies that $y_i = y_j = y$, say, for all $1 \leq i \leq j \leq k$. Let $Y = \bigcap_{i=1}^k Y_i$. Then $Y \cap X = \{u, v\}$ and $y \notin Y$. We can now use Lemma 3.1(a) and the fact that Case 3 does not occur to prove inductively that $i(Y) = 2|Y| - 3$.

Let $W = V \setminus X$. Since $i(W) \leq 2|W| - 2$ there is an integer t for which $i(W) = 2|W| - 2 - t$. Since $i(Y) = 2|Y| - 3$ and G is $(2, 2)$ -sparse, there are at least 3 edges from $Y \setminus \{u, v\}$ to $\{u, v\}$. Since $Y \setminus \{u, v\} \subseteq W$, $y \in W \setminus Y$ and there are two edges from y to $\{u, v\}$, we have at least five edges between $\{u, v\}$ and W . Note that the definition of X tells us that all degree 3 vertices in W are bad.

Suppose that no (bad) degree three vertex $z \in W$ is contained in a set $X \subseteq V$ which satisfies alternative (c) of Lemma 4.9 and has $i(X) \geq 2|X| - 3$, or a subgraph $H \cong K_4$ that satisfies alternative (b) of Lemma 4.11. Then every (bad) degree three vertex in W is adjacent to both u and v . Since Case 2 does not occur we have at most one degree three vertex in W . Since $i(Y) \geq 2|Y| - 4$, we have $|E| - |E(Y)| - |E(W)| \leq 4 + t$. The total degree of the vertices in W is at most $2(2|W| - 2 - t) + 4 + t = 4|W| - t$. Since there is at most one degree three vertex in W , $t \leq 1$. If $t = 0$, then W is tight and $W + u + v$ violates sparsity. Hence $t = 1$ and $W + u + v$ is a proper tight set which contradicts the fact that Case 3 does not occur.

Now consider the case when there is a (bad) degree three vertex $z \in W$ which is contained in a set $X \subseteq V$ which satisfies alternative (c) of Lemma 4.9 and has $i(X) \geq 2|X| - 3$, or a subgraph $H_1 \cong K_4$ that satisfies alternative (b) of Lemma 4.11. Then $H = G[N(z) \cup \{u, v, z\}]$ is isomorphic to one of the graphs shown in Figure 2, with $v_1 = z$. Since Case 2

does not occur we have $v_3 = y$. The facts that $y \notin Y$ and no $Z \subseteq V(H) - z$, with $\{u, v\} \subset Z$, has $i(Z) = 2|Z| - 3$ imply that $Y \cap V(H)$ is a proper subset of both Y and $V(H)$. Lemma 3.1(a) now implies that $Y \cup V(H)$ is tight. Since $Y \cup V(H) \neq V$ this contradicts the fact that Case 3 does not occur. \square

4.1. A deletion-contraction characterisation of uv -rigidity.

Theorem 4.14. *Let $G = (V, E)$ be a graph and let $u, v \in V$ be distinct vertices. Then G is uv -rigid if and only if $G - uv$ and G/uv are both rigid.*

Proof. Necessity follows from the fact that an infinitesimally rigid uv -coincident realisation of G is an infinitesimally rigid realisation of $G - uv$, and also gives rise to an infinitesimally rigid realisation of G/uv by (4.1).

To prove sufficiency, suppose, for a contradiction, that $G - uv$ and G/uv are both rigid but G is not uv -rigid. By Theorems 3.8 and 4.13 this implies that there is a thin cover \mathcal{K} of $G - uv$ with $\text{val}(\mathcal{K}) \leq 2|V| - 3$. If \mathcal{K} consists of subsets of V only, then $r(G - uv) \leq 2|V| - 3$ follows, which contradicts the fact that $G - uv$ is rigid.

Hence $\mathcal{K} = \{\mathcal{H}, H_1, \dots, H_k\}$, where $\mathcal{H} = \{X_1, \dots, X_l\}$ is a uv -compatible family. Contract the vertex pair u, v in G into a new vertex z_{uv} . This gives rise to a cover

$$\mathcal{K}' = \{X'_1, \dots, X'_l, H_1, \dots, H_k\}$$

of G/uv , where X'_j is obtained from X_j by replacing u, v by z_{uv} , for $1 \leq j \leq l$. Then we obtain

$$\begin{aligned} & \sum_{i=1}^k (2|H_i| - t_{H_i}) + \sum_{j=1}^l (2|X'_j| - t(X'_j)) \leq \sum_{i=1}^k (2|H_i| - t_{H_i}) + \\ & + \sum_{j=1}^l (2|X_j| - t(X_j)) - 2l = \text{val}(\mathcal{K}) - 2 \leq 2|V| - 3 - 2 = 2(|V| - 1) - 3, \end{aligned}$$

which implies that G/uv is not rigid, a contradiction. This completes the proof. \square

A similar proof can be used to verify the following more general result:

Theorem 4.15. *Let $G = (V, E)$ be a graph and let $u, v \in V$ be distinct vertices. Then $r_{uv}(G) = \min\{r(G - uv), r(G/uv) + 2\}$.*

Theorems 4.14 and 4.15 show that the polynomial-time algorithms for computing the rank of a count matroid (see e.g. [1, 7]) can be used to test whether G is uv -rigid, or more generally, to compute $r_{uv}(G)$.

5. VERTEX SPLITTING AND GLOBAL RIGIDITY

Suppose $G = (V, E)$ is a graph with $V = \{v_1, v_2, \dots, v_n\}$ and (G, p) is a realisation of G on a family of (not necessarily distinct) concentric cylinders $\mathcal{Y} = \mathcal{Y}_1 \cup \mathcal{Y}_2 \cup \dots \cup \mathcal{Y}_n$ such that $p(v_i) \in \mathcal{Y}_i$ for $1 \leq i \leq n$. We say that (G, p) is *globally rigid* if every equivalent framework (G, q) on \mathcal{Y} , with $q(v_i) \in \mathcal{Y}_i$ for all $1 \leq i \leq n$, is congruent to (G, p) .

Let $G = (V, E)$ be a graph and v_1 be a vertex of G with neighbours v_2, v_3, \dots, v_t . A *vertex split* of G at v_1 is a graph \tilde{G} which is obtained from G by deleting the edges $v_1v_2, v_1v_3, \dots, v_1v_k$ and adding a new vertex v_0 and new edges $v_0v_1, v_0v_2, \dots, v_0v_k$, for some $2 \leq k \leq t$. We will refer to the new edge v_0v_1 as the *bridging edge* of the vertex split. We will show in this section that a vertex splitting operation, in which the bridging edge is redundant, preserves generic global rigidity on the cylinder.

Given a map $p : V \rightarrow \mathbb{R}^{3n}$, there is a unique family of concentric cylinders \mathcal{Y} with $p(v_i) \in \mathcal{Y}_i$ for all $1 \leq i \leq n$ as long as $p(v_i)$ does not lie on the z -axis for all $1 \leq i \leq n$. We will refer to \mathcal{Y} as the family of concentric cylinders induced by p and denote it by \mathcal{Y}^p . We shall need the following analogue of [2, Theorem 13].

Lemma 5.1. *If (G, p) is infinitesimally rigid and globally rigid on \mathcal{Y} , then there exists an open neighbourhood N_p of p on \mathcal{Y} such that for any $q \in N_p$ the framework (G, q) is infinitesimally rigid and globally rigid on \mathcal{Y} .*

Proof. Suppose $|V| \geq 5$ and that for any open neighbourhood N_p , there is a $p^* \in N_p$ such that the framework (G, p^*) is not globally rigid on \mathcal{Y} . Then there is a convergent sequence (G, p^k) of non-globally rigid frameworks converging to (G, p) . For each framework (G, p^k) , let (G, q^k) be an equivalent but non-congruent realisation on \mathcal{Y} . We may assume that (G, p^k) and (G, q^k) are in standard position (that is $p^k(v_1) = q^k(v_1) = (0, 1, 0)$ assuming, without loss of generality, that $r_1 = 1$). By the compactness of $\mathbb{R}^{3|V|}$, there is a convergent subsequence (G, q^m) converging to a limiting framework (G, q) . As the limits of the respective sequences, (G, q) must be equivalent to (G, p) .

If (G, q) is not congruent to (G, p) then we contradict the global rigidity of (G, p) . So (G, p) and (G, q) are congruent, i.e. we can transform q to p by a reflection in the plane $x = 0$, a reflection in the plane $z = 0$ or a combination of the two. We apply this same congruence to all the (G, q^m) to obtain a sequence (G, r^m) converging to (G, p) with (G, r^m) being equivalent but not congruent to (G, p^m) for each m .

We next show that $p^m - r^m$ gives an infinitesimal motion of $(G, \frac{p^m + r^m}{2})$ on $\mathcal{Y}^{\frac{p^m + r^m}{2}}$. For each edge $v_i v_j$ we have

$$\begin{aligned} & \left(\frac{p^m(v_i) + r^m(v_i)}{2} - \frac{p^m(v_j) + r^m(v_j)}{2} \right) \cdot ((p^m(v_i) - r^m(v_i)) - (p^m(v_j) - r^m(v_j))) \\ &= \frac{1}{2} ((p^m(v_i) - p^m(v_j)) + (r^m(v_i) - r^m(v_j))) \cdot ((p^m(v_i) - p^m(v_j)) - (r^m(v_i) - r^m(v_j))) \\ &= \frac{1}{2} ((p^m(v_i) - p^m(v_j))^2 - (r^m(v_i) - r^m(v_j))^2) = 0. \end{aligned}$$

Recall that $\bar{p}_m(v_i)$ and $\bar{r}_m(v_i)$ denote the projections of $p_m(v_i)$ and $r_m(v_i)$ onto the plane $z = 0$. Since $p_m(v_i)$ and $r_m(v_i)$ both lie on \mathcal{Y}_i , we have $\bar{p}_m(v_i) \cdot \bar{p}_m(v_i) = \bar{r}_m(v_i) \cdot \bar{r}_m(v_i)$. Hence for each vertex v_i ,

$$(\bar{p}_m(v_i) + \bar{r}_m(v_i)) \cdot (\bar{p}_m(v_i) - \bar{r}_m(v_i)) = 0.$$

Since p^m and r^m are not congruent, $p^m - r^m$ is a nontrivial infinitesimal motion. This means that the rank of the rigidity matrix for each framework $(G, \frac{p^m + r^m}{2})$ is less than maximal. Since both p^m and r^m converge to p , so does $\frac{p^m + r^m}{2}$. Thus (G, p) is a limit of a sequence of infinitesimally flexible frameworks and hence itself is infinitesimally flexible, a contradiction. (The fact that (G, p) is infinitesimally rigid implies that the rank of $R_{\mathcal{Y}^q}(G, p)$ is maximum for all $q \in \mathbb{R}^{3|V|}$ sufficiently close to p .) \square

We can use this lemma and our main result to show that vertex splitting preserves global rigidity on \mathcal{Y} under the additional assumption that the new edge is redundant.

Theorem 5.2. *Let (G, p) be a generic globally rigid framework on a family of concentric cylinders \mathcal{Y} . Let \hat{G} be a vertex split of G at the vertex v_1 with new vertex v_0 and suppose that $\hat{G} - v_0 v_1$ is rigid on \mathcal{Y} . Let $\hat{p}(v) = p(v)$ for all $v \neq v_0$ and $\hat{p}(v_0) = p(v_1)$. Then for any q on \mathcal{Y} which is sufficiently close to \hat{p} , (\hat{G}, q) is globally rigid on \mathcal{Y} .*

Proof. Since $(\hat{G}/v_0v_1, p) = (G, p)$ is globally rigid on \mathcal{Y} and p is generic, \hat{G}/v_0v_1 is rigid on \mathcal{Y} . Since $G - v_0v_1$ is also rigid on \mathcal{Y} , Theorem 4.14 implies that \hat{G} has a v_0v_1 -coincident generic rigid realisation (\hat{G}, \hat{p}) , where $\hat{p}(v) = p(v)$ for all $v \neq v_0$ and $\hat{p}(v_0) = p(v_1)$. Since (G, p) is globally rigid on \mathcal{Y} , (\hat{G}, \hat{p}) is also globally rigid on \mathcal{Y} . We can now use Lemma 5.1 to deduce that (\hat{G}, q) is globally rigid on \mathcal{Y} for all q sufficiently close to \hat{p} . \square

6. CONCLUDING REMARKS

Similarly to our definition of a framework (G, p) on \mathcal{Y} we can define a framework on a family of concentric spheres $\mathcal{S} = \mathcal{S}_1 \cup \mathcal{S}_2 \cup \dots \cup \mathcal{S}_k$ where $\mathcal{S}_i = \{(x, y, z) \in \mathbb{R}^3 : x^2 + y^2 + z^2 = r_i^2\}$ and $r = (r_1, \dots, r_k)$ is a vector of positive real numbers. We can map a framework on \mathcal{S} to a framework in the union of parallel (affine) planes $P_1 \cup P_2 \cup \dots \cup P_k$, where P_i is the plane $z = r_i$ in \mathbb{R}^3 , by central projection. In [10, 11] this process was shown to preserve infinitesimal rigidity for any framework on \mathcal{S} . Since the projection also preserves the property that u and v are coincident, the problem of characterising generic rigidity for frameworks with two coincident points on concentric spheres is equivalent to the problem of characterising generic rigidity for frameworks with two coincident points on parallel planes. This latter problem can be characterised using the proof technique of Theorem 1.1. This gives us the following result.

Theorem 6.1. *Let $G = (V, E)$ be a graph and let $u, v \in V$ be distinct vertices. Then G is uv -rigid on \mathcal{S} if and only if $G - uv$ and G/uv are both rigid on \mathcal{S} .*

Note that a graph $G = (V, E)$ is rigid on \mathcal{S} if and only if it has rank $2|V| - 3$ in the $(2, 3)$ -sparse matroid by [8, Theorem 5.1].

We can also replace \mathcal{Y} with other surfaces. In particular if we choose a surface with 1 ambient rigid motion (such as the cone, hyperboloid or torus) then the analogue of Theorem 2.1 requires the graph to be $(2, 1)$ -tight [9]. In the uv -coincident case we would define the value as $\text{val}(H) = 2|H| - t_H$ where $t_H = 3$ if $|H| \in \{2, 3\}$ and $H \neq \{u, v\}$, $t_H = 2$ if $|H| \in \{0, 4\}$ or $H = \{u, v\}$ and $t_H = 1$ if $|H| \geq 5$. We expect that, using similar techniques to Section 3, the appropriate count matroid can be established. However we do not know how to prove an analogue of Theorem 4.13. To make a start on this problem would require dealing with the case when the only vertices of degree less than 4 are u and v .

7. ACKNOWLEDGEMENTS

The second author was supported by the EPSRC First Grant EP/M013642/1 and by the Hungarian Scientific Research Fund (OTKA, grant number K109240).

REFERENCES

- [1] A. BERG AND T. JORDÁN, Algorithms for graph rigidity and scene analysis, Proc. 11th Annual European Symposium on Algorithms (ESA) 2003, (G. Di Battista and U. Zwick, eds) Springer Lecture Notes in Computer Science 2832, pp. 78-89, 2003.
- [2] R. CONNELLY AND W. WHITELEY, Global rigidity: the effect of coning, *Discrete and computational geometry* 43 (2010), 717-735.
- [3] ZS. FEKETE, T. JORDÁN AND V. E. KASZANITZKY, Rigid two-dimensional frameworks with two coincident points, *Graphs and Combinatorics* 31 (2014), no. 3, 585-599.
- [4] B. JACKSON, T. MCCOURT AND A. NIXON, Necessary conditions for the generic global rigidity of frameworks on surfaces, *Discrete and computational geometry* 52 (2014), no. 2, 344-360.
- [5] B. JACKSON AND A. NIXON, Stress matrices and generic global rigidity of frameworks on surfaces, *Discrete and computational geometry* 54 (2015), no. 3, 586-609.
- [6] G. LAMAN, On graphs and rigidity of plane skeletal structures, *J. Engineering Math.* 4 (1970), 331-340.

- [7] A. LEE AND I. STREINU, Pebble game algorithms and sparse graphs, *Discrete Mathematics* 308 (2008), no.8, 1425–1437.
- [8] A. NIXON, J. OWEN AND S. POWER, Rigidity of frameworks supported on surfaces, *SIAM Journal on Discrete Mathematics* 26 (2012), no. 4, 1733–1757.
- [9] A. NIXON, J. OWEN AND S. POWER, A characterization of generically rigid frameworks on surfaces of revolution, *SIAM Journal on Discrete Mathematics* 28 (2014), no. 4, 2008–2028.
- [10] F. SALIOLA, W. WHITELEY, Some notes on the equivalence of first-order rigidity in various geometries, Preprint (2007) arXiv:0709.3354.
- [11] B. SCHULZE, W. WHITELEY Coning, symmetry and spherical frameworks, *Discrete Compu. Geom.*, Vol. 48, (2012)., 622-657
- [12] W. WHITELEY, Some matroids from discrete applied geometry, in *Matroid Theory*, J. E. Bonin, J. G. Oxley, and B. Servatius, Eds. American Mathematical Society, Contemporary Mathematics, 1996, vol. 197, pp 171-313.

SCHOOL OF MATHEMATICAL SCIENCES, QUEEN MARY, UNIVERSITY OF LONDON, E1 4NS, U.K.
E-mail address: `b.jackson@qmul.ac.uk`

DEPARTMENT OF MATHEMATICS AND STATISTICS, LANCASTER UNIVERSITY, LA1 4YF, U.K.
E-mail address: `viktoria@cs.elte.hu`

DEPARTMENT OF MATHEMATICS AND STATISTICS, LANCASTER UNIVERSITY, LA1 4YF, U.K.
E-mail address: `a.nixon@lancaster.ac.uk`