

New anomaly observed in ^{12}C supports the existence and the vector character of the hypothetical X17 boson

A.J. Krasznahorkay,* A. Krasznahorkay,† M. Begala, M. Csatlós,
L. Csige, J. Gulyás, A. Krakó, J. Timár, I. Rajta, and I. Vajda
Institute for Nuclear Research (ATOMKI), P.O. Box 51, H-4001 Debrecen, Hungary

N.J. Sas
University of Debrecen, 4010 Debrecen, PO Box 105, Hungary

Employing the $^{11}\text{B}(p,\gamma)^{12}\text{C}$ nuclear reaction, the angular correlation of e^+e^- pairs was investigated in the angular range of $40^\circ \leq \Theta \leq 175^\circ$ for five different proton energies between $E_p = 1.50 - 2.5$ MeV. At small angles ($\Theta \leq 120^\circ$), the results can be well interpreted by the internal pair creation process of electromagnetic radiations with E1 and M1 multipolarities and by the external pair creation in the target backing. However, at angles greater than 120° , additional count excesses and anomalies were observed, which could be well accounted for by the existence of the previously suggested hypothetical X17 particle. Our results suggest that the X17 particle was generated mainly in E1 radiation. The derived mass of the particle is $m_{\text{X}17}c^2 = 17.03 \pm 0.11(\text{stat}) \pm 0.20(\text{syst})$ MeV. According to the mass, and to the derived branching ratio ($B_x = 3.6(3) \times 10^{-6}$), this is likely the same X17 particle that we recently suggested for describing the anomaly observed in the decay of ^8Be and ^4He .

PACS numbers: 23.20.Ra, 23.20.En, 14.70.Pw

INTRODUCTION

Very challenging nuclear physics experiments were initiated in 1978 to detect a new particle, the axion, predicted by Weinberg [1, 2]. The quantum chromodynamics axion is one of the most compelling solutions to the strong CP (charge conjugation parity symmetry) problem. Donnelly [3] proposed to study the angular correlation of the e^+e^- pairs created in $1^+ \rightarrow 0^+$ nuclear transitions as a signature for the decay of the axion. However, it was quickly ruled out in the MeV/ c^2 mass regime.

The later introduced dark photon [4, 5] is proposed as a force carrier connected to dark matter. In a minimal scenario, this new force can be introduced by extending the gauge group of the Standard Model with a new abelian U(1) gauge symmetry.

Pierre Fayet suggested a generalized dark photon model that would produce a light gauge boson with an extra U(1) gauge group already in 1980 [6–8]. Such a generalized dark photon may act as a mediator of light dark matter annihilations, possibly allowing for lighter than GeV/ c^2 dark matter particles [9, 10]. More recent discussions on the light U boson as the mediator of a new force, coupled to a combination of Q, B, L and dark matter can be found in Ref [11, 12].

Searches for light particles, especially a 9 MeV/ c^2 particle suggested by Fokke de Boer and coworkers, have been performed in Frankfurt [13]. These results, although with little confidence, appeared to confirm the existence of the 9 MeV/ c^2 particle. However, due to the closure of the accelerator, these experiments could not be continued.

Recently, we studied electron-positron angular correlations for the 17.6 MeV and 18.15 MeV tran-

sitions in ^8Be and an anomalous angular correlation was observed for the 18.15 MeV transition [14]. This was interpreted as the creation and decay of an intermediate bosonic particle with a mass of $m_{\text{X}17}c^2 = 16.70 \pm 0.35(\text{stat}) \pm 0.5(\text{sys})$ MeV, which is now called X17. The possible relation of the X17 boson to the dark matter problem triggered an enormous interest in the wider physics community [15, 16].

The first theoretical interpretation of the experimental results was performed by Feng et al. [17, 18]. They explained the anomaly with a 16.7 MeV, vector gauge boson X17, which may mediate a fifth fundamental force with some coupling to Standard Model (SM) particles. The theory of the dark photon has been generalized to the fact that the new particle is coupled not only to electric charges but also to quarks.

Constraints on the coupling constants of such a new particle, notably from searches for $\pi_0 \rightarrow Z' + \gamma$ by the NA48/2 experiment [19] was also taken into account by Feng and co-workers [17, 18]. Based on their results, the X17 particle couples much more strongly to neutrons than to protons, so the particle was named protophobic.

Zhang and Miller [20] investigated the nuclear transition form factor as a possible origin of the anomaly, but they found the concluded form factor unrealistic for the ^8Be nucleus.

Ellwanger and Moretti suggested another interpretation of the experimental results in view of a light, pseudoscalar particle [21]. They predicted about ten times smaller branching ratio in case of the 17.6 MeV transition compared to the 18.15 MeV one, which is in nice agreement with our results.

Subsequently, many studies with different models have been performed including an extended two Higgs doublet

model [22–24]. They showed that the anomaly can be described with a very light Z_0 bosonic state, with significant axial couplings.

In parallel to these recent theoretical studies, we re-investigated the ^8Be anomaly with an improved experimental setup [25–27].

Recently, we also observed a similar anomaly in ^4He [26, 28]. The signal could be described by the creation and subsequent decay of a light particle during the proton capture process on ^3H to the ground state of the ^4He nucleus. The derived mass of the particle ($m_{\text{X}17}c^2 = 16.94 \pm 0.12(\text{stat.}) \pm 0.21(\text{syst.})$ MeV) agreed well with that of the proposed X17 particle. It was also shown, that the branching ratios of the X17 particle compared to the γ -decay are identical within uncertainties for three beam energies, proving that the X17 particle was most likely formed in direct proton capture, which has a dominant multipolarity of E1. Our results obtained for ^4He at different beam energies agree well with the present theoretical results of Viviani et al. [29].

Referring to our manuscript [28], Feng and co-workers have communicated a work very recently with the title of „Dynamical evidence for a fifth force explanation of the ATOMKI nuclear anomalies” [30], in which they propose to study the E1 ground state decay of the 17.2 MeV $J^\pi = 1^-$ state in ^{12}C in order to determine if X17 has a vector or axial-vector character.

In the present work, we investigated the 17.2 MeV $1^- \rightarrow 0^+$ transition of ^{12}C to search for signatures of the creation of the X17 particle.

EXPERIMENTAL METHODS

The experiments were performed in Debrecen, Hungary at the 2 MV Tandem accelerator of ATOMKI. The $E_x=17.23$ MeV ($J^\pi = 1^-$) [31] state of ^{12}C was excited by the $^{11}\text{B}(p,\gamma)^{12}\text{C}$ nuclear reaction. Due to its large cross section, the reaction is also widely used for detector calibrations. The resonant proton energy is $E_p = 1.388$ MeV [31, 32].

Owing to the rather large level width ($\Gamma = 1.15$ MeV [31]), a 2 mg/cm² thick ^{11}B target was applied to maximize the yield of the e^+e^- pairs. The target was evaporated onto a 5 μm thick Ta foil. The average energy loss of the protons in the target was ≈ 300 keV. To compensate for the energy losses, the energy of the protons was chosen to be $E_p = 1.50, 1.70, 1.88, 2.10$ and 2.50 MeV. The proton beam was impinging on a ^{11}B target with a typical current of 2 μA for about 50 hours at each beam energy. To achieve a more efficient cooling of the target and thus to reduce its degradation, we replaced the previously used [14, 33] Plexiglass support rods by Al rods. However, in turn, our data suffered a bit larger signal background from Al induced by the γ rays as shown in Fig. 1 than before (Fig. 9 of Ref. [17]).

Our previous detector setup [14, 33] has recently been

upgraded. The details of the upgrade are described in [28]. In the present experiment, the time and energy signals of the scintillators, as well as the time, energy and position signals of the DSSD detectors were recorded.

The energy calibration of the telescopes, the energy and position calibrations of the DSSD detectors, the Monte Carlo (MC) simulations as well as the acceptance calibration of the whole e^+e^- coincidence pair spectrometer were explained in Ref. [28]. Reasonably good agreement was obtained between the experimental acceptance and results of the MC simulations, as presented in Fig. 1. The average difference is within $\approx 3.0\%$ in the angular range of $40^\circ - 170^\circ$.

In order to validate the accuracy of the MC simulations, we performed measurements also on the $^7\text{Li}(p,\gamma)^8\text{Be}$ reaction. The experimental results for the angular correlations from this data taking on the $E_p = 441$ keV resonance (red dots with error bars) are shown in Fig. 1 b), together with the corresponding Monte Carlo simulation (histogram) of the IPC process stemming mostly from the M1 nuclear transition. The contribution of the external pair creation (EPC) process of the 17.6 MeV γ -rays is also shown by a black histogram. We note here that the direct-capture contribution is negligible compared to the M1 IPC due to the large resonance capture cross section and the thin target.

As it can be seen in Fig. 1, the simulation of the IPC process manages to describe the shape of the data distribution accurately, and the contribution of EPC created on the different parts of the spectrometer is reasonably low.

In order to search for the assumed X17 particle, both the sum-energy spectrum of the e^+e^- pairs measured by the telescopes, and their angular correlations, determined by the DSSD detectors, have been analyzed. Since the counting rates in the detectors were low (≈ 150 Hz in the scintillators and ≈ 25 Hz in the DSSD detectors) and the coincidence time window was sharp (≈ 10 ns) the effect of random coincidences was negligible. In the followings we show only the real coincidence gated spectra.

In the GEANT simulations, both e^+e^- pairs generated by internal pair creation in the target and the e^+e^- pairs generated by external pair creation in the Ta backing were taken into account. A more detailed description of the simulations can be found in Ref. [28].

EXPERIMENTAL RESULTS AND DISCUSSION

The total energy spectrum of the e^+e^- -pairs produced in the decay of the $E_x=17.2$ MeV ($J^\pi = 1^-$) state of ^{12}C at $E_p=1.7$ MeV is presented in Fig. 2. In addition to the E1 ground state transition, this state decays to the $E_x=4.44$ MeV ($J^\pi = 2^+$) level with an $E = 12.76$ MeV E1 transition, which is also present the energy spectrum.




FIG. 1. a) Detector response for the setup as a function of correlation angle (θ) for isotropic emission of e^+e^- pairs (red crosses) compared with the results of the Monte Carlo simulations (black line histogram) as explained in the text. b) e^+e^- angular correlations obtained for the 17.6 MeV transition of ^8Be by using thin target backing compared to the simulations performed for E1 and M1 IPC, as well as for the EPC created by the γ -rays on the different materials around the target.

The intense, 4.44 MeV ground state transition was discarded by setting a proper hardware threshold in order to reduce the high count rate of the DAQ.

The experimental efficiency of the e^+e^- spectrometer was determined with uncorrelated e^+e^- -pairs by taking the e^- and e^+ data from consecutive events as previously described in Refs. [14, 25, 27, 28]. The gated and efficiency-corrected e^+e^- angular correlation in the 17.2 MeV ($J^\pi = 1^- \rightarrow 0^+$) transition is shown in Fig. 3 for proton energies of $E_p = 1.5, 1.7, 1.88, 2.1$ and 2.5 MeV.




FIG. 2. Total energy spectrum of the e^+e^- -pairs from the $^{11}\text{B}(p, e^+e^-)^{12}\text{C}$ nuclear reaction.

As show in Fig. 3, a combination of the MC simulated IPC distributions of E1 and M1 radiations together with a small contribution of simulated external pair creation (EPC) in the Ta backing can describe the experimental distributions below $\Theta = 140^\circ$ reasonable well. However, we observe significant deviations at large angles ($> 140^\circ$) at each proton energy.

To derive the invariant mass of the decaying particle, we carried out a fitting procedure for both the mass value and the amplitude of the observed peak.

The fit was performed with RooFit [34] by describing the e^+e^- angular correlation with the following intensity function (INT):

$$INT(e^+e^-) = N_{Bg} * PDF(exp) + N_{Sig} * PDF(sig), \quad (1)$$

where $PDF(exp)$ was determined from a separate fit for the background region, $PDF(sig)$ was simulated by GEANT for the two-body decay of an X particle as a function of its mass, and N_{Bg} and N_{Sig} are the fitted numbers of background and signal events, respectively.

The signal PDF was constructed as a 2-dimensional model function of the e^+e^- opening angle and the mass of the simulated particle. To construct the mass dependence, the PDF linearly interpolates the e^+e^- opening angle distributions simulated for discrete particle masses.

Using the intensity function described in Equation 1, we first performed a list of fits by fixing the simulated particle mass in the signal PDF to a certain value, and letting RooFit estimate the best values for N_{Sig} and N_{Bg} . Allowing the particle mass to vary in the fit, the best fitted mass is calculated and the corresponding fit is shown for each studied beam energy in Fig. 4.




FIG. 3. a) Experimental angular correlations of the e^+e^- pairs measured in the $^{11}\text{B}(p, e^+e^-)^{12}\text{C}$ reaction at the vicinity of the $J^\pi = 1^+$ resonance for different proton energies.

A significant background is obtained from the E1 transition, but the contribution from the assumed particle decay is also significant at large angles.

The measured invariant masses of the hypothetical X17 and the branching ratios (B_x) of its e^+e^- decay to the γ decay, as derived from the fits are summarized in Table I. The values are compatible for each fitted parameter within 1σ error bars. Their average values are also highlighted.

TABLE I. X17 branching ratios (B_x), masses, and confidences derived from the fits.

E_p (MeV)	B_x $\times 10^{-6}$	Mass (MeV/ c^2)	Confidence
1.50	1.1(6)	16.81(15)	3σ
1.70	3.3(7)	16.93(8)	7σ
1.88	3.9(7)	17.13(10)	8σ
2.10	4.9(21)	17.06(10)	3σ
Averages	3.6(3)	17.03(11)	
Previous [14]	5.8	16.70(30)	
Previous [28]	5.1	16.94(12)	
Predicted [30]	3.0		




FIG. 4. Experimental angular correlations of the e^+e^- pairs measured at different proton energies. The full curves for each proton energy shows the results of the fit, using simulated angular distributions.

In Table I, only the statistical errors are indicated. The systematic uncertainties were estimated to be $\Delta m_X c^2(\text{syst.}) = \pm 0.20$ MeV by employing a series of MC simulations as presented in one of our previous works [28]. It mostly represents the uncertainty of the position of the beam spot, which was found to be shifted by about ± 2 mm in one measurement run.

The extracted invariant mass agrees well with the values published earlier for the ^8Be [14] and the ^4He [28] experiments, which provides a convincing kinematic verification of the existence of the X17 particle. The branching ratio of the X17 decay differs from the previous data, but, on the other hand, agrees well with the theoretically predicted value [30].

SUMMARY

We have studied the E1 ground state decay of the 17.2 MeV $J^\pi = 1^-$ state in ^{12}C . The energy-sum and the angular correlation of the e^+e^- pairs produced in the $^{11}\text{B}(p,e^+e^-)^{12}\text{C}$ reaction were measured at proton energies of $E_p = 1.50, 1.70, 1.88, 2.10$ and 2.50 MeV. The gross features of the distributions of these quantities can be described well by the IPC process following the decay of the 1^- resonance and by considering a small contribution of external pair creation process induced by the high-energy γ rays. However, on top of the smooth, monotonic distribution of the angular correlation of e^+e^- pairs, we observed significant peak-like anomalous excess around $155\text{-}160^\circ$ at four different beam energies.

The e^+e^- excess can be well-described by the creation and subsequent decay of the X17 particle, which we have recently suggested [14, 28]. The invariant mass of the particle was derived to be ($m_{Xc^2} = 17.03 \pm 0.11(\text{stat.}) \pm 0.20(\text{syst.})$ MeV), which agrees well with our previously published values. The branching ratio of the e^+e^- decay of X17 to the γ decay was found to be $3.6(3) \times 10^{-6}$. The present observation of the X17 particle in an E1 transition supports its vector character, as suggested by Feng et al. [30].

Given the present results on the X17 creation in E1 transitions, we consider to search for X17 in the decay of Giant Dipole Resonance (GDR) excitations of different nuclei.

ACKNOWLEDGEMENTS

We wish to thank Z. Pintye for the mechanical and J. Molnar for the electronic design of the experiment. This work has been supported by the Hungarian NKFI Foundation No. K124810 and by the GINOP-2.3.3-15-2016-00034 and GINOP-2.3.3-15-2016-00005 grants.

* kraszna@atomki.hu

† Currently working at CERN, Geneva, Switzerland

- [1] S. Weinberg, Phys. Rev. Lett., **40**, 223 (1978).
- [2] F. Wilczek, Phys. Rev. Lett., **40**, 278 (1978).
- [3] T. W. Donnelly, S. J. Freedman, R. S. Lytel, R. D. Peccei, and M. Schwartz, Phys. Rev. **D18**, 1607 (1978).
- [4] P. Galison, A. Manohar, Phys. Lett. B **136** 279 (1984).
- [5] B. Holdom, Phys. Lett. B. **166**, 196 (1986).
- [6] P. Fayet, Phys. Lett. B **95** (1980), 285-289
- [7] P. Fayet, Nucl. Phys. B **187** (1981), 184-204
- [8] P. Fayet, Phys. Lett. B **96** (1980), 83-88
- [9] C. Boehm and P. Fayet, Nucl. Phys. B **683** (2004), 219-263
- [10] P. Fayet, Phys. Rev. D **70** (2004), 023514
- [11] Pierre Fayet, Eur. Phys. J. C **77** 53 (2017).
- [12] Pierre Fayet, Phys. Rev. D **103**, 035034 (2021).
- [13] F. W. N. de Boer et al., J. Phys. **G27**, L29 (2001).
- [14] A.J. Krasznahorkay et al., Phys. Rev. Lett. **116**, 042501 (2016).
- [15] <https://inspirehep.net/search?ln=en&p=refersto%3A13582>
- [16] P. Agrawal et al., Eur. Phys. J. C **81**, 1015 (2021).
- [17] J. L. Feng, B. Fornal, I. Galon, S. Gardner, J. Smolinsky, T.M.P. Tait, P. Tanedo, Phys. Rev. Lett. **117**, 071803 (2016).
- [18] J.L. Feng, B. Fornal, I. Galon, S. Gardner, J. Smolinsky, T.M.P. Tait, P. Tanedo, Phys. Rev. **D 95**, 035017 (2017).
- [19] J. Batley et al., for the NA48/2 Collaboration, Phys. Lett. **B 746**, 178 (2015).
- [20] Xilin Zhang, Gerald A Miller Phys. Lett. **B773** 159 (2017).
- [21] U. Ellwanger and S. Moretti, JHEP **11** 39 (2016).
- [22] Luigi Delle Rose et al., Phys. Rev. **D 96**, 115024 (2017) and references therein.
- [23] Luigi Delle Rose et al., Phys. Rev. **D 99** 055022 (2019).
- [24] Luigi Delle Rose et al., Frontiers in Physics **7** 73 (2019).
- [25] A.J. Krasznahorkay et al., J. Phys.: Conf. Series **1056**, 012028 (2017).
- [26] A.J. Krasznahorkay et al., Acta Phys. Pol. **B 50**, 675 (2019).
- [27] A.J. Krasznahorkay et al., Proceedings of Science, (Bormio2017) 036 (2017).
- [28] A.J. Krasznahorkay et al., Phys. Rev. C **104**, 044003 (2021).
- [29] M. Viviani et al., Phys. Rev. C **105**, 014001 (2022)
- [30] J.L. Feng, T.M.P. Tait and C.B. Verharen, Phys. Rev. **D 102**, 036016 (2020).
- [31] F Ajzenberg-Selove, Nucl. Phys. **A506**, 1 (1990).
- [32] R.E. Segel, S.S. Hanna, R.G. Allas, Phys. Rev. **139**, B818 (1965).
- [33] J. Gulyás et al., Nucl. Instr. and Meth. in Phys. Res. A **808**, 21 (2016).
- [34] W. Verkerke and D. P. Kirkby, "The RooFit toolkit for data modeling," eConf C **0303241** (2003) MOLT007 [physics/0306116].