



EXISTENCE OF WEAK SOLUTIONS FOR A DOUBLE PHASE VARIABLE EXPONENT PROBLEM WITH A GRADIENT DEPENDENT REACTION TERM

MOHAMED EL OUAARABI, CHAKIR ALLALOU, AND SAID MELLIANI

Received 09 February, 2022

Abstract. In the present paper, we study the existence of at least one weak solution for a class of double phase variable exponent problem with a reaction term depending on the gradient and on two real parameters. By using the topological degree theory for a class of demicontinuous operators of generalized (S_+) and the theory of the variable exponent Sobolev spaces, we obtain the existence of at least one weak solution of this problem.

2010 *Mathematics Subject Classification:* 35J60; 35J70; 35D30; 47H11

Keywords: double phase variable exponent problem, $(p(x), q(x))$ -Laplacian operators, topological degree theory, variable exponent Sobolev spaces

1. INTRODUCTION AND MOTIVATION

The study of differential equations with with nonstandard $p(x)$ -Laplacian operator ($(p(x), q(x))$ -Laplacian operator) is an attractive topic and has been the object of considerable attention in recent years. Perhaps the impulse for this comes from the new search field that reflects a new type of physical phenomenon is a class of nonlinear problems with variable exponents. In the subject of fluid mechanics, for example, Rajagopal and M. Ruzicka recently developed a very interesting model for these fluids in [21] (see also [22]). Other applications relate to image processing [1], elasticity problems [18–20, 24, 26], the flow in porous media [3], and problems in the calculus of variations involving variational integrals with nonstandard growth [2].

Here and in the sequel, we will assume that Ω is a bounded domain in $\mathbb{R}^n (n > 1)$, with a Lipschitz boundary denoted by $\partial\Omega$, and let μ and λ be two real parameters.

In this paper, we consider the following double phase variable exponent problem:

$$\begin{cases} -\Delta_{p(x),q(x)}(u) = \mu g(x, u) + \lambda f(x, u, \nabla u) & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases} \quad (1.1)$$

where

$$\Delta_{p(x),q(x)}(u) := \operatorname{div} \left(|\nabla u|^{p(x)-2} \nabla u + a(x) |\nabla u|^{q(x)-2} \nabla u \right).$$

In this problem, the coefficient $a: \overline{\Omega} \rightarrow \mathbb{R}^+$ is Lipschitz continuous function, $g: \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ and $f: \Omega \times \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}$ are Carathéodory functions that satisfy the assumption of growth, and the variables exponents $p(\cdot), q(\cdot) \in C_+(\overline{\Omega})$ are assumed to satisfy the following assumption:

$$1 < q^- \leq q \leq q^+ < p^- \leq p \leq p^+ < +\infty.$$

The double phase operator has been used in the modelling of strongly anisotropic materials [26] and in Lavrentiev's phenomenon [27]. In the one hand, we have the physical motivation; since the double phase operator has been used to model the steady-state solutions of reaction-diffusion problems, that arise in biophysic, plasma-physic and in the study of chemical reactions. In the other hand, these operators provide a useful paradigm for describing the behaviour of strongly anisotropic materials, whose hardening properties are linked to the exponent governing the growth of the gradient change radically with the point, where the coefficient $a(\cdot)$ determines the geometry of a composite made of two different materials (see [4, 28] and the references given there).

Let us recall some known results on problem (1.1). For example, Fan and Zhang [12], based on the theory of the spaces $L^{p(x)}(\Omega)$ and $W_0^{1,p(x)}(\Omega)$, present several sufficient conditions for the existence of solutions for the problem (1.1) with $\mu = 1$ and $a = \lambda = 0$.

R. Alsaedi [9] establishes sufficient conditions for the existence of nontrivial weak solutions for the problem (1.1) when $\mu = 1$, $a = 0$, $g(x, u) = |u|^{q(x)-2}u$, $\lambda > 0$ and $f(x, u) = |u|^{p(x)-2}u$.

Problems related to (1.1) in case $p(x) \equiv p$ and $q(x) \equiv q$ have been studied by many scholars, for example, Liu et al. [17] study the problem (1.1) when $\lambda = 0$ and $\mu = 1$, and Wang et al. [23] showed, by using the topological degree theory for a class of demicontinuous operators, the existence of at least one weak solution of (1.1) with $\mu = 0$ and $\lambda = 1$.

We would like to draw attention to the fact that the $p(x)$ -laplacian operator has more complicated nonlinearity than the p -laplacian operator. For example, they are non-homogeneous, which prove that our problem is more difficult than the operators p -Laplacian type.

Motivated by the aforementioned works, in the present paper, we will generalize these works. By using a topological degree for a class of demicontinuous operators of generalized (S_+) type of [5] and the theory of the generalized Sobolev spaces, we establish the existence of weak solutions u in $W_0^{1,p(x)}(\Omega)$ for the problem (1.1).

The paper is designed as follows: in Section 2, we review some fundamental preliminaries about the functional framework where we will treat our problems. In Section 3, we introduce some classes of operators, as well as the topological degree theory for a class of demicontinuous operators of generalized (S_+) . Finally, Section 4 is devoted to discussing the existence of weak solutions to (1.1).

2. PRELIMINARIES

In this section, we recall the most important and relevant properties and notations about generalized Sobolev spaces $W^{1,p(x)}(\Omega)$, that we will need in our analysis of the problem (1.1), by that, referring to [6–8, 10, 11, 13, 16] for more details.

Let Ω be a smooth bounded domain in $\mathbb{R}^n (n > 1)$, with a Lipschitz boundary denoted by $\partial\Omega$. Set

$$C_+(\overline{\Omega}) = \left\{ p : p \in C(\overline{\Omega}) \text{ such that } p(x) > 1 \text{ for any } x \in \overline{\Omega} \right\}.$$

For each $p \in C_+(\overline{\Omega})$, we define

$$p^+ := \max \left\{ p(x), x \in \overline{\Omega} \right\} \text{ and } p^- := \min \left\{ p(x), x \in \overline{\Omega} \right\}.$$

For every $p \in C_+(\overline{\Omega})$, we define

$$L^{p(x)}(\Omega) = \left\{ u : \Omega \rightarrow \mathbb{R} \text{ is measurable such that } \int_{\Omega} |u(x)|^{p(x)} dx < +\infty \right\},$$

equipped with the Luxemburg norm

$$|u|_{p(x)} = \inf \left\{ \lambda > 0 : \rho_{p(x)} \left(\frac{u}{\lambda} \right) \leq 1 \right\},$$

where

$$\rho_{p(x)}(u) = \int_{\Omega} |u(x)|^{p(x)} dx, \forall u \in L^{p(x)}(\Omega).$$

Proposition 1 ([13, Theorem 1.3 and Theorem 1.4]). *Let (u_n) and $u \in L^{p(\cdot)}(\Omega)$, then*

$$|u|_{p(x)} < 1 \text{ (resp. } = 1; > 1) \Leftrightarrow \rho_{p(x)}(u) < 1 \text{ (resp. } = 1; > 1)$$

$$|u|_{p(x)} > 1 \Rightarrow |u|_{p(x)}^{p^-} \leq \rho_{p(x)}(u) \leq |u|_{p(x)}^{p^+}, \tag{2.1}$$

$$|u|_{p(x)} < 1 \Rightarrow |u|_{p(x)}^{p^+} \leq \rho_{p(x)}(u) \leq |u|_{p(x)}^{p^-}, \tag{2.2}$$

$$\lim_{n \rightarrow \infty} |u_n - u|_{p(x)} = 0 \Leftrightarrow \lim_{n \rightarrow \infty} \rho_{p(x)}(u_n - u) = 0. \tag{2.3}$$

Remark 1. According to (2.1) and (2.2), we have

$$|u|_{p(x)} \leq \rho_{p(x)}(u) + 1, \tag{2.4}$$

$$\rho_{p(x)}(u) \leq |u|_{p(x)}^{p^-} + |u|_{p(x)}^{p^+}. \tag{2.5}$$

Proposition 2 ([16, Theorem 2.5 and Corollary 2.7]). *The spaces $L^{p(x)}(\Omega)$ is a separable and reflexive Banach spaces.*

Proposition 3 ([16, Theorem 2.1]). *The conjugate space of $L^{p(x)}(\Omega)$ is $L^{p'(x)}(\Omega)$ where $\frac{1}{p(x)} + \frac{1}{p'(x)} = 1$ for all $x \in \Omega$. For any $u \in L^{p(x)}(\Omega)$ and $v \in L^{p'(x)}(\Omega)$, we have the following Hölder-type inequality*

$$\left| \int_{\Omega} uv \, dx \right| \leq \left(\frac{1}{p^-} + \frac{1}{p'^-} \right) \|u\|_{p(x)} \|v\|_{p'(x)} \leq 2 \|u\|_{p(x)} \|v\|_{p'(x)}. \tag{2.6}$$

Remark 2 ([13, Theorem 1.11]). If $p_1, p_2 \in C_+(\overline{\Omega})$ with $p_1(x) \leq p_2(x)$ for any $x \in \overline{\Omega}$, then there exists the continuous embedding $L^{p_2(x)}(\Omega) \hookrightarrow L^{p_1(x)}(\Omega)$.

Now, let $p \in C_+(\overline{\Omega})$ and we define $W^{1,p(x)}(\Omega)$ as

$$W^{1,p(x)}(\Omega) = \left\{ u \in L^{p(x)}(\Omega) \text{ such that } |\nabla u| \in L^{p(x)}(\Omega) \right\},$$

equipped with the norm

$$\|u\| = \|u\|_{p(x)} + \|\nabla u\|_{p(x)}.$$

We also define $W_0^{1,p(\cdot)}(\Omega)$ as the subspace of $W^{1,p(\cdot)}(\Omega)$, which is the closure of $C_0^\infty(\Omega)$ with respect to the norm $\|\cdot\|$.

Proposition 4 ([14, Theorem 4.3]). *If the exponent $p(\cdot)$ satisfies the log-Hölder continuity condition, i.e. there is a constant $b > 0$ such that for every $x, y \in \Omega, x \neq y$ with $|x - y| \leq \frac{1}{2}$ one has*

$$|p(x) - p(y)| \leq \frac{b}{-\log|x - y|}, \tag{2.7}$$

then we have the Poincaré inequality, i.e. there exists a constant $C > 0$ depending only on Ω and the function p such that

$$\|u\|_{p(x)} \leq C \|\nabla u\|_{p(x)}, \quad \forall u \in W_0^{1,p(\cdot)}(\Omega). \tag{2.8}$$

In this paper we will use the following equivalent norm on $W_0^{1,p(\cdot)}(\Omega)$

$$\|u\|_{1,p(x)} = \|\nabla u\|_{p(x)},$$

which is equivalent to $\|\cdot\|$.

Furthermore, we have the compact embedding $W_0^{1,p(\cdot)}(\Omega) \hookrightarrow L^{p(\cdot)}(\Omega)$ (see [16]).

Proposition 5 ([16, Theorem 3.1]). *The spaces $(W^{1,p(x)}(\Omega), \|\cdot\|)$ and $(W_0^{1,p(x)}(\Omega), \|\cdot\|_{1,p(x)})$ are separable and reflexive Banach spaces.*

Remark 3. The dual space of $W_0^{1,p(x)}(\Omega)$ denoted $W^{-1,p'(x)}(\Omega)$, is equipped with the norm

$$\|u\|_{-1,p'(x)} = \inf \left\{ \|u_0\|_{p'(x)} + \sum_{i=1}^N \|u_i\|_{p'(x)} \right\},$$

where the infimum is taken on all possible decompositions $u = u_0 - \operatorname{div}F$ with $u_0 \in L^{p'(x)}(\Omega)$ and $F = (u_1, \dots, u_N) \in (L^{p'(x)}(\Omega))^N$.

3. A REVIEW ON SOME CLASSES OF MAPPINGS AND TOPOLOGICAL DEGREE THEORY

We start by defining some classes of mappings. In what follows, let X be a real separable reflexive Banach space and X^* be its dual space with dual pairing $\langle \cdot, \cdot \rangle$ and given a nonempty subset Ω of X . Strong (weak) convergence is represented by the symbol \rightarrow (\rightharpoonup).

Definition 1. Let Y be another real Banach space. A operator $F : \Omega \subset X \rightarrow Y$ is said to be :

- (1) bounded, if it takes any bounded set into a bounded set.
- (2) demicontinuous, if for any sequence $(u_n) \subset \Omega$, $u_n \rightarrow u$ implies that $F(u_n) \rightharpoonup F(u)$.
- (3) compact, if it is continuous and the image of any bounded set is relatively compact.

Definition 2. A mapping $F : \Omega \subset X \rightarrow X^*$ is said to be :

- (1) of class (S_+) , if for any sequence $(u_n) \subset \Omega$ with $u_n \rightharpoonup u$ and $\limsup_{n \rightarrow \infty} \langle Fu_n, u_n - u \rangle \leq 0$, we have $u_n \rightarrow u$.
- (2) quasimonotone, if for any sequence $(u_n) \subset \Omega$ with $u_n \rightharpoonup u$, we have $\limsup_{n \rightarrow \infty} \langle Fu_n, u_n - u \rangle \geq 0$.

Definition 3. Let $T : \Omega_1 \subset X \rightarrow X^*$ be a bounded operator such that $\Omega \subset \Omega_1$. For any operator $F : \Omega \subset X \rightarrow X$, we say that

- (1) F is of class $(S_+)_T$, if for any sequence $(u_n) \subset \Omega$ with $u_n \rightharpoonup u$, $y_n := Tu_n \rightharpoonup y$ and $\limsup_{n \rightarrow \infty} \langle Fu_n, y_n - y \rangle \leq 0$, we have $u_n \rightarrow u$.
- (2) F has the property $(QM)_T$, if for any sequence $(u_n) \subset \Omega$ with $u_n \rightharpoonup u$, $y_n := Tu_n \rightharpoonup y$, we have $\limsup_{n \rightarrow \infty} \langle Fu_n, y - y_n \rangle \geq 0$.

In the sequel, we consider the following classes of operators:

$$\mathcal{F}_1(\Omega) := \left\{ F : \Omega \rightarrow X^* : F \text{ is bounded, demicontinuous and of class } (S_+) \right\},$$

$$\mathcal{F}_T(\Omega) := \left\{ F : \Omega \rightarrow X : F \text{ is demicontinuous and of class } (S_+)_T \right\},$$

$$\mathcal{F}_{T,B}(\Omega) := \left\{ F \in \mathcal{F}_T(\Omega) : F \text{ is bounded} \right\},$$

for any $\Omega \subset D(F)$, where $D(F)$ denotes the domain of F , and any $T \in \mathcal{F}_1(\Omega)$.

Now, let O be the collection of all bounded open sets in X and we define

$$\mathcal{F}(X) := \left\{ F \in \mathcal{F}_T(\bar{E}) : E \in O, T \in \mathcal{F}_1(\bar{E}) \right\},$$

where, $T \in \mathcal{F}_1(\overline{E})$ is called an essential inner map to F .

Lemma 1 ([15, Lemma 2.3]). *Let $T \in \mathcal{F}_1(\overline{E})$ be continuous and $S: D(S) \subset X^* \rightarrow X$ be demicontinuous such that $T(\overline{E}) \subset D(S)$, where E is a bounded open set in a real reflexive Banach space X . Then the following statements are true :*

- (1) *If S is quasimonotone, then $I + S \circ T \in \mathcal{F}_T(\overline{E})$, where I denotes the identity operator.*
- (2) *If S is of class (S_+) , then $S \circ T \in \mathcal{F}_T(\overline{E})$.*

Definition 4. Suppose that E is bounded open subset of a real reflexive Banach space X , $T \in \mathcal{F}_1(\overline{E})$ is continuous and $F, S \in \mathcal{F}_T(\overline{E})$. Then the affine homotopy $\mathcal{H}: [0, 1] \times \overline{E} \rightarrow X$ defined by

$$\mathcal{H}(t, u) := (1 - t)Fu + tSu, \quad \text{for } (t, u) \in [0, 1] \times \overline{E}$$

is called an admissible affine homotopy with the common continuous essential inner map T .

Remark 4 ([15, Lemma 2.5]). *The above affine homotopy is of class $(S_+)_T$.*

As in [15] we give the topological degree for the class $\mathcal{F}(X)$.

Theorem 1. *Let*

$$M = \left\{ (F, E, h) : E \in \mathcal{O}, T \in \mathcal{F}_1(\overline{E}), F \in \mathcal{F}_{T,B}(\overline{E}), h \notin F(\partial E) \right\}.$$

Then, there exists a unique degree function $d: M \rightarrow \mathbb{Z}$ that satisfies the following properties:

- (1) (Normalization.) *For any $h \in F(E)$, we have*

$$d(I, E, h) = 1.$$

- (2) (Additivity.) *Let $F \in \mathcal{F}_{T,B}(\overline{E})$. If E_1 and E_2 are two disjoint open subsets of E such that $h \notin F(\overline{E} \setminus (E_1 \cup E_2))$, then we have*

$$d(F, E, h) = d(F, E_1, h) + d(F, E_2, h).$$

- (3) (Homotopy invariance.) *If $\mathcal{H}: [0, 1] \times \overline{E} \rightarrow X$ is a bounded admissible affine homotopy with a common continuous essential inner map and $h: [0, 1] \rightarrow X$ is a continuous path in X such that $h(t) \notin \mathcal{H}(t, \partial E)$ for all $t \in [0, 1]$, then*

$$d(\mathcal{H}(t, \cdot), E, h(t)) = \text{const for all } t \in [0, 1].$$

- (4) (Existence.) *If $d(F, E, h) \neq 0$, then the equation $Fu = h$ has a solution in E .*

Definition 5 ([15, Definition 3.3]). The above degree is defined as follows:

$$d(F, E, h) := d_B(F|_{\overline{E}_0}, E_0, h),$$

where d_B is the Berkovits degree [5] and E_0 is any open subset of E with $F^{-1}(h) \subset E_0$ and F is bounded on \overline{E}_0 .

4. EXISTENCE OF WEAK SOLUTIONS

In this section, we will discuss the existence of weak solutions of (1.1). For this, we list our assumptions associated with our problem to show the existence result.

From now on, we always assume that $\Omega \subset \mathbb{R}^n (n > 1)$ is a bounded domain with a Lipschitz boundary $\partial\Omega$, $p \in C_+(\overline{\Omega})$ satisfy the log-Hölder continuity condition (2.7), $g: \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ and $f: \Omega \times \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}$ are functions such that :

(A₁): f is a Carathéodory function.

(A₂): There exists $\rho > 0$ and $\gamma \in L^{p'(x)}(\Omega)$ such that

$$|f(x, \zeta, \xi)| \leq \rho(\gamma(x) + |\zeta|^{q(x)-1} + |\xi|^{q(x)-1}).$$

(A₃): g is a Carathéodory function.

(A₄): There are $\sigma > 0$ and $v \in L^{p'(x)}(\Omega)$ such that

$$|g(x, \zeta)| \leq \sigma(v(x) + |\zeta|^{s(x)-1}),$$

for a.e. $x \in \Omega$ and all $(\zeta, \xi) \in \mathbb{R} \times \mathbb{R}^n$, where $k, s \in C_+(\overline{\Omega})$ with $2 \leq s^- \leq s(x) \leq s^+ < p^-$.

Remark 5.

- Note that for all $\varphi \in W_0^{1,p(x)}(\Omega)$

$$\int_{\Omega} \left(|\nabla u|^{p(x)-2} \nabla u \nabla \varphi + a(x) |\nabla u|^{q(x)-2} \nabla u \nabla \varphi \right) dx$$

is well defined (see [12, 17]).

- $\mu g(x, u)$ and $\lambda f(x, u, \nabla u)$ are belongs to $L^{p'(x)}(\Omega)$ under $u \in W_0^{1,p(x)}(\Omega)$, the assumptions (A₂) and (A₄) and the given hypotheses about the exponents p, α, q and s because: $r(x) = (q(x) - 1)p'(x) \in C_+(\overline{\Omega})$ with $r(x) < p(x)$, and $\kappa(x) = (s(x) - 1)p'(x) \in C_+(\overline{\Omega})$ with $\kappa(x) < p(x)$.

Then, by Remark 2 we can conclude that

$$L^{p(x)} \hookrightarrow L^{r(x)} \quad \text{and} \quad L^{p(x)} \hookrightarrow L^{\kappa(x)}.$$

Hence, since $\varphi \in L^{p(x)}(\Omega)$, we have

$$\left(\mu g(x, u) + \lambda f(x, u, \nabla u) \right) \varphi \in L^1(\Omega).$$

This implies that, the integral

$$\int_{\Omega} \left(\mu g(x, u) + \lambda f(x, u, \nabla u) \right) \varphi dx$$

exist.

Then, we shall use the definition of weak solution for (1.1) in the following sense:

Definition 6. We say that a function $u \in W_0^{1,p(x)}(\Omega)$ is a weak solution of (1.1), if for any $\varphi \in W_0^{1,p(x)}(\Omega)$, it satisfies the following:

$$\int_{\Omega} \left(|\nabla u|^{p(x)-2} \nabla u \nabla \varphi + a(x) |\nabla u|^{q(x)-2} \nabla u \nabla \varphi \right) dx = \int_{\Omega} \left(\mu g(x, u) + \lambda f(x, u, \nabla u) \right) \varphi dx.$$

Before giving the existence result for the problem (1.1), we first give two lemmas that will be used in the proof of this result.

Let us consider the following functional:

$$\mathcal{J}(u) := \int_{\Omega} \frac{1}{p(x)} |\nabla u|^{p(x)} dx + \int_{\Omega} \frac{a(x)}{q(x)} |\nabla u|^{q(x)} dx.$$

From [12, 17], it is obvious that the derivative operator of the functional \mathcal{J} in the weak sense at the point $u \in W_0^{1,p(x)}(\Omega)$ is the functional $\mathcal{T}(u) := \mathcal{J}'(u) \in W^{-1,p'(x)}(\Omega)$, given by

$$\langle \mathcal{T}u, \varphi \rangle = \int_{\Omega} \left(|\nabla u|^{p(x)-2} \nabla u \nabla \varphi + a(x) |\nabla u|^{q(x)-2} \nabla u \nabla \varphi \right) dx,$$

for all $u, \varphi \in W_0^{1,p(x)}(\Omega)$ where $\langle \cdot, \cdot \rangle$ means the duality pairing between $W^{-1,p'(x)}(\Omega)$ and $W_0^{1,p(x)}(\Omega)$. Furthermore, the properties of the operator \mathcal{T} are summarized in the following lemma (see [12, Theorem 3.1] and [17, Proposition 3.1]).

Lemma 2. *The operator $\mathcal{T} : W_0^{1,p(x)}(\Omega) \rightarrow W^{-1,p'(x)}(\Omega)$ defined by*

$$\langle \mathcal{T}u, \varphi \rangle = \int_{\Omega} \left(|\nabla u|^{p(x)-2} \nabla u \nabla \varphi + a(x) |\nabla u|^{q(x)-2} \nabla u \nabla \varphi \right) dx, \quad (4.1)$$

is a continuous, bounded, strictly monotone operator, and is of class (S_+) .

Lemma 3. *If $(A_1) - (A_4)$ hold, then the operator $\mathcal{S} : W_0^{1,p(x)}(\Omega) \rightarrow W^{-1,p'(x)}(\Omega)$ defined by*

$$\langle \mathcal{S}u, \varphi \rangle = - \int_{\Omega} \left(\mu g(x, u) + \lambda f(x, u, \nabla u) \right) \varphi dx, \quad (4.2)$$

for all $u, \varphi \in W_0^{1,p(x)}(\Omega)$, is compact.

Proof. In order to prove this lemma, we proceed in three steps.

Step 1 : Let $\Psi_1 : W^{1,p(x)}(\Omega) \rightarrow L^{p'(x)}(\Omega)$ be an operator defined by

$$\Psi_1 u(x) := -\mu g(x, u).$$

In this step, we prove that the operator Ψ_1 is bounded and continuous.

First, let $u \in W^{1,p(x)}(\Omega)$, bearing (A_4) in mind and using (2.4) and (2.5), we infer

$$\begin{aligned} |\Psi_1 u|_{p'(x)} &\leq \rho_{p'(x)}(\Psi_1 u) + 1 \\ &= \int_{\Omega} |\mu g(x, u(x))|^{p'(x)} dx + 1 \\ &= \int_{\Omega} |\mu|^{p'(x)} |g(x, u(x))|^{p'(x)} dx + 1 \\ &\leq \left(|\mu|^{p'^-} + |\mu|^{p'^+} \right) \int_{\Omega} |\sigma(v(x) + |u|^{s(x)-1})|^{p'(x)} dx + 1 \\ &\leq \text{const} \left(|\mu|^{p'^-} + |\mu|^{p'^+} \right) \int_{\Omega} \left(|v(x)|^{p'(x)} + |u|^{k(x)} \right) dx + 1 \\ &\leq \text{const} \left(|\mu|^{p'^-} + |\mu|^{p'^+} \right) \left(\rho_{p'(x)}(v) + \rho_{k(x)}(u) \right) + 1 \\ &\leq \text{const} \left(|v|_{p(x)}^{p'^+} + |u|_{k(x)}^{k^+} + |u|_{k(x)}^{k^-} \right) + 1. \end{aligned}$$

Then, we deduce from (2.8) and $L^{p(x)} \hookrightarrow L^{k(x)}$, that

$$|\Psi_1 u|_{p'(x)} \leq \text{const} \left(|v|_{p(x)}^{p'^+} + |u|_{1,p(x)}^{k^+} + |u|_{1,p(x)}^{k^-} \right) + 1,$$

that means Ψ_1 is bounded on $W^{1,p(x)}(\Omega)$.

Second, we show that the operator Ψ_1 is continuous. To this purpose let $u_n \rightarrow u$ in $W^{1,p(x)}(\Omega)$. We need to show that $\Psi_1 u_n \rightarrow \Psi_1 u$ in $L^{p'(x)}(\Omega)$. We will apply the Lebesgue's theorem. Note that if $u_n \rightarrow u$ in $W^{1,p(x)}(\Omega)$, then $u_n \rightarrow u$ in $L^{p(x)}(\Omega)$. Hence there exist a subsequence (u_k) of (u_n) and ϕ in $L^{p(x)}(\Omega)$ such that

$$u_k(x) \rightarrow u(x) \text{ and } |u_k(x)| \leq \phi(x), \tag{4.3}$$

for a.e. $x \in \Omega$ and all $k \in \mathbb{N}$.

Hence, from (A_2) and (4.3), we have

$$|g(x, u_k(x))| \leq \sigma(v(x) + |\phi(x)|^{s(x)-1}),$$

for a.e. $x \in \Omega$ and for all $k \in \mathbb{N}$.

On the other hand, thanks to (A_3) and (4.3), we get, as $k \rightarrow \infty$

$$g(x, u_k(x)) \rightarrow g(x, u(x)) \text{ a.e. } x \in \Omega.$$

Seeing that $v + |\phi|^{s(x)-1} \in L^{p'(x)}(\Omega)$ and

$$\rho_{p'(x)}(\Psi_1 u_k - \Psi_1 u) = \int_{\Omega} |g(x, u_k(x)) - g(x, u(x))|^{p'(x)} dx,$$

then, from the Lebesgue's theorem and the equivalence (2.3), we have

$$\Psi_1 u_k \rightarrow \Psi_1 u \text{ in } L^{p'(x)}(\Omega),$$

and consequently

$$\Psi_1 u_n \rightarrow \Psi_1 u \text{ in } L^{p'(x)}(\Omega),$$

that is, Ψ_1 is continuous.

Step 2 :: Let us define the operator $\Psi_2 : W^{1,p(x)}(\Omega) \rightarrow L^{p'(x)}(\Omega)$ by

$$\Psi_2 u(x) := -\lambda f(x, u(x), \nabla u(x)).$$

We will show that Ψ_2 is bounded and continuous. Let $u \in W^{1,p(x)}(\Omega)$. According to (A_2) and the inequalities (2.4) and (2.5), we obtain

$$\begin{aligned} |\Psi_2 u|_{p'(x)} &\leq \rho_{p'(x)}(\Psi_2 u) + 1 \\ &= \int_{\Omega} |\lambda f(x, u(x), \nabla u(x))|^{p'(x)} dx + 1 \\ &= \int_{\Omega} |\lambda|^{p'(x)} |f(x, u(x), \nabla u(x))|^{p'(x)} dx + 1 \\ &\leq \left(|\lambda|^{p'^-} + |\lambda|^{p'^+} \right) \int_{\Omega} |\rho(\gamma(x) + |u|^{q(x)-1} + |\nabla u|^{q(x)-1})|^{p'(x)} dx + 1 \\ &\leq \text{const} \left(|\lambda|^{p'^-} + |\lambda|^{p'^+} \right) \int_{\Omega} \left(|\gamma(x)|^{p'(x)} + |u|^{r(x)} + |\nabla u|^{r(x)} \right) dx + 1 \\ &\leq \text{const} \left(|\lambda|^{p'^-} + |\lambda|^{p'^+} \right) \left(\rho_{p'(x)}(\gamma) + \rho_{r(x)}(u) + \rho_{r(x)}(\nabla u) \right) + 1 \\ &\leq \text{const} \left(|\gamma|_{p(x)}^{p'^+} + |u|_{r(x)}^{r^+} + |u|_{r(x)}^{r^-} + |\nabla u|_{r(x)}^{r^+} + |\nabla u|_{r(x)}^{r^-} \right) + 1. \end{aligned}$$

Taking into account that $L^{p(x)} \hookrightarrow L^{r(x)}$ and (2.8), we have then

$$|\Psi_2 u|_{p'(x)} \leq \text{const} \left(|\gamma|_{p(x)}^{p'^+} + |u|_{1,p(x)}^{r^+} + |u|_{1,p(x)}^{r^-} \right) + 1,$$

and consequently Ψ_2 is bounded on $W^{1,p(x)}(\Omega)$.

It remains to show that Ψ_2 is continuous. Let $u_n \rightarrow u$ in $W^{1,p(x)}(\Omega)$, we need to show that $\Psi_2 u_n \rightarrow \Psi_2 u$ in $L^{p'(x)}(\Omega)$. We will apply the Lebesgue's theorem.

Note that if $u_n \rightarrow u$ in $W^{1,p(x)}(\Omega)$, then $u_n \rightarrow u$ in $L^{p(x)}(\Omega)$ and $\nabla u_n \rightarrow \nabla u$ in $(L^{p(x)}(\Omega))^N$. Hence, there exist a subsequence (u_k) and ϕ in $L^{p(x)}(\Omega)$ and ψ in $(L^{p(x)}(\Omega))^N$ such that

$$u_k(x) \rightarrow u(x) \text{ and } \nabla u_k(x) \rightarrow \nabla u(x), \tag{4.4}$$

$$|u_k(x)| \leq \phi(x) \text{ and } |\nabla u_k(x)| \leq |\psi(x)|, \tag{4.5}$$

for a.e. $x \in \Omega$ and all $k \in \mathbb{N}$.

Hence, thanks to (A_1) and (4.4), we get, as $k \rightarrow \infty$

$$f(x, u_k(x), \nabla u_k(x)) \rightarrow f(x, u(x), \nabla u(x)) \text{ a.e. } x \in \Omega.$$

On the other hand, from (A_2) and (4.5), we can deduce the estimate

$$|f(x, u_k(x), \nabla u_k(x))| \leq \rho(\gamma(x) + |\phi(x)|^{q(x)-1} + |\psi(x)|^{q(x)-1}),$$

for a.e. $x \in \Omega$ and for all $k \in \mathbb{N}$.

Seeing that

$$\gamma + |\phi|^{q(x)-1} + |\psi(x)|^{q(x)-1} \in L^{p'(x)}(\Omega),$$

and taking into account the equality

$$\rho_{p'(x)}(\Psi_2 u_k - \Psi_2 u) = \int_{\Omega} |f(x, u_k(x), \nabla u_k(x)) - f(x, u(x), \nabla u(x))|^{p'(x)} dx,$$

then, we conclude from the Lebesgue's theorem and (2.3) that

$$\Psi_2 u_k \rightarrow \Psi_2 u \text{ in } L^{p'(x)}(\Omega),$$

and consequently

$$\Psi_2 u_n \rightarrow \Psi_2 u \text{ in } L^{p'(x)}(\Omega),$$

and then Ψ_2 is continuous.

Step 3: Let $I^* : L^{p'(x)}(\Omega) \rightarrow W^{-1,p'(x)}(\Omega)$ be the adjoint operator of the operator $I : W^{1,p(x)}(\Omega) \rightarrow L^{p(x)}(\Omega)$.

We then define

$$I^* \circ \Psi_1 : W^{1,p(x)}(\Omega) \rightarrow W^{-1,p'(x)}(\Omega),$$

and

$$I^* \circ \Psi_2 : W^{1,p(x)}(\Omega) \rightarrow W^{-1,p'(x)}(\Omega).$$

On another side, taking into account that I is compact, then I^* is compact. Thus, the compositions $I^* \circ \Psi_1$ and $I^* \circ \Psi_2$ are compact, that means $\mathcal{S} = I^* \circ \Psi_1 + I^* \circ \Psi_2$ is compact. With this last step the proof of Lemma 3 is completed. □

We are now in the position to get the existence result of weak solution for (1.1).

Theorem 2. *Assume that the assumptions $(A_1) - (A_4)$ hold, then the problem (1.1) possesses at least one weak solution u in $W_0^{1,p(x)}(\Omega)$.*

Proof. The basic idea of our proof is to reduce the problem (1.1) to a new one governed by a Hammerstein equation, and apply the theory of topological degree introduced in Section 3 to show the existence of weak solutions to the state problem.

First, for all $u, \varphi \in W_0^{1,p(x)}(\Omega)$, we define the operators \mathcal{T} and \mathcal{S} , as defined in (4.1) and (4.2) respectively. Consequently, the problem (1.1) is equivalent to the equation

$$\mathcal{T}u + \mathcal{S}u = 0, \quad u \in W_0^{1,p(x)}(\Omega). \tag{4.6}$$

Taking into account that, by Lemma 2, the operator \mathcal{T} is a continuous, bounded, strictly monotone and of class (\mathcal{S}_+) , then, by [25, Theorem 26 A], the inverse operator

$$\mathcal{L} := \mathcal{T}^{-1} : W^{-1,p'(x)}(\Omega) \rightarrow W_0^{1,p(x)}(\Omega),$$

is also bounded, continuous, strictly monotone and of class (S_+) .

On another side, according to Lemma 3, we have that the operator \mathcal{S} is bounded, continuous and quasimonotone. Consequently, following Zeidler’s terminology [25], the equation (4.6) is equivalent to the following abstract Hammerstein equation

$$u = \mathcal{L}\varphi \text{ and } \varphi + \mathcal{S} \circ \mathcal{L}\varphi = 0, \quad u \in W_0^{1,p(x)}(\Omega) \text{ and } \varphi \in W^{-1,p'(x)}(\Omega). \quad (4.7)$$

Seeing that (4.6) is equivalent to (4.7), then to solve (4.6) it is thus enough to solve (4.7). In order to solve (4.7), we will apply the Berkovits topological degree introducing in Section 3. First, let us set

$$\mathcal{B} := \left\{ \varphi \in W^{-1,p'(x)}(\Omega) : \exists t \in [0, 1] \text{ such that } \varphi + t\mathcal{S} \circ \mathcal{L}\varphi = 0 \right\}.$$

Next, we show that \mathcal{B} is bounded in $W^{-1,p'(x)}(\Omega)$. Let us put $u := \mathcal{L}\varphi$ for all $\varphi \in \mathcal{B}$. Taking into account that $|\mathcal{L}\varphi|_{1,p(x)} = |\nabla u|_{p(x)}$, then we have the following two cases:

First case: If $|\nabla u|_{p(x)} \leq 1$, then $|\mathcal{L}\varphi|_{1,p(x)} \leq 1$, that means $\left\{ \mathcal{L}\varphi : \varphi \in \mathcal{B} \right\}$ is bounded.

Second case: If $|\nabla u|_{p(x)} > 1$, then, we deduce from (2.1), (A_2) and (A_4) , the inequalities (2.6) and (2.5) and the Young’s inequality that

$$\begin{aligned} |\mathcal{L}\varphi|_{1,p(x)}^{p^-} &= |\nabla u|_{p(x)}^{p^-} \\ &\leq \rho_{p(x)}(\nabla u) \\ &\leq \langle \mathcal{T}u, u \rangle \\ &\leq \langle \varphi, \mathcal{L}\varphi \rangle \\ &= -t \langle \mathcal{S} \circ \mathcal{L}\varphi, \mathcal{L}\varphi \rangle \\ &= t \int_{\Omega} \left(\mu g(x, u) + \lambda f(x, u, \nabla u) \right) u dx \\ &\leq \text{const} \left(\int_{\Omega} |v(x)u(x)| dx + \int_{\Omega} |u(x)|^{s(x)} dx + \int_{\Omega} |\gamma(x)u(x)| dx \right. \\ &\quad \left. + \int_{\Omega} |u(x)|^{q(x)} dx + \int_{\Omega} |\nabla u|^{q(x)-1} |u| dx \right) \\ &\leq \text{const} \left(\int_{\Omega} |v(x)u(x)| dx + \int_{\Omega} |\gamma(x)u(x)| dx + \rho_{s(x)}(u) + \rho_{q(x)}(u) \right. \\ &\quad \left. + \int_{\Omega} |\nabla u|^{q(x)-1} |u| dx \right) \\ &\leq \text{const} \left(|v|_{p'(x)} |u|_{p(x)} + |\gamma|_{p'(x)} |u|_{p(x)} + |u|_{s(x)}^{s^+} + |u|_{s(x)}^{s^-} + |u|_{q(x)}^{q^+} + |u|_{q(x)}^{q^-} \right. \\ &\quad \left. + \frac{1}{q^-} \rho_{q(x)}(\nabla u) + \frac{1}{q^-} \rho_{q(x)}(u) \right) \\ &\leq \text{const} \left(|u|_{p(x)} + |u|_{s(x)}^{s^+} + |u|_{s(x)}^{s^-} + |u|_{q(x)}^{q^+} + |u|_{q(x)}^{q^-} + |\nabla u|_{q(x)}^{q^+} \right), \end{aligned}$$

then, according to $L^{p(x)} \hookrightarrow L^{s(x)}$ and $L^{p(x)} \hookrightarrow L^{q(x)}$, we get

$$|\mathcal{L}\varphi|_{1,p(x)}^{p^-} \leq \text{const} \left(|\mathcal{L}\varphi|_{1,p(x)} + |\mathcal{L}\varphi|_{1,p(x)}^{s^+} + |\mathcal{L}\varphi|_{1,p(x)}^{q^+} \right),$$

what implies that $\{\mathcal{L}\varphi : \varphi \in \mathcal{B}\}$ is bounded.

On the other hand, we have that the operator is \mathcal{S} is bounded, then $\mathcal{S} \circ \mathcal{L}\varphi$ is bounded. Thus, thanks to (4.7), we have that \mathcal{B} is bounded in $W^{-1,p'(x)}(\Omega)$. However, $\exists R > 0$ such that

$$|\varphi|_{-1,p'(x)} < R \text{ for all } \varphi \in \mathcal{B},$$

which leads to

$$\varphi + t\mathcal{S} \circ \mathcal{L}\varphi \neq 0, \quad \varphi \in \partial\mathcal{B}_R(0) \text{ and } t \in [0, 1],$$

where $\mathcal{B}_R(0)$ is the ball of center 0 and radius R in $W^{-1,p'(x)}(\Omega)$. Moreover, by Lemma 1, we conclude that

$$I + \mathcal{S} \circ \mathcal{L} \in \mathcal{F}_L(\overline{\mathcal{B}_R(0)}) \text{ and } I = \mathcal{T} \circ \mathcal{L} \in \mathcal{F}_L(\overline{\mathcal{B}_R(0)}).$$

On another side, taking into account that I, \mathcal{S} and \mathcal{L} are bounded, then $I + \mathcal{S} \circ \mathcal{L}$ is bounded. Hence, we infer that

$$I + \mathcal{S} \circ \mathcal{L} \in \mathcal{F}_{L,B}(\overline{\mathcal{B}_R(0)}) \text{ and } I = \mathcal{T} \circ \mathcal{L} \in \mathcal{F}_{L,B}(\overline{\mathcal{B}_R(0)}).$$

Next, we define the homotopy

$$\mathcal{H}: [0, 1] \times \overline{\mathcal{B}_R(0)} \rightarrow W^{-1,p'(x)}(\Omega) \tag{4.8}$$

$$(t, \varphi) \mapsto \mathcal{H}(t, \varphi) := \varphi + t\mathcal{S} \circ \mathcal{L}\varphi. \tag{4.9}$$

Applying the homotopy invariance and normalization property of the degree d seen in Theorem 1, we obtain

$$d(I + \mathcal{S} \circ \mathcal{L}, \mathcal{B}_R(0), 0) = d(I, \mathcal{B}_R(0), 0) = 1 \neq 0.$$

Since $d(I + \mathcal{S} \circ \mathcal{L}, \mathcal{B}_R(0), 0) \neq 0$, then by the existence property of the degree d stated in Theorem 1, we conclude that there exists $\varphi \in \mathcal{B}_R(0)$ which verifies

$$(I + \mathcal{S} \circ \mathcal{L})(\varphi) = 0 \Leftrightarrow \varphi + \mathcal{S} \circ \mathcal{L}\varphi = 0 \Leftrightarrow \mathcal{T} \circ \mathcal{L}\varphi + \mathcal{S} \circ \mathcal{L}\varphi = 0.$$

Finally, we infer that $u = \mathcal{L}\varphi$ is a weak solution of (1.1). The proof is completed.

□

REFERENCES

- [1] R. Aboulaich, D. Meskine, and A. Souissi, “New diffusion models in image processing,” *Computers & Mathematics with Applications*, vol. 56, no. 4, pp. 874–882, 2008, doi: [10.1016/j.camwa.2008.01.017](https://doi.org/10.1016/j.camwa.2008.01.017).
- [2] E. Acerbi and G. Mingione, “Regularity results for a class of quasiconvex functionals with non-standard growth,” *Annali della Scuola Normale Superiore di Pisa-Classe di Scienze*, vol. 30, no. 2, pp. 311–339, 2001.
- [3] S. N. Antontsev and S. I. Shmarev, “A model porous medium equation with variable exponent of nonlinearity: existence, uniqueness and localization properties of solutions,” *Nonlinear Analysis: Theory, Methods & Applications*, vol. 60, no. 3, pp. 515–545, 2005, doi: [10.1016/j.na.2004.09.026](https://doi.org/10.1016/j.na.2004.09.026).
- [4] A. Bahrouni, V. D. Rădulescu, and D. D. Repovš, “Double phase transonic flow problems with variable growth: nonlinear patterns and stationary waves,” *Nonlinearity*, vol. 32, no. 7, p. 2481, 2019, doi: [10.1088/1361-6544/ab0b03](https://doi.org/10.1088/1361-6544/ab0b03).
- [5] J. Berkovits, “Extension of the Leray–Schauder degree for abstract hammerstein type mappings,” *Journal of differential equations*, vol. 234, no. 1, pp. 289–310, 2007, doi: [10.1016/j.jde.2006.11.012](https://doi.org/10.1016/j.jde.2006.11.012).
- [6] M. El Ouaarabi, C. Allalou, and S. Melliani, “Existence of weak solutions for $p(x)$ -Laplacian-like problem with $p(x)$ -Laplacian operator under Neumann boundary condition,” *São Paulo Journal of Mathematical Sciences*, pp. 1–19, 2022, doi: [10.1007/s40863-022-00321-z](https://doi.org/10.1007/s40863-022-00321-z).
- [7] M. El Ouaarabi, C. Allalou, and S. Melliani, “On a class of $p(x)$ -Laplacian-like Dirichlet problem depending on three real parameters,” *Arabian Journal of Mathematics*, vol. 11, no. 2, pp. 227–239, 2022, doi: [10.1007/s40065-022-00372-2](https://doi.org/10.1007/s40065-022-00372-2).
- [8] M. El Ouaarabi, C. Allalou, and S. Melliani, “Weak solution of a Neumann boundary value problem with $p(x)$ -Laplacian-like operator,” *Analysis*, vol. 42, no. 4, pp. 271–280, 2022, doi: [10.1515/anly-2022-1063](https://doi.org/10.1515/anly-2022-1063).
- [9] M. El Ouaarabi, C. Allalou, and S. Melliani, “Existence result for Neumann problems with $p(x)$ -Laplacian-like operators in generalized Sobolev spaces,” *Rendiconti del Circolo Matematico di Palermo Series 2*, vol. 72, no. 2, pp. 1337–1350, 2023, doi: [10.1007/s12215-022-00733-y](https://doi.org/10.1007/s12215-022-00733-y).
- [10] M. El Ouaarabi, C. Allalou, and S. Melliani, “On a class of nonlinear degenerate elliptic equations in weighted Sobolev spaces,” *Georgian Mathematical Journal*, vol. 30, no. 1, pp. 81–94, 2023, doi: [10.1515/gmj-2022-2191](https://doi.org/10.1515/gmj-2022-2191).
- [11] M. El Ouaarabi, C. Allalou, and S. Melliani, “Weak solutions for double phase problem driven by the $(p(x), q(x))$ -Laplacian operator under Dirichlet boundary conditions,” *Boletim da Sociedade Paranaense de Matemática*, vol. 41, pp. 1–14, 2023, doi: [10.5269/bspm.62182](https://doi.org/10.5269/bspm.62182).
- [12] X.-L. Fan and Q.-H. Zhang, “Existence of solutions for $p(x)$ -laplacian Dirichlet problem,” *Nonlinear Analysis: Theory, Methods & Applications*, vol. 52, no. 8, pp. 1843–1852, 2003, doi: [10.1016/S0362-546X\(02\)00150-5](https://doi.org/10.1016/S0362-546X(02)00150-5).
- [13] X. Fan and D. Zhao, “On the spaces $L^{p(x)}(\Omega)$ and $W^{m,p(x)}(\Omega)$,” *J. Math. Anal. Appl.*, vol. 263, no. 2, pp. 424–446, 2001, doi: [10.1006/jmaa.2000.7617](https://doi.org/10.1006/jmaa.2000.7617).
- [14] P. Harjulehto, P. Hästö, M. Koskenoja, and S. Varonen, “The Dirichlet energy integral and variable exponent sobolev spaces with zero boundary values,” *Potential Analysis*, vol. 25, no. 3, pp. 205–222, 2006, doi: [10.1007/s11118-006-9023-3](https://doi.org/10.1007/s11118-006-9023-3).
- [15] I.-S. Kim and S.-J. Hong, “A topological degree for operators of generalized (S_+) type,” *Fixed Point Theory and Applications*, vol. 2015, no. 1, pp. 1–16, 2015, doi: [10.1186/s13663-015-0445-8](https://doi.org/10.1186/s13663-015-0445-8).
- [16] O. Kováčik and J. Rákosník, “On spaces $L^{p(x)}(\Omega)$ and $W^{1,p(x)}(\Omega)$,” *Czechoslovak Math. J.*, vol. 41, pp. 592–618, 1991, doi: [10.21136/CMJ.1991.102493](https://doi.org/10.21136/CMJ.1991.102493).

- [17] W. Liu and G. Dai, “Existence and multiplicity results for double phase problem,” *Journal of Differential Equations*, vol. 265, no. 9, pp. 4311–4334, 2018, doi: [10.1016/j.jde.2018.06.006](https://doi.org/10.1016/j.jde.2018.06.006).
- [18] M. A. Ragusa and A. Tachikawa, “On continuity of minimizers for certain quadratic growth functionals,” *Journal of the Mathematical Society of Japan*, vol. 57, no. 3, pp. 691–700, 2005, doi: [10.2969/jmsj/1158241929](https://doi.org/10.2969/jmsj/1158241929).
- [19] M. A. Ragusa and A. Tachikawa, “Regularity of minimizers of some variational integrals with discontinuity,” *Zeitschrift für Analysis und ihre Anwendungen*, vol. 27, no. 4, pp. 469–482, 2008, doi: [10.4171/ZAA/1366](https://doi.org/10.4171/ZAA/1366).
- [20] M. A. Ragusa and A. Tachikawa, “Boundary regularity of minimizers of $p(x)$ -energy functionals,” *Annales de l’Institut Henri Poincaré C, Analyse non linéaire*, vol. 33, no. 2, pp. 451–476, 2016, doi: [10.1016/j.anihpc.2014.11.003](https://doi.org/10.1016/j.anihpc.2014.11.003).
- [21] K. R. Rajagopal and M. Ruzicka, “Mathematical modeling of electrorheological materials,” *Continuum mechanics and thermodynamics*, vol. 13, no. 1, pp. 59–78, 2001, doi: [10.1007/s001610100034](https://doi.org/10.1007/s001610100034).
- [22] M. Ruzicka, “Electrorheological fluids: modeling and mathematical theory,” *Lecture Notes in Math*, vol. 1748, pp. 16–38, 2000, doi: [10.1007/BFb0104029](https://doi.org/10.1007/BFb0104029).
- [23] B.-S. Wang, G.-L. Hou, and B. Ge, “Existence of solutions for double-phase problems by topological degree,” *Journal of Fixed Point Theory and Applications*, vol. 23, no. 1, pp. 1–10, 2021, doi: [10.1007/s11784-021-00847-3](https://doi.org/10.1007/s11784-021-00847-3).
- [24] J. Yang, H. Chen, and S. Liu, “Multiplicity of nodal solutions for a class of double-phase problems,” *Journal of Function Spaces*, vol. 2020, 2020, doi: [10.1155/2020/3805803](https://doi.org/10.1155/2020/3805803).
- [25] E. Zeidler, *Nonlinear Functional Analysis and its Applications: II/B: Nonlinear Monotone Operators*. Springer, New York, 1990. doi: [10.1007/978-1-4612-0981-2](https://doi.org/10.1007/978-1-4612-0981-2).
- [26] V. V. Zhikov, “Averaging of functionals of the calculus of variations and elasticity theory,” *Izvestiya Rossiiskoi Akademii Nauk. Seriya Matematicheskaya*, vol. 50, no. 4, pp. 675–710, 1986, doi: [10.1070/IM1987v029n01ABEH000958](https://doi.org/10.1070/IM1987v029n01ABEH000958).
- [27] V. V. Zhikov, “On Lavrentiev’s phenomenon,” *Russian journal of mathematical physics*, vol. 3, pp. 249–269, 1995.
- [28] V. V. Zhikov, “On variational problems and nonlinear elliptic equations with nonstandard growth conditions,” *Journal of Mathematical Sciences*, vol. 173, no. 5, pp. 463–570, 2011, doi: [10.1007/s10958-011-0260-7](https://doi.org/10.1007/s10958-011-0260-7).

Authors’ addresses

Mohamed El Ouaarabi

(Corresponding author) Laboratory LMACS, FST of Beni Mellal, Sultan Moulay Slimane University, Beni Mellal, BP 523, 23000, Morocco

E-mail address: mohamed.elouaarabi@usms.ma

Chakir Allalou

Laboratory LMACS, FST of Beni Mellal, Sultan Moulay Slimane University, Beni Mellal, BP 523, 23000, Morocco

E-mail address: c.allalou@usms.ma

Said Melliani

Laboratory LMACS, FST of Beni Mellal, Sultan Moulay Slimane University, Beni Mellal, BP 523, 23000, Morocco

E-mail address: s.melliani@usms.ma