

EI Eridani: a star under the influence[★]

The effect of magnetic activity in the short and long term

L. Kriskovics^{1,2}, Zs. Kővári^{1,2}, B. Seli^{1,2,3}, K. Oláh^{1,2}, K. Vida^{1,2,3}, G. W. Henry⁴, T. Granzer⁵, and A. Görgei^{1,2,3}

¹ Konkoly Observatory, Research Centre for Astronomy and Earth Sciences, Konkoly Thege út 15-17., H-1121, Budapest, Hungary
e-mail: kriskovics.levente@csfk.org

² CSFK, MTA Centre of Excellence, Budapest, Konkoly Thege Miklós út 15-17., H-1121, Hungary

³ Eötvös Loránd University, Budapest, Hungary

⁴ Center of Excellence in Information Systems, Tennessee State University, Nashville, TN 37209, USA

⁵ Leibniz-Institut für Astrophysik Potsdam (AIP), An der Sternwarte 16, D14482 Potsdam, Germany

Received ...; accepted ...

ABSTRACT

Context. Homogeneous photometric time series spanning decades provide a unique opportunity to study the long-term cyclic behaviour of active spotted stars such as our target EI Eridani. In addition, with ultra-precise space photometry data, it is possible to investigate the accompanying flare activity in detail. However, the rotation period of ≈ 2 days for EI Eri makes it impossible to achieve time-resolved surface images from a single ground-based observing site. Therefore, for this purpose, spectroscopic data from a multi-site observing campaign are needed.

Aims. We use our photometric time series of more than forty years to analyze the long-term behaviour of EI Eri. Flare activity is investigated using space-borne photometric data obtained with the Transiting Exoplanet Survey Satellite (TESS). The MULTI-Site COntinuous Spectroscopy (MUSICOS) campaign in 1998 aimed to achieve high-resolution, multi-wavelength spectroscopic observations from many sites around the globe, so that uninterrupted phase coverage of EI Eri became available. We use these data to reconstruct successive surface temperature maps of the star in order to study the changes of starspots on a very short timescale.

Methods. We use long-term, seasonal period analysis of our photometric time series to study changes in the rotational period. Short-term Fourier-transform is also applied to look for activity cycle-like changes. We also study the phase and frequency distribution of hand-selected flares. We apply our multi-line Doppler imaging code to reconstruct four consecutive Doppler images. These images are also used to measure surface differential rotation by our cross-correlation technique. In addition, we carry out tests to demonstrate how Doppler imaging is affected by the fact that the data came from several different instruments with different spectral resolutions.

Results. Seasonal period analysis of the light curve reveals a smooth, significant change in period, possibly indicating the evolution of active latitudes. Temperature curves from $B - V$ and $V - I$ show slight differences, indicating the activity of EI Eri is spot dominated. Short-term Fourier transform reveals smoothly changing cycles between 4.5–5.5 and 8.9–11.6 years. The time-resolved spotted surface of EI Eri from Doppler imaging enabled us to follow the evolution of the different surface features. Cross-correlating the consecutive Doppler maps reveal surface shear of $\alpha = 0.036 \pm 0.007$. Our tests validate our approach and show that the surface temperature distribution is adequately reconstructed by our method. The tests also indicate how accurately the cross-correlation method can reproduce the surface shear as a function of the spectral resolution.

Key words. stars: activity – stars: imaging – starspots – stars: individual: EI Eridani

1. Introduction

EI Eridani = HD 26337 (G5 IV, $P_{\text{rot}} = 1.945$ days, $V = 7^{\text{m}}1$) is a well-known, rapidly rotating ($v \sin i = 51$ km/s) active RS CVn type, non-eclipsing, single-lined spectroscopic binary. It has been identified as an active star by Bidelman & MacConnell (1973) based on its Ca II H&K emission, which was later confirmed by Fekel (1980). Later on, Fekel et al. (1982) identified it as an RS CVn type variable. They also detected photometric variability with an amplitude of $V \approx 0^{\text{m}}2$ and a period of roughly two days. Fekel et al. (1986) derived an orbital period of 1.9472 days, while Hall et al. (1987) detected a photometric period of 1.945 ± 0.005 days from UBV photometry.

Its long-term photometric variability and behavior were extensively studied in the last four decades. Hall et al. (1987) reported seasonal changes in the photometric period on the order of one percent. Strassmeier et al. (1989) found a seasonal change of 0.043 days in the rotational period throughout three seasons. Oláh et al. (2012) also reported a $\pm 2\%$ deviation from the orbital period on a three-decade-long dataset.

It was Rodono & Cutispoto (1992) who first suggested a long-term cycle of roughly 10 years. Strassmeier et al. (1997) also found an 11 ± 1 years-long cycle in the sinusoidal change of a 16 years long photometric time-series, without any apparent relation between the cycle and the change of the seasonal periods. Oláh & Strassmeier (2002) also found photometric cycles of roughly 2.4 and 12.2 years, which were later confirmed and refined by Oláh et al. (2009) with values of $\approx 2.9 - 3.1$ and ≈ 14 years. Moreover, they reported a cycle with a length of roughly 4.1–4.9 years. EI Eri is also known to exhibit flares. Pandey &

[★] Original spectroscopic data acquisition was coordinated by Albert Washuettl (former researcher at AIP) and was carried out within the framework of the MULTI-Site COntinuous Spectroscopy (MUSICOS) campaign in 1998.




Fig. 1. Observing sites and phase distribution of the data. *Left panel:* MUSICOS sites involved in this work, namely Observatoire Haute-Provence (OHP), Xinglong Observatory (BXO), Kitt Peak National Observatory (KPNO), ESO La Silla (ESO), La Palma (INT) and Mt. Stromlo Observatory (MSO). *Right panel:* Phases of the observations plotted against the reduced Julian Date. Different colors denote different instruments (see legend). Observations plotted with filled circles were used for Doppler imaging, while the empty circles denote omitted observations. The vertical dashed lines separate the four subsets used for Doppler imaging.

Singh (2012) estimates the peak energy during flares in the 0.3–10 keV energy band to be $\approx 10^{31} - 10^{32}$ erg using observations obtained by the XMM-Newton X-ray observatory.

EI Eri has been among the prime targets of Doppler imaging in the past decades since the first application of the technique. Strassmeier (1990), Strassmeier et al. (1991) and Hatzes & Vogt (1992) reported a permanent polar spot with a radically changing appendage. The presence of the polar feature was confirmed by Washuettl et al. (2001) and later, Washuettl et al. (2009) and Kóvári et al. (2009). They also reported constantly changing low latitude features over 6–10 rotations (approximately two-three weeks). However, since the rotational period of EI Eri is very close to two days, covering a rotation with optimal phase coverage from a singular observing site takes around a month, but reaching adequate coverage also requires at least two weeks of time series. On images reconstructed using these datasets, shorter-lived features are suppressed and smeared out, which renders the resulting Doppler images unfeasible to show rapid changes. It also makes differential rotation measurements less reliable. As conventional, ground-based optical observing runs are regularly interrupted by daybreak, the only solution for this issue is a multi-site campaign. Consequently, EI Eri was chosen as one of the main targets for the MUSICOS campaign in 1998 organized at ESTEC/ESA by Bernard Foing and Joana Oliveira (for general details on the campaign, see Catala & Foing 1988). MUSICOS stands for MUlti-Site COntinuous Spectroscopy and aims to achieve high-resolution, multi-wavelength spectroscopic observations from many sites around the globe so that uninterrupted phase coverage of selected objects can be obtained.

The MUSICOS dataset presented in this paper was used to create preliminary Doppler images in order to measure surface differential rotation by Kóvári et al. (2009). They also confirm the presence of a stable polar spot with a slightly changing shape, and short-lived spots at lower latitudes, even though the rapid changes are somewhat masked by the artifacts that could be the result of the single-line inversion approach that was used. They also derived a surface differential rotation parameter of

$\alpha = \Delta\Omega/\Omega_{\text{eq}} = 0.0037$. This is congruent with the estimation of Oláh et al. (2012) based on long-term photometry.

This paper is structured as follows. After presenting the observations in Sect. 2, we study the long-term spot activity, including photometric cycles, and the seasonal change of rotational periods of EI Eri using a photometric time series of unprecedented length, covering four decades (see Sect. 3.1). Photometric data from the Transiting Exoplanet Survey Satellite (TESS) are used to investigate the flaring activity in Sect. 3.2. The rapid changes of the spot configuration are investigated using spectral time series from the MUSICOS 98 campaign in Sect. 4. We apply a multi-line Doppler imaging code on the MUSICOS dataset to reconstruct four Doppler images of EI Eri in order to investigate the short-term changes of the spotted surface. In the Appendix Sect. A, we perform tests to verify that the spectra with different spectral resolutions used for Doppler imaging do not introduce artifacts and bias the results.

2. Observations

2.1. Photometric observations

Photometric observations in Johnson V and I_C colors were carried out with the Wolfgang–Amadeus twin Automatic Photoelectric Telescope (APT), two robotic 75-cm telescopes for photoelectric photometry at Fairborn Observatory in the Sonoran desert near Tucson, Arizona (Strassmeier et al. 1997). The observations were made differentially with HD 25852 as the comparison star ($V = 7^m83$, $V - I_C = 1^m03$) and HD 26409 as check star ($V = 5^m44$, $B - V = 0^m94$).

Johnson B and V data were gathered with the Tennessee State University’s T3 0.4 m Automatic Photoelectric Telescope (APT) located also at Fairborn Observatory. For this dataset HD 26409 was used as comparison star. Part of the V dataset was published in Jetsu et al. (2017). All photometry was transformed to the Johnson–Cousins $BV(RI)_C$ system. In addition, older photometric observations since 1980 were used from the literature (see Oláh et al. 2009; Strassmeier et al. 1997, Table 5). The light curves of the ground-based photometry are shown in Fig. 2.




Fig. 2. Available ground-based photometry, calculated temperature curves and yearly rotational periods for EI Eri. *Top panel:* Ground-based Johnson V light curve of EI Eri from the Potsdam Wolfgang APT at Fairborn Observatory and from the literature (grey dots) and from Jetsu et al. (2017) and present paper (black dots). *Middle panel:* Temperatures from $B-V$ (blue) and $V-I$ (red). The dashed line and the grey area denote the effective temperature from spectral synthesis with 1σ error ($T_{\text{eff}} = 5580 \pm 70$ K, see Sect. 4.1). Black dot shows the averaged surface temperature value from the four Doppler maps ($\langle T \rangle = 5467 \pm 25$ K, see Sect. 4.3). *Bottom panel:* Yearly rotational periods from the V light curves with grey and black points, as on the upper panel and the two available TESS sectors (grey triangles). The dashed lines show the maximal (pole) and minimal (equator) rotational periods calculated using the differential rotation from the Doppler maps (see Sect. 4.4), while the dotted line denotes the orbital period.

TESS 120 s cadence data from sectors 5 and 32 are also available for EI Eri and were included in the photometric analysis, see Sect. 3.2.

2.2. Spectroscopic observations

The MUSICOS 1998 campaign involved eight northern and southern sites and ten telescopes to fulfill the needs of six scientific programs and took place from November 20 to December 14, 1998. For this work, we used data from seven instruments at six observing sites. These instruments are summarized in Table 1. Fig. 1 shows the distribution of these observatories: Observatoire Haute-Provence (OHP), Xinglong Observatory (BXO), Kitt Peak National Observatory (KPNO), ESO La Silla (ESO), La Palma (INT) and Mt. Stromlo Observatory (MSO).

For this specific program, a total of 90 high-resolution spectra of EI Eri were obtained within a time of 21 days (November 23 – December 13). The observing log is summarized in Table B.1. The various observing sites made use of different spectrographs (long slit, echelle), which leads to data sets of different quality, different wavelength ranges, and spectral resolutions. Fortunately, the region around 6400–6450 Å, which contains frequently used Doppler imaging lines (Fe I 6400, Fe I 6408, Fe I 6421, Fe I 6430 and Ca I 6439), is present in all of them. Several spectra were discarded due to low signal-to-noise or contaminated line profiles, resulting 59 selected spectra. The data were reduced using standard reduction techniques for echelle spectra in the NOAO packages of IRAF: bias subtraction, flat-field correction to remove pixel-to-pixel variations, and the curvature of the blaze function. For OHP, HEROS and Xinglong

Table 1. Sites and telescopes involved in this work.

Site	Telescope	Instrument	Resolving Power
OHP	1.93 m	ELODIE	43 000
ESO90	0.9 m	HEROS	20 000
ESO152	1.52 m	FEROS	48 000
KPNO	0.9 m	Echelle	65 000
BXO	2.2 m	Echelle	35 000
MSO	1.9 m	Echelle	35 000
INT	2.5 m	MUSICOS	35 000

Telescope abbreviations are as follows: the 1.93-m telescope at Observatoire Haute-Provence (OHP), the 90-cm and 152-cm telescopes at ESO La Silla (ESO90 and ESO152 respectively), the McMath-Pierce telescope at Kitt Peak National Observatory, Xinglong Observatory, Beijing (BXO), Mt. Stromlo, Australia (MSO) and the Isaac Newton Telescope at La Palma (INT).

observations, the MIDAS package was used. Background subtraction and flat-field correction using exposures of a tungsten lamp were applied. Wavelength calibration was performed by taking the spectra of a Thorium–Argon lamp. The spectra were normalized by a low-order polynomial fit or a cubic spline fit to the observed continuum.

Since the dataset is comprised of spectra from seven different instruments with different spectral resolutions, which can cause problems during the Doppler inversions, we decided to decrease the resolution of all spectra to the lowest available resolution of 20 000. In Appendix A we present a series of tests to validate this approach.

The observations are phased with the following ephemeris (Kóvári et al. 2009):

$$HJD = 2448054.7130 + 1.9472287 \times E. \quad (1)$$

Table B.1 shows the observing log, while Fig. 1 shows the phase distribution of the different measurements.


3. Photometric analysis

3.1. Analysis of the long-term photometry

Due to differential rotation, the period measured from photometry can slightly change, depending on the latitude where the dominant spots reside. Fig. 2 shows the measured rotational period of EI Eri in seasonal blocks, using the peak of the Lomb-Scargle periodogram (Lomb 1976; Scargle 1982) with uncertainties calculated from the width of the peaks at 90% power. The periods from the literature and Potsdam APT data (grey dots) are consistent with the data from Jetsu et al. (2017) and present paper (black dots) and with the two points from the TESS light curves (grey triangles) within one season. There is a smooth and

Table 2. Astrophysical parameters from spectral synthesis.

T_{eff}	5580 ± 70 K
$\log g$	3.75 ± 0.35
[Fe/H]	-0.27 ± 0.18
v_{mic}	2.0 ± 0.5 km/s
$v \sin i$	51 ± 1 km/s


Fig. 3. Short-term Fourier transform of all available V observations of EI Eri. A smoothly changing cycle is seen between 4.5–5.5 years (and its half), and a longer one which moves between 8.9–11.6 years.

significant change in the period, indicating the evolution of active latitudes.

The middle panel of Fig. 2 shows the temperatures calculated from $B - V$ and $V - I$ color indices by linearly interpolating the grid of Worthey & Lee (2011) at the given $\log g$ and metallicity. The dashed line and the grey area denote $T_{\text{eff}} = 5580 \pm 70$ K from spectral synthesis with 1σ error (for details, see Sect. 4.1). The average temperature calculated from the Doppler maps is plotted with a black dot, and the 1σ error is computed from the four different values for the four different maps (for details on the inversion, see Sect. 4.3).

To look for activity cycles, we used Short-term Fourier transform (Kolláth & Oláh 2009) for the present dataset, which is 41 years long, 13 years longer than the one used in Oláh et al. (2009). Since the algorithm requires uniform sampling in time, a cubic spline interpolation was used. We find that the previously identified structures on the STFT diagram are still present, indicating roughly the same cyclic behaviour. The signal of the longest, high-amplitude feature in Fig. 3 is suppressed, but above about 13.6 years, which is one-third of the length of the data, all signals have the same amplification. Smoothly changing cycles are seen between 4.5–5.5, and 8.9–11.6 years, which are not harmonics, since the speed and the direction of their changes are different. A weak cycle-like feature of about 2.5 years is also seen at the beginning of the time span.

3.2. Flares

We manually identified 41 flares in the two available TESS sectors (see Fig. 4). To calculate their energies in the TESS band, we followed similar procedures as in Oláh et al. (2022), using a BT-NextGen model spectrum (Hauschildt et al. 1999) convolved




Fig. 4. TESS observations from Sector 5 (left) and Sector 32 (right). The identified flares are marked with black.




Fig. 5. Flare frequency distribution from the TESS data. The dashed line denotes a fit for all of the points with a power-law index of 1.66 ± 0.04 , while the two solid lines show a two-component fit (1.41 ± 0.02 and 2.26 ± 0.08), see Sect. 3.2.

with the TESS transmission curve to get the quiescent stellar luminosity of $L_{\text{TESS}} = 4.69 \cdot 10^{33} \text{ erg s}^{-1}$, and integrating over the flare light curves to get the equivalent durations. The flare frequency distribution (FFD) in Fig. 5 shows a broken power law shape. Two-component fits of the FFD yield power law indices of 1.41 ± 0.02 and 2.26 ± 0.08 (one minus the slope of the fitted line) for the first and second parts, respectively, while a fit for all datapoints gives 1.66 ± 0.04 (for further discussion, see Sect. 5).

Fig. 6 shows the phase distribution of flares, in the reference frame of the orbit, using the ephemeris from Eq. 1. While the spot configuration of EI Eri changes quickly, the orbiting secondary component provides a solid reference frame. To test whether there is a significant increase of flares at given phases, we run a two-sided Kuiper’s test (Kuiper 1960). It compares the measured data to a given distribution, similar to the Kolmogorov–Smirnov test, but it is invariant under a cyclic transformation, which makes it applicable to compare cyclic phase distributions (the Kolmogorov–Smirnov test would give different results for different starting epochs). The test rejects the null hypothesis of a uniform flare phase distribution with a p -value of 0.006. The increase in flare rate appears on the side facing the secondary component.




Fig. 6. Phase distribution of the TESS flares on the primary component in the reference frame of the binary orbit. The relative radii and separation of the circles are for scale. The colors are realistic digital colors for the given spectral types from Harre & Heller (2021). The grey area shows a Gaussian kernel density estimation with the bandwidth of 2° . Black ticks denote the orbital phase values of the flare observations.

4. Doppler imaging

4.1. Astrophysical parameters for the inversion

Precise astrophysical parameters are fundamental for Doppler inversion. Therefore we carried out a detailed spectroscopic analysis based on spectral synthesis using the code SME (Piskunov & Valenti 2017) and MARCS atmospheric models (Gustafsson et al. 2008). Atomic line parameters were taken from the VALD database (Kupka et al. 1999). Macroturbulence was estimated using the following equation (Valenti & Fischer 2005):

$$v_{\text{mac}} = \left(3.98 - \frac{T_{\text{eff}} - 5770\text{K}}{650\text{K}} \right) \text{ km s}^{-1}. \quad (2)$$

Astrophysical parameters were fitted one-by-one iteratively as described in Kriskovics et al. (2019). The results are summarized in Table 2.

4.2. The Doppler imaging code *iMap*

The Doppler imaging code *iMap* (Carroll et al. 2012), which is used in this work, carries out multi-line Doppler inversion on a list of photospheric lines. Since the wavelength coverage of our dataset is limited, we used five (Fe I 6400, Fe I 6408, Fe I 6421,




Fig. 7. Four consecutive Doppler images of EI Eri obtained for the MUSICOS 1998 data. The corresponding time intervals for the four images from top to bottom are 2451141.3758–2451144.9045, 2451145.6007–2451148.7618, 2451148.9391–2451150.7474 and 2451150.9827–2451156.5520.

Fe I 6430 and Ca I 6439) virtually unblended lines with well-defined continuum, suitable line depth, and temperature sensitivity. The stellar surface was divided into $6^\circ \times 6^\circ$ segments. For each local line profile, the code utilizes a full radiative solver (Carroll et al. 2008).

Then the local line profiles are disk integrated, and the individually modeled, disk-integrated lines are averaged. Atomic line data are taken from VALD (Kupka et al. 1999). Model atmospheres are taken from Castelli & Kurucz (2003) and are interpolated for the necessary temperature, gravity, or metallicity values. Due to the high computational capacity requirements,

LTE radiative transfer is used instead of spherical model atmospheres, but imperfections in the fitted line shapes are well compensated by the multi-line approach. Additional input parameters are micro- and macroturbulence, and $v \sin i$. Table 2 summarizes the astrophysical parameters used during the inversion. Inclination was set to $i = 56^\circ$ (adopted from Washuettl et al. 2009).

For the surface reconstruction, an iterative regularization method based on a Landweber algorithm is used (Carroll et al. 2012), meaning no additional constraints are imposed in the image domain.




Fig. 8. Measured surface differential rotation of EI Eri and its comparison to other stars. *Left panel:* Average cross-correlation map for EI Eri. The correlation peaks (white circles) are fitted with a quadratic differential rotation law. The resulting fit (continuous line) indicates a solar type differential rotation with a shear parameter $\alpha = 0.036 \pm 0.007$. *Right panel:* An extended version of the surface shear coefficient versus rotational period plot from Kóvári et al. 2017. Squares denote results from the cross-correlation technique, while circles show differential rotational coefficients from the sheared image method. White symbols correspond to single, greys to binary stars. Symbol size increases from dwarfs to subgiants to giants. EI Eri is represented by the blue square at $P_{\text{rot}} \approx 2$ days. The Sun is plotted with a dotted circle. The dotted and dash-dotted lines denote the linear fits to the points corresponding to singles and binaries, respectively, with slopes of $|\alpha| \propto (0.0049 \pm 0.00001)P_{\text{rot}}$ and $|\alpha| \propto (0.0014 \pm 0.00003)P_{\text{rot}}$.

4.3. Time-resolved surface evolution

The 59 selected spectra were divided into four subsets (hereafter DI1, DI2, DI3 and DI4) corresponding to the following Julian Date intervals: 2451141.3758–2451144.9045, 2451145.6007–2451148.7618, 2451148.9391–2451150.7474 and 2451150.9827–2451156.5520, with lengths of roughly 1.8, 1.6, 1.0 and 2.9 rotations, respectively. Imaging lines were selected from the spectral region that was covered by all of the spectra in order to make the resulting maps comparable. This gives us a unique opportunity to study the short-term spot evolution of EI Eri. The selected spectral lines are Fe I 6400, Fe I 6408, Fe I 6421, Fe I 6430 and Ca I 6439.

The resulting four Doppler images (DI1–DI4) are shown in Fig. 7. The corresponding line profile fits are shown in Fig. C.1, with corresponding root mean square deviations of 0.0030, 0.0030, 0.0036 and 0.0040, respectively. The four maps show a persistent polar feature (with a spot temperature of $\Delta T \approx 1100$ K below the temperature of the unspotted surface. (Here we refer to our tests presented in Sect. A of the Appendix, during which we also cover the accuracy of the spot temperature determination.) Associated with the dominant polar spot, a less prominent appendage appears in DI2 around $\phi \approx 0.35$ phase value, which is further strengthened in DI3. In DI4, the appendage starts to fade, but it is still present. The overall contrast of the polar feature seems to increase from DI1 towards DI4. With all this, several low-latitude features show rapid evolution as well. In DI1, the barely visible spot at $\phi \approx 0.45$ in the lower hemisphere becomes stronger in DI2, and after a slight retrograde shift seen in DI3, it disappears completely in DI4, being present for ~ 4.5 rotations. Around 0.2 phase, a low latitude feature emerges in DI2 and becomes more and more prominent. In DI2 at $\phi \approx 0.7$, a low latitude spot forms and strengthens along DI3 and DI4, while gradually shifting toward the visible pole.

In the end, it is worth emphasizing that the evolution of the different surface features can be followed quite nicely through the four *independently* reconstructed maps. Therefore we believe

that most of the recovered surface features are real and not imaging artifacts.

4.4. Differential rotation

A usual way of measuring surface differential rotation is cross-correlating consecutive Doppler images (Donati & Collier Cameron 1997). However, rapid spot evolution can easily hinder the correlation pattern, e.g., through the formation or dimming of spots or interaction between nearby spots, all of which can be seen in the Doppler images of EI Eri. To overcome this, and strengthen the signal of differential rotation, we use our code ACCORD (e.g. Kóvári et al. 2012), which uses the average of the cross-correlations of consecutive Doppler images. The latitudinal correlation peaks in the resulting average correlation map are fitted with a quadratic rotational law in the form

$$\Omega(\beta) = \Omega_{\text{eq}} - \Delta\Omega \sin^2 \beta, \quad (3)$$

where $\Omega(\beta)$ is the angular velocity at β latitude, Ω_{eq} is the angular velocity of the equator, and $\Delta\Omega = \Omega_{\text{eq}} - \Omega_{\text{pole}}$ gives the difference between the equatorial and polar angular velocities. With these, the dimensionless surface shear parameter α is defined as $\alpha = \Delta\Omega/\Omega_{\text{eq}}$.

For the average cross-correlation we formed three pairs of consecutive Doppler images, that is DI1–DI2, DI2–DI3, and DI3–DI4. After the cross-correlation of the image pairs, we normalized each resulting correlation map to the same time difference so that they can be averaged. For a more detailed description of the ACCORD method see Kóvári et al. (2015) and their references. In the left panel of Fig. 8 the average correlation pattern is fitted with a quadratic differential rotation law which yielded $\alpha = 0.036 \pm 0.007$ ($\Omega_{\text{eq}} = 186.922 \pm 0.384$ degree/day, $\Delta\Omega = 6.768 \pm 1.349$ degree/day). The errors are estimated from the amplitudes and the full width at half maximum values (FWHMs) of the Gaussian functions fitted to each longitudinal strip.

In the right panel of Fig. 8 we put the surface shear of EI Eri against the rotation period together with other well-known active stars as an updated plot of Kóvári et al. (2017). Our resulting shear value of 3.6% for EI Eri is in good agreement with the value expected according to the plot. We note, however, that the differential rotation pattern can be somewhat diminished by the relatively low spectral resolution; see our tests to investigate this question in Appendix A.

5. Discussion

5.1. Long term photometric behaviour

Kolláth & Oláh (2009) found shorter cycles of 2.9–3.1 and 4.1–4.9 years and a longer cycle of roughly 14 years on EI Eri. Our cycle lengths derived from a 13 years longer dataset of roughly 2.5, 4.5–5.5 and 8.9–11.6 years are in good agreement with these previous findings (in Fig. 3, the longest 13.6 year cycle is suppressed since the dataset is only three times longer than this value). The cycle lengths are smoothly changing, which is a well-documented behavior of activity cycles, see examples in e.g. Oláh et al. (2009), or the solar case in Kolláth & Oláh (2009). Visual inspection of the curve of seasonal rotation period changes (bottom panel of Fig. 2) hints at a cycle-like behavior of around 3500–4000 days (roughly 9.5–11 years). The similarity between the lengths of the cycle-like features of the long-term photometry and the change of the seasonal rotational periods around 9–11 years is indeed remarkable. However, there are problems in interpreting the photometric results due to the huge spots and the inclination of the stellar disk to the line of sight, which makes the tracing of the spot’s position and sizes complicated. Therefore, one-by-one correspondence of the brightness changes and rotational periods (like a butterfly diagram) cannot be drawn.

The temperatures calculated from $B - V$ and $V - I$ color indices show slightly different amplitudes. The higher temperature of the facular/plage regions contributes to the signal in the B bandpass, although the spots from the photosphere also have a lower contribution in B , whereas the signal in the I filter has no or negligible contribution from the higher temperature areas. The spot–plage ratio can be different star by star, and also within spot cycles of the same star. The difference between the $B - V$ and $V - I$ color indices could indicate that the activity of EI Eri is dominated by spots, rather than plages. This is also supported by the large spotted areas present on the Doppler maps and the fact that the temperature from $B - V$ correlates with the effective temperature from the spectral synthesis, while the average temperature from the Doppler images is in the interval of the temperatures calculated from $V - I$. We note, however, that the photometric accuracy of the color indices may hinder this result. Uncertainties in the order of 0.01 magnitude translate to an error of about 30 K in temperature. Nevertheless, average temperature coming from summing up the temperature elements of the surface of the Doppler images agree very well with the temperature values of the same season resulting from $V - I$, which indicates a reasonable accuracy of the results from the color indices.

5.2. Flare activity

The possible origin of breakpoints in FFDs was discussed by Mullan & Paudel (2018) for flare stars showing a critical energy value on the FFD when the flaring loop size exceeds the local scale height, which depend on local field strengths and densities. Above this value, the flares have higher energy and the power

law index is steeper, and the opposite stands below this energy value for smaller loops.

Extrapolating their results, Oláh et al. (2021) gave examples of broken power-law FFDs for two giant stars with different breakpoints of their FFDs but similar energy ranges suggesting different atmospheric properties and strengths of magnetic fields. In the case of the subgiant EI Eri this rationale may hold as well. The median flare duration of about 1.5 hours and the energy range shown in Fig. 5 are fully consistent with the general picture of flare energy–duration diagram given by Maehara et al. (2021, see their Fig. 14), placing EI Eri somewhere between G dwarfs and giants on the flare energy–flare duration diagram.

When an active star is in a close binary system, the tidal effects may play a role in the position of the emerging magnetic flux tubes that are supposed to be the origin of activity on stars. Such effects supposing different physical circumstances (like the strength of the magnetic field, the depth of the convection zone where the flux tubes originate, local density, etc.) were investigated in two papers by Holzwarth & Schüssler (2003a,b) showing that clustering of flux tube emergence positions in the orbital reference frame of close binaries may appear. Our results plotted in Fig. 6, where the observed flare phases are shown in the reference frame of the orbit, show a clustering on the hemisphere facing the secondary and may originate from such an effect.

5.3. Rapid spot evolution and differential rotation

Rapid spot evolution is not unheard of on subgiants. Strassmeier & Bartus (2000) found fast-evolving, low latitude features on V711 Tau, a similar binary subgiant (although we note that V711 Tau is somewhat cooler and its secondary is more massive). They found that in several cases, low-latitude features changed shape and became cooler on a scale of a few days – similar to what we found on EI Eri. Gradual strengthening of the polar feature is also observed on V711 Tau and EI Eri. It might indicate spot migration towards the poles. Moreover, Harutyunyan et al. (2016) reported rapid spot evolution on another subgiant, HU Vir, with a prominent, high-latitude, but not polar feature, which drastically changes shape from one rotation to the next. We note, however, that due to the nature of Doppler imaging, we have no information about the spot evolution on time scales shorter than the ≈ 10 -d rotational period of HU Vir.

Polar features persisting for years have been previously observed on subgiants (see e.g. Hussain 2002, and references therein), also it is a fairly common and well-explained phenomenon (even from a theoretical viewpoint) on fast-rotating active stars (e.g. Işık et al. 2007). However, rapid spot changes on the scale of a few days are much more difficult to explain with models. Işık et al. (2007) even point out that the lifetime of their simulated bipolar magnetic regions (BMRs) are longer on a star similar to V711 Tau than on a solar analog. According to their simulation, a large BMR above $\beta = 70^\circ$ latitude can be present for more than two years. This may even be consistent with the permanent polar features of EI Eri.

However, BMRs and spots can behave differently: BMRs consist of spots, plages, and short-lived structures. It is also debated whether large spots on Doppler images are single magnetic structures or clusters of smaller spots. Accordingly, it is possible that different evolution timescales are applicable to a homogeneous, large spot than to several smaller spots. Işık et al. (2007) showed that a cluster of spots (that still cannot be resolved by Doppler imaging) can dissipate the fastest in their simulation, especially when large-scale flows (e.g., meridional circulation) are introduced, with a decay time of at least a few tens of days.

Strassmeier & Bartus (2000) theorized that magnetic reconnection could play an important role in this phenomenon: it may induce magnetic field diffusion from the stronger flux tubes (i.e., the cooler spots) towards the weaker (warmer) ones, resulting in increased magnetic pressure and thus more suppressed convection and rapid cooling in the warmer spots. They also estimated that on V711 Tau, the Alfvén-velocity for the photosphere region corresponding to $\tau=1$ optical depth results in a magnetic interaction timescale between spots, which is on the order of a day. Thus, energy transport by Alfvén-waves may contribute to the rapid spot changes.

Rempel & Cheung (2014) carried out three-dimensional numerical simulations to study spot emergence and decay. They found that in the first phase of spot decay, the process is dominated by downward convective motion, while in the second phase, plasma intrusions fragment larger spots. The latter could be consistent with the assumptions of Strassmeier & Bartus (2000). Rempel & Cheung (2014) pointed out that strong, sub-surface large-scale convective motions can play a significant role in spot decay (however, their simulation box was limited to the upper part of the convective zone).

Namekata et al. (2019, and their Fig. 6) compares observed stellar spot lifetimes on different solar-type stars. Their result suggests that if spot evolution is driven by the same mechanism in all evolutionary stages, the typical spot lifetime on EI Eri should be in the range of 10–30 days. Our findings (and previous ones in the literature, see above) are in agreement with these results, since most of the active nests are present on all of the Doppler images covering altogether 15 days, although with changing shape and contrast (in agreement with Strassmeier & Bartus 2000). The TESS light curves (Fig. 2) also support this, e.g. a new nest seems to form around JD=2457143 which is persistent for at least ≈ 10 days, although with different amplitude. We note, however, that datasets spanning a much longer time period is required to confirm or disprove this.

The surface differential rotation parameter of $\alpha = 0.036 \pm 0.007$ fits well to the observation that rapidly rotating subgiants exhibit relatively low surface shear. Harutyunyan et al. (2016) measured $\alpha = -0.029 \pm 0.005$ on HU Vir, another K subgiant in a binary. On the K subgiant IL Hya, Kóvári & Weber (2004) reported $\alpha \approx 0.03$. Our results are also in good agreement with the empirical relation of $|\alpha| \approx 0.013 P_{\text{rot}}$ for binaries originally suggested by Kóvári et al. (2017) (see also the right panel of our Fig. 8).

6. Summary

Our study of the main component of the single lined RS CVn binary EI Eri based on a 4 decades long photometric time series and a multi-site high resolution spectroscopic campaign yielded the following results:

- Based on a 41 year long photometric dataset, we derive a roughly 2.5, a 4.5–5.5 and a 8.9–11.6 year photometric cycle. The apparent periodicity in the seasonal changes of the rotational period coincides with the longest cycle. The long term $B - V$ and $V - I$ multicolor photometry indicates spot dominated activity, rather than plage dominated.
- The FFD of 41 flares shows a broken power law shape with power law indices of 1.41 ± 0.02 and 2.26 ± 0.08 . According to Mullan & Paudel (2018), this might indicate that there is a critical energy break point on the FFD, where the flaring loop size exceeds the local scale height, resulting in higher energy flares and steeper power law indices. Phase distribution of

the flares indicate that magnetic flux emergence is affected by tidal effects (Holzwarth & Schüssler 2003a,b), and flares are concentrated on the hemisphere facing the secondary.

- Using our MUSICOS spectral time series, we derived four consecutive, independent Doppler-images of EI Eri, covering 1.8, 1.6, 1.0 and 2.9 rotations, enabling us to study very short term spot evolution on the rapidly evolving surface. The resolution of the spectra were blurred down to the lowest available resolution in order to avoid artifacts caused by different resolution. This approach was validated by a series of tests. The Doppler images revealed a strong, evolving, but always present polar cap and several emerging and disappearing low latitude features of constantly evolving shape and contrast. The approximate lifetime of these structures coincide with the lower limit proposed by Namekata et al. (2019), although datasets covering longer time periods are needed in order to conclusively verify this.
- Cross correlating the consecutive Doppler images revealed a weak, solar-like surface differential rotational pattern of $\alpha = 0.036 \pm 0.007$, which is in good agreement with the empirical law for binaries proposed by Kóvári et al. (2017).

Our paper also includes tests to show that our multi-instrument time-series can be used for Doppler imaging without introducing artifacts, if treated properly.

Acknowledgements. The authors would like to thank Albert Washuettl and the original MUSICOS team for organizing and gathering the data used in this work. The authors acknowledge the Hungarian National Research, Development and Innovation Office grants OTKA K-131508, KKP-143986 (Élvonal), and 2019-2.1.11-TÉT-2019-00056. LK acknowledges the Hungarian National Research, Development and Innovation Office grant OTKA PD-134784. LK and KV are Bolyai János research Fellows. KV is supported by the Bolyai+ grant ÚNKP-22-5-ELTE-1093, BS is supported by the ÚNKP-22-3 New National Excellence Program of the Ministry for Culture and Innovation from the source of the National Research, Development and Innovation Fund. GWK acknowledges long-term support from NASA, NSF, and the State of Tennessee through its Centers of Excellence Program.

References

- Bidelman, W. P. & MacConnell, D. J. 1973, *AJ*, 78, 687
 Carroll, T. A., Kopf, M., & Strassmeier, K. G. 2008, *A&A*, 488, 781
 Carroll, T. A., Strassmeier, K. G., Rice, J. B., & Künstler, A. 2012, *A&A*, 548, A95
 Castelli, F. & Kurucz, R. L. 2003, in *Modelling of Stellar Atmospheres*, ed. N. Piskunov, W. W. Weiss, & D. F. Gray, Vol. 210, A20
 Catala, C. & Foing, B. H. 1988, in *Comptes-rendus du premier atelier MUSICOS: Multi-Site Continuous Spectroscopy; Spectroscopie Multi-Sites en Continu*
 Donati, J. F. & Collier Cameron, A. 1997, *MNRAS*, 291, 1
 Fekel, F. C., Hall, D. S., Henry, G. W., Landis, H. J., & Renner, T. R. 1982, *Information Bulletin on Variable Stars*, 2110, 1
 Fekel, F. C., Moffett, T. J., & Henry, G. W. 1986, *ApJS*, 60, 551
 Fekel, F. C., J. 1980, *SAO Special Report*, 389, 133
 Gustafsson, B., Edvardsson, B., Eriksson, K., et al. 2008, *A&A*, 486, 951
 Hall, D. S., Osborn, S. A. G., Seufert, E. R., et al. 1987, *AJ*, 94, 723
 Harre, J.-V. & Heller, R. 2021, *Astronomische Nachrichten*, 342, 578
 Harutyunyan, G., Strassmeier, K. G., Künstler, A., Carroll, T. A., & Weber, M. 2016, *A&A*, 592, A117
 Hatzes, A. P. & Vogt, S. S. 1992, *MNRAS*, 258, 387
 Hauschildt, P. H., Allard, F., & Baron, E. 1999, *ApJ*, 512, 377
 Holzwarth, V. & Schüssler, M. 2003a, *A&A*, 405, 291
 Holzwarth, V. & Schüssler, M. 2003b, *A&A*, 405, 303
 Hussain, G. A. J. 2002, *Astronomische Nachrichten*, 323, 349
 Işık, E., Schüssler, M., & Solanki, S. K. 2007, *A&A*, 464, 1049
 Jetsu, L., Henry, G. W., & Lehtinen, J. 2017, *ApJ*, 838, 122
 Kóvári, Z., Kriskovics, L., Künstler, A., et al. 2015, *A&A*, 573, A98
 Kóvári, Zs., Korhonen, H., Kriskovics, L., et al. 2012, *A&A*, 539, A50
 Kóvári, Zs., Oláh, K., Kriskovics, L., et al. 2017, *Astronomische Nachrichten*, 338, 903

- Kővári, Zs., Washuettl, A., Foing, B. H., et al. 2009, in American Institute of Physics Conference Series, Vol. 1094, 15th Cambridge Workshop on Cool Stars, Stellar Systems, and the Sun, ed. E. Stempels, 676–679
- Kővári, Zs. & Weber, M. 2004, Publications of the Astronomy Department of the Eötvös Loránd University, 14, 221
- Kolláth, Z. & Oláh, K. 2009, A&A, 501, 695
- Krskovics, L., Kővári, Zs., Vida, K., et al. 2019, A&A, 627, A52
- Kuiper, N. H. 1960, in Nederl. Akad. Wetensch. Proc. Ser. A, Vol. 63, 38–47
- Kupka, F., Piskunov, N., Ryabchikova, T. A., Stempels, H. C., & Weiss, W. W. 1999, A&AS, 138, 119
- Lomb, N. R. 1976, Ap&SS, 39, 447
- Maehara, H., Notsu, Y., Namekata, K., et al. 2021, PASJ, 73, 44
- Mullan, D. J. & Paudel, R. R. 2018, ApJ, 854, 14
- Namekata, K., Maehara, H., Notsu, Y., et al. 2019, ApJ, 871, 187
- Oláh, K., Kővári, Z., Günther, M. N., et al. 2021, A&A, 647, A62
- Oláh, K., Kővári, Zs., Vida, K., & Strassmeier, K. G. 2012, in From Interacting Binaries to Exoplanets: Essential Modeling Tools, ed. M. T. Richards & I. Hubeny, Vol. 282, 478–479
- Oláh, K., Kolláth, Z., Granzer, T., et al. 2009, A&A, 501, 703
- Oláh, K., Seli, B., Kővári, Z., Krskovics, L., & Vida, K. 2022, arXiv e-prints, arXiv:2210.09710
- Oláh, K. & Strassmeier, K. G. 2002, Astronomische Nachrichten, 323, 361
- Pandey, J. C. & Singh, K. P. 2012, MNRAS, 419, 1219
- Piskunov, N. & Valenti, J. A. 2017, A&A, 597, A16
- Rempel, M. & Cheung, M. C. M. 2014, ApJ, 785, 90
- Rodono, M. & Cutispoto, G. 1992, A&AS, 95, 55
- Scargle, J. D. 1982, ApJ, 263, 835
- Strassmeier, K. G. 1990, ApJ, 348, 682
- Strassmeier, K. G. & Bartus, J. 2000, A&A, 354, 537
- Strassmeier, K. G., Bartus, J., Cutispoto, G., & Rodonó, M. 1997, A&AS, 125, 11
- Strassmeier, K. G., Boyd, L. J., Epan, D. H., & Granzer, T. 1997, PASP, 109, 697
- Strassmeier, K. G., Hall, D. S., Boyd, L. J., & Genet, R. M. 1989, ApJS, 69, 141
- Strassmeier, K. G., Rice, J. B., Wehlau, W. H., et al. 1991, A&A, 247, 130
- Valenti, J. A. & Fischer, D. A. 2005, ApJS, 159, 141
- Washuettl, A., Strassmeier, K. G., & Collier-Cameron, A. 2001, in Astronomical Society of the Pacific Conference Series, Vol. 223, 11th Cambridge Workshop on Cool Stars, Stellar Systems and the Sun, ed. R. J. Garcia Lopez, R. Rebolo, & M. R. Zapaterio Osorio, 1308
- Washuettl, A., Strassmeier, K. G., & Weber, M. 2009, Astronomische Nachrichten, 330, 366
- Worthey, G. & Lee, H.-c. 2011, ApJS, 193, 1

Appendix A: Resolution tests

MUSICOS 1998 data of EI Eri were obtained from seven different instruments with different spectral resolutions (and hence, there are spectra with different resolutions in each subset), which can induce artifacts on the Doppler maps due to the different levels of detail of the spectral features and the different instrumental profiles. To counteract this, we used a Gaussian kernel to decrease the resolution of the spectra to our lowest available resolution. Since this has not been done before, we carry out tests to see the effect of this treatment on the high resolution dataset of V1358 Ori (Kriskovics et al. 2019). We created datasets of decreased resolution corresponding to $R=20\,000$, $30\,000$, $40\,000$, $50\,000$, $60\,000$ and $70\,000$, and compared it to the original maps. For the details on the original dataset, astrophysical parameters and inversion, we refer to Kriskovics et al. (2019).

Doppler images for the datasets of decreased resolutions were derived with the same inversion settings and astrophysical parameters. Hotter features were not suppressed on purpose, to see if artifacts are introduced on lower resolutions. Fig. A.1 shows examples for both rotations corresponding to $R=60\,000$,

$40\,000$ and $20\,000$, along with the originals. Visual comparison of the test maps and the original ones reveal that the overall structure did not change on the maps corresponding to lower spectral resolutions, except for a few smaller features that gradually faded as the resolution decreased. Spot temperatures however can significantly change in the case of the strongest active regions, especially on the maps corresponding to $R=20\,000$. This is expected, since the blurring makes the bumps caused by the spots on the line profiles shallower. Fig. A.2 shows the difference maps of the original and the $R=20\,000$ map. It is apparent that in the case of the strong polar feature, there is about 600 K difference in the recovered spot temperature, while smaller features did not change much. Finally, it is important to note that artifacts are not introduced by the lower resolution.

Based on these tests we conclude that degrading the MUSICOS spectra to a reduced spectral resolution of $R=20\,000$ is still sufficient to investigate the rapid spot evolution and differential rotation of EI Eri, despite the fact that there is some fading of the spectral features due to a decrease in contrast, which affects the measurement of the differential rotation at some level.




Fig. A.1. Doppler imaging test for two consecutive rotations of V1358 Ori. Images on the left correspond to the first rotation, while images on the right correspond to the second one. The four rows from top to bottom correspond to data of different spectral resolutions: $R = 80\,000$ (original data taken from Kriskovics et al. 2019), and reduced levels of $R = 60\,000$, $R = 40\,000$ and $R = 20\,000$, respectively.




Fig. A.2. Difference images of the original V1358 Ori Doppler images (Kriskovics et al. 2019) and those that correspond to a spectral resolution reduced to $R=20\,000$. The left panel corresponds to the first rotation and the right panel for the second, cf. Fig. A.1.




Fig. A.3. Example cross-correlation test maps of V1358 Ori with their fitted surface differential rotation functions obtained for different spectral resolutions of $R=20\,000$ (left) and $R=70\,000$ (right). The derived surface shear coefficients are $\alpha = 0.010 \pm 0.0025$ and $\alpha = 0.015 \pm 0.003$, respectively. For the original cross-correlation map see Kriskovics et al. (2019).




Fig. A.4. The α surface differential rotation parameter values obtained for spectroscopic test datasets with different levels of R spectral resolution.

The reliability of measuring differential rotation by cross-correlation of consecutive Doppler maps is based on the reliability of the spot reconstruction (for the details of the cross-correlation technique, see Sect. 4.4). Therefore, here we test how the reduced spectral resolution affects the surface shear parameter derived from cross-correlation. For the test the two consecutive rotations of V1358 Ori (see Fig. A.1) were cross-correlated with different levels of spectral resolution. Two example cross-correlation maps are shown in Fig. A.3 which correspond to $R=20\,000$ ($\alpha = 0.01 \pm 0.0025$) and $R=70\,000$ ($\alpha = 0.015 \pm 0.003$). We note that the original cross-correlation map presented in Kriskovics et al. (2019) with a spectral resolution of $R=80\,000$ resulted in $\alpha=0.016$. The derived α surface shear values are plotted against the spectral resolution in Fig. A.4. It is apparent that the resolution affects the derived shear coefficient in a unique way, that is, at low resolution the dependence has some linear trend, while at mid to high resolution ($R>40\,000$) there is no real trend, the derived α values agree with each other within the estimated error bars. Nevertheless, on our lowest resolution ($R=20\,000$), the measured surface differential rotation underestimates the actual value by $\approx 60\%$.

Appendix B: Observing log

Table B.1. Observing log of the MUSICOS 1998 data.

<i>HJD</i> [d]	Telescope	Rotational phase	Used	<i>HJD</i> [d]	Telescope	Rotational phase	Used
2451141.3758	OHP	0.157	yes	2451148.7618	ESO152	0.950	yes
2451141.4856	OHP	0.213	yes	2451148.9391	MSO	0.041	yes
2451141.6050	OHP	0.274	yes	2451149.0746	MSO	0.111	no
2451141.6302	KPNO	0.287	yes	2451149.1508	MSO	0.150	yes
2451141.6797	KPNO	0.313	yes	2451149.2564	BXO	0.204	no
2451141.7891	KPNO	0.369	no	2451149.5683	ESO90	0.364	yes
2451141.9259	KPNO	0.439	yes	2451149.7026	ESO152	0.433	yes
2451142.6193	KPNO	0.795	yes	2451149.7849	ESO152	0.475	yes
2451142.7057	KPNO	0.840	yes	2451150.1065	BXO	0.640	no
2451142.8025	KPNO	0.889	no	2451150.2155	BXO	0.696	yes
2451142.9096	KPNO	0.944	yes	2451150.2784	BXO	0.729	yes
2451142.9849	KPNO	0.983	no	2451150.5648	ESO90	0.876	yes
2451143.7458	ESO152	0.374	no	2451150.6669	ESO152	0.928	yes
2451143.8030	KPNO	0.403	yes	2451150.7474	ESO152	0.970	yes
2451143.8529	ESO152	0.429	no	2451150.9827	MSO	0.090	yes
2451143.8823	KPNO	0.444	no	2451151.0459	MSO	0.123	yes
2451143.8973	KPNO	0.452	yes	2451151.1061	MSO	0.154	no
2451143.9122	KPNO	0.459	yes	2451151.5633	ESO90	0.389	yes
2451144.3730	OHP	0.696	yes	2451151.7115	ESO152	0.465	yes
2451144.5975	ESO152	0.811	no	2451151.7969	ESO152	0.509	yes
2451144.7046	KPNO	0.866	yes	2451152.0938	BXO	0.661	yes
2451144.7232	KPNO	0.876	yes	2451152.1989	BXO	0.715	yes
2451144.7419	KPNO	0.885	yes	2451152.2799	BXO	0.757	yes
2451144.8528	ESO152	0.942	no	2451152.5011	INT	0.870	no
2451144.8673	KPNO	0.950	yes	2451152.5616	ESO90	0.901	yes
2451144.8859	KPNO	0.959	yes	2451153.5985	ESO90	0.434	no
2451144.9045	KPNO	0.969	yes	2451154.5383	INT	0.916	no
2451145.3744	OHP	0.210	yes	2451154.5934	ESO90	0.945	yes
2451145.6007	ESO152	0.326	yes	2451155.1511	BXO	0.231	no
2451145.8042	ESO152	0.431	yes	2451155.2465	BXO	0.280	no
2451146.3699	OHP	0.722	yes	2451155.3879	INT	0.353	yes
2451146.5807	ESO90	0.830	yes	2451155.5990	ESO90	0.461	yes
2451146.7466	ESO152	0.915	yes	2451155.6447	INT	0.485	no
2451146.8321	ESO152	0.959	yes	2451156.3677	INT	0.856	no
2451147.0404	MSO	0.066	yes	2451156.4359	INT	0.891	yes
2451147.0592	MSO	0.076	no	2451156.5991	ESO90	0.975	yes
2451147.1641	MSO	0.129	yes	2451156.6333	INT	0.992	no
2451147.4508	OHP	0.277	yes	2451157.3997	INT	0.386	no
2451147.5618	ESO90	0.334	yes	2451157.4703	INT	0.422	no
2451147.7280	ESO152	0.419	yes	2451157.5991	ESO90	0.488	no
2451147.8123	ESO152	0.462	yes	2451157.6475	INT	0.513	no
2451147.9404	MSO	0.528	no	2451158.5358	INT	0.969	no
2451148.0240	MSO	0.571	yes	2451158.5984	ESO90	0.001	no
2451148.1543	MSO	0.638	no	2451159.5998	ESO90	0.516	no
2451148.6718	ESO152	0.904	yes	2451160.6313	ESO90	0.045	no

Columns from left to right: the Heliocentric Julian Dates (*HJD*), the observing facilities, the rotational phases (according to Eq. 1), and whether the measurement was used or not.

Appendix C: Line profile fits




Fig. C.1. Fitted line profile subsets (S1-S4) corresponding to the four Doppler images (DI1-DI4) presented in Fig. 7. The phase distributions of each subset are shown in the right panel of Fig. 1.