


INVITED REVIEW

Epigenomic regulation of macrophage polarization: Where do the nuclear receptors belong?

Zsolt Czimmerer¹ | Laszlo Nagy² 

¹Institute of Genetics, Biological Research Centre, Eotvos Lorand Research Network, Szeged, Hungary

²Departments of Medicine and Biological Chemistry, Johns Hopkins University School of Medicine, and Institute for Fundamental Biomedical Research, Johns Hopkins All Children's Hospital, St. Petersburg, Florida, USA

Correspondence

Laszlo Nagy, Johns Hopkins University, School of Medicine, Johns Hopkins All Children's Hospital, 600 5 Street South, St. Petersburg, FL 33701, USA.
Email: lnagy@jhmi.edu

Funding information

National Institutes of Health, Grant/Award Number: R01-DK115924 and R01-DK124782

Summary

Our laboratory has a long-standing research interest in understanding how lipid-activated transcription factors, nuclear hormone receptors, contribute to dendritic cell and macrophage gene expression regulation, subtype specification, and responses to a changing extra and intracellular milieu. This journey in the last more than two decades took us from identifying target genes for various RXR heterodimers to systematically mapping nuclear receptor-mediated pathways in dendritic cells to identifying hierarchies of transcription factors in alternative polarization in macrophages to broaden the role of nuclear receptors beyond strictly ligand-regulated gene expression. We detail here the milestones of the road traveled and draw conclusions regarding the unexpectedly broad role of nuclear hormone receptors as epigenomic components of dendritic cell and macrophage gene regulation as we are getting ready for the next challenges.

KEYWORDS

EGR2, epigenome, IL-4, macrophage, nuclear receptors, polarization, PPAR γ , STAT6, transcriptional regulation

1 | INTRODUCTION

Macrophages and dendritic cells are essential cellular components of the innate immune system. Despite sharing common functional characteristics, including their phagocytic, antigen-presenting, and cytokine-producing capacities or their ability to respond rapidly to the changing microenvironment, they still have a well-defined and cell-type-specific role in the body under various physiological and pathological conditions. Macrophages are critical contributors in maintaining normal tissue homeostasis, the defense against various pathogens, from bacteria to multicellular parasites, and important regulators of inflammation. Dendritic cells are critical in initiating and regulating pathogen-specific adaptive immune responses and contribute to the development of immunologic memory and tolerance.¹⁻³

Macrophages are found in almost all tissues, and although they have common functions, including phagocytotic and antimicrobial capacities, their homeostatic function strongly depends on the tissue microenvironment. Among others, the different macrophage subtypes participate in development, metabolism, wound healing, tissue remodeling, and angiogenesis.^{1,4,5} Besides, their maladaptive or altered functional properties also contribute to the development and progression of different chronic inflammatory disorders and cancers.⁵⁻⁷ This remarkable phenotypic heterogeneity and plasticity of macrophages are tightly determined by their origin and molecular microenvironment. Based on the origin of tissue-resident macrophages, yolk sack macrophages, fetal liver monocytes, and bone marrow monocytes-derived macrophages can be distinguished. In adult tissues in the steady-state, microglia in the brain, alveolar macrophages in the lung, Hofbauer cell in the placenta, and Kupffer cells in the liver are of embryonic origin,

This article is part of a series of reviews covering Lipids as Modulators of Immunity appearing in Volume 317 of *Immunological Reviews*.

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2023 The Authors. *Immunological Reviews* published by John Wiley & Sons Ltd.

while both embryonic and bone marrow monocytes-derived macrophages are present alongside one another in the intestine.⁸⁻¹⁰ The bone marrow monocyte-derived macrophages also play a critical role in the different organs of the body after tissue injury or infections.^{11,12} The macrophage microenvironment consists of a wide range of metabolites, nutrients, immunomodulatory factors, and pathogen-derived molecules, determining various phenotypic and functional features of the distinct macrophage subsets under different circumstances. The two endpoints of functional macrophage polarization induced by the microenvironmental activating signals are the Th1-type cytokine interferon-gamma (IFN γ) or gram-negative bacterial cell wall component lipopolysaccharide (LPS)-dependent classical and the Th2-type cytokines interleukin-4 (IL-4) and IL-13-dependent alternative macrophage polarization. These polarization states have well-distinguishable gene expression profiles and functional characteristics. The classically polarized macrophages have anti-bacterial capacities, while the alternatively polarized macrophages contribute to the protection against nematode infections and tissue regeneration.^{6,7,13,14} Nevertheless, the complex in vivo molecular milieu, often simultaneously containing different pathogen-derived molecules, cytokines, and metabolites, can result in various transient macrophage polarization forms associated with unique functional characteristics under normal and pathological conditions (Figure 1).¹⁵⁻¹⁷ The superfamily of nuclear hormone receptors, in particular, the retinoid X receptor (RXR) heterodimers such as peroxisome proliferator activated receptor gamma (PPAR γ), retinoic acid receptor (RAR), vitamin D receptor (VDR), and liver X receptor (LXR), have been implicated at multiple levels of gene regulation in macrophages and dendritic cells, including lineage determination to epigenomic bookmarking to conventional ligand activation. Here, we will summarize our contributions to this field.

Dendritic cells were discovered by Steinman and Cohn 50 years ago and their findings opened up an entire new field linking the until disparate innate and adaptive immunity fields in a highly mechanistic manner.¹⁸ This generated a flurry of investigations using multiple models. The human monocyte-derived cells generated by GM-CSF and IL-4 from CD14+ monocytes became the standard approach

for studying human DCs and for harnessing their therapeutic potential in the early 2000s.^{19,20} It has been suggested that these cells (MCs=monocyte-derived cells) represent the in vitro counterparts of CD14+ CD11c+ inflammatory cells generated by immune response.²¹ Human conventional dendritic cells however are generated in vitro from CD34+ cord blood HSPCs containing CD115 expressing progenitors and their differentiation is driven by Flt3L.²² Therefore, MCs should not be considered conventional DCs. Our work in this domain has focused on MCs, which could be considered immature monocyte-derived dendritic cells. We have carried out systematic work on profiling the transcriptional landscapes and changes in MCs upon the activation of the RAR, VDR, PPAR γ , and LXR. This led to the identification of new targets and established a role for these receptors in subtype specification including tissue specificity and specialized immune function (i.e. RAR and PPAR γ regulated CD1D expression and iNKT cell generation) summarized in a review.²³ By now, most of the human and mouse DC subtypes have been identified and their origins clarified using careful profiling and lineage tracing,¹⁸ presenting new opportunities to revisit the role of some of these nuclear hormone receptors for their roles in differentiation as well as in functional specification.

2 | THE TRANSCRIPTIONAL BASES OF MACROPHAGE AND DENDRITIC CELL RESPONSE TO THE IMMUNOMODULATORY AND ACTIVATING SIGNALS

The phenotypic and functional features of macrophages and dendritic cells are tightly controlled at the transcriptional level by the microenvironmental milieu. Several mechanistic studies demonstrated that the macrophage or dendritic cell subtype-specific and the extrinsic or intrinsic activation signals-promoted transcriptional programs are based on the complex collaborative interactions between the cis-regulatory elements, including promoters and enhancers, and trans-acting DNA-binding transcription factors (TFs).^{24,25} In general, cis-regulatory elements contain multiple, distinct transcription




FIGURE 1 General scheme of signal response and integration in macrophages.

factor-binding DNA motifs and have characteristic, partially overlapping histone modification patterns. Promoters are marked by a high level of H3K4 trimethylation, while enhancers are associated with H3K4 mono- and dimethylation instead. Additionally, both cis-regulatory elements exhibit high levels of H3K27 acetylation in the activated state and H3K27 trimethylation in the repressed state.²⁶⁻²⁸ The TFs-orchestrated transcriptional regulation shows a multi-layered organization with at least three distinct levels, including (i) the determination of the available enhancer platform by lineage-determining transcription factors (LDTFs), (ii) the development of a primary transcriptional response to microenvironmental signals and tissue demand by signal-dependent transcription factors (SDTFs), and (iii) the formation of a secondary or long-term transcriptional response to microenvironmental changes by SDTFs-regulated TFs or TF networks.²⁹⁻³¹ The core enhancer repertoire in the macrophages and dendritic cells is tightly determined by the cooperative binding of the general cell type-specific LDTFs, including ETS domain, transcription factor PU.1, activator protein 1 (AP1), and CCAAT-enhancer-binding proteins (CEBPs) in macrophages or PU.1 and CEBP β in dendritic cells.^{30,32-34} It has also been described that additional, specific transcription factors, such as GATA binding protein 6 (GATA6) in peritoneal macrophages or Spalt-like transcription factor 1 (SALL1) in microglial cells, can also act as LDTFs, determining the tissue-resident macrophage subtype-specific enhancer sets.^{35,36} The available enhancer repertoires serve as a binding platform for the environmental signals-activated SDTFs. Based on the invoked immunological consequences, the SDTFs and their activating signals can be divided into a minimum of two distinct groups. On the one hand, the homeostatic tissue and immunomodulatory signals-activated SDTFs, including immunomodulatory cytokines-activated signal transducer and activator of transcription (STAT) family members or the different hormones and lipids-activated nuclear hormone receptors, contribute to the development of a tissue-specific macrophage and dendritic cell phenotype and modulate their response to danger signals. On the other hand, the pathogen-derived or endogenous danger signals and inflammatory cytokines-activated SDTFs, such as nuclear factor kappa-light-chain-enhancer of activated B cells (NF- κ B) and AP-1 TF complexes, orchestrate the macrophage and dendritic cell activation.^{25,29} The microenvironmental signals-activated long-term transcriptional programs are provided by a wide range of SDTFs-induced transcription factors, among others, interferon regulatory factor (IRF), basic helix-loop-helix (BHLH), and early growth response factor (EGR) family members.^{31,37-40}

3 | NUCLEAR HORMONE RECEPTORS HETERODIMERIZING WITH THE RETINOID X RECEPTOR AS MULTIFACETED HORMONE AND LIPID-SENSING TRANSCRIPTION FACTORS

The nuclear hormone receptor superfamily includes 48 members in humans and 49 in mice, respectively. Based on their ligands and

mode of action, the nuclear receptors can be divided into steroid and nonsteroid receptors.⁴¹ The latter includes the RXR heterodimerizing nuclear hormone receptors. This subgroup of the nuclear hormone receptors is quite heterogenous, containing both classical endocrine and adopted orphan receptors. The classical endocrine receptors are activated by high-affinity ligands such as thyroid hormone for thyroid hormone receptor (TR), all-trans-retinoic acid for RARs, and $1\alpha,25$ -dihydroxy vitamin D₃ for VDR. In contrast, the adopted orphan receptors usually sense various lipid metabolites, including oxysterols for LXRs, fatty acids for PPARs, and bile acids for farnesoid X receptor (FXR), with low affinity.^{23,41,42} In the pre-epigenomic era, several general principles for a mode of action of RXR heterodimers were described using combined molecular biological, pharmacological, and genetic approaches. First, RXR heterodimers are bound to the DNA containing their directly repeated response elements with half-site sequence AGGTCA or a variant of it in the nucleus independently from the presence of ligands. Second, RXR heterodimers interact with corepressor complexes and act as transcriptional repressors without an appropriate ligand. Third, the ligand binding induces corepressor-coactivator complex exchange leading to transcription activation.²³ Besides, the phenomenon termed transrepression was also identified. In this case, the ligand-bound nuclear receptor heterodimer modulates transcription of a target gene without direct DNA binding interfering with the activity of other transcription factors through protein-protein interactions and post-translational modifications.⁴³ However, in the post-epigenomic era, the study of the transcription factor binding and function at the whole-genome level by many next-generation sequencing (NGS)-based methods allowed expanding and clarifying our knowledge about the activities of RXR heterodimers. Among others, these approaches significantly contributed to identifying new functional properties for the RXRs and their heterodimerizing partners, such as ligand-independent bookmarking or cell subtype-specific LDTF activities.⁴⁴⁻⁴⁶ In this review, we aimed to focus on the complex role of a retinoid X receptor heterodimerizing nuclear hormone receptors in macrophage and dendritic cell biology, especially in the context of our work in this field over the past two decades.

4 | INTEGRATED NGS-BASED EPIGENOMIC AND TRANSCRIPTOMIC APPROACHES TO BETTER UNDERSTAND THE REGULATORY FUNCTION OF RXR HETERODIMERS IN MACROPHAGES

Several studies demonstrated by the early 2010s that the lipid-sensing nuclear receptors can form a bridge between metabolism and innate immune system by directly activating the genome and regulating macrophage functions from oxLDL uptake to inflammatory response^{47,48} (Figure 2). However, the technologies available until then were biased and limited to discover ligand-regulated events and did not allow studying the regulatory mechanism of RXR heterodimers at the whole genome level and in an unbiased manner.




FIGURE 2 The classical ligand-dependent regulatory role of RXR/PPAR γ , RXR/LXR, RXR/RAR heterodimers, and RXR/RXR homodimers in macrophage biology and the proposed RXR-mediated enhancer promoter interaction of the *Vegfa* gene.

The advancement of NGS-based transcriptomic and epigenomic technologies has fundamentally changed the opportunities for studying transcriptional regulation enabling a detailed understanding of the already-known nuclear receptor-mediated regulatory mechanisms and identification of novel functions genome-wide. To take advantage of this technological breakthrough and better understand the regulatory role of RXR heterodimers in the macrophages, first we studied RXR signaling in murine bone marrow-derived macrophages combining chromatin immunoprecipitation sequencing (ChIP-seq), global run-on sequencing (GRO-seq), and RNA sequencing (RNA-seq) methods. By combining these methods, we developed a pipeline by which we called (i) directly regulated genes by utilizing a time course of nascent RNA production (GRO-Seq), (ii) active enhancers by the changing enhancer RNA (GRO-Seq) and active histone marks, (iii) RXR binding sites by ChIP-seq, and (iv) co-activator binding by p300 recruitment in the proximity of the regulated genes.⁴⁹ This rigorously applied comprehensive genome-wide approach provided a large amount of information about liganded RXR-mediated transcriptional changes, active binding sites, and cistromic interactions in the context of the 3D genome structure once validated using 3C-Q-RT-PCR. Our results confirmed many elements of our prior knowledge about RXRs and their heterodimeric partners, including nuclear localization and DNA binding capacity of RXRs in the unliganded state or their binding to receptor-specific hormone response elements as NR half sites, DR1, and DR4 motifs. It also established that the RXR cistrome in macrophages is not regulated by exogenously added synthetic ligands to any significant degree. To provide genome-wide evidence for the biological relevance of ligand-dependent transcriptional activation, we identified 387 liganded RXR-activated enhancers characterized by RXR and PU.1 binding,

as well as ligand-induced P300 recruitment and enhancer RNA expression. These RXR ligand-activated genomic elements are linked to 226 genes, such as previously identified RXR target genes *Abca1*, *Angptl4*, or *Tgm2*, validating the approach and allowed identifications of novel ones such as *Vegfa*⁴⁹ (Figure 2). *Vegfa* is a particularly notable example because the identified complex enhancer is almost 300 kilobase downstream of the transcription start site and thus we were able to call it only using the rigorous criteria detailed above. However, our approach could identify more than 5000 RXR-bound genomic regions without any evidence for ligand-mediated effects, nearly half of which are transcriptionally inactive.⁴⁹ Intriguingly, the ligand-insensitive fraction of LXR α and β cistromes could also be identified in non-polarized murine immortalized BMDMs indicating that this phenomenon is not restricted to the RXR cistrome.⁵⁰ These findings suggested that RXRs and their heterodimeric partners may also have an important ligand-insensitive or non-conventional transcriptional regulatory role in the macrophages. We could confirm the existence of this mechanism in two independent studies in recent years. On the one hand, we identified the elevated expression of several prometastatic genes in lung-derived RXR-deficient myeloid cells associated with their enhanced cancer cell migration and invasion-promoting activities in vitro and increased lung metastasis formation in vivo. The affected prometastatic genes were mainly insensitive to synthetic ligand activation. Still, they are associated with RXR, nuclear receptor corepressor (NCoR), and silencing mediator of retinoic acid and thyroid hormone receptor (SMRT) co-bound enhancers suggesting that RXRs regulate their expression through direct ligand-independent transcriptional repression.⁵¹ On the other hand, by studying the potential interactions between the alternative polarizing signal IL-4-activated STAT6 TF and the RXR signaling pathways

in human monocyte-derived differentiating macrophages, we could identify an atypical so-called potentiating effect of ligand-activated RXRs on the transcriptional regulation of IL-4 inducible genes. The IL-4-activated STAT6 cistrome showed extensive overlap with RXR-bound genomic sites following short-term (30min) IL-4 polarization without influencing the RXR occupancies at the overlapping regulatory regions. The synthetic RXR agonist (LG268) had distinct activities at the selected IL-4-activated and STAT6/RXR co-bound enhancers modulating the IL-4 responsiveness of their target genes regulated by these genomic elements. Based on the role of the RXR ligand, we could divide into three groups the selected IL-4-activated enhancers and genes, including RXR activation independent, synergistically activated, and liganded RXR potentiated categories. In the case of the latter group, the genes and enhancers were insensitive to RXR ligand activation in the non-polarized macrophages. Still, the applied synthetic RXR agonist further increased their IL-4-dependent induction.⁵² These findings raised the possibility that the liganded RXR plays a much more complex role in transcriptional regulation than we previously thought, especially in the case of a complex microenvironmental milieu where endogenous RXR ligands may be present. These new mechanisms represent unique opportunities for modulating inflammatory gene regulation.

5 | IL-4-ORCHESTRATED STAT6, EGR2, AND PPAR γ CONTAINING TF NETWORK IN THE DETERMINATION OF THE ALTERNATIVE MACROPHAGE POLARIZATION SPECIFIC EPIGENETIC AND TRANSCRIPTIONAL PROGRAM

After spending years to characterize primarily nuclear receptor-mediated events in human monocyte-derived immature dendritic-like cells and mouse bone marrow-derived macrophages, we came to the realization that the origin and state of the observed cell has an oversized and non-negligible influence on nuclear receptor activity both ligand dependent and also ligand independent and thus, we decided to focus on and exhaustively characterize a particular polarization pathway, alternative polarization, and examine and position RXR heterodimer and in particular PPAR γ /RXR signaling in its context and hierarchy.

Th2-type immunomodulatory cytokine IL-4 can turn on a specific transcriptional program through the direct activation of its SDF STAT6 in human and mouse macrophages resulting in their alternative polarization. This macrophage phenotype is characterized by a partially overlapping gene expression signature between mice and humans. Despite the differences in the IL-4-STAT6 signaling pathway-dependent gene expression programs, human and mouse alternatively polarized macrophages participate in the protection against nematode infections and have high tissue regenerative capacities.^{15,53,54} The alternatively polarized macrophages are also observed in Th2-type allergic airway inflammation and asthma, breast cancer, fibrotic processes, and the chronic phase of different

bacterial infections.⁵⁵ Although many features of direct STAT6-mediated transcription activation, including its target genes and its binding DNA motif, were already known in the pre-epigenomic era, the complex NGS-based methodology we developed for mapping nuclear receptor signaling in our prior works contributed to a better understanding of the IL-4-STAT6 signaling pathway-activated transcriptional program in macrophages. Applying comprehensive and integrated ChIP-seq, GRO-seq, and assay for transposase accessible chromatin sequencing (ATAC-seq) methods, we and others have made several important contributions in the context of direct STAT6-regulated epigenetic and transcriptional events. First, STAT6 binds to more than 20,000 genomic regions with specific dynamics. Second, it rapidly activates thousands of constitutively H3K4 mono and demethylated and PU.1-bound enhancers. Third, it can also bind to enhancers lacking LDTF binding and enhancer-specific histone marks in nonpolarized macrophages, leading to the latent or de novo enhancer formation and activation. Fourth, it can also act as a direct transcriptional repressor associating with reduced RNA Polymerase II binding and H3K27 acetylation as well as attenuated eRNA expression. These STAT6-repressed enhancers are also characterized by diminished PU.1 and co-activator P300 binding and decreased chromatin accessibility following short-term IL-4 exposure.^{56,57} Intriguingly, it has recently been demonstrated that many direct IL-4-STAT6 target genes show distinct IL-4 inducibility in different cell cycle phases of murine bone marrow-derived macrophages, raising the possibility that the STAT6 binding or its transcriptional activator capacity is also regulated in a cell cycle-dependent manner at different enhancer sets.⁵⁸

In addition to the direct transcriptional regulator activity of STAT6, it has also been demonstrated in the last two decades that additional transcription factors downstream from STAT6 are also required for proper alternative macrophage polarization, including IRF4, cMYC, KLF4, BHLH40, and PPAR γ .^{59–66} However, the full complement of this TF network and the relationship including the hierarchy between the individual TFs were not fully understood. To identify the key regulators during the transition between the early direct STAT6-dependent and the late stable alternative macrophage polarization-specific transcriptional and epigenetic events, we further studied the early and late enhancer activation following IL-4 polarization in murine bone marrow-derived macrophages. Our P300 and H3K27 acetylation-specific ChIP-seq-based approach could identify three IL-4-activated enhancer clusters with distinct dynamics, including “early transient,” “early sustained,” and “late” enhancers. As expected, STAT6 was responsible for the early activation at the “early transient” and “early sustained” enhancer sets. Besides, we could identify the EGR2 transcription factor as a new regulator of the late alternative macrophage polarization, directly and indirectly controlling the activation of “early sustained” and “late” enhancers following 24 hours length IL-4 exposure. The direct EGR2-dependent enhancer activation is associated with elevated chromatin openness and chromatin remodeling factor BRM binding, as well as increased co-activator P300 and BRD4 bindings. The indirect EGR2-mediated effects on the enhancer activity are probably caused by the activity

of an alternative macrophage polarization-specific TF network containing the previously described KLF4, BHLH40, or PPAR γ . Overall, the consequence of the direct and indirect regulatory roles of EGR2 is the altered expression of 77% of IL-4-induced and 64% of IL-4-repressed genes in the EGR2-deficient macrophages indicating that EGR2 is essential for the late stable epigenetic program of alternative macrophage polarization³⁷ (Figure 3). This was confirmed in a recent publication describing the role of natural genetic variations in the alternative macrophage polarization program also confirmed the contribution of EGR2 to the IL-4-induced gene expression and enhancer activation in murine bone marrow-derived macrophages.⁶⁷ A potential EGR2 autoregulatory loop can provide stability to the epigenomic state and gene expression. This is a particularly compelling example of how a transient epigenomic signal (STAT6 binding to chromatin) can be converted to a stable epigenome (EGR2 and downstream TF cistromes) providing transcriptional memory.

As noted above, the lipid-sensing nuclear hormone receptor PPAR γ is one of the best-known alternative macrophage polarization-specific TFs. Its induction by IL-4 in macrophages was described very early in 1999, and since then, it has been confirmed several times in different macrophage subtypes and species by us and others.^{60,68-70} However, the role of PPAR γ during alternative macrophage polarization was quite controversial for a long time. It has been previously described that PPAR γ is required for the proper IL-4-mediated alternative polarization, and the myeloid cell-specific disruption of PPAR γ sensitizes these mice to the development of diet-induced obesity, insulin resistance, and glucose intolerance.⁶⁰ Despite these facts, the activation of PPAR γ by the synthetic agonist rosiglitazone (RSG) could

not be linked to the induction of alternative polarization-specific gene signature,⁷⁰ raising the possibility of a minimum of two plausible explanations, including (i) the role of endogenous PPAR γ ligands or (ii) ligand-independent action of PPAR γ in this process. According to the first hypothesis, IL-4 can induce many enzymes with endogenous PPAR γ ligand-producing potential in murine and human macrophages, including 13-HETE and 15-HODE-producing ALOX15, lysophosphatidic acid-producing ENPP2, and serotonin metabolites-producing MAOA.^{68,69,71,72} The second hypothesis was also supported by the fact that PPAR γ heterodimeric partner RXR also has a significant genome-bound, ligand-insensitive, and transcriptionally inactive fraction in nonpolarized murine macrophages.⁴⁹ To systematically evaluate the synthetic and/or endogenous ligand sensitivity of the PPAR γ /RXR heterodimers at the whole genome level in long- and short-term polarization experiments using murine bone marrow-derived macrophages, we performed CHIP-seq and GRO-seq analyses in these systems in the absence or presence of synthetic PPAR γ agonist and antagonist. Using this approach, we could draw conclusions about the PPAR γ and RXR cistromes and their ligand sensitivity. On the one hand, IL-4 exposure can cause a large expansion of PPAR γ cistrome (binding) at the whole genome level associated with the redistribution of RXR cistrome in both polarization systems with minimal loss of binding. On the other hand, the formation of polarization-induced PPAR γ /RXR heterodimers is endogenous ligand-independent. Thirdly, the limited number of PPAR γ /RXR-bound enhancers annotated to the classical PPAR γ target genes, such as *Angptl4* or *Fabp4*, can be influenced by synthetic PPAR γ and RXR agonists, PPAR γ antagonist. Overall, these findings indicate that




FIGURE 3 The STAT6-EGR2 transcription factor axis-induced PPAR γ expression during IL-4-mediated alternative macrophage polarization in murine BMDMs.

alternative macrophage polarization is associated with the PPAR γ /RXR heterodimer cistromic expansion at the whole genome level. Still, PPAR γ activation by endogenous and synthetic ligands is negligible in volume and probably does not contribute to alternative macrophage polarization per se.^{44,73} The further study of ligand insensitive PPAR γ /RXR heterodimers-bound enhancers in wild-type and PPAR γ deficient alternatively polarized murine macrophages by ATAC-seq, as well as P300 and RAD21-specific ChIP-seq demonstrated these distal regulatory regions are associated with IL-4-induced P300 and RAD21 binding as well as chromatin accessibility in a PPAR γ dependent manner. These observations indicate that most PPAR γ /RXR heterodimers in the alternatively polarized macrophages act as epigenetic bookmarking factors contributing to the entrenchment of the alternative phenotype by keeping chromatin open and/or allow additional factors to act (Figure 4.) rather than the simple sensors of the extracellular or intracellular lipid microenvironment.⁴⁴

6 | IL-4-INDUCED TF NETWORK SUPPORTS THE UNIQUE RESPONSIVENESS OF ALTERNATIVELY POLARIZED MACROPHAGES TO THEIR SURROUNDING MOLECULAR MILIEU

Once we mapped the transcriptional cascade supporting alternative polarization and positioned PPAR γ /RXR signaling in it we extended our investigations to signal interactions with the alternative polarization and nuclear receptor pathways.

The immunomodulatory signals and exogenous pathogen-derived or endogenous danger signals may be present in the environment of macrophages simultaneously or consecutively. In general, the complex interactions between individual signals can be antagonistic, synergistic, or result in the de novo expression of a specific gene set leading to the development of more specialized macrophage phenotypes and functions aggravating or alleviating the disease outcome.

Transcriptional memory-like features associated with the first-stimulus epigenetic remodeling and metabolic changes are well-known in the macrophages following repeated exposure to the same or different stimuli. Transcriptional memory can result in attenuated (such as LPS tolerance) or increased (such as trained innate immunity) responsiveness to the second stimulus.^{74,75} To investigate whether the suspected bookmarking role of PPAR γ /RXR heterodimers can influence transcription memory in alternatively polarized macrophages, we applied IL-4 re-stimulation-based experimental system. Surprisingly, we identified an extracellular matrix-associated gene set that is de novo induced by the second IL-4 stimuli. These memory-like features were completely PPAR γ -dependent showing a novel epigenomic role of the ligand insensitive PPAR γ /RXR heterodimers⁴⁴ (Figure 4).

It has been demonstrated previously that Th1-type cytokine IFN γ can reinforce the Toll-like receptor (TLR) ligands-induced

inflammatory response in macrophages at multiple levels. IFN γ and TLR ligand co-stimulation results in the so-called super-induction of many canonical inflammatory genes for which STAT1 and IRF1 transcription factors are responsible.⁷⁶ Additionally, IFN γ -directed repression of feedback inhibitory and metabolic components of TLR responses are regulated by STAT3, further enhancing the inflammatory response in macrophages.⁷⁷ Finally, IFN γ can also impede and reverse TLR ligands-induced macrophage tolerance, leading to exacerbated inflammatory phenotype in autoimmune diseases.^{78,79}

The alternative macrophage polarization signal IL-4 may be present simultaneously with the opposing signal IFN γ and different pathogen-derived or endogenous danger signals at the same time in the macrophage microenvironment in various pathological conditions, including helminth-bacterial or helminth-viral coinfections, cancers, or exacerbated Th2-type airway inflammation and asthma.^{80–86} The mutual antagonistic effects have long been a known feature of IL-4 and Th1-type cytokines or TLR ligands, and recent studies explored the molecular and epigenetic bases of these interactions. It has been described that IFN γ could repress the basal homeostatic expression of many alternative macrophage polarization-associated genes through the inactivation and disassembly of MAF transcription factor binding enhancers.⁸⁷ Nevertheless, the co-stimulation of murine bone marrow-derived macrophages by IL-4 and IFN γ showed that IFN γ -activated enhancers associated with STAT1 and IRF binding are highly resistant to IL-4-dependent inhibition, but the AP-1 and C/EBP β -bound regulatory regions are more sensitive the inhibitory effects of IL-4.⁶⁴ To investigate whether the attenuated IFN γ responsiveness in alternatively polarized macrophages shows a progressive nature, we restimulated the macrophages two, three, and four times with IL-4 followed by IFN γ exposure. We applied washout after each stimulation and rested the cells for 24 hours before the following stimulation. We reported that the attenuated response of the selected genes, including *Ccl5*, *Irg1*, and *Irf8*, to IFN γ in the alternatively polarized macrophages proved quite progressive, resulting in almost complete desensitization after the fourth IL-4 restimulation and thus IFN γ resistance.⁴⁴ It has been recently published that macrophage responsiveness to IFN γ markedly reduced in S and G2/M cell cycle phases, while IL-4 induces the cell cycle phase distribution toward the G2/M phase resulting in the subpopulation-specific reduced IFN γ responsiveness in the IL-4 polarized macrophage populations.⁵⁸ Besides, we observed that the direct transcriptional repressor activity of the STAT6 transcription factor also affects the specific subset of the inflammatory enhancers attenuating both their basal activity and TLR their inducibility by TLR ligand LPS. This partially reduced inflammatory responsiveness results in blunted NLRP3 inflammasome activation, IL-1 β production, and pyroptosis in IL-4-primed and LPS-activate murine bone marrow-derived macrophages⁵⁷ (Figure 5). Intriguingly, the IL-4-enhanced inflammatory responsiveness of certain genes was also detected in macrophages suggesting that the interactions between IL-4-STAT6 and TLR signaling pathways are not restricted to antagonism.^{57,88,89} Based on these observations, we could identify a specific gene set containing

FIGURE 4 The ligand-independent epigenomic ratchet activity of PPAR γ leads to transcriptional memory and enhanced gene-specific responsiveness to IL-4 re-stimulation in alternatively polarized macrophages.





FIGURE 5 The direct transcriptional repressor activity of STAT6 at a specific subset of inflammatory enhancers limits the inflammatory responsiveness in the alternatively polarized macrophages resulting in blunted inflammasome activation and pyroptosis.

more than 1300 genes showing elevated LPS responsiveness, which we termed extended synergy, in IL-4-primed murine bone marrow-derived macrophages compared to the nonpolarized counterparts. Our systematic analyses resulted in the following important findings about extended synergism: (i) "de novo" and "enhanced" NF- κ B-p65 binding and synergistic activation are detected at the distal regulatory regions annotated to the synergistically activated genes in IL-4-primed, and LPS-activated macrophages, (ii) both elevated NF- κ B-p65 binding and synergistically activated gene expression

is completely STAT6 dependent, (iii) the synergistic enhancer activation is associated with increased chromatin openness and BRD4 binding, (iv) EGR2 is an important but not exclusive regulatory factor of this process, (v) the genetic variance can influence the extended synergism, (vi) the extended synergism is observable in different murine tissue-resident and human monocyte-derived macrophages, and (vii) this phenomenon is present in alveolar macs during allergic airway inflammation leading to enhanced LPS-induced inflammation⁹⁰ (Figure 6).




FIGURE 6 The IL-4-induced EGR2 and PPAR γ transcription factors contribute to the gene subset-specific enhanced responsiveness to various environmental signals, including pathogen-derived molecules and nuclear receptor ligands, in the alternatively polarized macrophages.




FIGURE 7 The potential connection between the phenomenon of extended synergism between IL4 and TLR signaling and the RXR signaling pathway in the alternatively polarized macrophages. (A) Partial overlap is observed between the LPS-activated de novo/enhanced NF- κ B-p65 binding and the RXR cistrome at the synergistically activated enhancers in the alternatively polarized macrophages. Venn diagrams showing the overlap of RXR peaks with de novo and enhanced NF κ B-p65 genomic regions, (B) Read distribution plot visualization of NF- κ B-p65 (purple), RNAPII-pS2 (red), and RXR (blue) binding at the de novo/enhanced NF- κ B-p65 and RXR co-binding-associated synergistically activated enhancers (+/-2.5kb from peak summits).




FIGURE 8 Both ligand-dependent and ligand-insensitive epigenomic regulatory roles of nuclear hormone receptors modulate the macrophage responsiveness to various microenvironmental signals at the transcriptional level.

RXRs and their heterodimerization partners can play diverse functions in macrophages, directly or indirectly modulating the inflammatory response. Including but not limited to, LXR agonists can inhibit the inflammatory gene expression through direct repression or transrepression,^{43,91} the liganded PPAR γ also has an anti-inflammatory effect,⁴³ while RXR ligand activation leads to the elevated expression of Ccl9 and Ccl6 in macrophages and RXR α deficiency in myeloid cells leads to the lower susceptibility to sepsis in murine CLP model.⁹² In addition, many nuclear hormone receptors play in subset-specific LDTF functions in different tissue-resident macrophages, including PPAR γ in alveolar macrophages,⁹³ LXRs in splenic marginal zone macrophages, and Kupffer cells,^{45,94} and RXRs in large peritoneal macrophages,⁴⁶ potentially also influencing the inflammatory responsiveness of the given macrophage subtypes. However, it is completely unknown whether RXR heterodimerizing nuclear hormone receptors can influence the complex interactions between different immunomodulatory signals and inflammatory signals detailed above. To evaluate the possibility of the modulatory action of RXR heterodimers on extended synergism, we determined the overlap between the RXR cistrome-derived from IL-4-polarized murine bone marrow-derived macrophages and the synergistically activated “de

novo” or “enhanced” NF κ B-p65 binding associated enhancers. As shown in Figure 7 A and B, we could identify nearly 700 synergistically activated enhancers having RXR binding in the IL-4 polarized macrophages. These findings raised the possibility that RXRs or their heterodimerization partners can influence the synergistic enhancer activation at these sites. Nevertheless, our hypotheses about the role of RXR heterodimers in the complex immunomodulatory cytokine-TLR ligand interactions need further experimental confirmation.

7 | CONCLUSION

Macrophages and dendritic cells are present in almost all tissues in various physiological and pathological conditions associated with the complex microenvironmental milieu. The SDFs play an important role in creating connections between the continuously changing microenvironment and the cellular response through the transcriptional regulation of several 100–1000s of genes. The activation of the immunomodulatory and danger signal-responsive SDFs in different combinations leads to a broad spectrum of functional states of macrophage and dendritic cell activation,

often determining the disease onset, progression and outcome. For all these reasons, it is essential to learn about the transcription programs activated by different SDFs and their combinations. Initially, the lipid-sensing nuclear receptors were recognized in macrophages and dendritic cells as simply forming a bridge between metabolism and inflammatory processes through the ligand-dependent transcriptional activator and repressor activities. However, various *in vivo* and global NGS-based approaches made it possible to significantly broaden our understanding by identifying novel non-conventional nuclear hormone receptor-mediated epigenomic regulatory functions (Figure 8). In the last decade, it was discovered that nuclear hormone receptors could act as LDTF in various tissue-resident macrophage subsets. We also uncovered their ligand-dependent gene-specific potentiating effect to another signal, their ligand-insensitive repressor, or bookmarking activities. These new developments shed a different light on this group of transcription factors, suggesting that they are much more than mere mediators of lipid signaling. They apparently have epigenomic roles serving processes as signal amplification, inhibition and transcriptional memory. It also questions how these distinct functions developed during evolution. Which was first, the liganded activity or the epigenomic one? Thus, there are still many open questions waiting to be answered. Additional work is needed in the near future in the nuclear hormone receptor field, including investigating the molecular background and ligand sensitivity of their LDTF activities and the extent of their described non-conventional activities, determining their regulatory role in complex molecular microenvironment-activated transcriptional programs relative to their described conventional ligand-dependent and non-conventional activities. Deciphering these enigmas preferably at the single cell level, can contribute to developing better targeted therapies for diseases associated with altered macrophage and dendritic cell functions.

ACKNOWLEDGMENTS

The authors are indebted to Dr. Laszlo Halasz for the figures and data analyses and to members of the Nagy laboratory for discussions and comments on aspects of the manuscript. Z.C. is supported by Hungarian National Research, Development, and Innovation Office (FK132185), and by the New National Excellence Program of the Ministry for Innovation and Technology (ÚNKP-22-5 - SZTE-549), and by the János Bolyai Research Scholarship of the Hungarian Academy of Sciences BO/00594/22/8. L.N. is supported by the National Institutes of Health (R01-DK115924, R01-DK124782).

CONFLICT OF INTEREST STATEMENT

The authors declare no conflict of interest.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are openly available in the NCBI GEO at <http://doi.org/10.1093/nar/gky157>, reference number GSE110465 and at <http://doi.org/10.1016/j.immuni.2022.10.004>, reference number GSE181223.

ORCID

Laszlo Nagy  <https://orcid.org/0000-0001-6653-2155>

REFERENCES

- Mosser DM, Hamidzadeh K, Goncalves R. Macrophages and the maintenance of homeostasis. *Cell Mol Immunol*. 2021;18(3):579-587. doi:10.1038/s41423-020-00541-3
- O'Neill LAJ, Pearce EJ. Immunometabolism governs dendritic cell and macrophage function. *J Exp Med*. 2016;213(1):15-23. doi:10.1084/jem.20151570
- Geissmann F, Manz MG, Jung S, Sieweke MH, Merad M, Ley K. Development of monocytes, macrophages, and dendritic cells. *Science*. 2010;327(5966):656-661.
- Nobs SP, Kopf M. Tissue-resident macrophages: guardians of organ homeostasis. *Trends Immunol*. 2021;42(6):495-507. doi:10.1016/j.it.2021.04.007
- Wynn TA, Chawla A, Pollard JW. Macrophage biology in development, homeostasis and disease. *Nature*. 2013;496(7446):445-455. doi:10.1038/nature12034
- Liu YC, Zou XB, Chai YF, Yao YM. Macrophage polarization in inflammatory diseases. *Int J Biol Sci*. 2014;10(5):520-529. doi:10.7150/ijbs.8879
- Sica A, Erreni M, Allavena P, Porta C. Macrophage polarization in pathology. *Cell Mol Life Sci*. 2015;72(21):4111-4126. doi:10.1007/s00018-015-1995-y
- Epelman S, Lavine KJ, Randolph GJ. Origin and functions of tissue macrophages. *Immunity*. 2014;41(1):21-35. doi:10.1016/j.immuni.2014.06.013
- Ginhoux F, Williams M. Tissue-resident macrophage ontogeny and homeostasis. *Immunity*. 2016;44(3):439-449. doi:10.1016/j.immuni.2016.02.024
- Bain CC, Schridde A. Origin, differentiation, and function of intestinal macrophages. *Front Immunol*. 2018;9:2733. doi:10.3389/fimmu.2018.02733
- Shi C, Pamer EG. Monocyte recruitment during infection and inflammation. *Nat Rev Immunol*. 2011;11(11):762-774. doi:10.1038/nri3070
- Watanabe S, Alexander M, Misharin AV, Budinger GRS. The role of macrophages in the resolution of inflammation. *J Clin Invest*. 2019;129(7):2619-2628. doi:10.1172/JCI124615
- Wculek SK, Dunphy G, Heras-Murillo I, Mastrangelo A, Sancho D. Metabolism of tissue macrophages in homeostasis and pathology. *Cell Mol Immunol*. 2022;19(3):384-408. doi:10.1038/s41423-021-00791-9
- Murray PJ. Macrophage polarization. *Annu Rev Physiol*. 2017;79:541-566. doi:10.1146/annurev-physiol-022516-034339
- Murray PJ, Allen JE, Biswas SK, et al. Macrophage activation and polarization: nomenclature and experimental guidelines. *Immunity*. 2014;41(1):14-20. doi:10.1016/j.immuni.2014.06.008
- Mosser DM, Edwards JP. Exploring the full spectrum of macrophage activation. *Nat Rev Immunol*. 2008;8(12):958-969. doi:10.1038/nri2448
- Lawrence T, Natoli G. Transcriptional regulation of macrophage polarization: enabling diversity with identity. *Nat Rev Immunol*. 2011;11(11):750-761. doi:10.1038/nri3088
- Cabeza-Cabrerizo M, Cardoso A, Minutti CM, Pereira da Costa M, Reis e Sousa C. Dendritic Cells Revisited. *Annu Rev Immunol*. 2021;39(1):131-166. doi:10.1146/annurev-immunol-061020-053707
- Wculek SK, Cueto FJ, Mujal AM, Melero I, Krummel MF, Sancho D. Dendritic cells in cancer immunology and immunotherapy. *Nat Rev Immunol*. 2020;20(1):7-24. doi:10.1038/s41577-019-0210-z
- Sallusto F, Lanzavecchia A. Efficient presentation of soluble antigen by cultured human dendritic cells is maintained by

- granulocyte/macrophage colony-stimulating factor plus interleukin 4 and downregulated by tumor necrosis factor alpha. *J Exp Med*. 1994;179(4):1109-1118. doi:10.1084/jem.179.4.1109
21. Segura E, Amigorena S. Inflammatory dendritic cells in mice and humans. *Trends Immunol*. 2013;34(9):440-445. doi:10.1016/j.it.2013.06.001
 22. Balan S, Dalod M. In vitro generation of human XCR1+ dendritic cells from CD34+ hematopoietic progenitors. *Methods Mol Biol*. 2016;1423:19-37. doi:10.1007/978-1-4939-3606-9_2
 23. Nagy L, Szanto A, Szatmari I, Széles L. Nuclear hormone receptors enable macrophages and dendritic cells to sense their lipid environment and shape their immune response. *Physiol Rev*. 2012;92:739-789. doi:10.1152/physrev.00004.2011-A
 24. Amit I, Winter DR, Jung S. The role of the local environment and epigenetics in shaping macrophage identity and their effect on tissue homeostasis. *Nat Immunol*. 2016;17(1):18-25. doi:10.1038/ni.3325
 25. Hoeksema MA, Glass CK. Nature and nurture of tissue-specific macrophage phenotypes. *Atherosclerosis*. 2019;281:159-167. doi:10.1016/j.atherosclerosis.2018.10.005
 26. Neele AE, van den Bossche J, Hoeksema MA, de Winther MPJ. Epigenetic pathways in macrophages emerge as novel targets in atherosclerosis. *Eur J Pharmacol*. 2015;763:79-89. doi:10.1016/j.ejphar.2015.03.101
 27. Heintzman ND, Stuart RK, Hon G, et al. Distinct and predictive chromatin signatures of transcriptional promoters and enhancers in the human genome. *Nat Genet*. 2007;39(3):311-318. doi:10.1038/ng1966
 28. Creighton MP, Cheng AW, Welstead GG, et al. Histone H3K27ac separates active from poised enhancers and predicts developmental state. *Proc Natl Acad Sci U S A*. 2010;107(50):21931-21936. doi:10.1073/pnas.1016071107
 29. Glass CK, Natoli G. Molecular control of activation and priming in macrophages. *Nat Immunol*. 2016;17(1):26-33. doi:10.1038/ni.3306
 30. Garber M, Yosef N, Goren A, et al. A high-throughput chromatin immunoprecipitation approach reveals principles of dynamic gene regulation in mammals. *Mol Cell*. 2012;47(5):810-822. doi:10.1016/j.molcel.2012.07.030
 31. Platanitis E, Decker T. Regulatory networks involving STATs, IRFs, and NFκB in inflammation. *Front Immunol*. 2018;9:2542. doi:10.3389/fimmu.2018.02542
 32. Gosselin D, Glass CK. Epigenomics of Macrophages. *Immunol Rev*. 2014;262(1):96-112.
 33. Heinz S, Benner C, Spann N, et al. Simple combinations of lineage-determining transcription factors prime cis-regulatory elements required for macrophage and B cell identities. *Mol Cell*. 2010;38(4):576-589. doi:10.1016/j.molcel.2010.05.004
 34. Paul F, Amit I. Plasticity in the transcriptional and epigenetic circuits regulating dendritic cell lineage specification and function. *Curr Opin Immunol*. 2014;30(1):1-8. doi:10.1016/j.coi.2014.04.004
 35. Okabe Y, Medzhitov R. Tissue-specific signals control reversible program of localization and functional polarization of macrophages. *Cell*. 2014;157(4):832-844. doi:10.1016/j.cell.2014.04.016
 36. Buttgerit A, Lelios I, Yu X, et al. Sall1 is a transcriptional regulator defining microglia identity and function. *Nat Immunol*. 2016;17(12):1397-1406. doi:10.1038/ni.3585
 37. Daniel B, Czimmerer Z, Halasz L, et al. The transcription factor EGR2 is the molecular linchpin connecting STAT6 activation to the late, stable epigenomic program of alternative macrophage polarization. *Genes Dev*. 2020;34(21-22):1474-1492. doi:10.1101/gad.343038.120
 38. Kuznetsova T, Prange KHM, Glass CK, de Winther MPJ. Transcriptional and epigenetic regulation of macrophages in atherosclerosis. *Nat Rev Cardiol*. 2020;17(4):216-228. doi:10.1038/s41569-019-0265-3
 39. Zhao GN, Jiang DS, Li H. Interferon regulatory factors: at the crossroads of immunity, metabolism, and disease. *Biochim Biophys Acta Mol Basis Dis*. 2015;1852(2):365-378. doi:10.1016/j.bbadis.2014.04.030
 40. Chistiakov DA, Myasoedova VA, Revin VV, Orekhov AN, Bobryshev YV. The impact of interferon-regulatory factors to macrophage differentiation and polarization into M1 and M2. *Immunobiology*. 2018;223(1):101-111. doi:10.1016/j.imbio.2017.10.005
 41. Mangelsdorf DJ, Evans RM. The RXR heterodimers and orphan receptors. *Cell*. 1995;83(6):841-850.
 42. Shulman AI, Mangelsdorf DJ. Retinoid X Receptor Heterodimers in the Metabolic Syndrome. *N Engl J Med*. 2005;353(6):604-615.
 43. Glass CK, Saijo K. Nuclear receptor transrepression pathways that regulate inflammation in macrophages and T cells. *Nat Rev Immunol*. 2010;10(5):365-376. doi:10.1038/nri2748
 44. Daniel B, Nagy G, Czimmerer Z, et al. The nuclear receptor PPARγ controls progressive macrophage polarization as a ligand-insensitive epigenomic ratchet of transcriptional memory. *Immunity*. 2018;49(4):615-626.e6. doi:10.1016/j.immuni.2018.09.005
 45. Sakai M, Troutman TD, Seidman JS, et al. Liver-derived signals sequentially reprogram myeloid enhancers to initiate and maintain Kupffer cell identity. *Immunity*. 2019;51(4):655-670.e8. doi:10.1016/j.immuni.2019.09.002
 46. Casanova-Acebes M, Menéndez-Gutiérrez MP, Porcuna J, et al. RXRs control serous macrophage neonatal expansion and identity and contribute to ovarian cancer progression. *Nat Commun*. 2020;11(1):1655. doi:10.1038/s41467-020-15371-0
 47. Bensinger SJ, Tontonoz P. Integration of metabolism and inflammation by lipid-activated nuclear receptors. *Nature*. 2008;454(7203):470-477. doi:10.1038/nature07202
 48. Nagy ZS, Czimmerer Z, Nagy L. Nuclear receptor mediated mechanisms of macrophage cholesterol metabolism. *Mol Cell Endocrinol*. 2013;368(1-2):85-98. doi:10.1016/j.mce.2012.04.003
 49. Daniel B, Nagy G, Hah N, et al. The Active enhancer network operated by liganded rxr supports angiogenic activity in macrophages. *Genes Dev*. 2014;28(14):1562-1577. doi:10.1101/gad.242685
 50. Ramón-Vázquez A, de La Rosa JV, Tabraue C, et al. Common and differential transcriptional actions of nuclear receptors liver X receptors and in macrophages. *Mol Cell Biol Published Online*. 2019;39(5):e00376-e00318. doi:10.1128/MCB
 51. Kiss M, Czimmerer Z, Nagy G, et al. Retinoid X receptor suppresses a metastasis-promoting transcriptional program in myeloid cells via a ligand-insensitive mechanism. *Proc Natl Acad Sci U S A*. 2017;114(40):10725-10730. doi:10.1073/pnas.1700785114
 52. Czimmerer Z, Nagy ZS, Nagy G, et al. Extensive and functional overlap of the STAT6 and RXR cisomes in the active enhancer repertoire of human CD14+ monocyte derived differentiating macrophages. *Mol Cell Endocrinol*. 2018;471:63-74. doi:10.1016/j.mce.2017.07.034
 53. Gordon S, Martinez FO. Alternative activation of macrophages: mechanism and functions. *Immunity*. 2010;32(5):593-604. doi:10.1016/j.immuni.2010.05.007
 54. Martinez FO, Helming L, Milde R, et al. Genetic programs expressed in resting and IL-4 alternatively activated mouse and human macrophages: similarities and differences. *Blood*. 2013;121(9):e57-e69. doi:10.1182/blood-2012-06-436212
 55. Sica A, Mantovani A. Macrophage plasticity and polarization: In vivo veritas. *J Clin Invest*. 2012;122(3):787-795. doi:10.1172/JCI59643
 56. Ostuni R, Piccolo V, Barozzi I, et al. Latent enhancers activated by stimulation in differentiated cells. *Cell*. 2013;152(1-2):157-171. doi:10.1016/j.cell.2012.12.018
 57. Czimmerer Z, Daniel B, Horvath A, et al. The transcription factor STAT6 mediates direct repression of inflammatory enhancers and limits activation of alternatively polarized macrophages. *Immunity*. 2018;48(1):75-90.e6. doi:10.1016/j.immuni.2017.12.010
 58. Daniel B, Belk JA, Meier SL, et al. Macrophage inflammatory and regenerative response periodicity is programmed by cell cycle and

- chromatin state. *Mol Cell*. 2023;83(1):121-138.e7. doi:10.1016/j.molcel.2022.11.017
59. Bouhrel MA, Derudas B, Rigamonti E, et al. PPAR γ activation primes human monocytes into alternative M2 macrophages with anti-inflammatory properties. *Cell Metab*. 2007;6(2):137-143. doi:10.1016/j.cmet.2007.06.010
 60. Odegaard JI, Ricardo-Gonzalez RR, Goforth MH, et al. Macrophage-specific PPAR γ controls alternative activation and improves insulin resistance. *Nature*. 2007;447(7148):1116-1120. doi:10.1038/nature05894
 61. Satoh T, Takeuchi O, Vandenbon A, et al. The Jmjd3-Irf4 axis regulates M2 macrophage polarization and host responses against helminth infection. *Nat Immunol*. 2010;11(10):936-944. doi:10.1038/ni.1920
 62. Pello OM, de Pizzol M, Mirolo M, et al. Role of c-MYC in alternative activation of human macrophages and tumor-associated macrophage biology. *Blood*. 2012;119(2):411-421. doi:10.1182/blood-2011-02
 63. Liao X, Sharma N, Kapadia F, et al. Krüppel-like factor 4 regulates macrophage polarization. *J Clin Invest*. 2011;121(7):2736-2749. doi:10.1172/JCI45444
 64. Piccolo V, Curina A, Genua M, et al. Opposing macrophage polarization programs show extensive epigenomic and transcriptional cross-talk. *Nat Immunol*. 2017;18(5):530-540. doi:10.1038/ni.3710
 65. Huang SCC, Smith AM, Everts B, et al. Metabolic reprogramming mediated by the mTORC2-IRF4 signaling Axis is essential for macrophage alternative activation. *Immunity*. 2016;45(4):817-830. doi:10.1016/j.immuni.2016.09.016
 66. Jarjour NN, Schwarzkopf EA, Bradstreet TR, et al. Bhlhe40 mediates tissue-specific control of macrophage proliferation in homeostasis and type 2 immunity. *Nat Immunol*. 2019;20(6):687-700. doi:10.1038/s41590-019-0382-5
 67. Hoeksema MA, Shen Z, Holtman IR, et al. Mechanisms underlying divergent responses of genetically distinct macrophages to IL-4. *Sci Adv*. 2021;7:9808-9824. doi:10.1126/sciadv.abf9808
 68. Huang JT, Welch JS, Ricote M, et al. Interleukin-4-dependent production of PPAR- γ ligands in macrophages by 12/15-lipoxygenase. *Nature*. 1999;400:378-382.
 69. Czimmerer Z, Varga T, Poliska S, Nemet I, Szanto A, Nagy L. Identification of novel markers of alternative activation and potential endogenous PPAR γ ligand production mechanisms in human IL-4 stimulated differentiating macrophages. *Immunobiology*. 2012;217(12):1301-1314. doi:10.1016/j.imbio.2012.08.270
 70. Szanto A, Balint BL, Nagy ZS, et al. STAT6 transcription factor is a facilitator of the nuclear receptor PPAR γ -regulated gene expression in macrophages and dendritic cells. *Immunity*. 2010;33(5):699-712. doi:10.1016/j.immuni.2010.11.009
 71. Waku T, Shiraki T, Oyama T, Maebara K, Nakamori R, Morikawa K. The nuclear receptor PPAR γ individually responds to serotonin and fatty acid-metabolites. *EMBO J*. 2010;29(19):3395-3407. doi:10.1038/emboj.2010.197
 72. McIntyre TM, Pontsler AV, Silva AR, et al. Identification of an Intracellular Receptor for Lysophosphatidic Acid (LPA): LPA Is a Transcellular PPAR Agonist. *Proc Natl Acad Sci U S A*. 2003;100(1):131-136.
 73. Daniel B, Nagy G, Horvath A, et al. The IL-4/STAT6/PPAR γ signaling axis is driving the expansion of the RXR heterodimer cistrome, providing complex ligand responsiveness in macrophages. *Nucleic Acids Res*. 2018;46(9):4425-4439. doi:10.1093/nar/gky157
 74. Seeley JJ, Ghosh S. Molecular mechanisms of innate memory and tolerance to LPS. *J Leukoc Biol*. 2017;101(1):107-119. doi:10.1189/jlb.3mr0316-118rr
 75. Netea MG, Domínguez-Andrés J, Barreiro LB, et al. Defining trained immunity and its role in health and disease. *Nat Rev Immunol*. 2020;20(6):375-388. doi:10.1038/s41577-020-0285-6
 76. Qiao Y, Giannopoulou EG, Chan CH, et al. Synergistic activation of inflammatory cytokine genes by interferon- γ -induced chromatin remodeling and toll-like receptor signaling. *Immunity*. 2013;39(3):454-469. doi:10.1016/j.immuni.2013.08.009
 77. Kang K, Bachu M, Park SH, et al. IFN- γ selectively suppresses a subset of TLR4-activated genes and enhancers to potentiate macrophage activation. *Nat Commun*. 2019;10(1):3320. doi:10.1038/s41467-019-11147-3
 78. *Regulation of Interferon and Toll-like Receptor Signaling during Macrophage Activation by Opposing Feedforward and Feedback Inhibition Mechanisms*.
 79. Chen J, Ivashkiv LB. IFN- γ abrogates endotoxin tolerance by facilitating toll-like receptor-induced chromatin remodeling. *Proc Natl Acad Sci U S A*. 2010;107(45):19438-19443. doi:10.1073/pnas.1007816107
 80. Raundhal M, Morse C, Khare A, et al. High IFN- γ and low SLPI mark severe asthma in mice and humans. *J Clin Invest*. 2015;125(8):3037-3050. doi:10.1172/JCI80911
 81. Potian JA, Rafi W, Bhatt K, McBride A, Gause WC, Salgame P. Preexisting helminth infection induces inhibition of innate pulmonary anti-tuberculosis defense by engaging the IL-4 receptor pathway. *J Exp Med*. 2011;208(9):1863-1874. doi:10.1084/jem.20091473
 82. Goleva E, Jackson LP, Harris JK, et al. The effects of airway microbiome on corticosteroid responsiveness in asthma. *Am J Respir Crit Care Med*. 2013;188(10):1193-1201. doi:10.1164/rccm.201304-0775OC
 83. Harris J, de Haro SA, Master SS, et al. T helper 2 cytokines inhibit autophagic control of intracellular mycobacterium tuberculosis. *Immunity*. 2007;27(3):505-517. doi:10.1016/j.immuni.2007.07.022
 84. Osborne LC, Monticelli LA, Nice TJ, et al. Virus-helminth coinfection reveals a microbiota-independent mechanism of immunomodulation. *Science*. 2014;345(6196):578-582. 1979. doi:10.1126/science.1256942
 85. Gauthier M, Chakraborty K, Oriss TB, et al. Severe asthma in humans and mouse model suggests a CXCL10 signature underlies corticosteroid-resistant Th1 bias. *JCI Insight*. 2017;2(13):94580. doi:10.1172/jci.insight.94580
 86. Eisenbarth SC, Piggott DA, Huleatt JW, Visintin I, Herrick CA, Bottomly K. Lipopolysaccharide-enhanced, toll-like receptor 4-dependent T helper cell type 2 responses to inhaled antigen. *J Exp Med*. 2002;196(12):1645-1651. doi:10.1084/jem.20021340
 87. Kang K, Park SH, Chen J, et al. Interferon- γ represses M2 gene expression in human macrophages by disassembling enhancers bound by the transcription factor MAF. *Immunity*. 2017;47(2):235-250.e4. doi:10.1016/j.immuni.2017.07.017
 88. Varin A, Mukhopadhyay S, Herbein G, Gordon S. Alternative activation of macrophages by IL-4 impairs phagocytosis of pathogens but potentiates microbial-induced signalling and cytokine secretion. *Blood*. 2010;115(2):353-362. doi:10.1182/blood-2009-08-236711
 89. Major J, Fletcher JE, Hamilton TA. IL-4 pretreatment selectively enhances cytokine and chemokine production in lipopolysaccharide-stimulated mouse peritoneal macrophages. *J Immunol*. 2002;168(5):2456-2463. doi:10.4049/jimmunol.168.5.2456
 90. Czimmerer Z, Halasz L, Daniel B, et al. The epigenetic state of IL-4-polarized macrophages enables inflammatory cistrome expansion and extended synergistic response to TLR ligands. *Immunity*. 2022;55(11):2006-2026.e6. doi:10.1016/j.immuni.2022.10.004
 91. Thomas DG, Doran AC, Fotakis P, et al. LXR suppresses inflammatory gene expression and neutrophil migration through cis-repression and cholesterol efflux. *Cell Rep*. 2018;25(13):3774-3785. e4. doi:10.1016/j.celrep.2018.11.100
 92. Núñez V, Alamedaa D, Rico D, et al. Retinoid X receptor α controls innate inflammatory responses through the up-regulation of chemokine expression. *Proc Natl Acad Sci U S A*. 2010;107(23):10626-10631. doi:10.1073/pnas.0913545107

93. Schneider C, Nobs SP, Kurrer M, Rehrauer H, Thiele C, Kopf M. Induction of the nuclear receptor PPAR- γ 3 by the cytokine GM-CSF is critical for the differentiation of fetal monocytes into alveolar macrophages. *Nat Immunol.* 2014;15(11):1026-1037. doi:10.1038/ni.3005
94. A-Gonzalez N, Guillen JA, Gallardo G, et al. The nuclear receptor LXR α controls the functional specialization of splenic macrophages. *Nat Immunol.* 2013;14(8):831-839. doi:10.1038/ni.2622

How to cite this article: Czimmerer Z, Nagy L. Epigenomic regulation of macrophage polarization: Where do the nuclear receptors belong? *Immunol Rev.* 2023;317:152-165. doi:10.1111/imr.13209