



An analogue of a theorem of Steinitz for ball polyhedra in \mathbb{R}^3

SAMI MEZAL ALMOHAMMAD, ZSOLT LÁNGI, AND MÁRTON NASZÓDI 

Abstract. Steinitz's theorem states that a graph G is the edge-graph of a 3-dimensional convex polyhedron if and only if, G is simple, plane and 3-connected. We prove an analogue of this theorem for ball polyhedra, that is, for intersections of finitely many unit balls in \mathbb{R}^3 .

Mathematics Subject Classification. 52B10, 52A30, 52B05.

Keywords. Steinitz's theorem, Polyhedron, Ball polyhedron, Edge-graph.

1. Introduction

Our work takes place in Euclidean 3-space. For the closed ball of radius ρ centered at $x \in \mathbb{R}^3$, we use the notation $\mathbf{B}[x, \rho] := \{y \in \mathbb{R}^3 : d(x, y) \leq \rho\}$. The 2-dimensional sphere (the boundary of a closed ball) is denoted by $\mathbb{S}^2(x, \rho) := \{y \in \mathbb{R}^3 : d(x, y) = \rho\}$. For brevity, we set $\mathbf{B}[x] := \mathbf{B}[x, 1]$, $\mathbb{S}(x) := \mathbb{S}^2(x, 1)$ and for a set $X \subseteq \mathbb{R}^3$, we write $\mathbf{B}[X] := \bigcap_{x \in X} \mathbf{B}[x]$.

Let $X \subset \mathbb{R}^3$ be a finite, nonempty set contained in a ball of radius less than 1. The set $P = \mathbf{B}[X]$ is called a *ball polyhedron*. For any $x \in X$, we call $\mathbf{B}[x]$ a *generating ball* of P and $\mathbb{S}(x)$ a *generating sphere* of P . Unless we state otherwise, we will assume that X is a *reduced set of centers*, that is, that $\mathbf{B}[X] \neq \mathbf{B}[X \setminus \{x\}]$ for any $x \in X$.

The face structure of a 3-dimensional ball polyhedron $\mathbf{B}[X]$ is defined in a natural way: a point on the boundary of $\mathbf{B}[X]$ belonging to at least three generating spheres is called a *vertex*; a connected component of the intersection of two generating spheres and $\mathbf{B}[X]$ is called an *edge*, if it is a non-degenerate circular arc; and the intersection of a generating sphere and $\mathbf{B}[X]$ is called a *face*.

The face structure of a ball polyhedron, unlike that of a convex polyhedron, is not necessarily an algebraic lattice, with respect to containment, see [3].

Following [2], we call a ball polyhedron in \mathbb{R}^3 *standard*, if its vertex-edge-face structure is a lattice with respect to containment. This is the case if, and only if, the intersection of any two faces is either empty, or one vertex or one edge, and any two edges share at most one vertex. The paper [4] and Chapter 6 of the beautiful book [5] by Martini, Montejano and Oliveros provide further background on the theory of ball polyhedra.

A fundamental result of Steinitz (see, [6, 8] and [7]) states that a graph G is the edge-graph of a 3-dimensional convex polyhedron if and only if, G is simple (ie., it contains no loops and no parallel edges), plane and 3-*connected* (ie., removing any two vertices and the edges adjacent to them yields a connected graph). In [2], it is shown that the edge-graph of any standard ball polyhedron in \mathbb{R}^3 is simple, plane and 3-connected. Solving an open problem posed in [2] and [1], our main result shows that the converse holds as well.

Theorem 1. *Every 3-connected, simple plane graph is the edge-graph of a standard ball polyhedron in \mathbb{R}^3 .*

The proof of Steinitz’s theorem consists of two parts. First, it is shown that 3-connected, simple plane graphs can be “reduced” by a finite sequence of certain graph operations to the complete graph K_4 on four vertices. Second, in the geometric part, it is shown that if a graph G is obtained from another graph G' by such an operation and G is realizable as the edge-graph of a polyhedron, then G' is realizable as well. To prove Theorem 1, we use the first, combinatorial part without modification. Our contribution is the proof of the second, geometric part in the setting of ball polyhedra.

The structure of the paper is the following. First, in Sect. 2, we introduce these operations on graphs, and recall facts on the face structure of the dual of a ball polyhedron. In Sect. 3, we state our main contribution, Theorem 2, which shows the “backward inheritance” of realizability by ball polyhedra under these graph operations, and deduce Theorem 1 from it. Finally, in Sect. 4, we prove Theorem 2.

2. Preliminaries

2.1. Simple Δ -to- Y and Y -to- Δ reductions on a plane graph

Let G be a 3-connected plane graph and K_3 be a triangular face with vertices v_1, v_2 and v_3 (resp., $K_{1,3}$ be a subgraph consisting of a 3-valent vertex v of G , its neighbors v_1, v_2, v_3 , and the edges $\{v, v_i\}$ connecting v to its neighbors). A ΔY operation is defined as the graph operation which removes the edges $\{v_i, v_j\}$ of a triangular face K_3 , adds a new vertex v from the face, and connects it to v_i s, or vice versa, it takes a subgraph $K_{1,3}$ of G , removes the vertex v and the edges incident to it, then connects all pairs $v - I, v_j$ by an edge.




FIGURE 1. \implies : A Δ -to- Y transformation; \impliedby : A Y -to- Δ transformation




FIGURE 2. Examples of SP-reductions

To specify the direction of the transformation, we will distinguish between a Δ -to- Y transformation and a Y -to- Δ transformation (see Fig. 1).

A ΔY operation may create multiple edges or vertices of degree two. A graph with such objects is clearly not the edge-graph of a standard ball polyhedron. To fix these issues, we define the following notion. A *series-parallel reduction*, or *SP-reduction* is the replacement of a pair of edges incident to a vertex of degree 2 with a single edge or, the replacement of a pair of parallel edges with a single edge that connects their common endpoints, see Fig. 2.

Assume that a graph G contains $K_{1,3}$ as a subgraph whose degree 3 vertex is denoted by v , and its neighbors are v_1, v_2, v_3 (resp., K_3 with vertices v_1, v_2, v_3), see Fig. 1. We call edges of G that connect two vertices of $K_{1,3}$ (resp., K_3) *internal edges*. We define the *outer degree* of a neighbor of v (resp., a vertex of K_3), as the number of non-internal edges adjacent to it. A $K_{1,3}$ is called Y_0, Y_1, Y_2 , or Y_3 if it has zero, one, two, or three internal edges respectively. A K_3 is called $\Delta_0, \Delta_1, \Delta_2$, or Δ_3 if it has zero, one, two, or three vertices of outer degree one, respectively.

A *simple ΔY reduction* means any ΔY operation followed immediately by SP-reductions that are then possible. There are four different types of simple Δ -to- Y and Y -to- Δ reductions (cf. Corollary 4.7 of [8]), as shown in Fig. 3.

Proposition 2.1. *Every 3-connected plane graph G can be reduced to K_4 by a sequence of simple ΔY reductions.*




FIGURE 3. **a** Four types of simple Δ -to- Y reduction, and **b** four types of simple Y -to- Δ reduction, where the dotted lines denote edges that may or may not be present, and are not affected by the simple Δ -to- Y and Y -to- Δ reductions

2.2. Standard graphs

A planar graph with a fixed drawing on the plane is called a *plane graph*. It is well known that 3-connected planar graphs have only one drawing, that is, all plane drawings of such a graph have isomorphic face lattices [8, Section 4.1].

Definition 2.1. Let G be a plane graph. We call G *standard*, if

- (i) the intersection of any two faces is either empty, or one vertex or one edge, and
- (ii) any two edges share at most one vertex.

Remark 2.1. Let G be the edge-graph of a ball polyhedron. Then G is standard if and only if the ball polyhedron is a standard ball polyhedron.

We leave the proof of the following two lemmas to the reader as an exercise.

Lemma 2.1. *Let G be a 3-connected plane graph and let the graph G_0 be derived from G by a simple Δ -to- Y reduction. If G is a standard graph, then so is G_0 .*

The subdivision of an edge $\{t_1, t_2\}$ of a graph G is another graph obtained from G by removing the edge $\{t_1, t_2\}$, then adding a new vertex t' and, finally, adding the edges $\{t_1, t'\}$ and $\{t', t_2\}$.

Lemma 2.2. *Let G be a standard graph, E be a face of G and $\{u_1, u_3\}, \{u_3, u_2\}$ be two edges of E such that u_1 and u_2 are non-adjacent vertices.*

- I. *If the graph H is obtained from G by adding the edge $\{u_1, u_2\}$, then H is a standard graph.*
- II. *If the graph H' is obtained from G by adding the edge $\{u_2, u'\}$ where u' is a new vertex subdividing the edge $\{u_1, u_3\}$, then H' is a standard graph.*
- III. *If the graph H'' is obtained from G by adding the edge $\{u', u''\}$ where u' and u'' are two new vertices subdividing the edges $\{u_1, u_3\}$ and $\{u_3, u_2\}$ respectively, then H'' is a standard graph.*

2.3. Graph duality

We denote the dual of a plane graph G by G^* , see [8, Section 4.1]. It is well known that G^* is also a plane graph, and G^* is 3-connected if and only if, G is 3-connected.

According to the following fact, simple Δ -to- Y reductions and simple Y -to- Δ reductions are dual to each other, see [8, Section 4.2].

Proposition 2.2. *Let G and G' be 3-connected plane graphs. Then G' is obtained from G by a simple Δ -to- Y reduction if and only if, G'^* is obtained from G^* by a simple Y -to- Δ reduction.*

2.4. The dual of a ball polyhedron

In the following, $\mathcal{F}(\mathbf{B}[X])$ denotes the set of faces, and $\mathcal{V}(\mathbf{B}[X])$ denotes the set of vertices of the ball polyhedron $\mathbf{B}[X]$.

Let $\mathbf{B}[X]$ be a ball polyhedron in \mathbb{R}^3 all of whose faces contain at least three vertices. In [3], the dual of $\mathbf{B}[X]$ is introduced as the ball polyhedron $\mathbf{B}[\mathcal{V}(\mathbf{B}[X])]$, and a bijection, called the duality mapping between $\mathbf{B}[X]$ and $\mathbf{B}[\mathcal{V}(\mathbf{B}[X])]$, is given between the faces, edges and vertices of $\mathbf{B}[X]$ and $\mathbf{B}[\mathcal{V}(\mathbf{B}[X])]$, consisting of the following three mappings:

- (1) The vertex-face mapping is

$$\mathcal{V}(\mathbf{B}[X]) \ni v \mapsto V \in \mathcal{F}(\mathbf{B}[\mathcal{V}(\mathbf{B}[X])])$$
 where V is the face of $\mathbf{B}[\mathcal{V}(\mathbf{B}[X])]$ with v as its center.

- (2) The *face-vertex* mapping is $\mathcal{F}(\mathbf{B}[X]) \ni F \mapsto f \in \mathcal{V}(\mathbf{B}[\mathcal{V}(\mathbf{B}[X])])$ where f is the center of the sphere supporting the face F .
- (3) The *edge-edge* mapping is the following. Two vertices in $\mathbf{B}[\mathcal{V}(\mathbf{B}[X])]$ are connected by an edge if and only if, the corresponding faces of $\mathbf{B}[X]$ meet in an edge.

Note that every face of a standard ball polyhedron contains at least three edges. The relationship between graph duality and duality of ball polyhedra is described below.

Lemma 2.3. ([1], Theorem 6.6.5) *Let P be a standard ball polyhedron of \mathbb{R}^3 . Then the intersection P^* of the closed unit balls centered at the vertices of P is another standard ball polyhedron whose face lattice is dual to that of P .*

3. Proof of Theorem 1

Our main contribution follows.

Theorem 2. *Let G' be a 3-connected plane graph, and let the graph G be derived from G' by a simple Y -to- Δ reduction. If G is the edge-graph of a standard ball polyhedron in \mathbb{R}^3 , then so is G' .*

First, we show how Theorem 2 implies Theorem 1.


Proof of Theorem 1. Let G be a 3-connected simple plane graph. By Proposition 2.1, the graph G reduces to K_4 the edge-graph of the standard ball tetrahedron by a sequence of simple ΔY reductions.

Now we show that the standard ball tetrahedron can be gradually turned into a realization of G . Let H be the edge-graph of a standard ball polyhedron and assume that H is obtained from another edge-graph H' by a simple ΔY reduction. We want to show that H' is realized by a standard ball polyhedron. So we need to discuss two cases:

First, assume that H is obtained from H' by a simple Y -to- Δ reduction. Then by Theorem 2, H' is realized by a standard ball polyhedron.

Second, assume that H is obtained from H' by a simple Δ -to- Y reduction. Then by Proposition 2.2, we get that the edge-graph H^* is obtained from the edge-graph H'^* by a simple Y -to- Δ reduction. By Lemma 2.3, the edge-graph H^* is realized by a standard ball polyhedron, and by Theorem 2, the edge-graph H'^* is realized by a standard ball polyhedron. Again by Lemma 2.3, the edge-graph H' is realized by a standard ball polyhedron, and this completes the proof. □

By Lemma 2.3, H' is the edge-graph of a standard ball-polyhedron.



By assumption, H is the edge-graph of a standard ball-polyhedron.

By Theorem 2, H'^* is the edge-graph of a standard ball-polyhedron.

By Lemma 2.3, H^* is the edge-graph of a standard ball-polyhedron.

4. Proof of Theorem 2

Let $\emptyset \neq X \subset \mathbb{R}^3$ be a finite set and $\mathbf{B}[X]$ be a standard ball polyhedron with edge-graph G , and assume that G is obtained from a graph G' by a simple Y -to- Δ reduction. We need to show that G' is realized by a standard ball polyhedron.

Let Λ denote the triangular face of $\mathbf{B}[X]$ which realizes the triangle obtained in the Y -to- Δ reduction, let $\mathbb{S}(x_\Lambda)$ be its supporting unit sphere, and v_1, v_2, v_3 be the vertices of Λ and e_1, e_2, e_3 the edges. Let F_1, F_2 and F_3 denote the faces of $\mathbf{B}[X]$ distinct from Λ containing e_1, e_2 and e_3 respectively, and let $\mathbb{S}(x_1), \mathbb{S}(x_2)$ and $\mathbb{S}(x_3)$ be the unit spheres supporting these faces, see Fig. 4.

The starting point of the proof of Theorem 2 is the removal of the ball that generates the triangular face Λ . Thus, we obtain another ball polyhedron, $\mathbf{B}[X \setminus \{x_\Lambda\}]$. The following lemma (which we prove later) describes the edge-graph of $\mathbf{B}[X \setminus \{x_\Lambda\}]$ and, combined with Lemma 2.1 yields that it is a standard graph, and hence, by Remark 2.1, $\mathbf{B}[X \setminus \{x_\Lambda\}]$ is a standard ball polyhedron.




FIGURE 4. Face Λ and its neighbors




FIGURE 5. Reconstruction of reductions

Lemma 4.1. *The edge-graph of the ball polyhedron $\mathbf{B}[X \setminus \{x_\Lambda\}]$ is obtained by a simple Δ -to- Y reduction applied to Λ in the role of K_3 .*

The edge-graph of $\mathbf{B}[X \setminus \{x_\Lambda\}]$ described in Lemma 4.1 may be G' , in which case we are done. However, it may happen that this is not G' , more precisely, the graph G is derived from G' by a simple Y -to- Δ reduction, but the converse is not always true, it may happen that G' is not derived from G by a simple Δ -to- Y reduction. The reason is that when we do a Δ -to- Y reduction, the vertices of the triangle of outer degree one in G become degree two vertices in the graph obtained from G by a Δ -to- Y reduction. Next, we do the SP -reduction, and these vertices are lost, see Fig. 5(c) and (d). Moreover, the internal edges will be missing as well, see Fig. 5(b), (c) and (d).

The following lemma describes how the edge-graph of $\mathbf{B}[X \setminus \{x_\Lambda\}]$ is converted into G' by adding the missing vertices and edges. We achieve this by adding some extra balls. Lemma 2.2 yields that G' is a standard graph, and hence, by Remark 2.1, the ball polyhedron realizing G' is a standard ball polyhedron.

Lemma 4.2. *Let $\emptyset \neq Z \subset \mathbb{R}^3$ be a finite set and $\mathbf{B}[Z]$ be a ball polyhedron. If the edge-graph of $\mathbf{B}[Z]$ contains Y_0 as an induced subgraph whose vertices are u, u_1, u_2 and u_3 , and whose edges are a_1, a_2 and a_3 , see Fig. 6, left side, then*

- I. *there exists a center w such that the edge-graph of the ball polyhedron $\mathbf{B}[Z \cup \{w\}]$ is obtained from the edge-graph of $\mathbf{B}[Z]$ by adding the internal edge $\{u_1, u_2\}$.*
- II. *there exists a center w' such that the edge-graph of the ball polyhedron $\mathbf{B}[Z \cup \{w'\}]$ is obtained from the edge-graph of $\mathbf{B}[Z]$ by adding the edge $\{u_2, u'_1\}$, where u'_1 is a new vertex subdividing the edge a_1 .*
- III. *there exists a center w'' such that the edge-graph of the ball polyhedron $\mathbf{B}[Z \cup \{w''\}]$ is obtained from the edge-graph of $\mathbf{B}[Z]$ by adding the edge $\{u'_1, u'_2\}$, where u'_1 and u'_2 are two new vertices subdividing the edges a_1 and a_2 respectively.*

In summary, proving Lemmas 4.1 and 4.2, we prove Theorem 2, which in turn yields Theorem 1.

4.1. Proof of Lemma 4.1

In this section, we use the notation of Lemma 4.1 and Fig. 4.

The following claim is obvious.

Claim 4.1. *For any $i \in \{1, 2, 3\}$, $\Lambda \setminus e_i$ is contained in the interior of $\mathbf{B}[x_i]$.*

The following claim is the key to our proof. It states that the “new part” of the boundary of the new ball polyhedron $\mathbf{B}[X \setminus \{x_\Lambda\}]$ belongs to the union of $\mathbb{S}(x_1)$, $\mathbb{S}(x_2)$ and $\mathbb{S}(x_3)$.

Claim 4.2. $\text{bd}(\mathbf{B}[X']) \setminus \text{bd}(\mathbf{B}[X]) \subseteq \mathbb{S}(x_1) \cup \mathbb{S}(x_2) \cup \mathbb{S}(x_3)$, where $X' = X \setminus \{x_\Lambda\}$.

Proof. Consider a point $q \in \text{bd}(\mathbf{B}[X']) \setminus \text{bd}(\mathbf{B}[X])$. Then there exists a generating sphere $\mathbb{S}(x_q)$ of $\mathbf{B}[X']$ such that $q \in \mathbb{S}(x_q)$ and $x_q \in X'$, implying that $\mathbb{S}(x_q)$ is a generating sphere of $\mathbf{B}[X]$ as well. Let $F = \mathbb{S}(x_q) \cap \mathbf{B}[X]$ and $F' = \mathbb{S}(x_q) \cap \mathbf{B}[X']$. Then $F = F' \cap \mathbf{B}[x_\Lambda]$, $F \subseteq F'$, $q \in F'$, and $q \notin F$. This yields that $F' \cap \mathbb{S}(x_\Lambda)$ is a non-degenerate circular arc in F' that separates q from F . Thus F intersects Λ in a non-degenerate circular arc. That only happens if F intersects Λ in an edge of $\mathbf{B}[X]$, and hence, $x_q = x_1, x_2$, or x_3 . □

The following Claim is obvious.

Claim 4.3. *Let $\mathbf{B}_1, \mathbf{B}_2$ and \mathbf{B}_3 be closed unit balls in \mathbb{R}^3 such that $\mathbf{B}_1 \cap \mathbf{B}_2 \cap \mathbf{B}_3$ is a ball polyhedron with three faces. Then $\mathbf{B}_1 \cap \mathbf{B}_2 \cap \mathbf{B}_3$ is a ball polyhedron with two vertices connected by three edges.*

Finally, we are in the position to prove Lemma 4.1. By Claim 4.3, the boundary of $\mathbf{B}[x_1] \cap \mathbf{B}[x_2] \cap \mathbf{B}[x_3]$ contains two vertices of degree three say q and \bar{q} , three edges, and three faces. We need to prove that the “new part” $\mathcal{N} := \text{bd}(\mathbf{B}[X \setminus \{x_\Lambda\}]) \setminus \text{bd}(\mathbf{B}[X])$ of the boundary of the ball polyhedron $\mathbf{B}[X \setminus \{x_\Lambda\}]$ contains either q or \bar{q} with part from each of the three edges, and part from each of the three faces (i.e., $K_{1,3}$). It means that when we remove the ball $\mathbf{B}[x_\Lambda]$, the triangular face Λ of G will be replaced by $Y_0 = K_{1,3}$ in the edge-graph of $\mathbf{B}[X \setminus \{x_\Lambda\}]$, i.e., the edge-graph of $\mathbf{B}[X \setminus \{x_\Lambda\}]$ is derived from G by a simple Δ -to- Y reduction.

Let $\Gamma := \mathbf{B}[x_1] \cap \mathbf{B}[x_2] \cap \mathbf{B}[x_3]$ and $\gamma := \text{bd}(\Lambda) = e_1 \cup e_2 \cup e_3$, see Fig. 4. By Claim 4.2, $\mathcal{N} \subseteq \mathbb{S}(x_1) \cup \mathbb{S}(x_2) \cup \mathbb{S}(x_3)$, which implies that $\mathcal{N} \subseteq \text{bd}(\Gamma)$. Clearly, $\text{bd}(\Gamma)$ has three edges, say e'_1, e'_2 and e'_3 such that $e'_1 \subseteq \mathbb{S}(x_2) \cap \mathbb{S}(x_3)$, $e'_2 \subseteq \mathbb{S}(x_1) \cap \mathbb{S}(x_3)$ and $e'_3 \subseteq \mathbb{S}(x_1) \cap \mathbb{S}(x_2)$.

By Claim 4.1, $v_i \in \text{int}(\mathbf{B}[x_i])$ for all $i = 1, 2, 3$. By Claim 4.3, $\mathbb{S}(x_1) \cap \mathbb{S}(x_2) \cap \mathbb{S}(x_3)$ is a set of two points q and \bar{q} . Clearly, $\{v_1, v_2, v_3\} \cap \{q, \bar{q}\} = \emptyset$.

Since $e'_1 \setminus \{q, \bar{q}\} \subseteq \text{int}(\mathbf{B}[x_1])$, $v_1 \in e'_1 \setminus \{q, \bar{q}\}$ and $\{e'_2, e'_3\} \subseteq \mathbb{S}(x_1)$, we have that v_1 belongs to $e'_1 \setminus \{q, \bar{q}\}$ and does not belong to e'_2 or to e'_3 . Similarly, v_i belongs to $e'_i \setminus \{q, \bar{q}\}$ ($i = 2, 3$) only, respectively. Thus, exactly one of v_1, v_2 and v_3 is contained on each of the three edges of Γ .

Observe that both $\mathbb{S}(x_\Lambda) \cap \Gamma$ and Λ are the intersections of three spherical disks on $\mathbb{S}(x_\Lambda)$, each smaller than a hemi-sphere: $\mathbb{S}(x_\Lambda) \cap \mathbf{B}[x_1]$, $\mathbb{S}(x_\Lambda) \cap \mathbf{B}[x_2]$ and $\mathbb{S}(x_\Lambda) \cap \mathbf{B}[x_3]$. Hence, $\mathbb{S}(x_\Lambda) \cap \Gamma = \Lambda$, and it follows that $\mathbb{S}(x_\Lambda) \cap \text{bd}(\Gamma) = \gamma$.

It follows that γ partitions $\text{bd}(\Gamma)$ into two components, q is in one component and \bar{q} is in the other, so we may assume that $\bar{q} \in \text{int}(\mathbf{B}[x_\Lambda])$. Claim 4.2 yields that $q \in \mathcal{N}$ as required. This completes the proof of Lemma 4.1.

4.2. Proof of Lemma 4.2

(I) Let g_1, g_2 and g_3 be the faces of $\mathbf{B}[Z]$ such that $a_1 = g_2 \cap g_3$, $a_2 = g_1 \cap g_3$ and $a_3 = g_1 \cap g_2$ and let $\mathbb{S}(z_1), \mathbb{S}(z_2)$ and $\mathbb{S}(z_3)$ be the spheres supporting these faces, see Fig. 6, left side.

To add an internal edge to Y_0 , we will add a rotated copy z'_3 of z_3 to the set Z , where the axis of the rotation is the line through u_1 and u_2 , the angle of the rotation is sufficiently small, and u is outside of $\mathbf{B}[z'_3]$. Thus, we obtain a new triangular face g'_3 supported by $\mathbb{S}(z'_3)$, see the dashed lines in Fig. 6(a), and remove the dotted lines in Fig. 6(a).

(II) To add a new vertex and a new edge to Y_0 , we use the same method as in (I), but we choose the rotation axis so that it passes through the vertex




FIGURE 6. The dotted (blue) vertices and edges are removed and the dashed (red) vertices and edges are introduced in the new graph (Color figure online)

u_2 and intersects the edge a_1 at a point, say u'_1 , distinct from its endpoints, see Fig. 6(b).

(III) To add two new vertices and a new edge to Y_0 , we use again the same method as in (I), but this time, we choose the rotation axis so that it intersects the edges a_1 and a_2 at two points, say u'_1 and u'_2 respectively, distinct from their endpoints, see Fig. 6(c). This finishes the proof of Lemma 4.2.

Acknowledgements

SMA would like to thank the Tempus Public Foundation (TPF), Stipendium Hungaricum program, and University of Thi-Qar, Iraq for the support for his PhD scholarship.

ZL was supported by Grants K119670 and BME Water Sciences & Disaster Prevention TKP2020 IE of the National Research, Development and Innovation Fund (NRDI), by the ÚNKP-20-5 New National Excellence Program of the Ministry for Innovation and Technology, and the János Bolyai Scholarship of the Hungarian Academy of Sciences.

MN was supported by the National Research, Development and Innovation Fund (NRDI) Grant K119670, by the ÚNKP-20-5 New National Excellence Program of the Ministry for Innovation and Technology from the source of the NRDI, as well as the János Bolyai Scholarship of the Hungarian Academy of Sciences.

Funding Open access funding provided by Eötvös Loránd University.

Open Access. This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

References

- [1] Bezdek, K.: Lectures on sphere arrangements-the discrete geometric side. Springer, Berlin (2013)
- [2] Bezdek, K., Lángi, Z., Naszódi, M., Papez, P.: Ball-polyhedra. *Discrete Comput. Geom.* **38**(2), 201–230 (2007)
- [3] Bezdek, K., Naszódi, M.: Rigidity of ball-polyhedra in Euclidean 3-space. *Eur. J. Comb.* **27**(2), 255–268 (2006)
- [4] Kupitz, Y.S., Martini, H., Perles, M.A.: Ball polytopes and the Vázsonyi problem. *Acta Math. Hungar.* **126**(1–2), 99–163 (2010)
- [5] Martini, H., Montejano, L., Oliveros, D.: Bodies of Constant Width. An Introduction to Convex Geometry with Applications, Birkhäuser/Springer, Cham (2019)
- [6] Steinitz, E., Rademacher, H.: Vorlesungen über die Theorie der Polyeder. Springer-Verlag, Berlin (1934)
- [7] Steinitz, E.: Polyeder und Raumeinteilungen. In: *Encyclopädie der mathematischen Wissenschaften*, Band 3 (Geometrie), vol. 3A12, pp. 1–139 (1922)
- [8] Ziegler, G.M.: Lectures on Polytopes, Graduate Texts in Mathematics. Springer-Verlag, Berlin (1995)

Sami Mezal Almohammad
Institute of Mathematics
Loránd Eötvös University
Budapest
Hungary
e-mail: sami85@cs.elte.hu;
sami.mezal@utq.edu.iq

and

Department of Mathematics
Faculty of Computer Science and Mathematics
University of Thi-Qar
Thi-Qar
Iraq

Zsolt Lángi
MTA-BME Morphodynamics Research Group and Department of Geometry
Budapest University of Technology
Budapest
Hungary
e-mail: zlangi@math.bme.hu

Márton Naszódi
Department of Geometry and MTA-ELTE Lendület Combinatorial Geometry Research
Group
Loránd Eötvös University
Budapest
Hungary
e-mail: marton.naszodi@math.elte.hu

Received: December 5, 2020

Revised: May 21, 2021

Accepted: May 24, 2021