

Algebraic models for shell-like quarteting of nucleons

J. Cseh

Institute for Nuclear Research, Hungarian Academy of Sciences, Debrecen, Pf. 51, Hungary-4001

(Dated: July 23, 2018)

Algebraic models are proposed for the description of the shell-like quarteting of the nucleons both on the phenomenologic and on the semimicroscopic levels. In the previous one the quartet is considered as a structureless object, while in the latter one its constituents are treated explicitly. The excitation spectrum is generated by the $SU(3)$ formalism in both cases. An application to the ^{20}Ne nucleus is presented.

PACS numbers: 21.60.Fw, 21.60.Cs, 27.30.+t

Quarteting is an important phenomenon in several branches of physics [1, 2]. In nuclear physics it appears in a straightforward way due to the fact that the exclusion principle allows two (spin $\frac{1}{2}$) protons and neutrons to occupy a single-particle state, and the short-range attractive nucleon-nucleon forces prefer this arrangement. Therefore, it has long been known, and recently a conjecture was put forward on the importance of quarteting also in nuclei away from the line of stability [3].

A well-known signature of quarteting is that the separation energy of a nucleon in an even-even $N = Z$ nucleus is much larger than that of an α -particle. The fact that the nuclear mass of $4n$ nuclei is approximately a linear function of n , while the masses of $4n + x$ nuclei are quadratic function of x , was already the motivation for Wigner's supermultiplet theory [4]. (Much work has been done on the binding energies and quarteting later on, too, see e.g. [5].)

A nuclear quartet model was formulated in [6], (based on the stretched scheme), and then it was generalized in several steps. In [7] quartet excitations were considered from one major shell to the other, and the corresponding energies were determined from mass relationships. In this generalized interpretation a quartet is not related to a specific angular momentum coupling scheme: it is made of 2 protons and 2 neutrons, occupying a fourfold degenerate single particle state (l, m orbit in $L - S$ coupling, or j, m and $j, -m$ orbits in $j - j$ coupling). The internal binding of a quartet is strong, while the quartet-quartet interaction is relatively weak. Arima and Gillet took into account [8] also pairs of nucleons, as further building blocks, extending the description to even-even nuclei of different Z and N .

In [9] intrashell quartet excitations have been introduced in addition to the intershell excitations of [7]. This concept leads to a quartet shell model, i.e. one assumes the existence of a self-consistent quartet potential well, and its states are used to describe the quartet states in $4n$ nuclei. The $0s, 0p, 1s - 0d, \dots$ oscillator shells of the nucleon-shell model are replaced by $0s, 0p, 1s - 0d, \dots$ quartet shells, having 1, 3, 6, ... single quartet states, respectively. The corresponding energies were determined empirically, too.

A further extension was presented in [10] by incorporating any number of particle-hole excitations (in the

language of the nucleon-shell-model), contrary to the quartet-shell-model of [7, 9] which had only 0, 4, 8 ... excitation quanta (in terms of nucleon-shell-model). This considerable extension of the quartet model space appeared due to the conceptual generalization of a quartet. Harvey defined [10] it as 2 protons and 2 neutrons having a quartet-symmetry: permutational symmetry of [4], and spin-isospin symmetry of [1,1,1,1].

Interacting boson type quartet models were invented [11, 12] for the description of quarteting in heavy nuclei. In [11] the basic building block quartets are treated as $l = 0$ (s) and $l = 2$ (d) bosons, and the model has a $U(6)$ group structure, like the interacting boson model of the quadrupole collectivity [13]. This model describes a spectrum of positive parity states. In [12] the alpha-like correlation is treated in terms of bosons of nucleon-pairs, but in addition to the s and d bosons another set of basic building blocks of $l = 0$ (s^*) boson and $l = 1$ (p) boson is included, therefore, negative parity states are also involved. These phenomenological models have the efficiency and elegance of the algebraic methods in generating the spectrum. E.g. they have dynamical symmetries as limiting cases, which provide us with exact solutions for the eigenvalue problem.

In [14] a BCS-like study was carried out for bosons of the proton-neutron interacting boson model [13] and it was concluded that the superfluid condensate is more of a quartet type, rather than separate superfluid phases of proton and neutron pairs. Recent investigations [15, 16] show that the isovector pairing in self-conjugate nuclei are of quartet type and can be well described by a quartet condensation model.

Another condensate, namely the alpha-particle condensate attracts much attention these years [17]. For the first sight it is very different from the quartet condensate. The latter one was shown to be important in the ground state, while the Bose-Einstein condensate (BEC) of alpha particles was invented for the description of the gas-like dilute structure near the alpha-threshold. However, later calculations revealed that the the THSR wavefunction, which is applied in the alpha-condensate studies have a very large overlap with the (resonating group method) wavefunction of ground state [18], indicating that the overlap with the quartet condensate is considerable, too. The non-localized nature of cluster-

ing in the BEC also shows in this direction. The exact relation of these two condensates still remains to be understood. In the present work we do not investigate the condensates, rather we concentrate on the “individual” quartet-excitations, in the sense of [7, 9, 10].

We propose algebraic quartet models based on the concepts of shell-model-like quarteting of [7, 9, 10]. Our main purpose is the description of the excitation spectrum. We propose two models: the simpler one is called phenomenologic algebraic quartet model (PAQM), which has the building blocks very similar to that of the quartet-shell model of [9], i.e. the composite nature of the quartet do not appear explicitly. The second one is the semimicroscopic algebraic quartet model (SAQM), based on the quartet concept of [10], in which each of the four nucleons of the quartet is treated. The novel feature in comparison with the works [7, 9, 10] is that an algebraic framework is formulated for the description of the detailed spectrum, like in the group theoretical approach of the works [11–13]. On the other hand, the new models are different from the interacting boson type models of [11, 12], because of the nature of their building blocks, and shell-like structure of the model spaces.

We apply Elliott’s SU(3) scheme [19, 20] for generating the spectrum both in the phenomenological and in the semimicroscopic descriptions. In the former case structureless quartets are supposed to occupy the single-particle levels of the harmonic oscillator shells, while in the latter model nucleons do so. Therefore, the phenomenological model space has only a spatial part, while the semimicroscopic one contains a space and a spin-isospin components. In fact, this latter model space is a truncation of that of the $L-S$ coupled no-core shell model [21], based on the spin-isospin formalism. The physical operators are expressed in terms of the group generators, thus algebraic techniques can be applied in calculating the matrix elements.

The phenomenologic algebraic quartet model. In this approach an excitation quantum $(\hbar\omega)_q$ between the major shells is expected to be approximately 4 times that of the nucleon shell model: $(\hbar\omega)_q \approx 4(\hbar\omega)$. All the shells have positive parity, due to their quartet nature. If a single quartet state is occupied, then no other particle can be put there, therefore, the permutational symmetry of the quartets has to be that of a single-columned Young diagram: [1,1,1,1].

The building blocks of the description are the nine operators, $\hat{A}_{\alpha\beta} = \frac{1}{2}(\hat{a}_\alpha^\dagger \hat{a}_\beta + \hat{a}_\beta \hat{a}_\alpha^\dagger)$, $\alpha, \beta = x, y, z$, $\hat{a}_\alpha = \sum_j \hat{a}_\alpha(j)$, $\hat{a}_\alpha^\dagger = \sum_j \hat{a}_\alpha^\dagger(j)$, $j = 1, \dots, N$; (here N is the total number of particles), which are number-conserving bilinear products of the creation and annihilation operators of oscillator quanta. They can be rewritten into three spherical tensors: a scalar operator \hat{n} , which is the number of oscillator quanta, five components of the quadrupole momentum \hat{Q}_m (acting in a single major shell), and three components of the angular momentum \hat{L}_m . The nine operators $\hat{n}, \hat{Q}_m, \hat{L}_m$ generate the U(3)

group, the eight operators \hat{Q}_m, \hat{L}_m generate the SU(3) group, and the three \hat{L}_m are generators of the SO(3) group.

The basis states are characterized by the representation labels of the group-chain:

$$U(3) \supset SU(3) \supset SO(3) \supset SO(2) \\ |[n_1, n_2, n_3], (\lambda, \mu), K, L, M \rangle. \quad (1)$$

Here $n = n_1 + n_2 + n_3$ is the eigenvalue of the \hat{n} operator. The angular momentum content of a (λ, μ) representation is as follows [19, 20]: $L = K, K+1, \dots, K+max(\lambda, \mu)$, $K = min(\lambda, \mu), min(\lambda, \mu) - 2, \dots, 1$ or 0, with the exception of $K_L = 0$, for which $L = max(\lambda, \mu), max(\lambda, \mu) - 2, \dots, 1$ or 0. The SU(3) content is given by the U(k) \supset SU(3) decomposition [22], where $k = 3, 6, 10, \dots$ for the major shell with 1, 2, 3, ... quartet excitations. The irreducible representation (irrep) of U(k) is the same as that of the permutational group in the major shell in question. The U(3) symmetry of the whole nucleus is obtained as a direct product of the major shell U(3) irreps. The irreps of the spurious center of mass excitations can be determined easily, due to the fact that the c.m. excitation operator is fully symmetric in particle indices, and has an [1, 0, 0] U(3) irreducible tensor character [23–25]. We illustrate here the construction of the model space with the lowest-lying states of the ^{20}Ne nucleus. The ground state contains the filled-in 0 and 1 $\hbar\omega_q$ major shells, and 1 quartet in the 2 $\hbar\omega_q$ major shell: $(0)^1(1)^3(2)^1$. The permutational symmetries in the three subsequent major shells are: [1] \otimes [1, 1, 1] \otimes [1], which give the [1, 1, 1, 1, 1] symmetry of the five-quartet-state. The corresponding U(3) symmetries are unique and simple in these cases, and they result in a single U(3) irrep: $[0, 0, 0] \otimes [1, 1, 1] \otimes [2, 0, 0] = [3, 1, 1]$. The $1\hbar\omega_q$ excitations are obtained in two different ways: 1a: $(0)^1(1)^2(2)^2$, or 1b: $(0)^1(1)^3(3)^1$. The permutational symmetries are: 1a: [1] \otimes [1, 1] \otimes [1, 1] = [1, 1, 1, 1, 1] $\oplus \dots$, and 1b: [1] \otimes [1, 1, 1] \otimes [1] = [1, 1, 1, 1, 1] $\oplus \dots$. The corresponding U(3) symmetries, and their products are: 1a: $[0, 0, 0] \otimes [1, 1, 0] \otimes [3, 1, 0] = [4, 2, 0] \oplus [4, 1, 1] \oplus [3, 2, 1]$; 1b: $[0, 0, 0] \otimes [1, 1, 1] \otimes [3, 0, 0] = [4, 1, 1]$. In total the U(3) irreps are: $[4, 2, 0] \oplus [4, 1, 1]^2 \oplus [3, 2, 1]$. The spurious excitation of the centre of mass: $[3, 1, 1] \otimes [1, 0, 0] = [4, 1, 1] \oplus [3, 2, 1]$. Therefore, the real $1\hbar\omega_q$ excitations are: $[4, 2, 0] \oplus [4, 1, 1]$.

Table I. shows the model space of ^{20}Ne for the 0-1 major shells (both for the phenomenologic and for the semimicroscopic approach). Note here the small angular momentum content of the PAQM space (limited by the SU(3) quantum numbers).

The operators of physical quantities are obtained in this description in terms of the generators of the U(3) group. In particular the Hamiltonian can be expanded in terms of the generators of the U(3) group, coupled to spherical scalars. The general solution of the eigenvalue problem than involves two steps: *i*) calculation of matrix elements of the Hamiltonian between the basis states, and *ii*) numerical diagonalization of the energy matrix.

TABLE I. SU(3) quantum numbers of the states of ^{20}Ne for the 0 and 1 major shells in the phenomenologic and semimicroscopic algebraic quartet model. The superscripts indicate multiplicity.

model	$\hbar\omega$	SU(3)
PAQM	0	(2,0)
	1	(2,2),(3,0)
SAQM	0	(8,0),(4,2),(0,4),(2,0)
	1	(8,2),(9,0),(6,3),(7,1) ² ,(4,4),(5,2) ⁴ ,(2,5),(6,0) (3,3) ⁴ ,(1,4) ² ,(4,1) ³ ,(2,2) ⁴ ,(0,3) ² ,(3,0) ³ ,(1,1) ²




FIG. 1. The spectrum of the semimicroscopic algebraic quartet model in comparison with the experimental data of the ^{20}Ne nucleus. The experimental bands are labeled by the K^π , and the model states by the $n(\lambda, \mu)K^\pi$ quantum numbers. The spin-parity in parenthesis indicates uncertain band-assignment. The width of the arrow between the states is proportional to the strength of the $E2$ transition.

In the special case of the dynamical symmetry, i.e. when the Hamiltonian is expressed in terms of the invariant operators of the group-chain (1), an analytical solution is available.

The electromagnetic transition operators are obtained as Hermitian combinations of group generators with appropriate tensorial character. The lowest-order transition operators are:

$$\hat{T}^{(E0)} = e^{(0)}\hat{n}, \quad \hat{T}_m^{(E2)} = e^{(2)}\hat{Q}_m^{(2)}, \quad \hat{T}_m^{(M1)} = m^{(1)}\hat{L}_m^{(1)}. \quad (2)$$

These operators are diagonal in the SU(3) basis, i.e. they indicate no transitions between states of different SU(3)

irreps. This is a typical situation in the algebraic models in the dynamical symmetry limit. Transitions between e.g. different major shells can be obtained either i) by applying symmetry-breaking interactions, which mix the SU(3) basis states, or ii) by constructing more complex operators.

The PAQM states of Table I. correspond to very highly excited states (due to the large excitation quantum of the quartets: $(\hbar\omega)_q \approx 4(\hbar\omega)$). Therefore, it is hard to find a well-established correspondence between the experimental and model states. In case of the semimicroscopic description, on the other hand, it is much more straightforward.

The relation of the PAQM to the previous models is as follows. Its model space is identical with that of [9]. The main difference between the approach of [7, 9] and the present one is that in [7, 9] the interaction matrix elements are obtained empirically from the binding energies, while here we construct all the physical operators algebraically. This enables us to calculate the complete spectrum in an easy way.

The semimicroscopic algebraic quartet model. On the semimicroscopic level we take into account the composition of the quartets explicitly. They are considered [10] as 2 protons and 2 neutrons having permutational symmetry of [4], and spin-isospin symmetry of [1,1,1,1]. Therefore, in this case the nucleon shell model space is applied, and it is truncated according to these symmetries. Subsequent major shells have opposite parities.

The building blocks of this description are, again, the creation and annihilation operators of oscillator quanta (of the nucleon shell model).

The spectrum is determined by the U(3) spatial and the $U^{ST}(4)$ spin-isospin irreps. In this case, however, the particles are nucleons, not structureless quartets, as in the phenomenologic model. Therefore, the groups describe the symmetries of the many-nucleon-systems. Their relevant irreps are obtained in the following way. In each major shell those U(k) (or permutational) symmetries has to be taken into account which result in the required quartet symmetry $[4]^{N_q}$ when calculating their outer product with those of the other major shells. (Equivalently, those $U^{ST}(4)$ irreps are relevant, which result in the $[1, 1, 1, 1]^{N_q}$ quartet symmetry in the direct products.) These U(k) (or $U^{ST}(4)$) symmetries determine the relevant U(3) representations, and their direct products define the model space. The center of mass excitations can be removed in the same way, like in the previous case.

The model space is much richer than that of the phenomenological model, as shown by Table I. Especially remarkable is the angular momentum content of the model space; already in the lowest-lying major shell $L=8$ appears.

The physical operators are expressed in this case, too, in terms of the group-generators. Due to the restriction to the quartet symmetry only the scalar $U^{ST}(4)$ part of the spin-isospin sector gives contribution to the Hamil-

tonian and to the transition operators. Therefore, the formulae of the phenomenologic quartet model are valid here, too, but the oscillator quanta in this case refer to those of the nucleon shell model.

As an application we show here the result of the semimicroscopic model for the ^{20}Ne nucleus. The $U(3)$ dynamical symmetry approach is used; the interactions are written in terms of the invariant operators of group-chain (1), therefore, an analytical solution is available.

The experimental data are taken from [26], but for the band-assignment of the highly-excited alpha-cluster states also the conclusions of [27] are taken into account. The lower part of Figure 1. shows the states with definite band-assignment. All the bands with K^π values of [26] are included, except the one of the very uncertain (and somewhat contradictory) 0_7^+ band. In case of the 7^- state of the 0^- band, and the 6^+ and 8^+ states of the 0_6^+ band, which have more than one experimental candidates, the average energies are indicated. (In [26] there are only three states, which are not included here, for not having corresponding states in the model spectrum: a 6^+ state in the 0_2^+ band, a 9^- state in the 1^- band, and a 8^+ state in the 0_6^+ band. Each of them have uncertain band-assignment.)

We have tried a few phenomenological interactions, expressed in terms of the invariant operators of the $U(3) \supset SU(3) \supset SO(3)$ algebra-chain. Each of them contained a harmonic oscillator term (linear invariant of the $U(3)$), with a strenght obtained from the systematics [28] $\hbar\omega = 13.19$ MeV, and a rotational term with a parameter to fit. The remaining parts were written in terms of the second ($\hat{C}_{SU3}^{(2)}$) and third order ($\hat{C}_{SU3}^{(3)}$) invariant of the $SU(3)$. The former one accounts for the quadrupole-quadrupole interaction, and the latter one distinguishes between the prolate and oblate shapes. The simple linear combination $a\hat{C}_{SU3}^{(2)} + b\hat{C}_{SU3}^{(3)}$ contains two parameters. Another two-parameter term can be written as $\frac{g}{2c} \exp(-c\hat{C}_{SU3}^{(2)} - 1)$, which is very similar to the $\exp(-c\hat{Q}\hat{Q} - 1)$ term of the symplectic model Hamiltonian [29], accounting for a set of many-body interactions with well-defined relative weights. In order not to destroy the shell structure for the case of large excitations, the major-shell-average of the quadratic invariant ($\langle\hat{C}_{SU3}^{(2)}\rangle$) can be subtracted [30]. We have obtained the best description (from among the (2+1) parameter formulae) with

$$\hat{H} = (\hbar\omega)\hat{n} + a\hat{C}_{SU3}^{(2)} + (\hat{C}_{SU3}^{(2)} - \langle\hat{C}_{SU3}^{(2)}\rangle) + b\hat{C}_{SU3}^{(3)} + d\frac{1}{2\theta}\hat{L}^2, \quad (3)$$

where θ is the moment of inertia calculated classically for the rigid shape determined by the $U(3)$ quantum numbers (for a rotor with axial symmetry) [31]. Note that the third term does not introduce new fitting parameter, but a constant coefficient of 1 MeV is quietly understood here. The model spectrum of Figure 1 was obtained with the parameters: $a = -1,065$ MeV, $b = -0,000360$ MeV

$d = 0,808$ MeV.

We note here that the experimentally identified bands are described by the lowest-lying models bands with the appropriate spin-parity content, i.e. the other model bands of the same character are all higher-lying.

The intraband E2 transition rates were calculated with the operator of Eq. (4). The $B(E2)$ value is given by the formula [32]:

$$B(E2, I_i \rightarrow I_f) = \frac{2I_f + 1}{2I_i + 1} \alpha^2 |\langle(\lambda, \mu)KI_i, (11)2||(\lambda, \mu)KI_f\rangle|^2 C(\lambda, \mu), \quad (4)$$

where $\langle(\lambda, \mu)KI_i, (11)2||(\lambda, \mu)KI_f\rangle$ is the $SU(3) \supset SO(3)$ Wigner coefficient [33], and α is a parameter fitted to the the experimental value of the $2_1^+ \rightarrow 0_1^+$ transition of 20.3 W.u. The interband transition rate is zero.

The relation of the SAQM to the approach of [10] is similar to that between the PAQM and the previous models of [7, 9]. In particular, the two model spaces are identical, but the physical operators are not.

To sum up: In this paper we have introduced two algebraic models for the shell-like quarteting of nucleons. The simpler one is based on the quartet-concept of Arima et al. [7, 9], which does not treat explicitly the degrees of freedom of the constituent nucleons. Nevertheless, the Pauli-principle is not violated in this phenomenological description, either: the quartets of four nucleons occupy different single-particle space-states. The semimicroscopic model is more detailed. It is based on the definition of quartets in terms of two protons and two neutrons of [4] permutational symmetry [10]. This model is able to take into account 0, 1, 2, 3, 4, ... (nucleonic) major shell excitations, as opposed to the ‘‘giant’’ quartet excitations of the phenomenologic approach [7, 9], which correspond to $4q$, $q = 0, 1, 2, \dots$ nucleon excitation quanta. For both description the $U(3)$ formalism of Elliott [19, 20] is applied for the calculation of the spectrum. The semimicroscopic model is practically a symmetry-dictated truncation of the $L - S$ coupled no-core shell model, focusing on the spin-isospin-zero sector, and multiple excitations. It can be considered as an effective model in the sense of [34]: the bands of different quadrupole shapes are described by their lowest-grade $U(3)$ irreps without taking into account the giant-resonance excitations, built upon them, and the model parameters are renormalised for the subspace of the lowest $U(3)$ irreps.

From the viewpoint of their group-theoretical formalism these models are similar to the fully algebraic interacting-boson-like quartet models of the 1980'th [11, 12], but they are different concerning the physical nature of the quartets.

Both of these models are easy to apply, yet the semimicroscopic approach seems to be detailed enough to account for a considerable amount of the experimental spectrum, as illustrated by the application to the ^{20}Ne nucleus. We expect that in addition to its applicability to

the s-d shell nuclei it can also be extended to the mass region of $A=92-100$ of current experimental interest.

Further generalizations are possible by applying symmetry-breaking interactions, which result in e.g. nonvanishing interband transitions. Since the shell-truncations scheme of the semimicroscopic approach is based on the nucleonic degrees of freedom it could be exported also to non-alpha-like nuclei.

As for the *connection to other models*, the transparent symmetry properties of the present approach is very helpful. Via its obvious shell-model relation, the connection of the quartet model to the cluster and collective models is also well-defined (see e.g. [31, 35], and references therein for a recent discussion). In this respect the models with algebraic structure are relevant, in partic-

ular, the microscopic cluster model applying $U(3)$ basis [36, 37], and the semimicroscopic algebraic cluster model [38], as well as the symplectic shell model [39], and the contracted symplectic model [40] of the quadrupole collectivity.

Especially promising can be the application of the present semimicroscopic quartet model in combination with the concept of the multichannel dynamical symmetry [41], when the spectra of different cluster configurations are obtained from the quartet spectrum by simple projections.

This work was supported by the OTKA (Grant No K106035). The technical help of Mr. G. Riczu is kindly acknowledged.

-
- [1] C. Wu, Phys. Rev. Lett. 95 (2005) 266404.
 [2] S.K. Moskalenko *et al.*, in Bose-Einstein Condensation, edited by A. Griffin *et al.* Cambridge University Press, Cambridge, UK 1995.
 [3] B. Cederwall *et al.*, Nature 469 (2011) 68.
 [4] E.P. Wigner, Phys. Rev. 51 (1937) 106.
 [5] P. Van Isacker, D.D. Warner, D.S. Brenner, Phys. Rev. Lett. 74 (1995) 4607.
 [6] M. Danos, V. Gillet, Phys. Rev. 161 (1967) 1034.
 [7] A. Arima, V. Gillett, J. Ginocchio, Phys. Rev. Lett. 25 (1970) 1043.
 [8] A. Arima, V. Gillett, Ann Phys. 66 (1971) 117.
 [9] L. Satpathy, K.W. Schmid, A. Faessler, Phys. Rev. Lett. 28 (1972) 832.
 [10] M. Harvey, Nucl. Phys. A 202 (1973) 191.
 [11] J. Dukelsky, P. Federman, R.P.J. Perazzo, H.M. Sofia, Phys. Lett. 115B (1982) 359.
 [12] F. Iachello, A.D. Jackson, Phys. Lett. 108B (1982) 151.
 [13] F. Iachello, and A. Arima, The Interacting Boson Model, Cambridge University Press, Cambridge, 1987.
 [14] Y.K. Gambhir, P. Ring, P. Schuck, Phys. Rev. Lett. 51 (1983) 1235.
 [15] N. Sandulescu, D. Negrea, J. Dukelsky, C.W. Johnson, Phys. Rev. C 85 (2012) 061303(R).
 [16] M. Sambataro, N. Sandulescu, J. Phys. G 40 (2013) 055107.
 [17] A. Tohsaki, H. Horiuchi, P. Schuck, G. Ropke, Phys. Rev. Lett. 87 (2001) 192501.
 [18] H. Horiuchi, J. Phys. Conf. Ser. 436 (2013) 012003, and references therein.
 [19] J.P. Elliott, Proc. Roy. Soc. 245 (1958) 128; 562.
 [20] M. Harvey, Adv. Nucl. Phys. 1, 67 (1968).
 [21] T. Dytrych, K.D. Sviratcheva, J.P. Draayer, C. Bahri, and J.P. Vary, J. Phys. G 35 (2008) 123101.
 [22] J.P. Draayer, Y. Lebschber, S.C. Park and R. Lopez, Comput. Phys. Commun. 56 (1989) 279.
 [23] J.P. Elliott, T.H.K. Skyrme, Proc. Roy. Soc. A 232 (1955) 561.
 [24] D.M. Brink, G.F. Nash, Nucl. Phys. 40 (1963) 608.
 [25] K.T. Hecht, Nucl. Phys. A 170 (1971) 34.
 [26] D.R. Tilley *et al.*, Nucl. Phys. A 636 (1998) 249.
 [27] H.T. Richards, Phys. Rev. C 29 (1984) 276.
 [28] J. Blomqvist, A. Molinari, Nucl. Phys. A 106 (1968) 545.
 [29] G.K. Tobin *et al.*, Phys. Rev. C 89 (2014) 034312.
 [30] G. Rosensteel, J.P. Draayer, Nucl. Phys. A 436 (1985) 455.
 [31] J. Cseh, arXiv 1404.3500 [nucl-th] (2014).
 [32] Y. Sun *et al.*, Nucl. Phys. A 703 (2002) 130.
 [33] Y. Akiyama, J.P. Draayer, Comp. Phys. Comm. 5 (1973) 405.
 [34] D.J. Rowe, G. Thiamowa, J.L. Wood, Phys. Rev. Lett. 97 (2006) 202501.
 [35] J. Cseh, J. Darai, arXiv 1404.3503 [nucl-th] (2014).
 [36] H. Horiuchi, Prog. Theor. Phys. Suppl. 62 (1977) 90.
 [37] K.T. Hecht, E.J. Reske, T.H. Seligman, W. Zahn, Nucl. Phys. A 356 (1981) 146.
 [38] J. Cseh, Phys. Lett. B 281 (1992) 173 ; J. Cseh, G. Lévai, Ann. Phys. (NY) 230 (1994) 165.
 [39] G. Rosensteel, D.J. Rowe, Phys. Rev. Lett. 38 (1977) 10; Ann. Phys. (N.Y.) 126 (1980) 343; D.J. Rowe, Rep. Prog. Phys. 48 (1985) 1419.
 [40] D.J. Rowe, G. Rosensteel, Phys. Rev. C 25 (1982) 3236; O. Castanos, J.P. Draayer, Nucl. Phys. A 491 (1989) 349.
 [41] J. Cseh, Phys. Rev. C 50 (1994) 2240; J. Cseh, K. Katō, Phys. Rev. C 87 (2013) 067301.