

Existence of solutions for nonlinear impulsive higher order fractional differential equations*

Xuhuan Wang[†]

School of Mathematics and Computer Science, Guangxi University for Nationalities,

Nanning, Guangxi 530006, P.R. China

Abstract. In this paper, we prove the existence of solutions for nonlinear impulsive differential equations of fractional order $q \in (n - 1, n]$. The results are based on Altman's fixed point theorem and Leray-Schauder's fixed point theorem.

Keywords: Nonlinear fractional differential equations; Impulse; Initial value problem; Existence of Solutions; Fixed point theorem.

1 Introduction

In this paper, we consider the existence and uniqueness of solutions of the following impulsive problems:

$$\begin{cases} {}^C D_t^q u(t) = f(t, u(t)), n - 1 < q \leq n, t \in J_i = (t_i, t_{i+1}], \\ \Delta u^{(j)}(t_k) = I_{j,k}(u(t_k)) (j = 0, 1, \dots, n - 1), k = 1, 2, \dots, p, \\ u^{(j)}(0) = u_j (j = 0, 1, \dots, n - 1), \end{cases} \quad (1)$$

where ${}^C D^q$ is the Caputo fractional derivative, $f \in C(J \times R, R)$, $I_{j,k} \in C(R, R)$ ($j = 0, 1, \dots, n - 1$), $J = [0, 1]$, $0 = t_0 < t_1 < \dots < t_k < \dots < t_p < t_{p+1} = 1$, $J' = J \setminus \{t_1, t_2, \dots, t_p\}$, $\Delta u(t_k) = u(t_k^+) - u(t_k^-)$,

*The project is supported by NNSF of China, Grant No. 10971019 and Guangxi Key Laboratory of Hybrid Computation and IC Design Analysis.

[†]Email: wangxuhuan85@yahoo.cn

where $u(t_k^+)$ and $u(t_k^-)$ denote the right and the left limit of $u(t)$ at $t = t_k (k = 1, 2, \dots, p)$, respectively. $\Delta u^{(j)}(t_k)$ ($j = 1, 2, \dots, n - 1$) have a similar meaning for $u^{(j)}(t)$ ($j = 1, 2, \dots, n - 1$).

Recently, the subject of fractional differential equations has emerged as an important area of investigation. Indeed, we can find numerous applications in electrochemistry, control, electromagnetic, porous media, etc.[1-4]. Therefore, they have received much attention. For the most recent works for the existence and uniqueness of solutions of the initial and boundary value problems for nonlinear fractional differential equations, we mention [5-16, 25-30]. But, as far as we know, there have been few papers which have considered the multi-order fractional differential equations can be found in [18-24].

The organization of this paper is as follows: In Section 2, we give some basic definitions and properties. In Section 3, will be devoted to existence and uniqueness results for nonlinear impulsive differential equations of fractional order.

2 Preliminaries

Let us set $J_0 = [0, t_1]$, $J_1 = (t_1, t_2], \dots, J_{p-1} = (t_{p-1}, t_p]$, $J_p = (t_p, 1]$ and introduce the spaces:

$PC(J, R) = \{u : J \rightarrow R \mid u \in C(J_k), k = 0, 1, \dots, p, \text{ and } u(t_k^+) \text{ exist}, k = 1, 2, \dots, p\}$ with the norm $\|u\| = \sup_{t \in J} |u(t)|$, and

$PC^{n-1}(J, R) = \{u : J \rightarrow R \mid u \in C^{n-1}(J_k), k = 0, 1, \dots, p, \text{ and } u^{(j)}(t_k^+) \text{ exist}, j = 0, 1, \dots, n - 1, k = 1, 2, \dots, p\}$,

with the norm $\|u\|_{PC^{n-1}} = \max\{\|u\|, \|u'\|, \dots, \|u^{(n-1)}\|\}$. Obviously, $PC(J, R)$ and $PC^{n-1}(J, R)$ are Banach spaces.

Definition 2.1. A function $u \in PC^{n-1}(J, R)$ with its Caputo derivative of order q existing on J is a solution of (1) if it satisfies (1).

Theorem 2.1 ([17]). Let E be a Banach space. Assume that Ω is an open bounded subset of E with $\theta \in \Omega$ and let $T : \overline{\Omega} \rightarrow E$ be a completely continuous operator such that $\|Tu\| \leq \|u\|, \forall u \in \partial\Omega$. Then T has a fixed point in $\overline{\Omega}$.

Theorem 2.2 ([17]). Let E be a Banach space. Assume that $T : E \rightarrow E$ is a completely continuous operator and the set $V = \{u \in E \mid u = \mu Tu, 0 < \mu < 1\}$ is bounded. Then T has a fixed point in E .

Lemma 2.1 ([1,9]). For $q > 0$, the general solution of the fractional differential equation ${}^C D^q u(t) = 0$ is given by

$$u(t) = C_0 + C_1 t + C_2 t^2 + \dots + C_{n-1} t^{n-1}, C_i \in R, i = 0, 1, \dots, n - 1, n = [q] + 1,$$

where $[q]$ denotes the integer part of the real number q .

In view of Lemma 2.1, it follows that

$$I^q D^q u(t) = u(t) + C_0 + C_1 t + C_2 t^2 + \dots + C_{n-1} t^{n-1}, \quad (2)$$

for some $C_i \in R, i = 0, 1, \dots, n-1, n = [q] + 1$.

Lemma 2.2. For a given $y \in C[0, 1]$, the unique solution of the following impulsive boundary value problem

$$\begin{cases} {}^C D_{t_i}^q u(t) = y(t), n-1 < q \leq n, t \in J_i = (t_i, t_{i+1}], \\ \Delta u^{(j)}(t_k) = I_{j,k}(u(t_k)) (j = 0, 1, \dots, n-1), k = 1, 2, \dots, p, \\ u^{(j)}(0) = u_j (j = 0, 1, \dots, n-1), \end{cases} \quad (3)$$

is given by

$$u(t) = \begin{cases} \frac{1}{\Gamma(q)} \int_0^t (t-s)^{q-1} y(s) ds + \sum_{j=0}^{n-1} \frac{u_j}{\Gamma(j+1)} t^j, & t \in J_0; \\ \frac{1}{\Gamma(q)} \int_{t_k}^t (t-s)^{q-1} y(s) ds + \sum_{j=0}^{n-1} \frac{(t-t_k)^j}{\Gamma(j+1)\Gamma(q-j)} \int_{t_{k-1}}^{t_k} (t_k-s)^{q-j-1} y(s) ds \\ + \sum_{j=0}^{n-1} \sum_{i=1}^{k-1} \frac{[(t-t_k) + (t_k-t_i)]^j}{\Gamma(j+1)\Gamma(q-j)} \int_{t_{i-1}}^{t_i} (t_i-s)^{q-j-1} y(s) ds + \sum_{j=0}^{n-1} \frac{(t-t_k)^j}{\Gamma(j+1)} I_{j,k}(u(t_k)) \\ + \sum_{j=0}^{n-1} \sum_{i=1}^{k-1} \frac{[(t-t_k) + (t_k-t_i)]^j}{\Gamma(j+1)} I_{j,i}(u(t_i)) + \sum_{j=0}^{n-1} \frac{u_j}{\Gamma(j+1)} t^j, & t \in J_k, k = 1, 2, \dots, p. \end{cases} \quad (4)$$

Proof. Let u be a solution of (3). Then, by (2), we have

$$u(t) = I^q y(t) - \sum_{j=0}^{n-1} c_{j+1} t^j = \frac{1}{\Gamma(q)} \int_0^t (t-s)^{q-1} y(s) ds - \sum_{j=0}^{n-1} c_{j+1} t^j, \quad t \in J_0, \quad (5)$$

for some $c_1, c_2, \dots, c_n \in R$. Furthermore

$$\begin{aligned} u'(t) &= \frac{1}{\Gamma(q-1)} \int_0^t (t-s)^{q-2} y(s) ds - c_2 - 2c_3 t - 3c_4 t^2 - \dots - (n-1)c_n t^{n-2}, \quad t \in J_0, \\ u''(t) &= \frac{1}{\Gamma(q-2)} \int_0^t (t-s)^{q-3} y(s) ds - 2c_3 - 6c_4 t - \dots - (n-1)(n-2)c_n t^{n-3}, \quad t \in J_0, \\ &\dots \\ u^{(n-1)}(t) &= \frac{1}{\Gamma(q-n+1)} \int_0^t (t-s)^{q-n} y(s) ds - (n-1)!c_n, \quad t \in J_0, \end{aligned}$$

if $t \in J_1$, then

$$u(t) = \frac{1}{\Gamma(q)} \int_{t_1}^t (t-s)^{q-1} y(s) ds - \sum_{j=0}^{n-1} d_{j+1} (t-t_1)^j,$$

$$\begin{aligned}
u'(t) &= \frac{1}{\Gamma(q-1)} \int_{t_1}^t (t-s)^{q-2} y(s) ds - d_2 - 2d_3(t-t_1) - 3d_4(t-t_1)^2 - \dots - (n-1)d_n(t-t_1)^{n-2}, \\
u''(t) &= \frac{1}{\Gamma(q-2)} \int_{t_1}^t (t-s)^{q-3} y(s) ds - 2d_3 - 6d_4(t-t_1) - \dots - (n-1)(n-2)d_n(t-t_1)^{n-3}, \\
&\dots \\
u^{(n-1)}(t) &= \frac{1}{\Gamma(q-n+1)} \int_{t_1}^t (t-s)^{q-n} y(s) ds - (n-1)d_n,
\end{aligned}$$

for some $d_1, d_2, \dots, d_n \in R$. Thus we have

$$\begin{aligned}
u(t_1^-) &= \frac{1}{\Gamma(q)} \int_0^{t_1} (t_1-s)^{q-1} y(s) ds - c_1 - c_2 t_1 - c_3 t_1^2 - c_4 t_1^3 - \dots - c_n t_1^{n-1}, \quad u(t_1^+) = -d_1, \\
u'(t_1^-) &= \frac{1}{\Gamma(q-1)} \int_0^{t_1} (t_1-s)^{q-2} y(s) ds - c_2 - 2c_3 t_1 - 3c_4 t_1^2 - \dots - (n-1)c_n t_1^{n-2}, \quad u'(t_1^+) = -d_2, \\
u''(t_1^-) &= \frac{1}{\Gamma(q-2)} \int_0^{t_1} (t_1-s)^{q-3} y(s) ds - 2c_3 - 6c_4 t_1 - \dots - (n-1)(n-2)c_n t_1^{n-3}, \quad u''(t_1^+) = -2d_3, \\
&\dots \\
u^{(n-1)}(t_1^-) &= \frac{1}{\Gamma(q-n+1)} \int_0^{t_1} (t_1-s)^{q-n} y(s) ds - (n-1)c_n, \quad u^{(n-1)}(t_1^+) = -(n-1)d_n,
\end{aligned}$$

In view of $\Delta u(t_1) = u(t_1^+) - u(t_1^-) = I_{0,1}(u(t_1))$, $\Delta u'(t_1) = u'(t_1^+) - u'(t_1^-) = I_{1,1}(u(t_1))$, $\Delta u''(t_1) = u''(t_1^+) - u''(t_1^-) = I_{2,1}(u(t_1))$, \dots , $\Delta u^{(n-1)}(t_1) = u^{(n-1)}(t_1^+) - u^{(n-1)}(t_1^-) = I_{n-1,1}(u(t_1))$, we have

$$\begin{aligned}
-d_1 &= \frac{1}{\Gamma(q)} \int_0^{t_1} (t_1-s)^{q-1} y(s) ds - c_1 - c_2 t_1 - c_3 t_1^2 - c_4 t_1^3 - \dots - c_n t_1^{n-1} + I_{0,1}(u(t_1)), \\
-d_2 &= \frac{1}{\Gamma(q-1)} \int_0^{t_1} (t_1-s)^{q-2} y(s) ds - c_2 - 2c_3 t_1 - 3c_4 t_1^2 - \dots - (n-1)c_n t_1^{n-2} + I_{1,1}(u(t_1)), \\
-2d_3 &= \frac{1}{\Gamma(q-2)} \int_0^{t_1} (t_1-s)^{q-3} y(s) ds - 2c_3 - 6c_4 t_1 - \dots - (n-1)(n-2)c_n t_1^{n-3} + I_{2,1}(u(t_1)), \\
&\dots \\
-(n-1)d_n &= \frac{1}{\Gamma(q-n+1)} \int_0^{t_1} (t_1-s)^{q-n} y(s) ds - (n-1)c_n + I_{n-1,1}(u(t_1)).
\end{aligned}$$

Consequently

$$\begin{aligned}
u(t) &= \frac{1}{\Gamma(q)} \int_{t_1}^t (t-s)^{q-1} y(s) ds + \sum_{j=0}^{n-1} \frac{(t-t_1)^j}{\Gamma(j+1)\Gamma(q-j)} \int_0^{t_1} (t_1-s)^{q-j-1} y(s) ds \\
&\quad + \sum_{j=0}^{n-1} \frac{(t-t_1)^j}{\Gamma(j+1)} I_{j,1}(u(t_1)) - \sum_{j=0}^{n-1} c_{j+1} t^j, \quad t \in J_1.
\end{aligned}$$

Similarly, we get

$$\begin{aligned}
 u(t) &= \frac{1}{\Gamma(q)} \int_{t_k}^t (t-s)^{q-1} y(s) ds + \sum_{j=0}^{n-1} \frac{(t-t_k)^j}{\Gamma(j+1)\Gamma(q-j)} \int_{t_{k-1}}^{t_k} (t_k-s)^{q-j-1} y(s) ds \\
 &+ \sum_{j=0}^{n-1} \sum_{i=1}^{k-1} \frac{[(t-t_k) + (t_k-t_i)]^j}{\Gamma(j+1)\Gamma(q-j)} \int_{t_{i-1}}^{t_i} (t_i-s)^{q-j-1} y(s) ds + \sum_{j=0}^{n-1} \frac{(t-t_k)^j}{\Gamma(j+1)} I_{j,k}(u(t_k)) \\
 &+ \sum_{j=0}^{n-1} \sum_{i=1}^{k-1} \frac{[(t-t_k) + (t_k-t_i)]^j}{\Gamma(j+1)} I_{j,i}(u(t_i)) - \sum_{j=0}^{n-1} c_{j+1} t^j, \quad t \in J_k, \quad k = 1, 2, \dots, p. \quad (6)
 \end{aligned}$$

By $u^{(j)}(0) = u_j$, we have

$$c_{j+1} = \frac{-u_j}{\Gamma(j+1)}.$$

It follows that the solution given by (4) satisfies (3). This completes the proof.

3 Main Results

Define an operator $T : PC(J, R) \rightarrow PC(J, R)$ as

$$\begin{aligned}
 Tu(t) &= \frac{1}{\Gamma(q)} \int_{t_k}^t (t-s)^{q-1} f(s, u(s)) ds \\
 &+ \sum_{j=0}^{n-1} \frac{(t-t_k)^j}{\Gamma(j+1)\Gamma(q-j)} \int_{t_{k-1}}^{t_k} (t_k-s)^{q-j-1} f(s, u(s)) ds \\
 &+ \sum_{j=0}^{n-1} \sum_{i=1}^{k-1} \frac{[(t-t_k) + (t_k-t_i)]^j}{\Gamma(j+1)\Gamma(q-j)} \int_{t_{i-1}}^{t_i} (t_i-s)^{q-j-1} f(s, u(s)) ds \\
 &+ \sum_{j=0}^{n-1} \frac{(t-t_k)^j}{\Gamma(j+1)} I_{j,k}(u(t_k)) + \sum_{j=0}^{n-1} \sum_{i=1}^{k-1} \frac{[(t-t_k) + (t_k-t_i)]^j}{\Gamma(j+1)} I_{j,i}(u(t_i)) \\
 &+ \sum_{j=0}^{n-1} m_{j+1} t^j, \quad t \in J_k, \quad k = 1, 2, \dots, p. \quad (7)
 \end{aligned}$$

where $m_{j+1} = -c_{j+1}$ ($j = 0, 1, \dots, n-1$). Using Lemma 2.2 with $y(t) = f(t, u(t))$, problem (1) reduces to a fixed point problem $u = Tu$, where T is given by (7). Thus problem (1) has a solution if the operator T has a fixed point.

Theorem 3.1. Let $\lim_{u \rightarrow 0} \frac{f(t,u)}{u} = 0$, $\lim_{u \rightarrow 0} \frac{I_{j,k}(u)}{u} = 0$, ($j = 0, 1, \dots, n-1$) then problem (1) has at least one solution.

Proof. Firstly, we show that the operator $T : PC(J, R) \rightarrow PC(J, R)$ is completely continuous. Note that T is continuous in view of continuity of f and $I_{j,k}$. Let $\Omega \subset PC(J, R)$ be bounded. Then, there exist

positive constants $L_i > 0 (i = 1, 2, \dots, n+1)$ such that $|f(t, u)| \leq L_1$, $|I_{j,k}(u)| \leq L_{j+2}, (j = 0, 1, \dots, n-1)$, $\forall u \in \Omega$. Thus, $\forall u \in \Omega$, we have

$$|m_j| \leq \frac{u_j}{\Gamma(j+1)}.$$

Therefore

$$\begin{aligned} |Tu(t)| &= \frac{1}{\Gamma(q)} \int_{t_k}^t (t-s)^{q-1} |f(s, u(s))| ds \\ &+ \sum_{j=0}^{n-1} \frac{(t-t_k)^j}{\Gamma(j+1)\Gamma(q-j)} \int_{t_{k-1}}^{t_k} (t_k-s)^{q-j-1} |f(s, u(s))| ds \\ &+ \sum_{j=0}^{n-1} \sum_{i=1}^{k-1} \frac{[(t-t_k) + (t_k-t_i)]^j}{\Gamma(j+1)\Gamma(q-j)} \int_{t_{i-1}}^{t_i} (t_i-s)^{q-j-1} |f(s, u(s))| ds \\ &+ \sum_{j=0}^{n-1} \frac{(t-t_k)^j}{\Gamma(j+1)} |I_{j,k}(u(t_k))| + \sum_{j=0}^{n-1} \sum_{i=1}^{k-1} \frac{[(t-t_k) + (t_k-t_i)]^j}{\Gamma(j+1)} |I_{j,i}(u(t_i))| + \sum_{j=0}^{n-1} |m_{j+1}| \\ &\leq \frac{L_1}{\Gamma(q+1)} + \sum_{j=0}^{n-1} \frac{(t-t_k)^j L_1}{\Gamma(j+1)\Gamma(q-j+1)} + \sum_{j=0}^{n-1} \sum_{i=1}^{k-1} \frac{[(t-t_k) + (t_k-t_i)]^j L_1}{\Gamma(j+1)} \\ &+ \sum_{j=0}^{n-1} \sum_{i=1}^{k-1} \frac{[(t-t_k) + (t_k-t_i)]^j L_1}{\Gamma(q-j+1)\Gamma(j+1)} + \sum_{j=0}^{n-1} \frac{(t-t_k)^j L_j}{\Gamma(j+1)} + \sum_{j=0}^{n-1} \frac{u_j}{\Gamma(j+1)} \\ &\leq \frac{L_1}{\Gamma(q+1)} + \sum_{j=0}^{n-1} \frac{L_1}{\Gamma(j+1)\Gamma(q-j+1)} + \sum_{j=0}^{n-1} \frac{(p-1)L_1}{\Gamma(j+1)} \\ &+ \sum_{j=0}^{n-1} \frac{(p-1)L_1}{\Gamma(q-j+1)\Gamma(j+1)} + \sum_{j=0}^{n-1} \frac{L_{j+2}}{\Gamma(j+1)} + \sum_{j=0}^{n-1} \frac{u_j}{\Gamma(j+1)}, \end{aligned} \tag{8}$$

which implies that

$$\begin{aligned} \|Tu\| &\leq \frac{L_1}{\Gamma(q+1)} + \sum_{j=0}^{n-1} \frac{L_1}{\Gamma(j+1)\Gamma(q-j+1)} + \sum_{j=0}^{n-1} \frac{(p-1)L_1}{\Gamma(j+1)} \\ &+ \sum_{j=0}^{n-1} \frac{(p-1)L_1}{\Gamma(q-j+1)\Gamma(j+1)} + \sum_{j=0}^{n-1} \frac{L_{j+2}}{\Gamma(j+1)} + \sum_{j=0}^{n-1} \frac{u_j}{\Gamma(j+1)} := L. \end{aligned}$$

On the other hand, for any $t \in J_k, 0 \leq k \leq p$, we have

$$\begin{aligned} |Tu'(t)| &= \frac{1}{\Gamma(q-1)} \int_{t_k}^t (t-s)^{q-2} |f(s, u(s))| ds \\ &+ \sum_{j=0}^{n-1} \frac{(t-t_k)^{j-1}}{\Gamma(j)\Gamma(q-j)} \int_{t_{k-1}}^{t_k} (t_k-s)^{q-j-1} |f(s, u(s))| ds \\ &+ \sum_{j=0}^{n-1} \sum_{i=1}^{k-1} \frac{[(t-t_k) + (t_k-t_i)]^{j-1}}{\Gamma(j)\Gamma(q-j)} \int_{t_{i-1}}^{t_i} (t_i-s)^{q-j-1} |f(s, u(s))| ds \end{aligned}$$

$$\begin{aligned}
& + \sum_{j=0}^{n-1} \frac{(t-t_k)^j}{\Gamma(j)} |I_{j,k}(u(t_k))| + \sum_{j=0}^{n-1} \sum_{i=1}^{k-1} \frac{[(t-t_k) + (t_k-t_i)]^{j-1}}{\Gamma(j)} |I_{j,i}(u(t_i))| + \sum_{j=0}^{n-1} \frac{u_j}{\Gamma(j)} \\
\leq & \frac{L_1}{\Gamma(q)} + \sum_{j=0}^{n-1} \frac{L_1}{\Gamma(j)\Gamma(q-j+1)} + \sum_{j=0}^{n-1} \frac{(p-1)L_1}{\Gamma(j)} + \sum_{j=0}^{n-1} \frac{(p-1)L_1}{\Gamma(q-j+1)\Gamma(j)} \\
& + \sum_{j=0}^{n-1} \frac{L_{j+2}}{\Gamma(j)} + \sum_{j=0}^{n-1} \frac{u_j}{\Gamma(j)} := \bar{L}.
\end{aligned}$$

Hence, for $t_1, t_2 \in J_k$ with $t_1 < t_2$, $0 \leq k \leq p$, we have

$$|(Tu)(t_2) - (Tu)(t_1)| \leq \int_{t_1}^{t_2} |(Tu)'(s)| ds \leq \bar{L}(t_2 - t_1).$$

This implies that T is equicontinuous on all the subintervals J_k , $k = 0, 1, \dots, p$. Thus, by Arzela-Ascoli Theorem, it follows that $T : PC(J, R) \rightarrow PC(J, R)$ is completely continuous. Now, in view of $\lim_{u \rightarrow 0} \frac{f(t, u)}{u} = 0$, $\lim_{u \rightarrow 0} \frac{I_{j,k}(u)}{u} = 0$ ($j = 0, 1, \dots, n-1$), there exists a constant $r > 0$ such that $|f(t, u)| \leq \delta_1|u|$, $|I_{j,k}(u)| \leq \delta_{j+2}|u|$ ($j = 0, 1, \dots, n-1$) for $0 < |u| < r$, where $\delta_i > 0$ ($i = 1, 2, \dots, n+1$) satisfy

$$\begin{aligned}
& \frac{\delta_1}{\Gamma(q+1)} + \sum_{j=0}^{n-1} \frac{\delta_{j+2}}{\Gamma(j+1)\Gamma(q-j+1)} + \sum_{j=0}^{n-1} \frac{(p-1)\delta_{j+2}}{\Gamma(j+1)} + \sum_{j=0}^{n-1} \frac{(p-1)\delta_1}{\Gamma(q-j+1)\Gamma(j+1)} \\
& + \sum_{j=0}^{n-1} \frac{\delta_{j+2}}{\Gamma(j+1)} + \sum_{j=0}^{n-1} \frac{u_j}{\Gamma(j+1)} \leq 1.
\end{aligned} \tag{9}$$

Define $\Omega = \{u \in PC(J, R) \mid \|u\| < r\}$ and take $u \in PC(J, R)$ such that $\|u\| = r$ so that $u \in \partial\Omega$. Then, by the process used to obtain (8), we have

$$\begin{aligned}
|Tu(t)| \leq & \left\{ \frac{L_1}{\Gamma(q+1)} + \sum_{j=0}^{n-1} \frac{L_1}{\Gamma(j+1)\Gamma(q-j+1)} + \sum_{j=0}^{n-1} \frac{(p-1)L_1}{\Gamma(j+1)} \right. \\
& \left. + \sum_{j=0}^{n-1} \frac{(p-1)L_1}{\Gamma(q-j+1)\Gamma(j+1)} + \sum_{j=0}^{n-1} \frac{L_j}{\Gamma(j+1)} + \sum_{j=0}^{n-1} \frac{u_j}{\Gamma(j+1)} \right\} \|u\|.
\end{aligned} \tag{10}$$

which implies that $\|Tu\| \leq \|u\|$, $u \in \partial\Omega$. Therefore, by Theorem 2.1, the operator T has at least one fixed point, which implies that (1) has at least one solution $u \in \bar{\Omega}$.

Theorem 3.2. Assume that

(H_1) there exist positive constants L_i ($i = 1, 2, \dots, n+1$) such that $|f(t, u)| \leq L_1$, $|I_{j,k}(u)| \leq L_{j+2}$, ($j = 0, 1, \dots, n-1$) for $t \in J$, $u \in R$ and $k = 1, 2, \dots, p$. Then problem (1) has at least one solution.

Proof. As shown in Theorem 3.1, the operator $T : PC(J, R) \rightarrow PC(J, R)$ is completely continuous. Now, we show the set $V = \{u \in PC(J, R) \mid u = \mu Tu, 0 < \mu < 1\}$ is bounded. Let $u \in V$, then

$u = \mu Au, 0 < \mu < 1$. For any $t \in J$, we have

$$\begin{aligned}
 u(t) &= \frac{\mu}{\Gamma(q)} \int_{t_k}^t (t-s)^{q-1} f(s, u(s)) ds \\
 &+ \sum_{j=0}^{n-1} \frac{\mu(t-t_k)^j}{\Gamma(j+1)\Gamma(q-j)} \int_{t_{k-1}}^{t_k} (t_k-s)^{q-j-1} f(s, u(s)) ds \\
 &+ \sum_{j=0}^{n-1} \sum_{i=1}^{k-1} \frac{\mu[(t-t_k) + (t_k-t_i)]^j}{\Gamma(j+1)\Gamma(q-j)} \int_{t_{i-1}}^{t_i} (t_i-s)^{q-j-1} f(s, u(s)) ds \\
 &+ \sum_{j=0}^{n-1} \frac{\mu(t-t_k)^j}{\Gamma(j+1)} I_{j,k}(u(t_k)) + \sum_{j=0}^{n-1} \sum_{i=1}^{k-1} \frac{\mu[(t-t_k) + (t_k-t_i)]^j}{\Gamma(j+1)} I_{j,i}(u(t_i)) \\
 &+ \sum_{j=0}^{n-1} \mu m_{j+1} t^j,
 \end{aligned} \tag{11}$$

where $m_{j+1} = -c_{j+1} (j = 0, 1, \dots, n-1)$. Combining (H_1) and (11), we obtain

$$\begin{aligned}
 |u(t)| &= \mu |Tu(t)| \\
 &\leq \frac{1}{\Gamma(q)} \int_{t_k}^t (t-s)^{q-1} |f(s, u(s))| ds \\
 &+ \sum_{j=0}^{n-1} \frac{(t-t_k)^j}{\Gamma(j+1)\Gamma(q-j)} \int_{t_{k-1}}^{t_k} (t_k-s)^{q-j-1} |f(s, u(s))| ds \\
 &+ \sum_{j=0}^{n-1} \sum_{i=1}^{k-1} \frac{[(t-t_k) + (t_k-t_i)]^j}{\Gamma(j+1)\Gamma(q-j)} \int_{t_{i-1}}^{t_i} (t_i-s)^{q-j-1} |f(s, u(s))| ds \\
 &+ \sum_{j=0}^{n-1} \frac{(t-t_k)^j}{\Gamma(j+1)} |I_{j,k}(u(t_k))| + \sum_{j=0}^{n-1} \sum_{i=1}^{k-1} \frac{[(t-t_k) + (t_k-t_i)]^j}{\Gamma(j+1)} |I_{j,i}(u(t_i))| + \sum_{j=0}^{n-1} |m_{j+1}| \\
 &\leq \frac{L_1}{\Gamma(q+1)} + \sum_{j=0}^{n-1} \frac{L_1}{\Gamma(j+1)\Gamma(q-j+1)} + \sum_{j=0}^{n-1} \frac{(p-1)L_1}{\Gamma(j+1)} \\
 &+ \sum_{j=0}^{n-1} \frac{(p-1)L_1}{\Gamma(q-j+1)\Gamma(j+1)} + \sum_{j=0}^{n-1} \frac{L_{j+2}}{\Gamma(j+1)} + \sum_{j=0}^{n-1} \frac{u_j}{\Gamma(j+1)},
 \end{aligned}$$

Thus, for any $t \in J$, it follows that

$$\begin{aligned}
 \|u\| &\leq \frac{L_1}{\Gamma(q+1)} + \sum_{j=0}^{n-1} \frac{L_1}{\Gamma(j+1)\Gamma(q-j+1)} + \sum_{j=0}^{n-1} \frac{(p-1)L_1}{\Gamma(j+1)} \\
 &+ \sum_{j=0}^{n-1} \frac{(p-1)L_1}{\Gamma(q-j+1)\Gamma(j+1)} + \sum_{j=0}^{n-1} \frac{L_{j+2}}{\Gamma(j+1)} + \sum_{j=0}^{n-1} \frac{u_j}{\Gamma(j+1)}.
 \end{aligned}$$

So, the set V is bounded. Therefore, by the conclusion of Theorem 2.2, the operator T has at least one fixed point. This implies that (1) has at least one solution.

Theorem 3.3. Assume that

(H₂) there exist positive constants K_i ($i = 1, 2, \dots, n + 1$) such that $|f(t, u) - f(t, v)| \leq K_1|u - v|$, $|I_{j,k}(u) - I_{j,k}(v)| \leq K_{j+2}|u - v|$, ($j = 0, 1, \dots, n - 1$) for $t \in J, u, v \in R$ and $k = 1, 2, \dots, p$. Then problem (1) has a unique solution if

$$\begin{aligned} \Lambda &= \frac{L_1}{\Gamma(q+1)} + \sum_{j=0}^{n-1} \frac{L_1}{\Gamma(j+1)\Gamma(q-j+1)} + \sum_{j=0}^{n-1} \frac{(p-1)L_1}{\Gamma(j+1)} \\ &\quad + \sum_{j=0}^{n-1} \frac{(p-1)L_1}{\Gamma(q-j+1)\Gamma(j+1)} + \sum_{j=0}^{n-1} \frac{L_{j+2}}{\Gamma(j+1)} < 1. \end{aligned} \tag{12}$$

Proof. For $u, v \in C(J)$, we have

$$\begin{aligned} |(Tu)(t) - (Tv)(t)| &\leq \frac{1}{\Gamma(q)} \int_{t_k}^t (t-s)^{q-1} |f(s, u(s)) - f(s, v(s))| ds \\ &\quad + \sum_{j=0}^{n-1} \frac{(t-t_k)^j}{\Gamma(j+1)\Gamma(q-j)} \int_{t_{k-1}}^{t_k} (t_k-s)^{q-j-1} |f(s, u(s)) - f(s, v(s))| ds \\ &\quad + \sum_{j=0}^{n-1} \sum_{i=1}^{k-1} \frac{[(t-t_k) + (t_k-t_i)]^j}{\Gamma(j+1)\Gamma(q-j)} \int_{t_{i-1}}^{t_i} (t_i-s)^{q-j-1} |f(s, u(s)) - f(s, v(s))| ds \\ &\quad + \sum_{j=0}^{n-1} \frac{(t-t_k)^j}{\Gamma(j+1)} |I_{j,k}(u(t_k)) - I_{j,k}(v(t_k))| \\ &\quad + \sum_{j=0}^{n-1} \sum_{i=1}^{k-1} \frac{[(t-t_k) + (t_k-t_i)]^j}{\Gamma(j+1)} |I_{j,i}(u(t_i)) - I_{j,i}(v(t_i))| \\ &\leq \left\{ \frac{L_1}{\Gamma(q+1)} + \sum_{j=0}^{n-1} \frac{L_1}{\Gamma(j+1)\Gamma(q-j+1)} + \sum_{j=0}^{n-1} \frac{(p-1)L_1}{\Gamma(j+1)} \right. \\ &\quad \left. + \sum_{j=0}^{n-1} \frac{(p-1)L_1}{\Gamma(q-j+1)\Gamma(j+1)} + \sum_{j=0}^{n-1} \frac{L_{j+2}}{\Gamma(j+1)} \right\} \|u - v\| \\ &\leq \Lambda \|u - v\|. \end{aligned}$$

where Λ is given by (12). Thus, $\|Tu - Tv\| \leq \Lambda \|u - v\|$. As $\Lambda < 1$, therefore, T is a contraction operator. Hence, by the contraction mapping principle, problem (1) has a unique solution.

References

- [1] A.A. Kilbas, H.M. Srivastava, J.J. Trujillo, Theory and Applications of Fractional Differential Equations, Elsevier, Amsterdam, 2006.
- [2] S.G. Samko, A.A. Kilbas, O.I. Marichev, Fractional Integrals and Derivatives, Theory and Applications, Gordon and Breach, Yverdon, 1993.
- [3] I. Podlubny, Fractional Differential Equations, Academic Press, San Diego, 1999.
- [4] V. Lakshmikantham, S. Leela, J. Vasundhara Devi, Theory of Fractional Dynamic Systems, Cambridge Scientific Pub, Cambridge, UK, 2009.
- [5] R.P. Agarwal, M. Benchohra, S. Hamani, A Survey on Existence Results for Boundary Value Problems of Nonlinear Fractional Differential Equations and Inclusions, Acta. Appl. Math. 109 (2010) 973-1033.
- [6] C.F. Li, X.N. Luo, Y. Zhou, Existence of positive solutions of the boundary value problem for nonlinear fractional differential equations, Comput. Math. Appl. 59 (2010) 1363-1375.
- [7] V. Lakshmikantham, A.S. Vatsala, Basic theory of fractional differential equations, Nonlinear Anal. TMA. 69 (2008) 2677-2682.
- [8] V. Lakshmikantham, S. Leela, Nagumo-type uniqueness result for fractional differential equations, Nonlinear Anal. 8 (2009) 2886-2889.
- [9] Z.B. Bai, H.S. Lü, Positive solutions of boundary value problems of nonlinear fractional differential equation, J. Math. Anal. Appl. 311 (2005) 495-505.
- [10] B. Ahmad, J.J. Nieto, Existence results for a coupled system of nonlinear fractional differential equations with three-point boundary conditions, Comput. Math. Appl. 58 (2009) 1838-1843.
- [11] Y.K. Chang, J.J. Nieto, Some new existence results for fractional differential inclusions with boundary conditions, Math. Comput. Modelling. 49 (2009) 605-609.
- [12] M. Belmekki, J.J. Nieto, R. Rodriguez-Lopez, Existence of periodic solution for a nonlinear fractional differential equation, Bound. Value. Probl. 2009 (2009) 18 pages. Article ID 324561.
- [13] B. Ahmad, J.J. Nieto, Existence results for nonlinear boundary value problems of fractional integrodifferential equations with integral boundary conditions, Bound. Value. Probl. 2009 (2009) 11 pages. Article ID 708576.

- [14] K. Balachandran, S. Kiruthika, J.J. Trujillo, Existence results for fractional impulsive integrodifferential equations in Banach spaces, *Commun Nonlinear Sci Numer Simulat* 16 (2011) 1970-1977.
- [15] Z.H.Liu, Anti-periodic solutions to nonlinear evolution equations, *J. Funct. Anal.* 258(6)(2010) 2026-2033.
- [16] Y. Zhou, F. Jiao, Nonlocal Cauchy problem for fractional evolution equations, *Nonlinear Anal.* 11 (2010) 4465-4475.
- [17] J.X. Sun, *Nonlinear Functional Analysis and its Application*, Science Press, Beijing, 2008.
- [18] V. Daftardar-Gejji, H. Jafari, Solving a multi-order fractional differential equation using adomian decomposition, *Appl. Math. Comput.* 189 (2007) 541-548.
- [19] V.S. Erturk, S. Momani, Z. Odibat, Application of generalized differential transform method to multi-order fractional differential equations, *Communications in Nonlinear Science and Numerical Simulation.* 13 (2008) 1642-1654.
- [20] R. Lin, F. Liu, Fractional high order methods for the nonlinear fractional ordinary differential equation, *Nonlinear Anal.* 66 (2007) 856-869.
- [21] V. Daftardar-Gejji, H. Jafari, Analysis of a system of nonautonomous fractional differential equations involving Caputo derivatives, *J. Math. Anal. Appl.* 328 (2007) 1026-1033.
- [22] J.Q. Deng, L.F. Ma, Existence and uniqueness of solutions of initial value problems for nonlinear fractional differential equations, *Appl. Math. Lett.* 23 (2010) 676-680.
- [23] D. Bugajewska, P. Kasprzak, On the existence uniqueness and topological structure of solution sets to a certain fractional differential equation, *Comput. Math. Appl.* 59 (2010) 1108-1116.
- [24] B. Ahmad, J.J. Nieto, Existence of solutions for nonlocal boundary value problems of higher-order nonlinear fractional differential equations, *Abstr. Appl. Anal.* 2009, art. No. 494720.
- [25] J.J. Nieto, S. Abbas, M. Benchohra, Global uniqueness results for fractional order partial hyperbolic functional differential equations, *Adv. Difference Equ.* 2011, art. No. 379876.
- [26] Rui A.C. Ferreira, Nontrivial solutions for fractional q -difference boundary value problems, *E. J. Qualitative Theory of Diff. Equ.*, No. 70 (2010), 1-10.
- [27] B. Ahmad, J.J. Nieto, J. Pimentel, Some boundary value problems of fractional differential equations and inclusions, *Comput. Math. Appl.* Article in Press, doi:10.1016/j.camwa.2011.02.035.

- [28] L.H. Zhang, G.T. Wang, Existence of solutions for nonlinear fractional differential equations with impulses and anti-periodic boundary conditions, *E. J. Qualitative Theory of Diff. Equ.*, No. 7 (2011) 1-11.
- [29] J.J. Nieto, Maximum principles for fractional differential equations derived from Mittag-Leffler functions, *Appl. Math. Lett.* 23 (2010) 1248-1251.
- [30] X.J. Wang, C.Z. Bai, Periodic boundary value problems for nonlinear impulsive fractional differential equation, *E. J. Qualitative Theory of Diff. Equ.*, No. 3 (2011) 1-15.

(Received June 16, 2011)