

Data on the parasitoid complexes of *Metallus pumilus* (Hymenoptera: Tenthredinidae) and *Emmetia heinemanni* (Lepidoptera: Tischeriidae) mining leaves of *Rubus* sp.*

L. SZÖCS^{1**}, G. MELIKA², Cs. THURÓCZY³ & Gy. CSÓKA¹

¹NARIC Forest Research Institute, Department of Forest Protection, H-3232 Mátrafüred, Hungary.

E-mails: szocsl@erti.hu, csokagy@erti.hu

²Directorate of Plant Protection, Soil Conservation and Agri-environment, Budapest Plant Pest Diagnostic Laboratory, H-1118 Budapest, Budaörsi út 141–145, Hungary.

E-mail: melikag@nebih.gov.hu

³H-9730 Kőszeg, Malomárok utca 27, Hungary. E-mail: thuroczy.cs@freemail.hu

Abstract – 251 *Rubus* leaves mined by three species of leaf miners, *Metallus pumilus* (Klug, 1816), *Emmetia heinemanni* (Wocke, 1871) and *Ectoedemia rubivora* (Wocke, 1860), were collected between 2011 and 2014 in order to rear out their parasitoids. No parasitoids have been reared out from *Ectoedemia rubivora*, but we have identified 3 parasitoid species from *Emmetia heinemanni* and 5 species from *Metallus pumilus*. With 2 tables.

Key words – *Ectoedemia rubivora*, *Emmetia heinemanni*, *Metallus pumilus*, parasitoid complexes, *Rubus*

INTRODUCTION

The genus *Rubus* (Rosaceae) contains more than 700 species all over Europe (KIRÁLY *et al.* 2013). Some of them are significant weeds of forest clearings and agricultural fields. Although several leaf miners live on them, their parasitoid complexes are poorly studied. Here, we present preliminary data for the parasitoid complexes of two of these leaf mining species.

Emmetia (syn.: *Coptotriche*) *heinemanni* (Wocke, 1871) (Lepidoptera: Tischeriidae) is a bivoltine species (Szöcs 1977) with mines develop in June–July and September–October. The autumn mines can be confused with those of *Tischeria marginea* (Braun, 1972). *E. heinemanni* produces less silk in the mine, and

* The paper is dedicated to Dr László Móczár, doyen of the Hungarian hymenopterists, celebrating his 100th birthday.

** Corresponding author.

thus the mines are not contracted (HUSIMAN *et al.* 2013). Food plants are various *Rubus* (BUHR 1939, SZÓCS 1977, NIEUKERKEN 2006) and *Agrimonia* species (SZÓCS 1977, NIEUKERKEN 2006). The mine is an irregular blotch mine on the upper side of the leaf. A single larva develops inside it. Only one species of parasitoid (*Chrysocharis nephereus*) is mentioned from this host by ERDŐS (1956). The Universal Chalcidoidea Database (NOYES 2014) lists 6 parasitoid species associated with *T. heinemanni*: *Chrysocharis budensis* (Erdős, 1954), *Ch. nephereus* (Walker, 1839), *Ch. pentheus* (Walker, 1839), *Pediobius saulius* (Walker, 1839), *Pnigalio pectinicornis* (Linnaeus, 1758) and *Pteromalus semotus* (Walker, 1834).

The leaf mining sawfly *Metallus pumilus* (Klug, 1816) is oligophagous on *Rubus* species (BUHR 1939). It has two generations per year (ALTENHOFER *et al.* 2003), and the mines are present from June to September (ZOMBORI 1990), and even into October (personal observation by L. Szócs). The solitary larvae develop in the blotch mines on the upper surface of the leaves. The Universal Chalcidoidea Database (NOYES 2014) lists only one *Eulophus* species under the synonymous name *Metallus rubi* (Boie, 1848).

MATERIAL AND METHODS

We have studied the parasitoid complexes of various leaf miners during four years (2011–2014). A total of 251 leaf mines (137 *Emmetia*, 60 *Metallus* and 54 *Ectoedemia*) were collected from *Rubus idaeus* and *Rubus caesius* from seven locations across Hungary (Table 1). The mines were cut out from the leaf in order to avoid contamination from other parasitoids (i.e. egg or aphid parasitoids) which parasitise other insects on leaves. After a short period of drying, the samples were placed individually into air ventilated plastic rearing tubes. The parasitoids that emerged were preserved in 95% ethanol, and later identified by Csaba Thuróczy and George Melika. The emerging leaf miner adults were also identified to make sure that the identification of the leaf mines was correct. After the identification process, the adult insects were kept in alcohol for further molecular analysis.

RESULTS AND DISCUSSION

From *Emmetia heinemanni* samples, we have identified three parasitoid species belonging to the family Eulophidae. These are *Chrysocharis budensis*, *Ch. nephereus* and *Ch. pentheus*. They all are solitary koinobiont endoparasitoids (Table 2). The species reared out were different at each sampling locations. The parasitisation rate was 27.6% at Várgesztes (n = 58 mines). The only parasitoid species reared out from the Várgesztes samples was *Ch. budensis* (68 parasitoid specimens). The parasitisation rate was 21.4% at Csongrád (n = 14 mines). At

Table 1. Collecting localities and collected sample sizes of our rearings between 2011 and 2014

Host plant species	Leaf miner species	Sample date	Sample place							Total samples
			Kötegyán	Mátrafüred	Nógrád	Csongrád	Tiszabura	Várgesztes		
<i>Rubus caesius</i>	<i>Emmetia heinemanni</i>	14.VII.2011		1	55	14		9	58	137
	<i>Metallus pumilus</i>	27.IX.2012			1				10	11
<i>Rubus idaeus</i>	<i>Metallus pumilus</i>	21.X.2013	33					16		49
	<i>Ectoedemia rubivora</i>	10.VII.2012							54	54
Total samples		09.VII.2014	33	1	1	55	14	16	9	122
		14.VII.2011								251
		06.VII.2014								
		19.X.2013								

Csongrád, also only one species, *Ch. pentheus*, emerged from our samples. Only *Ch. nephereus* emerged from one single mine from Tiszabura. Due to the low sample size, we cannot calculate the parasitisation rate.

All three parasitoids, reared from *E. heinemanni* are listed in the Universal Chalcidoidea Database (NOYES 2014). With the exception of *Ch. budensis*, all

Table 2. Main biological characteristics and parasitoid assemblages of the two leaf miners from our rearings

Parasitoid species	Parasitoid biology	Leaf miner species	
		<i>Emmetia heinemanni</i>	<i>Metallus pumilus</i>
<i>Achrysocharoides cilla</i>	Gregarious Koinobiont Endoparasitoid		7
<i>Aprostocetus</i> sp.	Solitary ? ?		1
<i>Chrysocharis budensis</i>	Solitary Koinobiont Endoparasitoid	68	
<i>Chrysocharis nephereus</i>	Solitary Koinobiont Endoparasitoid	1	
<i>Chrysocharis pentheus</i>	Solitary Koinobiont Endoparasitoid	17	
<i>Pnigalio pectinicornis</i>	Solitary Idiobiont Ectoparasitoid		9
<i>Sympiesis sericeicornis</i>	Solitary Idiobiont Ectoparasitoid		1
<i>Itopectis alternans</i>	Solitary Koinobiont Endoparasitoid		2
Total parasitoid specimens		86	20

species have a broad range of hosts. According to the literature, *Ch. budensis* develops as a solitary parasitoid, but even so, many adults (up to seven) emerged from a single leaf mine. The reared parasitoid species seem to be semi-specialized to the Tenthredinidae, Nepticulidae and Tischeriidae leaf miners (NOYES 2014, SZÖCS *et al.* unpublished). They seem to prefer leaf mines situated on the upper surface of the leaves.

From the *Metallus pumilus* mines ($n = 60$) collected from two host plants (*R. caesius* and *R. idaeus*) we reared 20 specimens from five parasitoid species. Four of the emerged parasitoid species belonged to the family Eulophidae, and one to the family Ichneumonidae.

The dominant parasitoid in the samples was a secondary, solitary ectoparasitoid, *Pnigalio pectinicornis*. The subdominant species in the samples was the *Achrysocharoides cilla* (Walker, 1839), the only gregarious koinobiont endoparasitoid found on the sawfly, with seven emerged specimens.

We identified three parasitoid species in the samples from Kötegyán, which were *A. cilla*, *P. pectinicornis* and an ichneumonid wasp *Itopectis alternans* (Gravenhorst, 1829). The attack rate was 36.4% ($n = 33$), with 16 parasitoid specimens emerged. Three species of parasitoids (*P. pectinicornis*, *A. cilla* and *Aprostocetus* sp.) also emerged from the samples from Tiszabura.

At Várgesztes, only one specimen of *Sympiesis sericeicornis* emerged from the samples, and therefore the parasitisation rate cannot be calculated.

The parasitoid complex of *M. pumilus*, based on our rearings, is composed mainly of generalist species. The parasitoid species of the complex can be reared from other members of the family Tenthredinidae, especially from *Profenusa pygmaea* (Klug, 1816) (NOYES 2014, SCHÖNROGGE & ALTENHOFER 1992, SZÖCS *et al.* 2013).

No parasitoid has emerged from the *Ectoedemia rubivora* (Wocke, 1860) (Lepidoptera: Nepticulidae) sample (54 mines) collected at Várgesztes.

Despite the low sample size, we managed to identify three parasitoid species of *E. heinemanni* and five parasitoid species of *M. pumilus*. We have not found any overlap between the complexes of these two leaf miners. Even in the Universal Chalcidoidea Database, the two leaf miner species have very few common parasitoids, despite the fact that they regularly occur on the same host plant.

Both leaf miners have two generations and are abundant (locally very common), therefore, we assume that these species might have more species-rich parasitoid assemblages. Considering the relatively small sample sizes, it is conceivable that further rearings will help provide more information regarding their parasitoid fauna.

*

Acknowledgements – This project was supported by the OTKA 91238 research grant. We are grateful to Csaba Szabóky (Budapest) for his help in identification of the leaf miner adults.

REFERENCES

- ALTENHOFER E. & PSCHORN-WALCHER H. 2003: Biologische Notizen über die Blattwespen-Gattungen *Metallus* Forbes, *Monostegia* A. Costa und *Phymatocera* Dahlbom (Hymenoptera: Tenthredinidae). – *Linzer Biologische Beiträge* 35(1): 405–417. Online: http://www.landesmuseum.at/pdf_frei_remote/LBB_0035_1_0405-0417.pdf [Accessed 29 December 2014.]
- BUHR H. 1939: Beobachtungen über Nahrungspflanzen, Verbreitung und Auftreten von minierenden Blattwespen. – *Mitteilungen der Münchner Entomologische Gesellschaft* 31: 903–926. Online: http://www.landesmuseum.at/pdf_frei_remote/NachBlBayEnt_052_0060-0070.pdf [Accessed 29 December 2014.]
- ERDŐS J. 1956: Additamenta ad cognitionem faunae Chalcidoidarum in Hungaria et regionibus finitimis. VI. 19. Eulophidae. – *Folia entomologica hungarica (series nova)* 9: 29.
- HUSIMAN K. J., KOSTER J. G., MUUS T. S. T. & NIEUKERKEN E. J. VAN 2013: Microlepidoptera in Nederland, vooral in 2007–2010. – *Entomologische Berichten* 73(3): 91–117. Online: http://www.microlepidoptera.nl/library/pdf/Huisman_etal2013EB73.pdf [Accessed 29 December 2014.]
- KIRÁLY G., BOHUMIL T. & VOJTĚCH Z. 2013: A szeder (*Rubus* L.) nemzetség modern taxonómiai koncepciója. (Modern *Rubus* taxonomy.) – *Erdészettudományi Közlemények* 3(1): 147–156. Online: http://www.erdtudkoz.hu/fileadmin/dokumentumok/emk/erfaved/ErdTudKozl/Szamok/2013/Kiraly_es_mtsai_2013.pdf [Accessed 29 December 2014.]
- NIEUKERKEN E. J. VAN 2006: Records of mining Lepidoptera in Belgium with nine species new to the country (Nepticulidae, Opostegidae, Tischeriidae, Lyonetiidae). – *Phegea* 34(4): 125–144. Online: <http://www.repository.naturalis.nl/document/50443> [Accessed 29 December 2014.]
- NOYES J. S. 2014: *Universal Chalcidoidea Database*. – Online: <http://www.nhm.ac.uk/research-curation/research/projects/chalcidoids/database/> [Accessed 29 December 2014.]
- SCHÖNRÖGGE K. & ALTENHOFER E. 1992: On the biology and larval parasitoids of the leaf-mining sawflies *Profenusa thomsoni* (Konow) and *P. pygmaea* (Konow) (Hym.: Tenthredinidae). – *Entomologist's Monthly Magazine* 128: 99–108. Online: http://www.nhm.ac.uk/resources/research-curation/projects/chalcidoids/pdf_Y/SchonrA1992.pdf [Accessed 29 December 2014.]
- SZÓCS J. 1977: Lepidoptera-aknák és -gubacsok. Hyponomia et Cecidia Lepidopterorum. – In: *Magyarország Állatvilága (Fauna Hungariae), XIV, 16*. Akadémiai Kiadó, Budapest, 424 pp.
- ZOMBORI L. 1990: Levéldarázs-alakúak III. – Tenthredinoidea III. – In: *Magyarország Állatvilága (Fauna Hungariae), XI, 3/B*. Akadémiai Kiadó, Budapest, 85 pp.