



## Nonradial solutions for semilinear Schrödinger equations with sign-changing potential

Dingyang Lv and Xuxin Yang 

Department of Mathematics, Hunan First Normal University,  
Fenglin Road, No. 1015, Changsha, 410205, PR China

Received 9 February 2014, appeared 23 March 2015

Communicated by Vilmos Komornik

**Abstract.** In this paper, we investigate the existence of infinite nonradial solutions for the Schrödinger equations

$$\begin{cases} -\Delta u + b(|x|)u = f(|x|, u), & x \in \mathbb{R}^N, \\ u \in H^1(\mathbb{R}^N), \end{cases}$$

where  $b$  is allowed to be sign-changing. Under some assumptions on  $b \in C([0, \infty), \mathbb{R})$  and  $f \in C([0, \infty) \times \mathbb{R}^N, \mathbb{R})$ , we obtain that the above system possesses infinitely many nonradial solutions. The method of proof relies on critical point theorem.

**Keywords:** Schrödinger equations, variational methods, critical point, Sign-changing potential, nonradial solution.

**2010 Mathematics Subject Classification:** 35J20, 35J25.

### 1 Introduction and statement of the main result

In this paper, we study the existence of infinitely many nonradial solutions for the following semilinear Schrödinger equation

$$\begin{cases} -\Delta u + b(|x|)u = f(|x|, u), & x \in \mathbb{R}^N, \\ u \in H^1(\mathbb{R}^N). \end{cases} \quad (1.1)$$

We suppose that  $b: [0, \infty) \rightarrow \mathbb{R}$  and  $f: [0, \infty) \times \mathbb{R}^N \rightarrow \mathbb{R}$  satisfy the following assumptions:

(B<sub>1</sub>)  $b \in C([0, \infty), \mathbb{R})$  and  $\inf_{x \in \mathbb{R}^N} b(|x|) > -\infty$ ;

(B<sub>2</sub>) there exists a constant  $a > 0$  such that

$$\lim_{|y| \rightarrow +\infty} \text{meas} \left\{ x \in \mathbb{R}^N : |x - y| \leq a, b(|x|) \leq M \right\} = 0, \quad \forall M > 0,$$

where  $\text{meas}(\cdot)$  denotes the Lebesgue measure in  $\mathbb{R}^N$ ;

---

 Corresponding author. Email: yangxx2002@sohu.com

(B<sub>3</sub>)  $f \in C([0, \infty) \times \mathbb{R}^N, \mathbb{R})$  and there exist constants  $a_1, a_2 > 0$  and  $p \in (1, \frac{N+2}{N-2})$  such that

$$|f(x, u)| \leq a_1|u| + a_2|u|^p, \quad \forall (x, u) \in \mathbb{R}^N \times \mathbb{R}; \quad (1.2)$$

(B<sub>4</sub>) there exist  $\mu > 2$  and  $R > 0$  such that

$$0 < \mu F(r, u) := \mu \int_0^u f(r, v) dv \leq uf(r, u), \quad \text{for any } r \geq 0 \text{ and } |u| \geq R; \quad (1.3)$$

(B<sub>5</sub>)  $f(|x|, -u) = -f(|x|, u)$ ,  $\forall (x, u) \in \mathbb{R}^N \times \mathbb{R}$ .

We say that a solution  $u: \mathbb{R}^N \rightarrow \mathbb{R}$  is a *radial solution* (see for instance in [4, 7–9]) if  $u(x) = u(|x|)$ , that is, solution  $u$  has spherical symmetry. In the present paper, we consider the solutions of (1.1) which are different from the radial ones.

The following theorems are the main results of the paper.

**Theorem 1.1.** *Under assumptions (B<sub>1</sub>)–(B<sub>5</sub>), if  $N = 4$  or  $N \geq 6$ , then system (1.1) possesses an unbounded sequence of solutions  $\pm u_k$ ,  $k \in \mathbb{N}$ , which are not radial. The solutions are classical if  $f$  is locally Lipschitz with respect to  $u$ .*

Recently, by using variational methods and critical point theory, many authors have studied the existence of solution for system (1.1) or the following general type:

$$-\Delta u + b(x)u = f(x, u), \quad x \in \mathbb{R}^N. \quad (1.4)$$

The interest in equation (1.1) or (1.4) originates from various problems in physics and mathematical physics. In cosmology and constructive field theory, system (1.1) or (1.4) is also called *nonlinear Euclidean scalar field equation* (see [8, 9]). As it was mentioned in [4], a solution of (1.1) can also be interpreted as a *stationary state* (see [8, 9]) of the reaction diffusion:

$$u_t = -\Delta u - b(|x|)u + f(|x|, u),$$

for more physics background of (1.1), we refer the readers to [8, 9] and the references therein.

In [28], professor W. A. Strauss did pioneering work for the autonomous case of (1.1), that is:

$$-\Delta u = g(u), \quad x \in \mathbb{R}^N, \quad (1.5)$$

where  $g: \mathbb{R} \rightarrow \mathbb{R}$  is continuous and odd in  $u$ . In [8, 9], Berestycki and Lions obtained the existence of infinitely many radial solutions of (1.5) under almost necessary growth conditions on  $g$ . The solutions they obtained have exponential decay at infinity. When  $N = 1$ , they obtained a necessary and sufficient condition for the existence of a solution of problem (1.5). Some open problems are also mentioned in [8, 9]. For more results of radial solutions of (1.1) or (1.2), we refer the readers to [5, 7, 32]. For more applications of critical point theory to PDE, we refer the readers to the work of Michel Willem [32], Strauss [30], Rabinowitz [26], Zou [35] and T. Bartsch, Z. Q. Wang, M. Willem [7].

We are motivated by [4] written by T. Bartsch and Michel Willem. They make the following assumptions.

(A<sub>1</sub>)  $b \in C([0, \infty), \mathbb{R})$  is bounded from below by a positive constant  $a_0$ .

(A<sub>2</sub>)  $f \in C([0, \infty) \times \mathbb{R}, \mathbb{R})$  and there are positive constants  $a_1, R$  and a constant  $1 < q < \frac{N+2}{N-2}$  such that

$$|f(r, u)| \leq a_1|u|^q \quad \text{for any } r \geq 0, |u| \geq R.$$

(A<sub>3</sub>) There exists  $\mu > 2$  such that

$$\mu F(r, u) := \mu \int_0^u f(r, v) dv \leq u f(r, u), \quad \text{for any } r \geq 0, u \in \mathbb{R}. \quad (1.6)$$

(A<sub>4</sub>) There exists  $K > 0$  such that  $\inf_{r>0, |u|=K} F(r, u) > 0$ .

(A<sub>5</sub>)  $f(r, u) = o(|u|)$  for  $u \rightarrow 0$  uniformly in  $r \geq 0$ .

(A<sub>6</sub>)  $f$  is odd in  $u$ :  $f(r, -u) = -f(r, u)$  for any  $r \geq 0, u \in \mathbb{R}$ .

They state the following result.

**Theorem 1.2.** *Suppose  $N = 4$  or  $N \geq 6$ . If the assumptions (A<sub>1</sub>)–(A<sub>6</sub>) hold, then there exists an unbounded sequence of solutions  $\pm u_k, k \in \mathbb{N}$ , of (1.1) which are not radial. The solutions are classical if  $f$  is locally Lipschitz with respect to  $u$ .*

**Remark 1.3.** (1) As it is mentioned in [4] that solutions of (1.1) always occur in pairs because of the oddness of  $f$ .

(2) Compared with Theorem 1.2, our result allows  $b$  to be sign-changing.

(3) Assumption (1.6) is known as *global A–R condition* which was introduced by A. Ambrosetti and R. H. Rabinowitz (see for instance in [26]). It is obvious that the second part of assumption (1.3) is weaker than (1.6).

(4) In our result, assumption (A<sub>5</sub>) is not necessary.

In [4], (A<sub>5</sub>) together with (A<sub>3</sub>) plays a key role while discussing the functional  $\varphi$  (see later) corresponding to the system (1.1) satisfying the (P.S.)-condition (see [26,30,32,35]). If a function  $f \in C(\mathbb{R}^N \times \mathbb{R}, \mathbb{R})$  satisfies (A<sub>3</sub>) and (A<sub>5</sub>), then for any  $\varepsilon > 0$  (in application, we only concern about sufficiently small positive  $\varepsilon$ , that is  $0 < \varepsilon \ll 1$ ), there exists a finite  $C_\varepsilon = C(\varepsilon) > 0$  such that

$$|f(x, u)| \leq \varepsilon |u| + C_\varepsilon |u|^q. \quad (1.7)$$

Though in (1.7),  $C_\varepsilon$  may change for different  $\varepsilon > 0$ , but by (A<sub>3</sub>) and (A<sub>5</sub>) one can easily show that we can always assume that

$$C_\varepsilon < \infty, \quad \text{uniformly for any } \varepsilon > 0, x \in \mathbb{R}^N \text{ and } u \in \mathbb{R}. \quad (1.8)$$

That is,  $C_\varepsilon$  is independent of  $x \in \mathbb{R}^N$  and  $u \in \mathbb{R}$ . But under our assumptions in Theorem 1.1, (1.7) does not work any more, for example, let  $f(r, u) = f(u) = u + |u|^{p-1}u$  ( $1 < p < \frac{N+2}{N-2}$ ), then one can easily check that  $f$  satisfies the conditions (B<sub>3</sub>) and (B<sub>5</sub>) in our result. Now we are going to prove that  $f$  also satisfies condition (B<sub>4</sub>): firstly, we have  $F(r, u) = \frac{u^2}{2} + \frac{|u|^{p+1}}{p+1}$  and  $u f(r, u) = u^2 + |u|^{p+1}$ , choose some  $\mu \in (2, p+1)$ .

From

$$0 < \mu F(r, u) \leq u f(r, u),$$

that is,

$$\begin{aligned} \frac{\mu u^2}{2} + \frac{\mu |u|^{p+1}}{p+1} &\leq u^2 + |u|^{p+1}, \\ \Leftrightarrow \frac{\mu - 2}{2} u^2 &\leq \frac{p+1 - \mu}{p+1} |u|^{p+1}, \end{aligned}$$

$$\begin{aligned}
&\Leftrightarrow \frac{\mu - 2}{2} \leq \frac{p + 1 - \mu}{p + 1} |u|^{p-1}, \\
&\Leftrightarrow \frac{(\mu - 2)(p + 1)}{2(p + 1 - \mu)} \leq |u|^{p-1}, \\
&\Leftrightarrow \left[ \frac{(\mu - 2)(p + 1)}{2(p + 1 - \mu)} \right]^{\frac{1}{p-1}} \leq |u|.
\end{aligned}$$

Let  $R = \left[ \frac{(\mu - 2)(p + 1)}{2(p + 1 - \mu)} \right]^{\frac{1}{p-1}}$ , then we know that  $f$  satisfies condition  $(B_4)$ . But for any given  $\varepsilon_0 > 0$ , there does not exist a finite  $C_{\varepsilon_0} > 0$  such that

$$|f(|x|, u)| \leq \varepsilon_0 |u| + C_{\varepsilon_0} |u|^p, \quad \text{uniformly for any } \varepsilon > 0, x \in \mathbb{R}^N \text{ and } u \in \mathbb{R}. \quad (1.9)$$

If not, we assume that for some  $\varepsilon_0 > 0$  (without loss of generality, we suppose that  $0 < \varepsilon_0 < 1$ ), there exists some finite  $C = C_{\varepsilon_0} > 0$  such that

$$|f(|x|, u)| = |u| + |u|^p \leq \varepsilon_0 |u| + C_{\varepsilon_0} |u|^p. \quad (1.10)$$

By (1.10) we have  $C_{\varepsilon_0} > 1$  and

$$(1 - \varepsilon_0) |u| \leq (C_{\varepsilon_0} - 1) |u|^p,$$

this implies that

$$C_{\varepsilon_0} \geq \frac{1 - \varepsilon_0}{|u|^{p-1}} + 1 \rightarrow +\infty \quad \text{as } |u| \rightarrow 0.$$

This is obviously a contradiction. That is, in this case  $C_{\varepsilon_0}$  in (1.10) depends on  $u$ . Also, one can easily show that  $f$  does not satisfy  $(A_3)$  and  $(A_5)$ . As far as we know, while using the *fountain theorem* [32, 34] to discuss the existence of solutions of second order elliptic partial differential equations, many authors always assume that  $(A_5)$ , or similar type:  $f(x, u) = o(|u|)$  for  $u \rightarrow 0$  uniformly in  $x \in \mathbb{R}^N$  holds (see for instance in [12, 25, 32]).

Finally, we recall an abstract critical point lemma which we shall use later. Let  $X$  be a Banach space. We say that  $I \in C^1(X, \mathbb{R})$  satisfies  $(C)_c$ -condition (or weak-(P.S.)-condition [35]) if any sequence  $\{u_n\}$  such that

$$I(u_n) \rightarrow c, \quad \|I'(u_n)\| (1 + \|u_n\|) \rightarrow 0 \quad (1.11)$$

has a convergent subsequence.

**Lemma 1.4** ([3, 26]). *Let  $X$  be an infinite dimensional Banach space,  $X = Y \oplus Z$ , where  $Y$  is finite dimensional. If  $I \in C^1(X, \mathbb{R})$  satisfies  $(C)_c$ -condition for all  $c > 0$ , and*

$$(I_1) \quad I(0) = 0, I(-u) = I(u) \text{ for all } u \in X;$$

$$(I_2) \quad \text{there exist constants } \rho, \alpha > 0 \text{ such that } I|_{\partial B_\rho \cap Z} \geq \alpha;$$

$$(I_3) \quad \text{for any finite dimensional subspace } \tilde{X} \subset X, \text{ there is } R = R(\tilde{X}) > 0 \text{ such that } I(u) \leq 0 \text{ on } \tilde{X} \setminus B_R;$$

then  $I$  possesses an unbounded sequence of critical values.

## 2 Variational setting and proof of Theorem 1.1

Our proof is divided into a sequence of lemmas. Throughout this section, we make the following assumption instead of  $(B_1)$ .

$(B'_1)$   $b \in C(\mathbb{R}^N, \mathbb{R})$  and  $\inf_{\mathbb{R}^N} b(|x|) > 0$ .

We work in the Hilbert space

$$X = \left\{ u \in H^1(\mathbb{R}^N) : \int_{\mathbb{R}^N} (|\nabla u|^2 + b(|x|)u^2) dx < +\infty \right\}$$

equipped with the inner product

$$(u, v) = \int_{\mathbb{R}^N} (\nabla u \cdot \nabla v + b(|x|)uv) dx, \quad u, v \in X,$$

the associated norm

$$\|u\| = \left\{ \int_{\mathbb{R}^N} (|\nabla u|^2 + b(|x|)u^2) dx \right\}^{1/2}, \quad u \in X.$$

Evidently,  $C_0^\infty(\mathbb{R}^N, \mathbb{R}) \subset X$  and  $X$  is continuously embedded into  $H^1(\mathbb{R}^N)$  and hence continuously embedded into  $L^r(\mathbb{R}^N)$  for  $2 \leq r \leq 2^*$ , (where  $2^* = \frac{2N}{N-2}$  for  $N \geq 3$  and  $2^* = \infty$  for  $N = 1, 2$ ), i.e., there exists  $S_r > 0$  such that

$$\|u\|_r \leq S_r \|u\|, \quad \forall u \in X, \quad (2.1)$$

where  $\|\cdot\|_r$  denotes the usual norm in  $L^r(\mathbb{R}^N)$  for all  $2 \leq r \leq 2^*$ . In fact we further have the following lemma due to [7].

**Lemma 2.1** ([7, Lemma 3.1]). *Under assumptions  $(B'_1)$  and  $(B_2)$ , the embedding from  $X$  into  $L^s(\mathbb{R}^N)$  is compact for  $2 \leq s < 2^*$ .*

Now we define a functional  $\Phi$  on  $X$  by

$$\Phi(u) = \frac{1}{2} \int_{\mathbb{R}^N} (|\nabla u|^2 + b(|x|)u^2) dx - \int_{\mathbb{R}^N} F(|x|, u) dx \quad (2.2)$$

for all  $u \in X$ . Then it is well known that  $u \in X$  is a solution of (1.1) if and only if  $u$  is a critical point of  $\Phi$  in  $X$ . By assumption  $(B_3)$ , we have

$$|F(x, u)| \leq \frac{a_1}{2}|u|^2 + \frac{a_2}{p+1}|u|^{p+1}, \quad \forall (x, u) \in \mathbb{R}^N \times \mathbb{R}. \quad (2.3)$$

Consequently, under assumptions  $(B'_1)$ ,  $(B_2)$  and  $(B_3)$ , the functional  $\Phi$  is of class  $C^1(X, \mathbb{R})$ . Moreover, we have

$$\Phi(u) = \frac{1}{2} \|u\|^2 - \int_{\mathbb{R}^N} F(|x|, u) dx, \quad \forall u \in X, \quad (2.4)$$

$$\langle \Phi'(u), v \rangle = (u, v) - \int_{\mathbb{R}^N} f(|x|, u)v dx, \quad \forall u, v \in X. \quad (2.5)$$

By (2.3), for  $|u| < R$  ( $R$  is the same as in  $(B_4)$ ), we have

$$|f(|x|, u)u| + \mu|F(|x|, u)| \leq d|u|^2, \quad (2.6)$$

where  $d = \frac{2+\mu}{2}a_1 + \frac{p+1+\mu}{p+1}a_2R^{p-1}$ .

Now, we shall show that  $\Phi$  defined as (2.4) in  $X$  satisfies all the conditions in Lemma 1.4. By  $(B_5)$ , it is obvious that  $\Phi(0) = 0$  and  $\Phi(-u) = \Phi(u)$  for all  $u \in X$ . That is,  $(I_1)$  is satisfied. In order to prove that  $\Phi$  satisfies the  $(C)_c$ -condition, we firstly introduce an inequality (see for instance in [1]) which we will use later: if  $1 \leq p < \infty$  and  $a, b \geq 0$ , then

$$(a + b)^p \leq 2^{p-1}(a^p + b^p). \quad (2.7)$$

**Lemma 2.2.** *Under assumptions  $(B'_1)$  and  $(B_2)$ – $(B_4)$ , any sequence  $\{u_n\} \subset X$  satisfying*

$$\Phi(u_n) \rightarrow c > 0, \quad \langle \Phi'(u_n), u_n \rangle \rightarrow 0 \quad (2.8)$$

*is bounded in  $X$ . Moreover,  $\{u_n\}$  contains a convergent subsequence.*

*Proof.* Let  $\{u_n\} \subset X$  be a sequence satisfying (2.8), for the sake of discussion below, we introduce an auxiliary function  $\mathcal{F}(|x|, u) = f(|x|, u)u - \mu F(|x|, u)$  and  $\Omega_n = \{x \in \mathbb{R}^N : |u_n(x)| < R\}$  where  $R$  is the same as in  $(B_4)$ . By  $(B_4)$  and (2.6), without loss of generality, we may assume that for all  $n \in \mathbb{N}$ , we have:

$$\begin{aligned} c + 1 &\geq \Phi(u_n) - \frac{1}{\mu} \langle \Phi'(u_n), u_n \rangle \\ &= \frac{\mu - 2}{2\mu} \|u_n\|^2 + \frac{1}{\mu} \int_{\mathbb{R}^N} [f(|x|, u_n)u_n - \mu F(|x|, u_n)] dx \\ &= \frac{\mu - 2}{2\mu} \|u_n\|^2 + \frac{1}{\mu} \int_{\Omega_n} \mathcal{F}(|x|, u_n) dx + \frac{1}{\mu} \int_{\mathbb{R}^N \setminus \Omega_n} \mathcal{F}(|x|, u_n) dx \\ &\geq \frac{\mu - 2}{2\mu} \|u_n\|^2 + \frac{1}{\mu} \int_{\Omega_n} \mathcal{F}(|x|, u_n) dx \\ &\geq \frac{\mu - 2}{2\mu} \|u_n\|^2 - \frac{1}{\mu} \int_{\Omega_n} [f(|x|, u_n)u_n + \mu |F(|x|, u_n)|] dx \\ &\geq \frac{\mu - 2}{2\mu} \|u_n\|^2 - \frac{d}{\mu} \int_{\mathbb{R}^N} u_n^2 dx \\ &= \frac{\mu - 2}{2\mu} \|u_n\|^2 - \frac{d}{\mu} \|u_n\|_2^2. \end{aligned} \quad (2.9)$$

By (2.9), we have

$$\frac{\|u_n\|_2^2}{\|u_n\|^2} \geq \frac{\mu - 2}{2d} - \frac{\mu(c + 1)}{d\|u_n\|^2}.$$

So for sufficiently large  $\|u_n\|^2$  (actually we only require  $\|u_n\|^2 \geq \frac{4(c+1)\mu}{(\mu-2)}$ ),

$$\frac{\|u_n\|_2^2}{\|u_n\|^2} \geq \frac{\mu - 2}{4d} > 0. \quad (2.10)$$

If  $\{u_n\} \subset X$  is an unbounded sequence in  $X$ , passing to a subsequence if necessary, we may assume that  $\|u_n\| \rightarrow \infty$  as  $n \rightarrow \infty$ . Let  $v_n = \frac{u_n}{\|u_n\|}$ , then (2.10) implies that

$$\|v_n\|_2^2 > 0. \quad (2.11)$$

Let  $A_n = \{x \in \mathbb{R}^N : v_n \neq 0\}$ , then  $\text{meas}(A_n) > 0$ . Furthermore, under the assumption that  $\|u_n\| \rightarrow \infty$  as  $n \rightarrow \infty$ , we obtain

$$|u_n(x)| \rightarrow \infty \quad \text{as } n \rightarrow \infty \text{ for } x \in A_n. \quad (2.12)$$

Hence  $A_n \subseteq \mathbb{R}^N \setminus \Omega_n$  for sufficiently large  $n \in \mathbb{N}$ . By  $(B_4)$ , there exists some  $d_1 > 0$  such that

$$F(|x|, u) \geq d_1 |u|^\mu \quad \text{for } x \in \mathbb{R}^N \text{ and } |u| \geq R.$$

Hence by  $\mu > 2$ , we obtain

$$\lim_{|u| \rightarrow \infty} \frac{F(|x|, u)}{|u|^2} = +\infty. \quad (2.13)$$

By (2.1), (2.3), (2.4), (2.11), (2.13) and Fatou's lemma [21], for sufficiently large  $n \in \mathbb{N}$ , we have

$$\begin{aligned} 0 &= \lim_{n \rightarrow \infty} \frac{\Phi(u_n)}{\|u_n\|^2} \\ &= \frac{1}{2} - \int_{\mathbb{R}^N} \frac{F(|x|, u_n)}{u_n^2} v_n^2 dx \\ &= \frac{1}{2} - \int_{\Omega_n} \frac{F(|x|, u_n)}{u_n^2} v_n^2 dx - \int_{\mathbb{R}^N \setminus \Omega_n} \frac{F(|x|, u_n)}{u_n^2} v_n^2 dx \\ &\leq \frac{1}{2} + \left( \frac{a_1}{2} + \frac{a_2}{p+1} R^{p-1} \right) S_2 - \liminf_{n \rightarrow \infty} \int_{\mathbb{R}^N \setminus \Omega_n} \frac{F(|x|, u_n)}{u_n^2} v_n^2 dx \\ &\leq \frac{1}{2} + \left( \frac{a_1}{2} + \frac{a_2}{p+1} R^{p-1} \right) S_2 - \int_{\mathbb{R}^N \setminus \Omega_n} \liminf_{n \rightarrow \infty} \frac{F(|x|, u_n)}{u_n^2} v_n^2 dx \\ &\leq \frac{1}{2} + \left( \frac{a_1}{2} + \frac{a_2}{p+1} R^{p-1} \right) S_2 - \int_{A_n} \liminf_{n \rightarrow \infty} \frac{F(|x|, u_n)}{u_n^2} [\chi_{A_n}(x)] v_n^2 dx \\ &\rightarrow -\infty, \quad \text{as } n \rightarrow \infty. \end{aligned} \quad (2.14)$$

This is an obvious contradiction. Hence  $\{u_n\} \subset X$  is bounded.

Now we shall prove  $\{u_n\}$  contains a convergent subsequence. Without loss of generality, by the Eberlein–Shmulyan theorem (see for instance in [33]), passing to a subsequence if necessary, there exists a  $u \in X$  such that  $u_n \rightharpoonup u$  in  $X$ . Again by Lemma 2.1,  $u_n \rightarrow u$  in  $L^r(\mathbb{R}^N)$  for  $2 \leq r < 2^*$ , that is

$$\|u_n - u\|_r \rightarrow 0 \quad \text{as } n \rightarrow \infty, \quad (2.15)$$

and  $u_n \rightarrow u$  a.e.  $x \in \mathbb{R}^N$ . Observe that

$$\|u_n - u\|^2 = \langle \Phi'(u_n) - \Phi'(u), u_n - u \rangle + \int_{\mathbb{R}^N} [f(|x|, u_n) - f(|x|, u)](u_n - u) dx. \quad (2.16)$$

It is clear that

$$\langle \Phi'(u_n) - \Phi'(u), u_n - u \rangle \rightarrow 0, \quad n \rightarrow \infty. \quad (2.17)$$

By  $(B_3)$ , (2.7), (2.15), Hölder's inequality and the fact  $2 < p+1 < 2^*$ ,

$$\begin{aligned} &\int_{\mathbb{R}^N} |(f(|x|, u_n) - f(|x|, u))(u_n - u)| dx \\ &\leq \int_{\mathbb{R}^N} (|f(|x|, u_n)| + |f(|x|, u)|) |u_n - u| dx \\ &\leq \int_{\mathbb{R}^N} [a_1(|u_n| + |u|) + a_2(|u_n|^p + |u|^p)] |u_n - u| dx \\ &\leq \int_{\mathbb{R}^N} [2a_1(|u_n - u| + |u|) + 2^p a_2(|u_n - u|^p + |u|^p)] |u_n - u| dx \\ &= \int_{\mathbb{R}^N} [2a_1(|u_n - u|^2 + |u| |u_n - u|) + 2^p a_2(|u_n - u|^{p+1} + |u|^p |u_n - u|)] dx \\ &\leq 2a_1(\|u_n - u\|_2^2 + \|u\|_2 \|u_n - u\|_2) + 2^p a_2 \left( \|u_n - u\|_{p+1}^{p+1} + \|u\|_{p+1}^{\frac{p}{p+1}} \|u_n - u\|_{p+1} \right) \\ &\rightarrow 0, \quad \text{as } n \rightarrow \infty. \end{aligned} \quad (2.18)$$

This together with (2.16), (2.17) implies  $u_n \rightarrow u$  in  $X$  as  $n \rightarrow \infty$ .  $\square$

Let  $\{e_j\}$  be an orthonormal basis of  $X$  and define  $X_j = \mathbb{R}e_j$ ,

$$Y_k = \oplus_{j=1}^k X_j, \quad Z_k = \oplus_{j=k+1}^{\infty} X_j, \quad k \in \mathbb{Z}. \quad (2.19)$$

**Lemma 2.3.** *Under assumptions  $(B'_1)$  and  $(B_2)$ , for  $2 \leq r < 2^*$ , we have*

$$\beta_k(s) := \sup_{u \in Z_k, \|u\|=1} \|u\|_s \rightarrow 0, \quad k \rightarrow \infty. \quad (2.20)$$

*Proof.* By Lemma 2.1,  $X \hookrightarrow L^r(\mathbb{R}^N)$  is compact, then Lemma 2.3 can be proved by a similar way as Lemma 3.8 in [28] or Lemma 3.3 in [4].  $\square$

By Lemma 2.3, we can choose an integer  $m \geq 1$  such that

$$\|u\|_2^2 \leq \frac{1}{2a_1} \|u\|^2, \quad \|u\|_{p+1}^{p+1} \leq \frac{p+1}{4a_2} \|u\|^{p+1}, \quad \forall u \in Z_m. \quad (2.21)$$

**Lemma 2.4.** *Under the assumptions  $(B'_1)$ ,  $(B_2)$  and  $(B_3)$ , there exist constants  $\rho, \alpha > 0$  such that  $\Phi|_{\partial B_\rho \cap Z_m} \geq \alpha$ .*

*Proof.* By (2.4) and (2.21), we have

$$\begin{aligned} \Phi(u) &= \frac{1}{2} \|u\|^2 - \int_{\mathbb{R}^N} F(x, u) dx \\ &\geq \frac{1}{2} \|u\|^2 - \frac{a_1}{2} \|u\|_2^2 - \frac{a_2}{p+1} \|u\|_{p+1}^{p+1} \\ &\geq \frac{1}{4} (\|u\|^2 - \|u\|^{p+1}). \end{aligned} \quad (2.22)$$

Let  $0 < \rho < 1$ , then  $\alpha = \frac{1}{4}(\rho^2 - \rho^{p+1}) > 0$  satisfies the conditions of the lemma.  $\square$

**Lemma 2.5.** *Under assumptions  $(B'_1)$ ,  $(B_2)$ – $(B_4)$ , for any finite dimensional subspace  $\tilde{X} \subset X$ , there holds*

$$\Phi(u) \rightarrow -\infty, \quad \|u\| \rightarrow \infty, \quad u \in \tilde{X}. \quad (2.23)$$

*Proof.* Arguing indirectly, assume that there exists a sequence  $\{u_n\} \subset \tilde{X}$  with  $\|u_n\| \rightarrow \infty$  and  $M > 0$  such that  $\Phi(u_n) \geq -M$  for all  $n \in \mathbb{N}$ . Set  $v_n = u_n / \|u_n\|$ , then  $\|v_n\| = 1$ . Passing to a subsequence, we may assume that  $v_n \rightharpoonup v$  in  $X$ . Since  $\tilde{X}$  is finite dimensional, then  $v_n \rightarrow v \in \tilde{X}$  in  $X$ ,  $v_n \rightarrow v$  a.e. on  $\mathbb{R}^N$ , and so  $\|v\| = 1$ . Hence, we can conclude a contradiction by a similar fashion as (2.14).  $\square$

By  $(B_1)$ , there exists a constant  $b_0 > 0$  such that  $\bar{b}(|x|) := b(|x|) + b_0 \geq 1$  for all  $x \in \mathbb{R}^N$ . Let  $\bar{f}(|x|, u) = f(|x|, u) + b_0 u$ . Then  $\bar{b}$  and  $\bar{f}$  satisfy  $(B'_1)$ ,  $(B_2)$ – $(B_5)$  and it is also easy to verify the following lemma.

**Lemma 2.6.** *Problem (1.1) is equivalent to the following problem*

$$\begin{cases} -\Delta u + \bar{b}(|x|)u = \bar{f}(|x|, u), & x \in \mathbb{R}^N, \\ u \in H^1(\mathbb{R}^N). \end{cases} \quad (2.24)$$

At last, to complete the proof of Theorem 1.1, we need the following result (see [4]).



**Lemma 2.7.** *Let  $G$  be a group acting on  $X$  via orthogonal maps  $\rho(g): X \rightarrow X$  and such that the following hold.*

- (i)  $\Phi: X \rightarrow \mathbb{R}$  is  $G$ -invariant.
- (ii) The inclusion  $X^G \hookrightarrow L^s(\mathbb{R}^N)$  is compact for every  $s \in (2, 2^*)$ .
- (iii)  $\dim X^G = \infty$ .

Here  $X^G = \{u \in X : \rho(g)u = u \text{ for all } g \in G\}$  is the  $G$ -fixed point set. Then  $\Phi$  has unbounded sequence of critical values with associated critical points lying in  $X^G$ .

*Proof of Theorem 1.1.* Firstly we shall find a group  $G$  and an action of  $G$  on  $X$  which satisfies the assumptions of Lemma 2.7. We should point out the main idea of the discussion below due to the work of T. Bartsch and M. Willem in [4].  $G \subset O(N)$  is defined as follows. Choose an integer  $2 \leq m \leq \frac{N}{2}$  satisfying  $2m \neq N - 1$ . This always holds for  $N = 4$  or  $N \geq 6$ . The action of

$$H = O(m) \times O(m) \times O(N - 2m)$$

on  $X$  is defined by

$$gu(x) = u(g^{-1}x).$$

Let  $\tau$  be the involution defined on  $\mathbb{R}^N = \mathbb{R}^m \oplus \mathbb{R}^m \oplus \mathbb{R}^{N-2m}$  by

$$\tau(x_1, x_2, x_3) \doteq (x_2, x_1, x_3),$$

where  $(x_1, x_2, x_3) \in \mathbb{R}^m \oplus \mathbb{R}^m \oplus \mathbb{R}^{N-2m}$ . Let  $G = \langle H \cup \{\tau\} \rangle \subset O(N)$ . Then elements of  $G$  can be represented uniquely as  $h$  or  $h\tau$  with  $h \in H$  and the action of  $G$  on  $X$  defined as

$$\begin{aligned} \rho(g)u(x) &:= hu(x) = u(h^{-1}x), & g = h \in H, \\ &:= -\tau u(x) = -u(\tau x), & g = h\tau. \end{aligned}$$

Then it is clear that 0 is the only radial function in  $X^G$ . By the work of T. Bartsh and M. Willem in [4] (also see in [32]) we know that  $G$  and the action  $\rho(g)$  satisfy all the assumptions in Lemma 2.7. Thus we obtain an unbounded sequence of critical values  $c_k$  of  $\Phi: X \rightarrow \mathbb{R}$ . By Lemma 2.7, we know the associated critical points  $u_k$  lie in  $X^G$ , from discussion above we know that  $u_k$  are of nonradial solutions of (2.24). By Lemma 2.6, we know that  $u_k$  are also of nonradial solutions of (1.1). When  $f$  is locally Lipschitz with respect to  $u$ , by [14] we know that  $u_k$  are classical.  $\square$

## Acknowledgement

This work is supported by Scientific Research Fund of Hunan Provincial Education Department (14C0253), Hunan Provincial Natural Science Foundation of China (14JJ7083) and a Key Project Supported by Scientific Research Fund of Hunan Provincial Education Department (14A028).

## References

- [1] R. A. ADAMS, J. J. F. FOURNIER, *Sobolev spaces*, second edition, Academic Press, Amsterdam, 2003. [MR2424078](#)
- [2] A. AMBROSETTI, P. H. RABINOWITZ, Dual variational methods in critical point theory and applications, *J. Funct. Anal.* **14**(1973), 349–381. [MR0370183](#)
- [3] T. BARTOLO, V. BENCI, D. FORTUNATO, Abstract critical point theorems and applications to some nonlinear problems with strong resonance at infinity, *Nonlinear Anal.* **7**(1983), 981–1012. [MR713209](#); [url](#)
- [4] T. BARTSCH, M. WILLEM, Infinitely many nonradial solutions of a Euclidean scalar field equation, *J. Funct. Anal.* **117**(1993), 447–460. [MR1244943](#); [url](#)
- [5] T. BARTSCH, M. WILLEM, Infinitely many radial solutions of a semilinear elliptic problem on  $\mathbb{R}^N$ , *Arch. Rational Mech. Anal.* **124**(1993), 261–276. [MR1237913](#); [url](#)
- [6] T. BARTSCH, Z. Q. WANG, Existence and multiplicity results for some superlinear elliptic problems on  $\mathbb{R}^N$ , *Comm. Partial Differential Equations* **20**(1995), 1725–1741. [MR1349229](#); [url](#)
- [7] T. BARTSCH, Z. Q. WANG, M. WILLEM, The Dirichlet problem for superlinear elliptic equations, in: M. Chipot, P. Quittner (Eds.), *Stationary partial differential equations*, Vol. 2, Handbook of Differential Equations, Elsevier, 2005, pp. 1–55 (Chapter 1). [MR2181481](#); [url](#)
- [8] H. BERESTYCKI, P. L. LIONS, Nonlinear scalar field equations I. Existence of a ground state, *Arch. Rational Mech. Anal.* **82**(1983), 313–345. [MR695535](#); [url](#)
- [9] H. BERESTYCKI, P. L. LIONS, Nonlinear scalar field equations II. Existence of infinitely many solutions, *Arch. Rational Mech. Anal.* **82**(1983), 347–375. [MR695536](#); [url](#)
- [10] H. BRÉZIS, L. NIRENBERG, Positive solutions of nonlinear elliptic equations involving critical Sobolev exponents, *Comm. Pure Appl. Math.* **36**(1983), 437–477. [MR709644](#); [url](#)
- [11] V. COTI ZELATI, P. H. RABINOWITZ, Homoclinic type solutions for a semilinear elliptic PDE on  $\mathbb{R}^N$ , *Comm. Pure Appl. Math.* **45**(1992), 1217–1269. [MR1181725](#); [url](#)
- [12] W. Y. DING, W.-M. NI, On the existence of positive entire solutions of a semilinear elliptic equation, *Arch. Rational Mech. Anal.* **91**(1986), 183–208. [MR807816](#); [url](#)
- [13] Y. DING, C. LEE, Multiple solutions of Schrödinger equations with indefinite linear part and super or asymptotically linear terms, *J. Differential Equations* **222**(2006), 137–163. [MR2200749](#); [url](#)
- [14] L. C. EVANS, *Partial differential equations*, Graduate Studies in Mathematics, Vol. 19, American Mathematical Society, Providence, RI, 1998. [MR1625845](#)
- [15] L. JEANJEAN, On the existence of bounded Palais–Smale sequences and application to a Landesman–Lazer type problem set on  $\mathbb{R}^N$ , *Proc. Roy. Soc. Edinburgh Sect. A* **129**(1999), 787–809. [MR1718530](#); [url](#)
- [16] L. JEANJEAN, K. TANAKA, Singularly perturbed elliptic problems with superlinear or asymptotically linear nonlinearities, *Calc. Var. Partial Differential Equations* **21**(2004), 287–318. [MR2094325](#); [url](#)

- [17] A. KRISTÁLY, Multiple solutions of a sublinear Schrödinger equation, *NoDEA Nonlinear Differential Equations Appl.* **14**(2007), 291–301. [MR2364894](#); [url](#)
- [18] G. B. LI, A. SZULKIN, An asymptotically periodic Schrödinger equation with indefinite linear part, *Commun. Contemp. Math.* **4**(2002), 763–776. [MR1938493](#); [url](#)
- [19] Y. LI, Remarks on a semilinear elliptic equation on  $\mathbb{R}^N$ , *J. Differential Equations* **74**(1988), 34–49. [MR949624](#); [url](#)
- [20] Y. Y. LI, Nonautonomous nonlinear scalar field equations, *Indiana Univ. Math. J.* **39**(1990), 283–301.
- [21] E. H. LIEB, M. LOSS, *Analysis*, Graduate Studies in Mathematics, Vol. 14, American Mathematical Society, Providence, RI, 1997. [MR1415616](#)
- [22] S. B. LIU, S. J. LI, Infinitely many solutions for a superlinear elliptic equation, *Acta Math. Sinica (Chin. Ser.)* **46**(2003), 625–630. [MR2010750](#)
- [23] D. MOSCHETTO, Existence and multiplicity results for a nonlinear stationary Schrödinger equation, *Ann. Polon. Math.* **99**(2010), 39–43. [MR2660590](#); [url](#)
- [24] K. PERERA, Nontrivial critical groups in  $p$ -Laplacian problems via the Yang index, *Topol. Methods Nonlinear Anal.* **21**(2003), 301–309. [MR1998432](#)
- [25] Q. LI, H. SU, Z. WEI, Existence of infinitely many large solutions for the nonlinear Schrödinger–Maxwell equations, *Nonlinear Anal.* **72**(2010), 4264–4270. [MR2606783](#); [url](#)
- [26] P. H. RABINOWITZ, Minimax methods in critical point theory with applications to differential equations, in: *CBMS Regional Conference Series in Mathematics*, Vol. 65, Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 1986. [MR845785](#)
- [27] P. H. RABINOWITZ, On a class of nonlinear Schrödinger equations, *Z. Angew. Math. Phys.* **43**(1992), 270–291. [MR1162728](#); [url](#)
- [28] W. A. STRAUSS, Existence of solitary waves in higher dimensions, *Comm. Math. Phys.* **55**(1977), 149–163. [MR0454365](#)
- [29] M. STRUWE, Multiple solutions of differential equations without the Palais–Smale condition, *Math. Ann.* **261**(1982), 399–412. [MR0679798](#)
- [30] M. STRUWE, *Variational methods. Applications to nonlinear partial differential equations and Hamiltonian systems*, third edition, Springer, Berlin, 2000. [MR1736116](#); [url](#)
- [31] X. H. TANG, Infinitely many solutions for semilinear Schrödinger equations with sign-changing potential and nonlinearity, *J. Math. Anal. Appl.* **401**(2013), 407–415. [MR3011282](#); [url](#)
- [32] M. WILLEM, *Minimax theorems*, Progress in Nonlinear Differential Equations and their Applications, Vol. 24, Birkhäuser, Boston, 1996. [MR1400007](#); [url](#)
- [33] K. YOSIDA, *Functional analysis*, Classics in Mathematics, Springer-Verlag, 1995. [MR1336382](#)

- [34] W. Zou, Variant fountain theorems and their applications, *Manuscripta Math.* **104**(2001), 343–358. [MR1828880](#); [url](#)
- [35] W. Zou, *Sign-changing critical point theory*, Springer, 2008. [MR2442765](#)