

Bile acids activate ryanodine receptors in pancreatic acinar cells via a direct allosteric mechanism

Nikolett Geyer^a, Gyula Diszházi^a, László Csénoch^a, István Jónai^{a,b}, János Almássy^{a,b,*}

^a Department of Physiology, University of Debrecen, Faculty of Medicine, 98. Nagyerdei krt. PO Box: 22, Debrecen 4012, Hungary

^b Research Center for Molecular Medicine, University of Debrecen, Faculty of Medicine, 98. Nagyerdei krt. PO Box: 72, Debrecen 4012, Hungary

ARTICLE INFO

Article history:

Received 22 December 2014

Received in revised form 10 March 2015

Accepted 30 March 2015

Available online xxx

Keywords:

Ryanodine receptor

SERCA

Calcium release

Pancreas

Pancreatitis

Bile acid

ABSTRACT

The earliest critical event of pancreatitis is a long lasting high amplitude rise of intracellular Ca^{2+} concentration of the acinar cell, which can be triggered by high concentration of bile acids. Although, Ca^{2+} -release through ryanodine receptors (RyR) is involved in the process, the significance and the exact mechanism of bile acid's action on RyR has not been fully elucidated yet. Therefore, we aimed to test with various techniques and aspects whether bile acids exert a direct effect on RyR and SERCA pump.

Our data show that taurocholic acid (TCA)-induced Ca^{2+} release in pancreatic acinar cells was significantly reduced by the RyR antagonist dantrolene. Further, we show that TCA enhanced RyR's ^3H -ryanodine binding and triggered robust Ca^{2+} -release from RyR-enriched vesicles in the pathologically relevant concentration range. RyR single channel current analysis demonstrated that 200 μM TCA induced a 5-fold increase in the channel's open probability and caused a significant lengthening of the mean open time. TCA also suppressed Ca^{2+} -uptake rate and ATP-ase activity of SERCA-enriched vesicles, but interestingly, failed to decrease Ca^{2+} elimination rate in intact cells.

Overall, our results strongly suggest that TCA opens RyR by an allosteric mechanism, which contribute significantly to bile acid-induced pathologic Ca^{2+} -leak from the endoplasmic reticulum in pancreatic acinar cells.

© 2015 Published by Elsevier Ltd.

1. Introduction

Pancreatitis is a painful and potentially fatal inflammatory disease with a high incidence rate of around 13–45/100,000 people/year [1]. Specific medical therapies are not available. The most common cause of pancreatitis are gallstones, which cause reflux of the bile into the pancreatic duct system by occluding the common bile duct [2,3]. Bile acid concentration therefore may reach millimolar levels in the pancreas, which triggers uncontrolled Ca^{2+}

release and prolonged elevation of the intracellular Ca^{2+} concentration ($[\text{Ca}^{2+}]_{\text{i}}$) in the acinar cell [4]. Impaired Ca^{2+} signaling is considered to be the critical molecular step of acute pancreatitis because it causes premature intraacinar protease activation and acinar cell damage that provokes the inflammation [5,6].

During physiological pancreatic acinar cell stimulation, exocytosis of zymogens is controlled by Ca^{2+} oscillations highly localized to the apical region of the cell [7]. The spatial localization of Ca^{2+} release is explained by high inositol trisphosphate receptor (IP₃R) density in the apical endoplasmic reticulum (ER) (also called Ca^{2+} release trigger zone) [8,9]. In contrast, overstimulation of the pancreatic acinar cell by secretagogues and high concentrations of bile acids induce whole-cell, high amplitude, long lasting, (peak-plateau type) Ca^{2+} signals. The mechanism underlying global Ca^{2+} signals is Ca^{2+} wave propagation, which involves the Ca^{2+} dependent activation of ryanodine receptor (RyR) Ca^{2+} release channels that show higher density in the basolateral-supranuclear region of the acinar cell. Since the ryanodine sensitive compartment is the major source of Ca^{2+} under pathological Ca^{2+} release, the role of RyR in the pathomechanism seems to be essential [4,10,11]. For instance, recent studies showed that Ca^{2+} signals in the basolateral areas induced by supramaximal concentrations of carbachol and that of

Abbreviations: RyR, ryanodine receptor; $[\text{Ca}^{2+}]_{\text{i}}$, intracellular Ca^{2+} concentration; IP₃R, inositol trisphosphate receptor; ER, endoplasmic reticulum; TC, terminal cisternae; LSR, longitudinal sarcoplasmic reticulum; CICR, Ca^{2+} induced Ca^{2+} release; SERCA, sarco-endoplasmic reticulum Ca^{2+} pump; TLCS, taurolithocholic acid-3-sulphate; TCA, taurocholate; TDC, taurodeoxycholate; CPA, cyclopiazonic acid; RR, ruthenium red; Po, open probability; FWHM, full widths at half maximum; cch, carbachol.

* Corresponding author at: Department of Physiology, University of Debrecen, Faculty of Medicine, 98. Nagyerdei krt. PO Box: 22, Debrecen 4012, Hungary. Tel.: +36 304331450.

E-mail addresses: geyer.nikoletta@med.unideb.hu (N. Geyer), diszgyuszi@gmail.com (G. Diszházi), csl@edu.unideb.hu (L. Csénoch), jona.istvan@med.unideb.hu (I. Jónai), almassy.janos@med.unideb.hu (J. Almássy).

<http://dx.doi.org/10.1016/j.ceca.2015.03.009>
0143-4160/© 2015 Published by Elsevier Ltd.

the subcellular distribution of RyR overlapped with the regions where caerulein-induced early zymogen activation occurred [10]. More importantly, the RyR inhibitor dantrolene diminished the carbachol-induced $[Ca^{2+}]_i$ elevation selectively in the basolateral region and reduced the severity of experimental pancreatitis in vivo [11]. These data provide strong evidence that RyR plays a significant role in the pathomechanism of pancreatitis and therefore it can be considered as an important therapeutic target.

The bile acid taurolithocholic acid-3-sulphate (TLCS) was also demonstrated to produce global Ca^{2+} release, which originated in the apical region and spread toward the basal end of the acinar cell [5]. In permeabilized cell assay, bile acid retained its effect when IP₃R or RyR were blocked, but failed when both IP₃R and RyRs were inhibited [12]. Moreover, both dantrolene and ryanodine converted bile acid-induced sustained intracellular $[Ca^{2+}]$ elevations to physiological Ca^{2+} oscillations in intact cells [4]. Although, these data suggest that bile acids trigger excessive Ca^{2+} release by directly opening RyRs too, this possibility remains to be tested, because alternatively, RyR might be activated indirectly via Ca^{2+} induced Ca^{2+} release (CICR) mechanism by the Ca^{2+} , released from acidic vesicles through NAADP receptor Ca^{2+} channels [6,12]. Nevertheless, this question has not been addressed yet. Furthermore, bile acids were reported to inhibit sarco-endoplasmic reticulum Ca^{2+} pump's (SERCA) activity which might result in an increased RyR and IP₃R mediated Ca^{2+} leak and CICR rate [13].

As pictured above, despite the fact that bile acid-induced Ca^{2+} signals in pancreatic acinar cells are intensively studied, the complex interplay between the Ca^{2+} transporters makes the role of RyR and SERCA difficult to evaluate in this process. Therefore, we aimed to test the action of different bile acids explicitly on RyR and SERCA function using the most direct methods available and in the simplest settings possible.

The major components of the bile, cholate and chenodeoxycholate are synthesized by the liver and converted to secondary bile acids including taurocholate, deoxycholate and lithocholates by the intestinal flora. Due to secondary bile acid recycling through the enterohepatic circulation, bile acids occur in taurine-conjugated form. Because taurine conjugated bile acids are the most effective Ca^{2+} mobilizers among bile acids, the major taurine-conjugated bile acid taurocholic acid's (TCA) effect was tested throughout the study. In addition, other secondary bile acids taurodeoxycholate (TDC) and taurolithocholic acid sulphate (TLCS) have also been tested in Ca^{2+} imaging and Ca^{2+} release assays. In order to selectively investigate RyR's or SERCA function, ³H ryanodine binding assay, Ca^{2+} release measurements, single channel recording, and an enzyme-linked NADH assay were performed using purified microsomes. These experiments all demonstrate that bile acids are potent RyR openers and SERCA inhibitors.

2. Material and methods

2.1. Chemicals

If not specified, chemicals were purchased from Sigma.

2.2. Pancreatic acinar cell isolation

Pancreatic acinar cells were freshly isolated from mouse pancreas as described previously [14]. Briefly, 4 months old NMRI mice of both genders were sacrificed by cervical dislocation and the pancreas was removed. The tissue was injected with F12/DMEM medium containing 100 U/ml collagenase P (Roche), 0.1 mg/ml trypsin inhibitor and 2.5 mg/ml BSA, then incubated in this solution in a 37 °C shaking water bath for 30 min, and continuously gassed with carbogen. The tissue was dissociated by pipetting with a

serological pipette 4–6 times, then filtered through mesh #60 (150 μ m). Cells were layered on the top of 400 mg/ml BSA and washed through the medium by gentle centrifugation. The pellet was resuspended in saline solution, collected by centrifugation. Acinar cell clumps were gently resuspended in F12/DMEM medium and kept gassed at room temperature until use.

2.3. Intracellular Ca^{2+} imaging

Acinar cell clumps were loaded with 2 μ M Fluo-4 AM Ca^{2+} sensitive dye for 30 min at room temperature. Acinar cells were plated on glass coverslips and mounted on a perfusion chamber. After perfusion with Ca^{2+} -free Tyrode's solution containing (in mM): 140 NaCl, 5 KCl, 2 MgCl and 10 HEPES, pH 7.2, taurocholate (1, 2 or 5 mM), tauro deoxycholate (1 mM) or taurolythocholic acid sulphate (0.25 mM) was applied by superfusing the cells for 3 or 12 min at room temperature. At the end of each measurement, 2 μ M carbachol was applied to obtain an internal control signal. In some experiments, the loading and imaging media was supplemented with 12 μ M dantrolene or 1.8 mM $CaCl_2$.

Fluorescence was monitored using a Zeiss LSM 5 LIVE confocal microscope equipped with a 40× objective. Fluo-4 was excited at 488 nm and the emitted light was collected through a 520 nm filter. Fluorescence emission data of single cells was analyzed and F/F₀ ratio was calculated after background subtraction using Zeiss ZEN 2009 software. Data were averaged from multiple cells in each experiment then, these values were further averaged to get the mean \pm SEM.

2.4. Microsome isolation

Microsomes from rat pancreas were isolated as described earlier [15]. A pancreas was homogenized using a glass homogenizer in 6 ml ice cold DMEM supplemented with Roche protease inhibitor cocktail. The homogenate was transferred into 1.5 ml microcentrifuge tubes and centrifuged at 11,000 × g for 15 min. The fluffy layer on the top of the pellet was collected and centrifuged at 240,000 × g in a sw 55 ti rotor for 30 min at 4 °C. The pellet was resuspended in a small volume of 300 mM sucrose, 10 mM K-PIPES (pH = 7.0), rapidly frozen in liquid nitrogen and stored at –70 °C.

Sarcoplasmic reticulum terminal cisternae (TC)- and longitudinal SR (LSR) vesicles were isolated from 50 g of rabbit skeletal muscle by differential centrifugation as described previously [16]. All the steps were carried out in cold room or on ice and in the presence of protease inhibitors (in μ M: 200 pepflobc SC, 0.1 aprotinin, 1 leupeptin, 0.2 pepstatin A, 500 benzamidine). After homogenization in 450 ml buffer (in mM: 100 NaCl, 20 EGTA, 20 Na-HEPES, pH = 7.5), cell debris was pelleted at 3500 × g, for 35 min using a tabletop centrifuge equipped with a swing-out rotor. Crude microsomes were collected from the supernatant by centrifugation in a Ti45 rotor at 40,000 × g, for 30 min. To dissolve the actomyosin content the pellet was resuspended in (in mM) 600 KCl, 10K-Pipes, 250 sucrose, 1 EGTA, 0.9 $CaCl_2$ (pH, 7.0). After incubation for 1 h at 4 °C, the microsome fraction was centrifuged at 109,000 × g, for 30 min, the pellet was resuspended and loaded onto a 20–45% linear sucrose gradient (in mM: 105 NaCl, 10 Pipes, 0.1 EGTA, 0.09 $CaCl_2$, pH = 7.0). After spinning overnight at 90,000 × g in a SW27 rotor, two visible rings, corresponding to LSR and TC vesicles were collected from the 30–32% and the 36–38% regions of the sucrose gradient, respectively. The microsomes were washed with a buffer, containing (in mM) 475 sucrose, 1 NaCl, 10 Pipes (pH = 7.0) and pelleted by centrifugation at 124,000 × g for 60 min in a Ti45 rotor. The pellet was resuspended at a final protein concentration of >20 mg/ml in a solution containing 300 mM sucrose and 10 mM K-PIPES (pH = 7.0). Vesicles were aliquoted and rapidly frozen in liquid nitrogen and stored at –70 °C until further use or were immediately

used for RyR purification. Because the LSR fraction is more abundant in SERCA pump than TC, but does not contain RyR, it was used in Ca^{2+} uptake- and in ATPase activity measurements. TC microsomal fraction contains both RyR and SERCA pump, so they are used in Ca^{2+} release assays and for RyR purification. Importantly, these skeletal muscle microsomal fractions do not contain IP_3Rs .

2.5. RyR purification

TC vesicles (3 mg protein/ml) were solubilized for 2 h in a buffer containing 1% CHAPS, 1 M NaCl, 100 μM EGTA, 150 μM CaCl_2 , 5 mM AMP, 0.45% phosphatidylcholine, 20 mM Na-PIPES (pH = 7.2) at 4°C. The sample was loaded onto a 10–28% linear sucrose gradient and centrifuged overnight at 90,000 $\times g$ in a SW27 rotor. In order to detect RyR in the sucrose gradient, a small portion of the sample was labeled with ^3H ryanodine during solubilization. RyR-containing fractions were identified, aliquoted, snap-frozen and stored at –70°C until use [16].

2.6. ^3H ryanodine binding assay

Equilibrium ryanodine-binding assay was carried out using ^3H ryanodine in a medium composed of (in mM) 130 NaCl, 20 Na-PIPES pH 7.1, 1 Pefabloc SC. The desired ionized Ca^{2+} concentrations were adjusted with EGTA [17–19]. Aliquots (32 μl) containing 16 μg protein were incubated at 37°C for 120 min, with various concentrations of radioligand, as indicated in the figure legends. The reaction was terminated by filtering the samples using a BIODOT 96-well filter apparatus and 0.45 μm nitrocellulose membrane (Millipore). After filtration, the membrane was washed with equal aliquots of washing medium. The washing buffer was identical to the incubation medium except that ryanodine and Ca^{2+} was omitted. Nonspecific binding was determined in the presence of 1000× excess of ryanodine, which had been added to the incubation mixture prior to the radioligand [20]. The membrane was cut into appropriate pieces and their radioactivity was determined using a liquid scintillation counter (Packard Tri-Carb) [21].

2.7. Calcium flux- and uptake measurements

TC vesicles were actively loaded with Ca^{2+} in the cuvette at 37°C by the addition of the appropriate amount of CaCl_2 in the solution containing (in mM) 92.5 KCl, 18.5 MOPS, 1 MgCl_2 , 1 ATP, pH = 7.0. In each experiment, 0.5 mg protein was used. Calcium efflux was determined by measuring the extravesicular $[\text{Ca}^{2+}]$ using the metallochromic dye antipyrylazo III (250 μM). Transmittance was monitored by a spectrofluorimeter (Spex Fluoromax) at 710 nm. The rate of Ca^{2+} release was obtained by measuring the slope of the initial segment of the process [16]. Absorbance was calculated off-line and was normalized to the average of the first 5 data points. In some experiments vesicles were pretreated with 12 μM dantrolene 30 min prior to be suspended in the imaging solution, which was also supplemented with dantrolene.

Ca^{2+} uptake rate of 250 μg longitudinal sarcoplasmic reticulum vesicles (LSR) was assessed by measuring the slope of absorbance change following an addition of Ca^{2+} under control conditions or in the presence of 250 μM TCA. To inhibit leak through RyRs in these experiments, 5 μM ruthenium red was included into the solution.

2.8. RyR reconstitution and single-channel current recording

Measurements of channel activity were carried out using purified RyR incorporated into planar lipid bilayers [16]. Bilayers were formed in a symmetrical buffer solution (250 mM KCl, 100 μM $\text{K}_2\text{H}_2\text{EGTA}$, 150 μM CaCl_2 , 20 mM HEPES, pH 7.2), across a 200 μm wide hole drilled into the wall of a delrin cap. The bilayer

chamber was manufactured by Warner Instruments Inc. (Hamden, CT, U.S.A.). The lipid mixture contained phosphatidylethanolamine, phosphatidylserine, and phosphatidylcholine in the ratio of 5:4:1 dissolved in *n*-decane in the final lipid concentration of 20 mg/ml. The lipids were obtained from Avanti Polar Lipids.

RyR fusion with the bilayer was induced by the addition of a small aliquot of purified RyR into the chamber, which was defined as *cis* (cytoplasmic) side. The other chamber was referred to *trans* (luminal) and was kept on ground potential. After successful incorporation of RyR, free $[\text{Ca}^{2+}]$ in the *cis* chamber was decreased from 50 μM , to 270 nM by the addition of EGTA. Then, TCA was applied to the cytoplasmic side of the RyR. The experiments were performed at a membrane potential of +60 or –60 mV. The holding potential of the membrane was adjusted, and the bilayer currents were recorded using an Axopatch 200 amplifier and pCLAMP 6.03 software (Axon Instruments, Foster City, CA, U.S.A.). Currents were filtered at 1 kHz through an eight-pole lowpass Bessel filter and digitized at 3 kHz. Open probabilities and mean open times were calculated with clampfit 10 software.

2.9. ATPase activity measurements

ATPase activity of LSR vesicles was determined at 37°C by a coupled enzyme assay in a medium containing (in mM) 100 KCl, 20 Tris-HCl, 5 MgCl_2 , 5 ATP, 0.42 phosphoenolpyruvate, 0.001 calcimycin (A23187, a Ca^{2+} ionophore), 0.2 NADH, 7.5 U/ml pyruvate kinase, and 18 U/ml lactate dehydrogenase (pH 7.5). The assay was performed at free $[\text{Ca}^{2+}]$ of 1 μM to allow maximum SERCA activity. Calcimycin was applied to prevent accumulation of Ca^{2+} in the vesicles (i.e. end-product inhibition). LSR was obtained as described above from rabbit skeletal muscle and the protein concentration was 5.5 $\mu\text{g}/\text{ml}$ in each experiment [22]. Total hydrolytic activity was measured as the decrease of optical density at the NADH absorbance wavelength (340 nm) and expressed in micromoles of inorganic phosphate per milligrams of protein per minute (abbreviated as I.U.) [16]. TCA was added to the medium 1 min before starting the enzyme activity measurements.

2.10. Statistical analysis

Analysis was made in Origin 7.0 (Microcal Software, Northampton, MA, USA) or in Microsoft Excel. Data are presented as the average of at least three independent experiments (*n*, number of experiments or cells measured). Averages are expressed as mean \pm SEM (standard error of the mean). Statistical analysis was performed by using Student's *t*-test. Threshold for statistically significant differences as compared to the respective control was set at **p* < 0.05.

The number of experiments are expressed as *n* = number of experimental repeats/total number of cells/responding cells.

3. Results

3.1. Intracellular calcium imaging

Our first series of experiments were performed to reproduce previous results and to further investigate the role of RyR in pancreatic acinar cells' intracellular Ca^{2+} signaling. To this end, Fluo-4 fluorescence data was collected from enzymatically isolated pancreatic acinar cell clumps. A bright field image of a typical cell group used in this work is shown in the top image of Fig. 1, which illustrates that polarized apical secretory granule localization is retained after the isolation.

In order to selectively investigate Ca^{2+} release from the intracellular compartments and to avoid Ca^{2+} entry from the extracellular space, cells were bathed in a Ca^{2+} -free saline solution throughout




Fig. 1. Taurocholic acid triggers Ca^{2+} release in pancreatic acinar cells. Ca^{2+} imaging was performed on enzymatically isolated mouse pancreatic acinar cells loaded with Fluo-4 AM. An example of an acinar cell clump is shown in the top image. Cells were bathed in Ca^{2+} -free physiological saline solution and treated with 1, 2, and 5 mM taurocholic acid (TCA) for 12 min. Further experiments were performed in 1.8 mM Ca^{2+} and in the presence of 12 μM dantrolene. Examples of normalized

the whole experiment. Examples of such experiments are illustrated in Fig. 1. Cells were treated with 1, 2 and 5 mM TCA for 12 min, respectively. 1 mM TCA failed to trigger Ca^{2+} release, whereas 2 and 5 mM TCA induced repetitive $[\text{Ca}^{2+}]_{\text{i}}$ oscillations. The effect of TCA was concentration dependent: 5 mM TCA induced significantly higher amplitude of Ca^{2+} -spikes, with earlier onset compared to 1 mM. The first spike developed approximately 3 min after cells have been started to be perfused with the bile acid. The lazy development of the effect can probably be attributed to the relatively slow transport rate of carrier mediated bile acid transporters. We never observed a sustained response, however, in some way the fact that (at 5 mM TCA) the baseline between the spikes tends to increase in about 38% of the cells can be interpreted as progressive Ca^{2+} accumulation. The effect of 5 mM TCA was also tested in 1.8 mM extracellular Ca^{2+} . Apparently, the presence of extracellular Ca^{2+} did not change the features of the Ca^{2+} signals (bottom trace). These data are in agreement with previous studies using TCA at the same concentrations for the same period of time [5]. Because, we were interested in the role of RyR in TCA-triggered Ca^{2+} release, we also tested the effect of the RyR inhibitor dantrolene. The fourth trace in Fig. 1 shows a representative record of a cell treated with 5 mM TCA together with 12 μM dantrolene in a Ca^{2+} -free solution. These raw data imply that dantrolene significantly inhibited the amplitude of $[\text{Ca}^{2+}]$ spikes through the whole length of TCA application.

Quantitative analysis of the effect of dantrolene was performed on experimental data, when 5 mM TCA was applied for 3 min. This treatment resulted in a single Ca^{2+} -transient (Fig. 2A). In order to assess the contribution of RyRs to Ca^{2+} release, cells were also treated with the RyR antagonist dantrolene before and during TCA application (Fig. 2B). At the end of each experiment, supramaximal dose (2 μM) of the parasympathomimetic drug carbachol (cch) was applied to induce maximal Ca^{2+} response. During data analysis, this cch-induced peak-fluorescence was used as a reference value to quantify the relative amplitude of the TCA-signal. In Fig. 2C, pooled data of TCA and cch induced Ca^{2+} -signal peak ratios of 39 and 27 cells are shown under control conditions (TCA-only) and in the presence of dantrolene, respectively. These results demonstrate that 12 μM dantrolene significantly suppressed the relative magnitude of the TCA-induced acinar cell response from 0.6 ± 0.08 to 0.29 ± 0.08 , but essentially the same proportion of cells responded (30 vs. 34%). The average peak latency was 195 ± 5 s which slightly shifted to 210 ± 3.9 s in the dantrolene treated cells. These data confirm earlier observations implying that RyR plays a significant role in bile acid-induced $[\text{Ca}^{2+}]$ transients. Next, using more direct methods we tested whether bile acids may act on RyRs.

3.2. Taurocholic acid enhances ^3H ryanodine binding to microsomes

To investigate TCA's action on RyR activity under more dissected conditions, next, we used pancreas microsomes in a radiolabeled ^3H ryanodine binding assay. The assay utilizes the feature of ryanodine that it binds to RyR in a use-dependent and irreversible manner. Therefore, the quantity of ^3H ryanodine bound to microsomes at unit length of time is proportional to RyR's mean open probability.

First, the amount of ryanodine specifically bound to pancreatic crude microsomes was determined under control conditions and in the presence of 200 μM TCA. Pancreatic microsomes showed

fluorescence-background ratios are shown. 2 and 5 mM TCA triggered repetitive fluctuations of the fluorescence signal, which was suppressed by the ryanodine receptor antagonist dantrolene. 1 mM TCA remained without effect during the whole application period. TCA: taurocholic acid.




Fig. 2. Dantrolene inhibits taurocholic acid-induced Ca^{2+} release in pancreatic acinar cells. Ca^{2+} imaging was performed on enzymatically isolated mouse pancreatic acinar cells loaded with Fluo-4 AM. In order to avoid the effect of interfering Ca^{2+} influx mechanisms, the experiments were performed in Ca^{2+} -free medium. The representative Ca^{2+} signal shown in (A) was recorded in an individual cell stimulated with 5 mM taurocholic acid (TCA) under control conditions, while the one shown in (B) was obtained after dantrolene (30 min, 12 μM) pretreatment and in the presence of dantrolene. 3 min after TCA perfusion was started, a robust, simultaneous elevation of intracellular $[\text{Ca}^{2+}]$ was observed in both cases, but in the presence of dantrolene the TCA-related fluorescence-change was significantly lower. TCA-responses were normalized to the amplitude of Ca^{2+} release induced by 2 μM carbachol (cch), which was used at the end of each measurement as a proof of retained functionality of the cells. The normalized values for control and dantrolene treated cells are summarized in (C) ($n=6/39$ and $4/27$, $*p<0.05$). TCA: taurocholic acid, cch: carbachol.

relatively small specific activity of $0.13 \pm 0.03 \text{ pM ryanodine/mg protein}$ at physiologic Ca^{2+} concentration (472 nM), which was increased by TCA treatment to 0.24 ± 0.05 ($n=8$ and 9). Fig. 3A shows pooled data of three independent experiments. These data support our basic hypothesis that TCA stimulates RyR in pancreatic acinar cells. However, this approach was limited by the high background noise and low specific activity of the pancreatic sample,

possibly due to the relatively low density of RyR in the pancreatic ER and the low yield of ER microsome isolation procedure.

Pancreatic acinar cells were shown to express the skeletal muscle type RyR isoform (RyR1) in PCR studies [23,11]. Our live cell Ca^{2+} imaging data using the RyR1 isoform specific antagonist dantrolene [24] also suggest that the primary RyR isoform in pancreatic acinar cells is the RyR type 1. Also, because skeletal muscle microsome preparations are of higher quality and quantity source of RyR1, they make a good substitute of pancreatic microsomes. Therefore, in order to improve our measurements' signal to background ratio, skeletal muscle terminal cisternae (TC) vesicles were used as a model to test further the effect of TCA. These experiments were designed to measure ^3H ryanodine binding to TC vesicles using different concentrations of radiolabeled ryanodine under control conditions or in the presence of 200 μM TCA and to test the Ca^{2+} dependence of TCA-RyR interaction in media containing 472 nM or 5 μM Ca^{2+} . We chose these $[\text{Ca}^{2+}]$ s because under our circumstances ryanodine binding is too low in the lower nanomolar range to measure accurately, whereas 5 μM is around the dissociation constant for Ca^{2+} . TCA was found to promote ^3H ryanodine binding to TC vesicles at both Ca^{2+} concentrations, as shown by Fig. 3B and C. These findings confirm the data, which were obtained with pancreatic microsomes, that TCA stimulates RyR activity and also verifies that TCA acts similarly on both preparations. Next, TCA's dose-response relationship was determined by incubating TC vesicles along with different concentrations of TCA and 18 nM ^3H ryanodine in a 472 nM or 1.3 μM Ca^{2+} containing solutions. Specific radioactivity significantly increased even in the presence of 25 μM TCA, which was further increased by higher concentrations of TCA at both $[\text{Ca}^{2+}]$ s (Fig. 3D and E). These results show that TCA activates RyR in the wide range of $[\text{Ca}^{2+}]$, therefore bile acid action neither depends on $[\text{Ca}^{2+}]$, nor use-dependent. Importantly, these results argue against the idea that bile acids trigger Ca^{2+} release through their ionophoretic activity on SR vesicles [25].

3.3. TCA triggers Ca^{2+} efflux from TC vesicles

Afterwards, Ca^{2+} release assay was used to further elucidate the effect of TCA on RyR. Fig. 4A shows a representative record of four independent experiments, in which extravesicular $[\text{Ca}^{2+}]$ was monitored by measuring the absorbance of the metallochromic dye antipyrylazo III at 710 nm wavelength. In each experiment, equal quantities of TC vesicles were actively loaded with identical amounts of Ca^{2+} using the vesicles' SERCA pump activity and MgATP. Extravesicular $[\text{Ca}^{2+}]$ was suddenly raised by two subsequent injections of small volumes of Ca^{2+} containing solution, which was instantly loaded into the intraluminal space by the ATPase, as reported by the rapid recovery of the absorbance. The newly established steady baseline is due to the dynamic balance between two opposing mechanisms: the intrinsic Ca^{2+} leak through RyRs and the Ca^{2+} uptake. Therefore, in order to selectively investigate RyR function, SERCA needs to be inhibited. For this reason, SERCA inhibitor cyclopiazonic acid (CPA) was added few seconds before TCA treatment. Experiments were also done with CPA alone. These records showed slow Ca^{2+} leak, which is due to the constant background RyR activity and the lack of SERCA activity. Therefore, they served as control. Representative records and averaged data (Fig. 4A right) clearly show that 500 μM TCA triggered a significantly stronger Ca^{2+} efflux compared to CPA only. More importantly, incubation of TC vesicles with therapeutic concentrations (12 μM) of RyR inhibitor dantrolene caused a 113 second delay in the Ca^{2+} efflux's latency (t was 112 ± 15 s for the control and 225 ± 31 s for the dantrolene treated vesicles) (Fig. 4B), whereas 20 μM ryanodine prevented the TCA-induced Ca^{2+} release. Thus, we conclude that TCA exerts its effect specifically through RyR by a direct allosteric mechanism and it is independent of the




Fig. 3. Taurocholic acid-treatment stimulates RyR's ^3H -ryanodine binding. Averaged data of three independent experiments is presented in panel (A), showing that ^3H ryanodine binding to pancreatic microsomes was higher in the presence of 200 μM taurocholic acid (TCA) at physiological $[\text{Ca}^{2+}]$ (472 nM) (* $p < 0.05$). (B–E) Pooled data of experiments, obtained by using skeletal muscle terminal cisternae vesicles. Different concentrations of ^3H ryanodine was incubated with or without 200 μM TCA in solutions, containing 472 nM and 5 μM Ca^{2+} . Data show that 200 μM TCA increased ^3H ryanodine binding of TC vesicles (B, C). Dose response curves using various TCA concentrations at 18 nM ^3H ryanodine and 472 nM or 1.3 μM Ca^{2+} are presented in panels (D) and (E), respectively. Data show that specific radioactivity was significantly increased by TCA in the 25–500 μM range. TCA: taurocholic acid.

detergent-properties of the bile acid. However, these measurements do not directly confirm whether the binding site for TCA is on the RyR itself, or it is on a closely associated regulatory protein.

3.4. TCA enhances RyR gating by increasing its open probability and mean open time

To investigate TCA's effect on RyR channels free of associated proteins, RyRs were solubilized from TC vesicles and purified on sucrose gradient, because most of the RyR's associated counterparts are believed to be lost during this procedure. In order to get deeper insights into the mechanism of action of TCA on RyR channel gating, the purified channels were reconstituted into artificial lipid bilayers and single channel currents were recorded under voltage clamp conditions at +60 mV membrane potential. Fusion of single RyR channels was initiated at 50 μM $[\text{Ca}^{2+}]$ on either side of the bilayer. Then, $[\text{Ca}^{2+}]$ on the cytoplasmic side was lowered

to 270 nM. The top trace in Fig. 5A presents a representative 3 s long segment of the single channel current recording. Under these conditions the open events were detected as positive (upward) deflections and the channel displayed low open probability (P_o) of 0.025 for a ~20 s record. When TCA was added in increasing concentrations to the cytoplasmic face of the RyR, P_o and mean open time (τ) of the channel increased, which was a consistent finding (Fig. 5A). The average P_o and τ of three independent experiments showed saturating concentration dependence, increasing P_o several fold at 500 μM TCA (Fig. 5B and C). The half effective concentration was $181.2 \pm 56.6 \mu\text{M}$ ($n_{\text{Hill}} = 1.54 \pm 0.42$). These data strongly suggest that TCA acts on the RyR protein itself too, and significantly stimulates RyR gating below the pathophysiologically relevant concentration range by increasing both the number and duration of channel openings. The conductance of the channel was not modified by TCA. Finally, the addition of ryanodine caused the characteristic transition to a half-conducting state, implying that




Fig. 4. Taurocholic acid triggers Ca²⁺ efflux from TC vesicles. The effect of taurocholic acid (TCA) was investigated on the function of RyR, using Ca²⁺ flux measurements. Terminal cisternae (TC) vesicles were suspended in a solution, containing the Ca²⁺ indicator dye antipyrylazo III. Extravesicular [Ca²⁺] was monitored by measuring the absorbance at 710 nm as the function of time. (A) Two representative records. To generate concentration gradient for Ca²⁺ efflux, vesicles were loaded with Ca²⁺, by utilizing SERCA's pump-activity. In order to establish similar driving force in each individual experiment, two subsequent injections of the same amount of Ca²⁺ was applied to load the intravesicular space. To prevent interference between Ca²⁺-uptake and -release mechanisms, 20 μM SERCA inhibitor cyclopiazonic acid (CPA) was also added into the solution, prior to the flux measurement. (Ca²⁺ and drug additions are marked by arrows.) SERCA inhibition (CPA) impaired the balance of Ca²⁺ handling mechanisms in TC vesicles, thus it caused a spontaneous Ca²⁺ release. This flux rate is attributed to the basal Ca²⁺ leak through RyRs, (in gray) and served as control. In different experiments, 500 μM TCA was added along with CPA (in black). We found that TCA together with CPA caused a steeper Ca²⁺ flux compared to control. Followed by calibration, Ca²⁺ release rate was calculated. Averaged data (±SE, n = 4, *p < 0.05) is shown on the right of (A). (B, C) Representative experiments, when 12 μM dantrolene, or 20 μM ryanodine (rya) was included in the cuvette respectively. Dantrolene caused a 113 s delay in Ca²⁺ release, while ryanodine prevented the effect of TCA. To demonstrate that ryanodine treated vesicles were also appropriately loaded with Ca²⁺, 1 μM Ca²⁺ ionophore (A23187) was added (marked by arrow), that resulted in a strong Ca²⁺ efflux. TCA: taurocholic acid, TC: terminal cisternae, CPA: cyclopiazonic acid, rya: ryanodine.

the pharmacological properties of RyR are retained after TCA treatment and ryanodine's and TCA's binding sites are distinct.

3.5. TDC and TLCS also cause Ca²⁺ release in pancreatic acinar cells and from TC vesicles

Because the effective 2–5 mM bile acid concentration might not be reached during bile duct obstruction in the pancreas, other bile acids were also investigated to confirm our basic observations made using TCA. As TLCS and TDC are also often used to investigate bile acid induced changes in the exocrine pancreas [4,5,12], we performed Ca²⁺ imaging and Ca²⁺ release experiments using these bile acids too. These results demonstrate that TLCS's and TDC's effect is qualitatively very similar to that we have observed for TCA. Representative experiments are shown in Fig. 6. Notably, TLCS and TDC worked even more potently compared to TCA, in as low as 1 mM (TDC) and 250 μM (TLCS) concentrations in intact acinar cells and in 100 μM and 50 μM on TC vesicles, respectively. Furthermore, TLCS-induced Ca²⁺ release developed much earlier when compared to Ca²⁺ signals triggered by TCA as the peak latency was 194 ± 5 s for TCA, 195 ± 12 s for TDC, and 49 ± 3 s for TLCS. These results suggest that the effect of bile acids on RyR is universal. In few experiments the cells were perfused with 250 μM TLCS in a Ca²⁺ free saline for prolonged period (9 min, Fig. 6B), which induced oscillating [Ca²⁺]i signals. This pattern is very similar to the one triggered by 5 mM TCA (Fig. 1). At the end of these experiments, 2 μM carbachol was immediately added after TLCS

treatment, while in the experiment shown in Fig. 6A, TLCS was applied for only 3 min and TLCS was washed out before carbachol was added. The carbachol-responses in the two cases were very different: they were significantly lower and returned slower in the latter case, when TLCS was applied for 9 min and was not washed out before carbachol. (B). Ca²⁺-depletion of the ER (smaller driving force) due to the repetitive Ca²⁺ release events could account for the smaller amplitude, while the slower Ca²⁺ clearance can be explained by lower Ca²⁺ pump (SERCA and/or PMCA) activities. The amplitude and the spatio-temporal characteristics of a Ca²⁺ signal is ultimately determined by the relative rate of Ca²⁺ release and Ca²⁺ elimination mechanisms. Therefore, suppressed Ca²⁺ reuptake should result in enhanced and long lasting Ca²⁺ release events and resting Ca²⁺ leak, which might also significantly contribute to Ca²⁺ overload in pancreatic acinar cells. However, enhanced ATP expenditure of Ca²⁺ elimination and mitochondrial depolarization during oscillatory Ca²⁺ signals could lead to severe ATP depletion and suppressed Ca²⁺ pump activity, inhibitory action of bile acids on SERCA might also contribute to the slow clearance rate [26–30].

3.6. TCA inhibits SERCA activity of LSR microsomes

Then, experiments were designed to test the idea that TLCS's and TCA's inhibitory action on the SERCA pump is partially responsible for the Ca²⁺ overload during bile acid induced Ca²⁺ release. SERCA pump's hydrolytic activity was measured using a coupled enzyme-NADH assay and LSR, because this skeletal muscle




Fig. 5. Taurocholic acid promotes RyR1 gating. Single RyR1 channels were reconstituted into lipid bilayers, and current traces were recorded under voltage clamp conditions, at +60 mV membrane potential. Representative single channel current traces, recorded under control conditions (270 nM Ca^{2+}) and after taurocholic acid (TCA) was added to the cytoplasmic side of the channel are shown in (A). The closed state level is marked by "C". Single openings of the channel are represented by upward deflections. Under control conditions, due to the relatively low $[\text{Ca}^{2+}]$, the channel stayed mostly in closed state. After TCA treatment, the channel opened more frequently. Detailed data analysis revealed that TCA increased both RyR's open probability (P_o) and mean open time, as shown by the graphs in (B) and (C) respectively ($n=3$). The apparent dissociation constant was $181.2 \pm 56.6 \mu\text{M}$. TCA: taurocholic acid.

vesicular fraction is the richest in SERCA. Graphs in Fig. 7A illustrate original records of these experiments, with increasing concentrations of TCA. Smaller slopes at gradually higher TCA concentrations represent lower SERCA ATPase activities. The specific Ca^{2+} -ATPase activity was identified as the portion of total hydrolytic activity, which was inhibited by 10 μM cyclopiazonic acid. During analysis, hydrolytic activity was calculated from the slopes and the averaged data of 3–5 independent experiments are shown in Fig. 7A, on the right. These results demonstrate that TCA and TLCS exert an inhibitory action on SERCA's hydrolytic activity and they are in agreement with results published previously [13].

The findings obtained by coupled enzyme assay have also been verified by using a different approach, when SERCA's activity was assessed by determining the Ca^{2+} uptake rate of LSR vesicles. The experimental setup was very similar to the one used in Ca^{2+} release experiments. Extravesicular $[\text{Ca}^{2+}]$ was monitored using antipyrylazo III by measuring the absorbance of the Ca^{2+} indicator. At a certain point extravesicular $[\text{Ca}^{2+}]$ was suddenly raised by 12.5 μM . The cuvette contained 250 μg protein (LSR)/ml imaging solution. The experiments were performed in the presence of the RyR inhibitor ruthenium red to prevent interference between RyR mediated background Ca^{2+} efflux and Ca^{2+} uptake. Ca^{2+} uptake rate was calculated using the slope of the best linear fit for the data points. Quantitative data analysis show, that Ca^{2+} uptake rate was significantly lower in the presence of 250 μM TCA, compared to control (9.57 ± 0.39 vs. $6.75 \pm 0.64 \text{ nmol Ca}^{2+}/\text{s}$).

Next, to test the effect of TCA on the Ca^{2+} removal mechanisms in native pancreatic acinar cells, Ca^{2+} release was triggered by using




Fig. 6. Tauro deoicholate and tauroolithocholic acid sulphate induces Ca^{2+} release in intact pancreatic acinar cells and from TC vesicles. Bile acids tauro deoicholate (TDC) and tauroolithocholic acid sulphate (TLCS) were tested using the approaches detailed above. Ca^{2+} imaging data of two cells treated with 1 mM TDC (gray) or 0.25 mM TLCS (black) are displayed in (A). Both bile acids triggered robust Ca^{2+} release in the acinar cells in these concentrations. Interestingly, the TLCS-related response developed earlier compared to TDC and TCA-induced Ca^{2+} signals. The average peak latency was 49 ± 3 , 195 ± 12 and 194 ± 5 s, respectively. At the end of all experiments, 2 μM carbachol (cch) was applied to test the functionality of the cells. (B) An example of repetitive Ca^{2+} spikes of a cell treated with 250 μM TLCS for prolonged period (9 min) in a Ca^{2+} -free medium. (C) Representative records of Ca^{2+} release experiments using terminal cisternae (TC) vesicles. They present that 100 μM TDC (gray line) and 50 μM TLCS (in black) triggered a stronger Ca^{2+} efflux compared to the release elicited by cyclopiazonic acid (CPA) only (arrow). Thus, these results demonstrate that TDC and TLCS are also effective Ca^{2+} releasing bile acids, with RyR activating abilities. TDC: tauro deoicholate, TLCS: tauroolithocholic acid sulphate, CPA: cyclopiazonic acid.

2 μM cch, under control conditions and in the presence of 250 μM TCA or 100 μM TLCS respectively. We chose to apply these bile acid concentrations, because they failed to induce Ca^{2+} release in acinar cells, but caused significant inhibition of SERCA pump hydrolytic activity. Ca^{2+} elimination rate was evaluated by measuring the full widths at half maximum (FWHM) of the fluorescence change after carbachol treatment. While carbachol transients' amplitude was not significantly different between the two groups, Ca^{2+} removal was quicker in cells treated with the bile acids compared to control, as suggested by smaller FWHM values (61.8 ± 2.3 s vs. 95.4 ± 5.1 s for TCA, data not shown). Because we did not observe SERCA activation at any bile acid concentration, these data suggest that




Fig. 7. Taurocholic acid inhibits SR Ca^{2+} pump's hydrolytic activity. Total hydrolytic activity of LSR vesicles was measured as the decrease of optical density at the NADH absorbance wavelength (340 nm) and expressed in micromoles of inorganic phosphate per milligrams of protein per minute. Examples of experiments under control conditions (1) and in the presence of 50, 250 and 500 μM taurocholic acid (TCA) (2, 3, and 4) are presented in (A) (left). Averaged data show that ATP-ase activity was significantly reduced by TCA in the 50–500 μM range (A, right). The inset in (A) shows the result of similar experiments using 25, 50, 100 and 250 μM taurolithocholic acid sulphate (TLCS). SERCA activity was also assessed by determining the Ca^{2+} uptake rate of LSR vesicles. The slope of absorbance-change that was measured after the extravesicular $[\text{Ca}^{2+}]$ was suddenly raised by 12.5 μM (B). Ruthenium red (RR) was included into the measuring solution in order to prevent TCA induced Ca^{2+} leak, therefore to selectively study SERCA function. Otherwise the same method was used as for the Ca^{2+} -release experiments. We found that Ca^{2+} uptake rate was significantly lower in the presence of 250 μM TCA, compared to the control. Original records and pooled data of three independent experiments are shown in (B) (* $p < 0.05$). TCA: taurocholic acid, TLCS: taurolithocholic acid sulphate, RR: ruthenium red.

bile acids also influence other Ca^{2+} handling proteins. We also performed experiments using 2 mM TCA (for 3 min), which resulted in qualitatively same results (data not shown), suggesting that the bile acid does not contribute to the magnitude of Ca^{2+} release by inhibiting SERCA activity in acinar cells, or that these bile acid concentrations are not high enough (or applied for long enough) to reach effective concentrations in the cytoplasm. Therefore, we performed similar experiments using carbachol-stimulation, followed by 250 μM TLCS treatment for 9 min (see Fig. 6B). Control cells were stimulated only with carbachol. Because cells treated with 250 μM TLCS developed $[\text{Ca}^{2+}]$ oscillations and significantly smaller carbachol response (possibly due to their lower ER $[\text{Ca}^{2+}]$ and poor ATP-status (also see Section 3.5)), the untreated cells did not serve as adequate controls any more. Therefore, analysis and comparison of Ca^{2+} clearance rates in the presence or absence of 250 μM TLCS cannot be performed. In summary, bile acids inhibit SERCA activity directly, and indirectly, by Ca^{2+} release induced ATP depletion, but their significance is technically difficult to evaluate in native acinar cells.

4. Discussion

The action of bile acids with particular interest to the effect of TLCS have been recently intensively investigated on pancreatic acinar cell's intracellular Ca^{2+} signaling and cytotoxicity. Cytotoxicity is related to the lipid solubility of the particular bile acid. In general, dehydroxylated and deconjugated bile acids are more toxic than others [31]. The bile mostly contains taurin- or glycine-conjugated secondary (dehydroxylated) bile acids. TCA is one of the most

abundant secondary bile acid constituent of the bile. The three most extensively investigated bile acids in the order of cytotoxicity are TLCS > TDC > TCA [5].

Bile acids attracted much attention after the link between persistent elevation of intraacinar $[\text{Ca}^{2+}]$ and pancreatitis has been established and TCA was demonstrated to trigger excessive Ca^{2+} release from intracellular stores in pancreatic acinar cells. TCA and TLCS were previously shown to most probably activate IP₃Rs [4,5,12]. Although, the enrollment of RyRs during the process was quite obvious, the way they were recruited was not completely understood. Our study addressed the question for the first time, whether bile acids release Ca^{2+} by activating RyR. We hypothesized that they directly activate the channel. The objective of our study was to test the action of bile acids on the calcium release channel using direct methods and to assess its significance in impaired Ca^{2+} signaling.

The data presented here clearly show that bile acids potently and directly open the channel. They thus highlight the role of RyR in bile acid-induced toxic Ca^{2+} release and propose that the contribution of RyR is direct and cannot only be through the CICR mechanism. Although, bile acid-induced Ca^{2+} release is initiated from the luminal end of ER, where IP₃Rs are localized at higher density than RyRs, our study raises the possibility that the pathological process is initiated by RyR activation. Although, answering these questions needs further investigations.

Recent studies have shown that 5–50 mg/kg dantrolene mitigated caerulein-induced pancreatitis *in vivo* in mice and suggested dantrolene in the therapy of acute pancreatitis [11]. Therefore, we aimed to assess the role of RyR in TCA-triggered Ca^{2+} release using dantrolene in live cell Ca^{2+} imaging experiments.

Dantrolene is the only RyR antagonist in human medicine, which is used as a muscle relaxant to treat malignant hyperthermia and muscle spasticity caused by stroke, spinal cord injury or multiple sclerosis. One of the most important finding of the present study is that Ca^{2+} release was significantly attenuated by the drug. Since dantrolene is selective for the skeletal muscle type RyR1 [24], these results provide evidence that the most important isoform involved in bile acid-induced Ca^{2+} release is RyR1. We used dantrolene in the concentration of 12 μM , because our group has previously shown that this concentration is sufficient to significantly suppress Ca^{2+} efflux rate and the amount of Ca^{2+} released from TC vesicles by thymol [31]. Interestingly, dantrolene inhibited the TCA-induced Ca^{2+} release by causing a delay in Ca^{2+} release instead of inhibiting Ca^{2+} flux rate. The dissimilarity of these data possibly due to the idea that dantrolene probably act on or requires a yet unidentified accessory protein. This is supported by dantrolene's ineffectiveness on solubilized RyR channel currents [32]. Importantly though, ryanodine treatment completely abolished the effect of TCA, which argues against the ionophoretic theory of bile acid action [25] and confirms the specificity of TCA's action on the RyR protein complex.

The most important result of this study is provided by our single channel current data. In these experiments, purified RyRs, – free of most of the accessory proteins – were incorporated into artificial lipid bilayers. They provide clear evidence that TCA directly acts on the RyR protein itself, most probably without modifying the interactions between RyR and its regulatory proteins. The fact that TCA's effect developed and reached steady state quickly after addition, implies that RyR activation does not rely on nonspecific mechanisms, such as TCA's detergent properties and increased membrane fluidity, which could have been due to the progressive accumulation of the bile acid in the bilayer.

Earlier studies have demonstrated that bile acids slow down Ca^{2+} clearance rate by inhibiting SERCA pump activity in stimulated pancreatic acinar cells. By doing so, they chronically deplete Ca^{2+} stores, activate store-operated Ca^{2+} entry that cause Ca^{2+} overload and cell death. In these experiments a bile acid mixture was used; but the effects of the components were not investigated specifically [13]. Therefore, we aimed to specify the action of TCA and TLCS on SERCA pump activity. Although, these bile acids significantly inhibited SERCA pump activity in LSR vesicles, they failed to decrease Ca^{2+} elimination rate in native acinar cells in relatively low concentrations (250 μM TCA and 100 μM TLCS). Instead, they slightly enhanced Ca^{2+} removal rate. This discrepancy suggests that the SERCA-inhibiting action might be masked by an unknown mechanism, or that these bile acid concentrations are not high enough or they are not applied for long enough to reach effective concentrations in the cytoplasm. Therefore, Ca^{2+} clearance rate after carbachol stimulation was evaluated after cells were treated with higher (250 μM) TLCS concentration and for more prolonged periods. TLCS at this concentration produces robust Ca^{2+} oscillation and mitochondrial membrane depolarization, which leads to high rate of ATP consumption by Ca^{2+} ATPases and interferes with ATP synthesis. As a consequence, Ca^{2+} depletion of the ER and severe ATP depletion develops, which results in significantly smaller amplitude of carbachol response with slow slope of recovery compared to control. Thus, because this approach to evaluate the effect of bile acids on SERCA pump in native cells is very limited by the above mentioned interfering circumstances, appropriate consequences cannot be drawn. We can only rely on SERCA ATPase activity measurements to assess the effect of bile acids on the Ca^{2+} pump [26–30].

In summary, our data supports the idea that RyR is allosterically gated by bile acids and RyR is a possible therapeutic target in pancreatitis, furthermore, they emphasize the therapeutic potential of dantrolene.

Although, most of the studies investigating the pathophysiology of acute pancreatitis focused on pancreatic acinar cells, the action of bile acids on ductal cell might be pathologically significant too, because during the biliary reflux bile acids reach the ductal cells first. The effect of bile acids on ductal cells was biphasic. Low concentration of chenodeoxycholic acid (CDC) stimulated HCO_3^- secretion in a Ca^{2+} dependent manner, which has a protective effect on the pancreas. In contrast, high concentrations of bile acids induce toxic $[\text{Ca}^{2+}]_{\text{i}}$ elevations, which leads to the loss of mitochondrial membrane potential and ATP depletion. At severely low [ATP] the $\text{Cl}^-/\text{HCO}_3^-$ exchanger function is suppressed. As a consequence, low fluid secretion rate favors ductal cell injury and the damage of ductal integrity, thus acute pancreatitis [31,33–36].

Conflict of interest

None.

Acknowledgements

This work was supported by a grant provided to IJ and JA from the Hungarian Scientific Research Fund (OTKA K81923 and OTKA PD 112199).

JA is supported by the Janos Bolyai Research Scholarship of the Hungarian Academy of Sciences and the Lajos Szodoray Scholarship of the University of Debrecen, Faculty of Medicine.

References

- [1] D. Yadav, A.B. Lowenfels, The epidemiology of pancreatitis and pancreatic cancer, *Gastroenterology* 144 (2013) 1252–1261.
- [2] E.L. Opie, The etiology of acute hemorrhagic pancreatitis, *Johns Hopkins Hosp. Bull.* 112 (1901) 182–188.
- [3] H.J. Aho, S.M. Koskensalo, T.J. Nevalainen, Experimental pancreatitis in the rat. Sodium taurocholate-induced acute haemorrhagic pancreatitis, *Scand. J. Gastroenterol.* 15 (1980) 411–416.
- [4] S.Z. Husain, A.I. Orabi, K.A. Muili, et al., Ryanodine receptors contribute to bile acid induced pathological calcium signaling and pancreatitis in mice, *Am. J. Physiol. Gastrointest. Liver Physiol.* 302 (2012) 1423–1433.
- [5] S. Voronina, R. Longbottom, R. Sutton, O. Petersen, A. Tepikin, Bile acids induce calcium signals in mouse pancreatic acinar cells: implications for bile-induced pancreatic pathology, *J. Physiol.* 540 (2002) 49–55.
- [6] J.V. Gerasimenko, G. Lur, M.W. Sherwood, et al., Pancreatic protease activation by alcohol metabolite depends on Ca^{2+} release via acid store IP_3 receptors, *Proc. Natl. Acad. Sci. U. S. A.* 106 (2009) 10758–10763.
- [7] L.R. Johnson, K.F. Ghishan, D.J. Kaunitz, L.J. Merchant, M.H. Said, D.J. Wood, *Physiology of the Gastrointestinal Tract* Academic Press, 2012.
- [8] O.H. Petersen, V.A. Tepikin, Polarized calcium signaling in exocrine gland cells, *Annu. Rev. Physiol.* 70 (2007) 273–299.
- [9] J.H. Won, W.J. Cottrell, T.H. Foster, D.J. Yule, Ca^{2+} release dynamics in parotid and pancreatic exocrine acinar cells evoked by spatially limited flash photolysis, *Am. J. Physiol. Gastrointest. Liver Physiol.* 293 (2007) 1166–1177.
- [10] S.Z. Husain, P. Prasad, W.M. Grant, T.R. Kolodkin, M.H. Nathanson, F.S. Gorelick, The ryanodine receptor mediates early zymogen activation in pancreatitis, *Proc. Natl. Acad. Sci. U. S. A.* 102 (2005) 14386–14391.
- [11] A.I. Orabi, A.U. Shah, M.U. Ahmad, et al., Dantrolene mitigates caerulein-induced pancreatitis *in vivo* in mice, *Am. J. Physiol. Gastrointest. Liver Physiol.* 299 (2010) 196–204.
- [12] V.J. Gerasimenko, G.S. Flowerdew, K.T. Voronina, et al., Bile acids induce Ca^{2+} release from both the endoplasmic reticulum and acidic intracellular calcium stores through activation of inositol triphosphate receptors and ryanodine receptors, *J. Biol. Chem.* 281 (2006) 40154–40163.
- [13] J.Y. Kim, K.H. Kim, J.A. Lee, et al., Transporter-mediated bile acid uptake causes Ca^{2+} -dependent cell death in rat pancreatic acinar cells, *Gastroenterology* 122 (2002) 1941–1953.
- [14] J.A. Williams, M. Korc, R.L. Dorner, Action of secretagogues on a new preparation of functionally intact, isolated pancreatic acini, *Am. J. Physiol.* 235 (1978) 517–524.
- [15] T. Ozawa, Ryanodine-sensitive Ca^{2+} release mechanism of rat pancreatic acinar cells is modulated by calmodulin, *Biochim. Biophys. Acta* 1452 (1999) 254–262.
- [16] S. Sárközi, J. Almássy, B. Lukács, N. Dobrosi, G. Nagy, I. Jóná, Effect of natural phenol derivatives on skeletal type sarcoplasmic reticulum Ca^{2+} -ATPase and ryanodine receptor, *J. Muscle Res. Cell Motil.* 28 (2007) 167–174.
- [17] X. Altafaj, J. France, J. Almássy, et al., Maurocalcine interacts with the cardiac ryanodine receptor without inducing channel modification, *Biochem. J.* 406 (2007) 309–315.

- [18] T.J. Lanner, D.K. Georgiou, A. Dagnino-Acosta, et al., AICAR prevents heat induced sudden death in RyR1 mutant mice independent of AMPK activation, *Nat. Med.* 18 (2012) 244–251.
- [19] G.G. Du, J.P. Imredy, D.H. MacLennan, Characterization of recombinant rabbit cardiac and skeletal muscle Ca^{2+} release channels (ryanodine receptors) with a novel [^3H]ryanodine binding assay, *J. Biol. Chem.* 273 (1998) 33259–33266.
- [20] J.M. Dias, C. Szegedi, I. Jóna, P.D. Vogel, Insights into the regulation of the ryanodine receptor: differential effects of Mg^{2+} and Ca^{2+} on ATP binding, *Biochemistry* 45 (2006) 9408–9415.
- [21] S. Sárközi, C. Szegedi, B. Lukács, M. Ronjat, I. Jóna, Effect of gadolinium on the ryanodine receptor/sarcoplasmic reticulum calcium release channel of skeletal muscle, *FEBS J.* 272 (2005) 464–471.
- [22] N. Szentandrásy, A. Szabó, J. Almásy, et al., Effects of articaine and ropivacaine on calcium handling and contractility in canine ventricular myocardium, *Eur. J. Anaesthesiol.* 27 (2010) 153–161.
- [23] T.J. Fitzsimmons, I. Gukovsky, J.A. McRoberts, E. Rodriguez, F.A. Lai, S.J. Pandol, Multiple isoforms of the ryanodine receptor are expressed in rat pancreatic acinar cells, *Biochem. J.* 351 (2000) 265–271.
- [24] F. Zhao, P. Li, S.R. Chen, C.F. Louis, B.R. Fruen, Dantrolene inhibition of ryanodine receptor Ca^{2+} release channels. Molecular mechanism and isoform selectivity, *J. Biol. Chem.* 276 (2001) 13810–13816.
- [25] P. Zimniak, J.M. Little, A. Radominska, D.G. Oelberg, M.S. Anwer, R. Lester, Taurine-conjugated bile acids act as Ca^{2+} ionophores, *Biochemistry* 30 (1991) 8598–8604.
- [26] J.I. Bruce, Plasma membrane calcium pump regulation by metabolic stress, *World J. Biol. Chem.* (2010) 221–228.
- [27] J.I. Bruce, The Pancreapedia: Exocrine Pancreas Knowledge Base, PMCA (2013), <http://dx.doi.org/10.3998/panc.2013.7>
- [28] D.N. Criddle, J. Murphy, G. Fistetto, et al., Fatty acid ethyl esters cause pancreatic calcium toxicity via inositol trisphosphate receptors and loss of ATP synthesis, *Gastroenterology* 130 (2006) 781–793.
- [29] S.L. Barrow, S.G. Voronina, G. da Silva Xavier, et al., ATP depletion inhibits Ca^{2+} release, influx and extrusion in pancreatic acinar cells but not pathological Ca^{2+} responses induced by bile, *Pflugers Arch.* 455 (2008) 1025–1039.
- [30] S.G. Voronina, S.L. Barrow, O.V. Gerasimenko, O.H. Petersen, A.V. Tepikin, Effects of secretagogues and bile acids on mitochondrial membrane potential of pancreatic acinar cells: comparison of different modes of evaluating DeltaPsim, *J. Biol. Chem.* 279 (2004) 27327–27338.
- [31] V. Venglovecz, Z. Rakonczay, P. Hegyi, The effects of bile acids on pancreatic ductal cells, The Pancreapedia: Exocrine Pancreas Knowledge Base (2012), <http://dx.doi.org/10.3998/panc.2012.8>
- [32] P. Szentesi, C. Collet, S. Sárközi, et al., Effects of dantrolene on steps of excitation-contraction coupling in mammalian skeletal muscle fibers, *J. Gen. Physiol.* 118 (2001) 355–375.
- [33] C. Alvarez, A. Fasano, B.L. Bass, Acute effects of bile acids on the pancreatic duct epithelium in vitro, *J. Surg. Res.* 74 (1998) 43–46.
- [34] J. Maléth, V. Venglovecz, Z. Rázga, L. Tiszavicz, Z. Rakonczay Jr., P. Hegyi, Non-conjugated chenodeoxycholate induces severe mitochondrial damage and inhibits bicarbonate transport in pancreatic duct cells, *Gut* 60 (2011) 136–138.
- [35] H.A. Reber, J.G. Mosley, The effect of bile salts on the pancreatic duct mucosal barrier, *Br. J. Surg.* 67 (1980) 59–62.
- [36] V. Venglovecz, Z. Rakonczay Jr., B. Ozsvári, T. Takács, J. Lonovics, A. Varró, M.A. Gray, B.E. Argent, P. Hegyi, Effects of bile acids on pancreatic ductal bicarbonate secretion in guinea pig, *Gut* 57 (2008) 1102–1112.