Percolation on the stationary distributions of the voter model

Balázs Ráth¹, Daniel Valesin²

February 5, 2015

Abstract

The voter model on \mathbb{Z}^d is a particle system that serves as a rough model for changes of opinions among social agents or, alternatively, competition between biological species occupying space. When $d \geq 3$, the set of (extremal) stationary distributions is a family of measures μ_{α} , for α between 0 and 1. A configuration sampled from μ_{α} is a strongly correlated field of 0's and 1's on \mathbb{Z}^d in which the density of 1's is α . We consider such a configuration as a site percolation model on \mathbb{Z}^d . We prove that if $d \geq 5$, the probability of existence of an infinite percolation cluster of 1's exhibits a phase transition in α . If the voter model is allowed to have sufficiently spread-out interactions, we prove the same result for d > 3.

Keywords: interacting particle systems, voter model, percolation

AMS MSC 2010: 60K35, 82C22, 82B43

1 Introduction

1.1 Model and results

Given integers $d \geq 1$ and $R \geq 1$, the voter model with range R on the d-dimensional lattice \mathbb{Z}^d is a Markov process, denoted here by $(\xi_t)_{t\geq 0}$, with configuration space $\{0,1\}^{\mathbb{Z}^d}$ and stochastic dynamics described informally as follows. Each vertex (or site) x of \mathbb{Z}^d updates its current state $\xi_t(x) \in \{0,1\}$ at rate one by copying the state $\xi_t(y)$ of a vertex y that is chosen uniformly among all vertices at $(\ell^1$ -norm) distance at most R from x.

In Section 3 we give the formal definition of the model and recall some of its relevant properties. In this Introduction, we will only very briefly present the concepts that are needed to state our main results.

The voter model was introduced independently by Clifford and Sudbury in [CS73] and Holley and Liggett in [HL75]. In the interpretation of the latter pair of authors, each site of \mathbb{Z}^d represents a voter which can have one of two possible opinions (corresponding to the states 0 and 1). The model thus represents the evolution of the opinions among the population. Clifford and Sudbury gave a biological interpretation for the model: there are two competing species, denoted 0 and 1, and each site is a region of space that can be occupied by an individual of one of the two species.

The set of stationary distributions of the voter model on \mathbb{Z}^d has been thoroughly studied; the following is a summary of known results. For fixed $d \geq 3$, $R \geq 1$ and $\alpha \in [0,1]$, one defines a probability measure μ_{α} on $\{0,1\}^{\mathbb{Z}^d}$ as the distributional limit (which is shown to exist), as time is taken to infinity, of the voter model with the random initial configuration in which the states of all sites are independent and Bernoulli(α). μ_{α} is then stationary for the voter model dynamics. Moreover, it is shown that the set of stationary distributions for the voter model dynamics that are extremal – i.e., that

¹Budapest University of Technology, Hungary. E-mail: rathb@math.bme.hu

²University of Groningen, The Netherlands. E-mail: d.rodrigues.valesin@rug.nl

cannot be expressed as non-trivial convex combinations of other stationary distributions – is precisely the family

$$\{ \mu_{\alpha} : \alpha \in [0,1] \}.$$

The measures μ_{α} can be obtained in a more constructive way with the aid of coalescing random walks. A realization of a system of coalescing random walks with range R on \mathbb{Z}^d induces a partition of \mathbb{Z}^d into coalescence classes: we say that x and y are in the same class if the walkers started at x and y are eventually joined. We then assign 0's or 1's to the coalescence classes independently with probabilities $1-\alpha$ and α , respectively, and the resulting configuration $\xi \in \{0,1\}^{\mathbb{Z}^d}$ has law μ_{α} . (Again, the sentences in this paragraph will be given a precise meaning in Section 3).

With the aid of this construction, it is not difficult to show that each μ_{α} is invariant and ergodic with respect to translations of \mathbb{Z}^d and satisfies $\mu_{\alpha}[\xi(0) = 1] = \alpha$, so that α is equal to the density of 1's. Moreover, the family $\{\mu_{\alpha}\}$ is stochastically increasing: in the partial order on $\{0,1\}^{\mathbb{Z}^d}$ induced by the order 0 < 1 on the coordinates, we have that μ_{α} is stochastically dominated by $\mu_{\alpha'}$ when $\alpha < \alpha'$.

The objective of this paper is to show that the measures μ_{α} exhibit non-trivial percolation phase transition. Loosely speaking, we want to show that if α is close to zero then the set of 1's only contains finite connected components and if α is close to one then the set of 1's contains an infinite component. Let us explain this concept more precisely. We define the event $\text{Perc} \subseteq \{0,1\}^{\mathbb{Z}^d}$ which consists of those voter configurations ξ for which the subgraph of the nearest-neighbour lattice \mathbb{Z}^d spanned by the set of occupied sites $\{x:\xi(x)=1\}$ has an infinite connected component. By ergodicity, $\mu_{\alpha}(\text{Perc})$ is either 0 or 1. If Perc occurs, we say that the set $\{x:\xi(x)=1\}$ percolates. We can then define α_c as the supremum of all the values of α for which $\mu_{\alpha}(\text{Perc})=0$. By the stochastic ordering mentioned in the previous paragraph, $\mu_{\alpha}(\text{Perc})$ is non-decreasing in α . Thus for any $\alpha < \alpha_c$ we have $\mu_{\alpha}(\text{Perc})=0$ and for any $\alpha > \alpha_c$ we have $\mu_{\alpha}(\text{Perc})=1$. Our aim is to show that the family of measures $\{\mu_{\alpha}: 0 \leq \alpha \leq 1\}$ exhibits non-trivial percolation percolation phase transition, i.e., that $0 < \alpha_c < 1$. Our main results are

Theorem 1.1. If $d \ge 5$ and $R \ge 1$, then the family of stationary distributions of the voter model exhibits non-trivial percolation phase transition.

Theorem 1.2. If d = 3 or 4 then there exists $R_0 = R_0(d) \in \mathbb{N}$ such that if $R \geq R_0$ then the family of stationary distributions of the voter model exhibits non-trivial percolation phase transition.

1.2 Context

Although it may at first seem intuitively clear that, similarly to the case of Bernoulli percolation, ξ should be non-percolative if α is close to zero, this statement is not obvious. As the dynamics of the voter model favours that voters synchronize their opinions, the measures μ_{α} present long-range dependences. In fact, it follows from (3.6) below that for any $\alpha \in (0,1)$, the configuration ξ under law μ_{α} has covariances given by

$$c(\alpha, d, R) \cdot |x - y|^{2 - d} \le \operatorname{Cov}_{\mu_{\alpha}}(\xi(x), \xi(y)) \le C(\alpha, d, R) \cdot |x - y|^{2 - d}, \qquad x \ne y \in \mathbb{Z}^d. \tag{1.1}$$

It is a priori possible that percolation models with strong correlations present no phase transition. It is easy to build artificial examples, but let us recall an example that arises "naturally". The random interlacement set \mathcal{I}^u at level u > 0, introduced in [Sz10] is a random subset of \mathbb{Z}^d : (a) the law of \mathcal{I}^u is stochastically dominated by the law of \mathcal{I}^u when u < u', (b) the correlations of \mathcal{I}^u decay like (1.1) (see [Sz10, (1.68)]) and (c) the density of \mathcal{I}^u can be taken arbitrarily small by making u small (see [Sz10, (1.58)]), yet the set \mathcal{I}^u is connected for any u > 0, (see [Sz10, (2.21)]).

On the other hand, in case one attempts to prove that phase transition does occur, then the slowly decaying correlations (1.1) pose a challenge, as many of the well-known tools that are used for Bernoulli percolation are not applicable. Additionally, since general criteria are lacking and (as mentioned above) phase transition may in principle fail to occur, one needs to envisage strategies of proof that are model-specific. The proof of non-degeneracy of the percolation threshold has been carried out recently for the vacant set $\mathcal{V}^u = \mathbb{Z}^d \setminus \mathcal{I}^u$ of random interlacements in [Sz10, S10] and the excursion sets of the Gaussian free field in [RS13]. Both of these percolation models exhibit a decay of correlations described by (1.1).

In the case of the voter model, the question of percolation has been considered before, in [LS86], [BLM87], [LM06] and [Ma07]. The main focus of these works is in the case where d=3 and R=1. Through simulations and numerical studies, the first, third and fourth of these references argue that there should be a non-trivial phase transition and that the predictions of [HW83, W84] regarding the critical behaviour of percolation models with correlations described by (1.1) should be correct. However, the problem of finding a rigorous proof of the non-triviality of percolation phase transition of the stationary state of the voter model remained open. This problem is (partially) settled by our Theorems 1.1 and 1.2.

Another investigation of geometric properties of the stationary distribution of the voter model has recently been carried out in [HMN14]. The object of interest there is the voter model on a finite rhombus of the triangular lattice; the boundary of the rhombus, composed of four segments, is frozen so that two adjacent segments are always in state 0 and the other two in state 1. In this finite setting, there is only one stationary distribution, which can be constructed with the aid of coalescing random walks and the resulting coalescence classes, similarly to the μ_{α} 's on \mathbb{Z}^d . The authors study the volume of the coalescing classes and the interface curve that appears as a consequence of the opposing boundaries.

1.3 Ideas and structure of proof

Let us now explain how the paper is organized and also the contents of each section.

In Section 2, we give a notation summary and also collect some facts regarding martingales and random walks that are needed in the rest of the exposition.

Section 3 contains an introduction to the voter model on \mathbb{Z}^d , including its graphical construction, duality properties and the construction of the extremal stationary distributions using a family of coalescing random walks.

We begin to prove our main results in Section 4. Our goal is to show (see (4.1)) that for sufficiently small values of α , the probability that a large annulus is crossed by a *-connected path of 1's in ξ is smaller than a stretched exponential function of the radius of the annulus. The condition (4.1) is then shown to imply $0 < \alpha_c < 1$. It is self-evident that if (4.1) holds, then there is no percolation for small enough α . We also show, through a classical argument using planar duality, that (4.1) implies that if α is close enough to 1, then there is percolation.

We were able to establish (4.1) for the two sets of assumptions that appear in our main theorems (namely: first for $d \geq 5$, $R \geq 1$ and second for $d \geq 3$ and R large enough). We prove both cases using a renormalization scheme inspired by [Sz12, Section 2,3], which involves embeddings of binary trees into \mathbb{Z}^d that are "spread-out on all scales". In Section 4.1, we present this renormalization scheme and some of its properties.

In Section 5 we establish (4.1) for $d \ge 3$ and R large, and in Section 6 we establish it for $d \ge 5$ and $R \ge 1$. For simplicity of notation, Section 6 only treats explicitly $d \ge 5$ and R = 1 (i.e., the case of nearest neighbour interactions), but it will be easy to see that

the proof given there applies for any value of R. In fact, the proof of Section 6 could also be adapted to cover the case of $d \geq 3$ and R large enough, so that Section 5 is (strictly speaking) redundant. We have nevertheless chosen to include it for three reasons: first, because it is quite short; second, because the method might find other applications; and third, the contents of Section 5 may be helpful for the reader to grasp the more involved arguments of Section 6.

A common point in the proofs of Section 5 and 6 is the need to provide an upper bound for probabilities of the form

$$\mu_{\alpha} \left[\xi(x) = 1 \text{ for all } x \in \mathcal{X} \right]$$
 (1.2)

for certain finite sets $\mathcal{X} \subseteq \mathbb{Z}^d$ that appear at the "bottom" scale of the renormalization construction. An immediate consequence (as we will explain in Section 3, up to equation (3.5)) of the construction of μ_{α} through "coalescence classes" is that (1.2) is equal to $\mathbb{E}\left[\alpha^{\mathcal{N}_{\infty}(\mathcal{X})}\right]$, where $\mathcal{N}_{\infty}(\mathcal{X})$ is the (random) terminal number of random walkers in a system of coalescing random walks started from the configuration in which there is one walker in each vertex of \mathcal{X} . Hence, in order to give a good upper bound for (1.2), one needs to argue that $\mathcal{N}_{\infty}(\mathcal{X})$ is comparable to $|\mathcal{X}|$ (the cardinality of \mathcal{X}). It is worth noting that $\alpha^{|\mathcal{X}|}$ is the probability of the event in (1.2) for independent, Bernoulli(α) percolation.

Our renormalization construction ensures that the set \mathcal{X} under consideration here is "sparse on all scales". Hence, one expects that walkers started from the vertices of \mathcal{X} tend to avoid other walkers, and the amount of loss due to coalescence, $|\mathcal{X}| - \mathcal{N}_{\infty}(\mathcal{X})$, is far from $|\mathcal{X}|$ with overwhelming probability. In order to make this precise, we use different strategies in Sections 5 and 6.

- $(d \geq 3, R \gg 1)$ In Section 5, we replace the system of coalescing random walks with a system of annihilating random walks and observe that annihilation events are "negatively correlated". This allows us to derive a useful explicit bound on (1.2) which is particularly effective if the range R of the walkers is big enough to guarantee that the expected number of annihilations is sufficiently small.
- $(d \ge 5, R = 1)$ The proof of Section 6 involves two important ideas. First, it turns out that under some carefully constructed circumstances one can run the walkers for some period of time independently from each other (i.e., without coalescence), which allows them to "wander away" from each other before they start to coalesce. Second, we reveal the paths of random walkers one by one and pre-emptively throw away those future walks that are too likely to coalesce with the ones already revealed. We can then control
 - (a) the number of walkers that we throw away and
 - (b) the number of coalescences occurring between the remaining walkers

in such a way that the sum of these two numbers (which is greater than or equal to $|\mathcal{X}| - \mathcal{N}_{\infty}(\mathcal{X})$) is not too big compared to $|\mathcal{X}|$.

To state the obvious, Theorems 1.1 and 1.2 leave open the cases of dimension 3 and 4 and range R small, even though, as mentioned above, simulations and numerical work suggest that non-trivial phase transition should also occur in these cases. In our final and brief Section 7, we give an heuristic explanation to the ineffectiveness of the method of Section 6 in treating d = 3, 4 and R = 1. We also explain there why the method of Section 5 is insufficient to prove Theorem 1.1, so that we could not do without the more involved method of Section 6.

2 Notation and preliminary facts on martingales and random walks

2.1 Summary of notation

Given a set or event A, we denote by $\mathbb{1}_A$ its indicator function and by |A| its cardinality. Given a vertex $x \in \mathbb{Z}^d$, we denote by |x| its ℓ^{∞} norm and by $|x|_1$ its ℓ^1 norm. We then write

$$B(L) = \{x \in \mathbb{Z}^d : |x| \le L\}, \qquad B(x,L) = x + B(L);$$

$$B_1(L) = \{x \in \mathbb{Z}^d : |x|_1 \le L\}, \quad B_1(x,L) = x + B_1(L);$$

$$S(L) = \{x \in \mathbb{Z}^d : |x| = L\}, \qquad S(x,L) = x + S(L).$$
(2.1)

If for $x, y \in \mathbb{Z}^d$ we have $|x - y|_1 = 1$, then these points are said to be neighbors, and we abbreviate this by $x \sim y$. They are *-neighbors if |x - y| = 1. For sets $A, B \subset \mathbb{Z}^d$, $\operatorname{dist}(A, B) = \min\{|x - y| : x \in A, y \in B\}$. The expression $A \subset \mathbb{Z}^d$ indicates that A is a finite subset of \mathbb{Z}^d .

A nearest-neighbor path in \mathbb{Z}^d is a (finite or infinite) sequence $\gamma(0), \gamma(1), \ldots$ so that $\gamma(i+1) \sim \gamma(i)$ for each i. A *-connected path is a sequence $\gamma(0), \gamma(1), \ldots$ so that $\gamma(i+1)$ and $\gamma(i)$ are *-neighbors for each i.

Let $\xi \in \{0,1\}^{\mathbb{Z}^d}$. Given two disjoint sets $A, B \subset \mathbb{Z}^d$, we say A and B are connected by an open path in ξ (and write $A \stackrel{\xi}{\Leftrightarrow} B$) if there exists a nearest-neighbor path $\gamma(0), \ldots, \gamma(n)$ such that $\gamma(0)$ is the neighbor of a point of A, $\gamma(n)$ is a neighbor of a point of B and $\xi(\gamma(i)) = 1$ for each i. Similarly, we write $A \stackrel{*\xi}{\longleftrightarrow} B$ if there exists a *-connected path $\gamma(0), \ldots, \gamma(n)$ so that $\gamma(0)$ is the *-neighbor of a point of A, $\gamma(n)$ is the *-neighbor of a point of B and $\xi(\gamma(i)) = 1$ for each i.

2.2 Martingale facts

We will need a concentration inequality involving continuous-time martingales. We start recalling two definitions. Consider a probability space with a filtration $(\mathcal{F}_t)_{t>0}$.

Definition 2.1. A process $(X_t)_{t>0}$ is predictable with respect to (\mathcal{F}_t) if

$$X_t \in \mathcal{F}_{t-} = \sigma\left(\bigcup_{s < t} \mathcal{F}_s\right)$$
 for all t .

Note that if (X_t) is continuous and adapted to (\mathcal{F}_t) , then it is predictable with respect to (\mathcal{F}_t) .

Definition 2.2. Let $(N_t)_{t\geq 0}$ be a square-integrable càdlàg martingale with respect to $(\mathcal{F}_t)_{t\geq 0}$. The predictable quadratic variation of (N_t) is the predictable process $(\langle N \rangle_t)_{t\geq 0}$ such that $(N_t^2 - \langle N \rangle_t)_{t\geq 0}$ is a martingale with respect to (\mathcal{F}_t) .

The almost sure uniqueness of the predictable quadratic variation follows from Doob-Meyer-Doléans decomposition ([Kal, Theorem 25.5]) applied to the submartingale (N_t^2) . Note that $\langle N \rangle_t$ is a non-decreasing function of t. We refer the reader to [Kal, Proposition 26.1] for elementary properties of $\langle N \rangle$. The result we will need, which follows from [Kal, Theorem 26.17], is:

Theorem 2.3. Let $S \in [0, +\infty]$. Let (N_t) be a càdlàg martingale with $\langle N \rangle_S \leq \sigma^2$ almost surely for some $\sigma^2 \in (0, +\infty)$. Assume that the jumps of N are almost surely bounded by $\Delta \in (0, \sigma]$. Then we have

$$\mathbb{P}\left(\max_{t\in[0,S]} N_t - N_0 \ge r\right) \le \exp\left(-\frac{1}{2}\frac{r}{\Delta}\ln\left(1 + \frac{r\Delta}{\sigma^2}\right)\right), \qquad r \ge 0. \tag{2.2}$$

Note that [Kal, Theorem 26.17] is only stated for the $\sigma = 1$ case; however, our version (2.2) follows from an application of that theorem to the martingale N_t/σ .

2.3 Random walk facts

Definition 2.4. Given $R \in \mathbb{N}_+$, we say that $(X_t^z)_{t\geq 0}$ is an R-spread-out random walk on \mathbb{Z}^d starting at $z \in \mathbb{Z}^d$ if $X_0^z = z$ and $(X_t^z)_{t\geq 0}$ is a continuous-time càdlàg Markov process on \mathbb{Z}^d with infinitesimal generator

$$(\mathcal{L}f)(x) = \sum_{\substack{y \in \mathbb{Z}^d: \\ 0 < |x-y|_1 < R}} \frac{f(y) - f(x)}{|B_1(R)| - 1},$$

where $f: \mathbb{Z}^d \to \mathbb{R}$. When R = 1, then we call (X_t^z) a (continuous-time) nearest-neighbour simple random walk on \mathbb{Z}^d .

In words: the holding times between jumps are i.i.d. with Exp(1) distribution and if a jump occurs at time t and $X_{t-}^z = x$ then X_t^z is uniformly distributed on $B_1(x,R) \setminus \{x\}$. If R = 1, then X_t^z is uniformly distributed on the set of nearest neighbours of x.

Let us formulate a useful corollary of Theorem 2.3 about random walks:

Corollary 2.5. Let X_t denote a d-dimensional continuous-time nearest-neighbour simple random walk with jump rate 1 starting at the origin. Then for any $S, r \geq 0$ we have

$$\mathbb{P}\left[\max_{0 \le t \le S} |X_t| > r\right] \le 2d \exp\left(-\frac{1}{2}r \ln\left(1 + \frac{d \cdot r}{S}\right)\right). \tag{2.3}$$

Proof. The d coordinates of X_t are 1-dimensional simple random walks with jump rate 1/d, hence after a union bound we only need to apply (2.2) with $\sigma^2 = S/d$ and $\Delta = 1$ to achieve (2.3).

Let us define the transition kernel and the Green function of R-spread-out random walk on \mathbb{Z}^d by

$$p_{R,t}(x,y) = \mathbb{P}\left[X_{s+t}^z = y \,|\, X_s^z = x\right], \qquad g_R(x,y) = \int_0^\infty p_{R,t}(x,y) \,\mathrm{d}t.$$

If R = 1 then we drop the R from the subscript and simply denote $p_t(x, y)$ and g(x, y). We have

$$p_{R,t}(x,y) = p_{R,t}(y,x), \quad p_{R,t}(x,y) = p_{R,t}(y-x,0), q_R(x,y) = q_R(y,x), \quad q_R(x,y) = q_R(y-x,0), \quad q_R(x,x) \ge 1.$$
 (2.4)

It follows from the Chapman-Kolmogorov equations for $p_{R,t}(\cdot,\cdot)$ that we have

$$\sum_{y \in \mathbb{Z}^d} p_{R,T}(x,y) \cdot g_R(y,z) = \int_T^\infty p_{R,t}(x,z) \,\mathrm{d}t. \tag{2.5}$$

It follows from the Local Central Limit Theorem (see [L96, Section 1.2]) that for any $d \ge 3$ there exist constants c = c(d, R) > 0 and $C = C(d, R) < +\infty$ such that

$$\frac{\int_{T}^{\infty} p_{R,t}(x,y) \, \mathrm{d}t}{(|x-y| \vee \sqrt{T} + 1)^{2-d}} \in [c,C], \qquad x,y \in \mathbb{Z}^d, \quad T \ge 0.$$
 (2.6)

It follows from the strong Markov property of random walks that we have

$$P[\exists t \ge 0 : X_t^x = y] = \frac{g_R(x, y)}{g_R(y, y)} \stackrel{(2.4)}{\le} g_R(x, y). \tag{2.7}$$

The distributions of the increments of our random walks are symmetric, therefore if the random walks (X_t^x) and (X_t^y) are independent, then

$$(X_t^y - X_t^x)_{t \ge 0}$$
 has the same law as $(X_{2t}^{y-x})_{t \ge 0}$. (2.8)

Let us define

$$h_R(x,y) = P[\exists t \ge 0 : X_t^x = X_t^y], \quad x, y \in \mathbb{Z}^d$$
 (2.9)

the probability that two independent R-spread-out random walks started from x and y ever meet. We have

$$h_R(x,y) \stackrel{(2.8),(2.7),(2.4)}{=} \frac{g_R(x,y)}{g_R(0,0)} \le g_R(x,y).$$
 (2.10)

We will make use of the following claim about spread-out random walks:

Claim 2.6. Given $d \geq 3$, there exists $C < +\infty$ and $f : \mathbb{N} \to \mathbb{R}_+$ such that

$$\forall R \in \mathbb{N}, \ x \neq y \in \mathbb{Z}^d : \quad h_R(x, y) \le f(R) \wedge C|x - y|^{2 - d}, \qquad \lim_{R \to \infty} f(R) = 0. \tag{2.11}$$

The proof of this claim follows from (2.10), [HvdHS03, Proposition 1.6] and the observation that the Green function of a continuous-time random walk with jump rate 1 is identical to the Green function of the corresponding discrete-time random walk.

We will also make use of the following bound on the difference of Green function values of nearest neighbour sites: there exists a C = C(d) such that

$$|g(x,y) - g(x,y+e)| \le C \cdot (|x-y|+1)^{1-d}, \quad x,y \in \mathbb{Z}^d, \ e \sim 0.$$
 (2.12)

This bound follows from the much stronger [L96, Theorem 1.5.5].

The following heat kernel bound follows from the Local Central Limit Theorem: there exist $C = C(d) < +\infty$ and c = c(d) > 0 such that

$$p_t(x,y) \le Ct^{-\frac{d}{2}} \exp\left(-c\frac{|x-y|^2}{t}\right), \quad x,y \in \mathbb{Z}^d, \ t \ge 1.$$
 (2.13)

The bounds (2.12) and (2.13) together imply that there exists a C = C(d) such that

$$\sum_{w \in \mathbb{Z}^d} p_t(y, w) |g(v + e, w) - g(v, w)| \le C t^{\frac{1}{2} - \frac{d}{2}}, \qquad y, v \in \mathbb{Z}^d, \ t \ge 1.$$
 (2.14)

3 Voter model: graphical construction, duality, stationary distributions

We will now define the voter model on \mathbb{Z}^d and present some well-known facts about it. We refer the reader to [Li85] for an introduction to the voter model and proofs of all the statements that we make in this section.

Fix $d, R \in \mathbb{N}$. The voter model on \mathbb{Z}^d with range R, denoted by $(\xi_t)_{t\geq 0}$, is the Markov process with state space $\{0,1\}^{\mathbb{Z}^d}$ and infinitesimal generator given by

$$(\mathcal{L}f)(\xi) = \sum_{\substack{x,y \in \mathbb{Z}^d:\\0 < |x-y|_1 < R}} \frac{f(\xi^{y \to x}) - f(\xi)}{|B_1(R)| - 1},\tag{3.1}$$

where $f: \{0,1\}^{\mathbb{Z}^d} \to \mathbb{R}$ is any function that only depends on finitely many coordinates, $\xi \in \{0,1\}^{\mathbb{Z}^d}$ and

$$\xi^{y \to x}(z) = \begin{cases} \xi(z) & \text{if } z \neq x, \\ \xi(y) & \text{if } z = x. \end{cases}$$

In words, each site $x \in \mathbb{Z}^d$ updates its state $\xi(x)$ with rate 1 by uniformly choosing a site $y \in B_1(R) \setminus \{x\}$ and adopting the state $\xi(y)$ of y. In case R = 1, we say that the model is nearest-neighbour.

Given $\xi \in \{0,1\}^{\mathbb{Z}^d}$, we denote by P_{ξ} a probability measure under which $(\xi_t)_{t\geq 0}$ is defined and satisfies $P_{\xi}[\xi_0 = \xi] = 1$. Likewise, given a probability distribution ν on $\{0,1\}^{\mathbb{Z}^d}$, we write $P_{\nu} = \int P_{\xi} d\nu(\xi)$.

The process (ξ_t) satisfies a duality relation with respect to a system of coalescing random walks. We will now explain what is meant by this – or rather, we will give a particularly simple formulation of duality that will be sufficient for our purposes.

For each $x,y \in \mathbb{Z}^d$ with $0 < |x-y|_1 \le R$, let $(D_t^{(x,y)})_{t\ge 0}$ be a Poisson process with rate $(|B_1(R)|-1)^{-1}$ on $[0,\infty)$, so that $D_0^{(x,y)}=0$ and $D_t^{(x,y)}-D_{t-}^{(x,y)}$ is equal to 0 or 1 for all t. One pictures $D_t^{(x,y)}-D_{t-}^{(x,y)}=1$ as an arrow pointing from x to y at time t. We denote by $\mathbb P$ a probability measure under which all these processes are defined and are independent. For each $x \in \mathbb Z^d$, we then define (on this same probability space) $(Y_t^x)_{t\ge 0}$ as the unique $\mathbb Z^d$ -valued process which is right-continuous with left limits and satisfies

$$Y_0^x = x, Y_t^x = Y_{t-}^x + \sum_{z \in B_1(R)} z \cdot \left(D_t^{(Y_{t-}^x, Y_{t-}^x + z)} - D_{t-}^{(Y_{t-}^x, Y_{t-}^x + z)} \right). (3.2)$$

One pictures Y_t^x as it moves along the time axis and follows the arrows that it encounters. The collection of processes $\{(Y_t^x)_{t\geq 0}: x\in \mathbb{Z}^d\}$ is what we refer to as a system of coalescing random walks. This terminology makes sense because, as is clear from the above definition, each $(Y_t^x)_{t\geq 0}$ is a continuous-time random walk on \mathbb{Z}^d with rate 1 which jumps to a uniformly distributed location in $Y_{t-}^x + (B_1(R) \setminus \{0\})$ and moreover, these walks move independently until they meet, after which they coalesce and remain together.

Now, the duality equation for the voter model is, for any $A \subset\subset \mathbb{Z}^d$, $t \geq 0$ and probability measure ν on $\{0,1\}^{\mathbb{Z}^d}$,

$$P_{\nu}\left[\xi_{t}(x) = 1 \text{ for all } x \in A\right] = \int \mathbb{P}\left[\left\{Y_{t}^{x} : x \in A\right\} \subset \left\{y : \xi(y) = 1\right\}\right] \, d\nu(\xi). \tag{3.3}$$

Note that by inclusion-exclusion the equation (3.3) characterizes the distribution of ξ_t for the process started with distribution ν . Of particular interest is the case when ν is equal to

$$\pi_{\alpha} := (\alpha \delta_{\{1\}} + (1 - \alpha) \delta_{\{0\}})^{\otimes \mathbb{Z}^d},$$

the product measure of Bernoulli(α) on \mathbb{Z}^d , for $\alpha \in [0,1]$. In order to discuss this case, let us introduce some notation. For $A \subset \mathbb{Z}^d$, we let

$$\mathcal{N}_t(A) = |\{Y_t^x : x \in A\}|, \quad t \ge 0 \quad \text{ and } \quad \mathcal{N}_\infty(A) = \lim_{t \to \infty} \mathcal{N}_t(A);$$
 (3.4)

the limit exists because $\mathcal{N}_t(A)$ decreases with t. Denoting by \mathbb{E} the expectation operator associated with \mathbb{P} , we can then rewrite (3.3) as

$$P_{\pi_{\alpha}}\left[\xi_{t}(x) = 1 \text{ for all } x \in A\right] = \mathbb{E}\left[\alpha^{\mathcal{N}_{t}(A)}\right].$$

By taking the limit on the right-hand side as $t \to \infty$, we can conclude that, under $P_{\pi_{\alpha}}$, as $t \to \infty$, ξ_t converges in distribution to a measure μ_{α} on $\{0,1\}^{\mathbb{Z}^d}$ characterized by

$$\mu_{\alpha}\left[\xi(x) = 1 \text{ for all } x \in A\right] = \mathbb{E}\left[\alpha^{\mathcal{N}_{\infty}(A)}\right]$$
 (3.5)

for every finite $A \subset \mathbb{Z}^d$. The measures μ_{α} are invariant and ergodic with respect to translations on \mathbb{Z}^d and satisfy

$$\mu_{\alpha}[\xi(x) = 1] = \alpha, \qquad \operatorname{Corr}_{\mu_{\alpha}}(\xi(x), \xi(y)) \stackrel{(2.9)}{=} h_{R}(x, y), \qquad x, y \in \mathbb{Z}^{d},$$
 (3.6)

thus (1.1) indeed holds by (2.6) and (2.10).

As these measures are obtained as distributional limits of (ξ_t) , they are also stationary with respect to the dynamics of the voter model. In fact, in Section V.1 of [Li85] it is shown that

- if $d \ge 3$, then the set of extremal stationary distributions of the voter model is equal to $\{\mu_{\alpha} : \alpha \in [0,1]\}$. Here, a measure is said to be extremal if it cannot be written as a nontrivial convex combination of other stationary distributions.
- if d=1 or 2, then there are only two extremal stationary distributions, namely the point masses on the constant configurations $\xi \equiv 1$ and $\xi \equiv 0$. (If d=1 or 2, by recurrence of the random walk we have $\mathcal{N}_{\infty}(A) = 1$ almost surely for any finite and non-empty A. We can then see from (3.5) that μ_{α} is a convex combination with weight α of the point masses on the constant configurations).

Finally, we give a useful construction, jointly on the same probability space, of the system of coalescing random walks and for each $\alpha \in [0,1]$, a random $\xi^{(\alpha)} \in \{0,1\}^{\mathbb{Z}^d}$ distributed as μ_{α} . To this end, we take the probability space in which the aforementioned measure \mathbb{P} and the processes $\{(Y_t^x)_{t\geq 0}: x\in\mathbb{Z}^d\}$ are defined, and enlarge it so that a sequence of random variables $\{\mathcal{U}_n: n\in\mathbb{N}\}$, all independent and uniformly distributed on [0,1], are also defined (and are independent of the Y_t^x 's). Next, fix an arbitrary enumeration x_1, x_2, \ldots of \mathbb{Z}^d . For any $n\geq 1$, define the random variables

$$\eta(n) = \min\{m : m \le n \text{ and } Y_t^{x_m} = Y_t^{x_n} \text{ for some } t \ge 0\}$$
(3.7)

and then set

$$\xi^{(\alpha)}(x_n) = \mathbb{1}_{\{\mathcal{U}_{\eta(n)} \le \alpha\}}, \qquad n \in \mathbb{N}, \ \alpha \in [0, 1].$$
(3.8)

It is then straightforward to check that $\xi^{(\alpha)}$ has law μ_{α} , as defined in (3.5), and moreover it satisfies

$$\xi^{(\alpha)}(x) = \xi^{(\alpha)}(y)$$
 if $Y_t^x = Y_t^y$ for some t .

Moreover, it follows from this construction that if $\alpha \leq \alpha'$, then $\xi^{(\alpha)}(x) \leq \xi^{(\alpha')}(x)$ for each $x \in \mathbb{Z}^d$, therefore μ_{α} is stochastically dominated by $\mu_{\alpha'}$, that is, if $f: \{0,1\}^{\mathbb{Z}^d} \to \mathbb{R}$ is increasing (with respect to the partial order on $\{0,1\}^{\mathbb{Z}^d}$ that is induced by the order 0 < 1 on the coordinates), then

$$\int f \, \mathrm{d}\mu_{\alpha} \le \int f \, \mathrm{d}\mu_{\alpha'} \tag{3.9}$$

We will also need the following consequence of the joint construction:

the law of
$$1 - \xi$$
 under μ_{α} is the same as law of ξ under $\mu_{1-\alpha}$. (3.10)

4 First facts about voter model percolation, $d \geq 3$

Throughout this section, we fix $d \geq 3$, $R \geq 1$ (see (3.1)) and $\alpha \in [0,1]$. ξ denotes an element of $\{0,1\}^{\mathbb{Z}^d}$ and μ_{α} denotes the extremal stationary distribution of the voter model with density α , as described in Section 3.

We will show that there exist $\alpha_0 > 0$ and a sequence $(L_N)_{N \geq 0}$ of form $L_N = L \cdot \ell^N$ (where $\ell \geq 6$, $L \geq 1$, see (4.3)) such that

$$\mu_{\alpha_0} \left[B(L_N) \stackrel{*\xi}{\longleftrightarrow} B(2L_N)^c \right] \le 2^{-2^N}, \qquad N \ge 0.$$
 (4.1)

In words: the probability under μ_{α_0} that an annulus with inner radius L_N and outer radius $2L_N$ is crossed by a *-connected path of 1's in ξ is less than or equal to 2^{-2^N} . We will prove (4.1) for $d \geq 3$ and $R \gg 1$ in Section 5 and for $d \geq 5$ and R = 1 in Section 6. Let us now deduce the main results of this paper from (4.1).

Proof of Theorems 1.1 and 1.2. As we have already discussed in Section 3, the measure μ_{α} is invariant and ergodic under spatial shifts of \mathbb{Z}^d . Therefore the probability under μ_{α} of the event

$$Perc = \{\{x : \xi(x) = 1\} \text{ has an infinite connected component}\}$$
 (4.2)

can only be zero or one for any α . Also, since the event in (4.2) is increasing, by (3.9) there indeed exists $0 \le \alpha_c \le 1$ such that μ_{α} (Perc) = 0 for any $\alpha < \alpha_c$ and μ_{α} (Perc) = 1 for any $\alpha > \alpha_c$. Our aim is to to prove that $0 < \alpha_c < 1$.

Let us now briefly explain how (4.1) implies

$$\alpha_0 \leq \alpha_c \leq 1 - \alpha_0$$
.

As soon as we prove these inequalities, the statements of Theorems 1.1 and 1.2 will follow. First, since a nearest neighbour path is also a *-path, we have

$$\mu_{\alpha_0}[\operatorname{Perc}] \leq \lim_{N \to \infty} \mu_{\alpha_0} \left[B(L_N) \stackrel{*\xi}{\longleftrightarrow} B(2L_N)^c \right] \stackrel{(4.1)}{=} 0,$$

from which $\alpha_0 \leq \alpha_c$ follows.

Now we will prove $\alpha_c \leq 1 - \alpha_0$. Let us define the plane $F \subset \mathbb{Z}^d$ by

$$F = \{x = (x_1, \dots, x_d) \in \mathbb{Z}^d : x_i = 0 \text{ for all } i \ge 3\}.$$

For $\xi \in \{0,1\}^{\mathbb{Z}^d}$, we denote by $\bar{\xi}$ the restriction of ξ to F. By planar duality (see [K86, Lemma 2.23]) $B(L_N) \cap F$ is not connected to infinity by a nearest neighbour path of 1's in $\bar{\xi}$ if and only if $B(L_N) \cap F$ is surrounded by a *-connected cycle of 0's in $\bar{\xi}$. Now if the latter event occurs, then there exists $M \geq N$ and an $x \in (L_M \cdot \mathbb{Z}^d) \cap F$ with $|x| \leq 2L_{M+1}$ such that the annulus $B(x, 2L_M) \setminus B(x, L_M)$ is crossed by a *-connected path of 0's in $\bar{\xi}$. Thus for any $N \in \mathbb{N}$ we can bound

$$\mu_{1-\alpha_0}[\operatorname{Perc}^c] \leq \sum_{M=N}^{\infty} \sum_{\substack{x \in L_M \cdot \mathbb{Z}^d \\ |x| \leq 2L_{M+1}}} \mu_{1-\alpha_0} \left[B(x, L_M) \stackrel{*(1-\xi)}{\longleftrightarrow} B(x, 2L_M)^c \right] \stackrel{(3.10), (4.1)}{\leq}$$

$$\sum_{M=N}^{\infty} (4\ell + 1)^d \cdot 2^{-2^M},$$

from which $\mu_{1-\alpha_0}[\text{Perc}] = 1$ follows by letting $N \to \infty$. This implies $\alpha_c \le 1 - \alpha_0$. The proof of Theorems 1.1 and 1.2 is complete, given (4.1).

4.1 Renormalization scheme for percolation, $d \geq 3$

We will borrow the renormalization scheme of [Ra05]. We are going to use multi-scale renormalization with geometrically growing scales.

Let us fix $d \geq 3$. We let ℓ and L be two integers describing the scales of renormalization:

$$L_N = L \cdot \ell^N, \qquad N \ge 0. \tag{4.3}$$

Using these scales we define the renormalized lattices

$$\mathcal{L}_N = L_N \cdot \mathbb{Z}^d, \qquad N \ge 0. \tag{4.4}$$

We want to consider certain embeddings of binary trees on \mathbb{Z}^d . First let $T_{(k)} = \{1,2\}^k$ for $k \geq 0$ (in particular, $T_{(0)} = \varnothing$) and then let

$$T_N = \cup_{k=0}^N T_{(k)}$$

be the binary tree of height N. If $0 \le k < N$ and $m = (\eta_1, \dots, \eta_k) \in T_{(k)}$, we let

$$m_1 = (\eta_1, \dots, \eta_k, 1), \qquad m_2 = (\eta_1, \dots, \eta_k, 2)$$

be the two children of m in $T_{(k+1)}$

Definition 4.1. $\mathcal{T}: T_N \to \mathbb{Z}^d$ is a proper embedding of T_N if

- 1. $\mathcal{T}(\emptyset) = 0$;
- 2. for all $0 \le k \le N$ and $m \in T_{(k)}$ we have $\mathcal{T}(m) \in \mathcal{L}_{N-k}$;
- 3. for all $0 \le k < N$ and $m \in T_{(k)}$ we have

$$|\mathcal{T}(m_1) - \mathcal{T}(m)| = L_{N-k}, \qquad |\mathcal{T}(m_2) - \mathcal{T}(m)| = 2L_{N-k}.$$

We denote by Λ_N the set of proper embeddings of T_N into \mathbb{Z}^d .

We now collect a few facts from [Ra05] about these embeddings. Although the lemmas in [Ra05] that correspond to our Lemmas 4.2, 4.3 and 4.4 below are stated for $\ell = 6$, their statements hold true (and have the same proof) for any integer $\ell \geq 6$.

Lemma 4.2.

$$|\Lambda_N| = \left[((4\ell+1)^d - (4\ell-1)^d) \cdot ((2\ell+1)^d - (2\ell-1)^d) \right]^{2^N - 1}. \tag{4.5}$$

This follows from [Ra05, Lemma 3.2].

Next is the statement that, given a crossing of the L_N -scale annulus $B(2L_N)\setminus B(L_N)$, we can find a proper embedding $\mathcal{T} \in \Lambda_N$ so that all L_0 -scale annuli $B(\mathcal{T}(m), 2L_0) \setminus B(\mathcal{T}(m), L_0) : m \in T_{(N)}$ are crossed. Recall the notion of S(x, L) from (2.1).

Lemma 4.3. If γ is a *-connected path in \mathbb{Z}^d with

$$\{\gamma\} \cap S(L_N - 1) \neq \emptyset, \quad \{\gamma\} \cap S(2L_N) \neq \emptyset,$$

then there exists $\mathcal{T} \in \Lambda_N$ such that

$$\{\gamma\} \cap S(\mathcal{T}(m), L_0 - 1) \neq \emptyset \text{ and } \{\gamma\} \cap S(\mathcal{T}(m), 2L_0) \neq \emptyset \text{ for all } m \in T_{(N)}.$$

This is [Ra05, Lemma 3.3] (in fact, the statement given here corresponds to equation (3.7) in the proof of that lemma).

Finally, given a proper embedding $\mathcal{T} \in \Lambda_N$, the set of images of the leaves $\{\mathcal{T}(m) : m \in T_{(N)}\}$ is very sparse.

Lemma 4.4. For any $T \in \Lambda_N$ and any $m_0 \in T_{(N)}$, we have

$$\left| \left\{ m \in T_{(N)} : \text{dist} \left(B(\mathcal{T}(m_0), 2L), \ B(\mathcal{T}(m), 2L) \right) \le \ell^k L/2 \right\} \right| \le 2^{k-1}, \quad k \ge 1.$$
 (4.6)

This is a consequence of our assumption $\ell \geq 6$ and [Ra05, Lemma 3.4]. In particular, (4.6) implies that the sets $B(\mathcal{T}(m), 2L)$ for $m \in T_{(N)}$ are disjoint.

5 Spread-out model, $d \geq 3$

In this section we work with the voter model with range R, thus we will denote the stationary distribution (see (3.5)) with density α by $\mu_{R,\alpha}$. The goal of this section is to prove Theorem 1.2. More specifically, we will show that (4.1) holds for any $d \geq 3$ if $R \geq R_0(d)$ for some large R_0 and some $\alpha_0 = \alpha_0(d) > 0$.

Recall the notion of $h_R(x, y)$ from (2.9). The key lemma in our proof of Theorem 1.2 is as follows.

Lemma 5.1. For any $\mathcal{X} = \{x_1, \dots, x_{|\mathcal{X}|}\} \subset \mathbb{Z}^d$ we have

$$\mu_{R,\alpha}\left[\xi(x) = 1 \text{ for all } x \in \mathcal{X}\right] \le \alpha^{|\mathcal{X}|} \prod_{1 \le i < j \le |\mathcal{X}|} \left(1 + h_R(x_i, x_j) \left(\alpha^{-2} - 1\right)\right). \tag{5.1}$$

Before we prove Lemma 5.1, let us see how it allows us to conclude.

Proof of (4.1) for $d \ge 3$ and $R \gg 1$. We use the renormalization scheme described in Section 4.1. In this proof we choose $\ell = 6$ and L = 1 in (4.3). Given $\mathcal{T} \in \Lambda_N$, we denote

$$\mathcal{X}_{\mathcal{T}} = \bigcup_{m \in T_{(N)}} \{ \mathcal{T}(m) \}.$$

Combining Lemmas 4.2 and 4.3 in a union bound, we get, for any N,

$$\mu_{R,\alpha} \left[B(L_N) \stackrel{*\xi}{\longleftrightarrow} B(2L_N)^c \right] \le \widehat{C}^{2^N} \max_{\mathcal{T} \in \Lambda_N} \mu_{R,\alpha} \left[\xi(x) = 1 \text{ for all } x \in \mathcal{X}_{\mathcal{T}} \right].$$
 (5.2)

Now we fix some N and $\mathcal{T} \in \Lambda_N$ with the aim of bounding the probability on the right-hand side of (5.2). Note that by Lemma 4.4 we have $|\mathcal{X}_{\mathcal{T}}| = 2^N$. Let us denote $\mathcal{X}_{\mathcal{T}} = \{x_1, \ldots, x_{2^N}\}$.

$$\mu_{R,\alpha} \left[\xi(x) = 1 \text{ for all } x \in \mathcal{X}_{\mathcal{T}} \right]^{(5.1)} \leq \alpha^{2^N} \prod_{1 \le i < j \le 2^N} \left(1 + h_R(x_i, x_j) \alpha^{-2} \right) \le \alpha^{2^N} \prod_{i=1}^{2^N} \exp \left(\alpha^{-2} \sum_{j \ne i} h_R(x_i, x_j) \right)$$
(5.3)

Now for any $1 \le i \le 2^N$ let us bound

$$\sum_{j \neq i} h_R(x_i, x_j) \stackrel{(2.11)}{\leq} \sum_{j \neq i} f(R) \wedge C|x_i - x_j|^{2-d} \stackrel{(4.6)}{\leq} \sum_{k=1}^{\infty} 2^k \cdot \left(f(R) \wedge C \cdot \left(6^k/2 \right)^{2-d} \right), \tag{5.4}$$

thus if we choose $\alpha = \frac{1}{4}\widehat{C}^{-1}$ (with \widehat{C} as in (5.2)) then by $\lim_{R\to\infty} f(R) = 0$ (see (2.11)) and (5.4) we can choose R big enough so that for any $1 \le i \le 2^N$ we have

$$\alpha^{-2} \sum_{j \neq i} h_R(x_i, x_j) \le \ln(2),$$
(5.5)

which implies the desired bound (4.1):

$$\mu_{R,\alpha} \left[B(L_N) \stackrel{*\xi}{\longleftrightarrow} B(2L_N)^c \right] \stackrel{(5.2),(5.3)}{\le} \widehat{C}^{2^N} \alpha^{2^N} \prod_{i=1}^{2^N} \exp \left(\alpha^{-2} \sum_{j \neq i} h_R(x_i, x_j) \right)$$

$$\stackrel{(5.5)}{\le} \widehat{C}^{2^N} \left(\frac{1}{4} \widehat{C}^{-1} \right)^{2^N} 2^{2^N} = 2^{-2^N}.$$

The rest of this section is devoted to the proof of Lemma 5.1.

Recall the graphical construction of coalescing random walks Y_t^x , $x \in \mathbb{Z}^d$, $t \in \mathbb{R}_+$ defined on the probability space of the Poisson point processes $(D_t^{(x,y)})_{t\geq 0}$ from Section 3. Given $\mathcal{X} \subset\subset \mathbb{Z}^d$, denote by $\mathcal{X}_t = \{Y_t^x : x \in \mathcal{X}\}$, so that $\mathcal{N}_t(\mathcal{X}) = |\mathcal{X}_t|$. If $D_t^{(x,y)} - D_{t-}^{(x,y)} = 1$ for some $x \in \mathcal{X}_{t-}$, $y \in \mathbb{Z}^d$ and $t \in \mathbb{R}_+$, then the graphical construction (3.2) of coalescing random walks implies

$$\mathcal{X}_t = (\mathcal{X}_{t-} \setminus \{x\}) \cup \{y\}. \tag{5.6}$$

Let us introduce another set-valued stochastic process \mathcal{X}'_t , annihilating random walks, also defined on the probability space of the Poisson point processes $(D_t^{(x,y)})_{t\geq 0}$. Starting also from $\mathcal{X}'_0 := \mathcal{X}$ these particles also perform independent R-spread-out continuous-time random walks until one of the walkers tries to jump on a site occupied by another walker, in which case both of them disappear immediately. The formal definition is as follows. If $D_t^{(x,y)} - D_{t-}^{(x,y)} = 1$ for some $x \in \mathcal{X}'_{t-}, y \in \mathbb{Z}^d$ and $t \in \mathbb{R}_+$, then

$$\mathcal{X}_{t}' = \left(\mathcal{X}_{t-}' \setminus \{x\}\right) \Delta\{y\},\tag{5.7}$$

where $A\Delta B$ denotes the symmetric difference of the sets A and B. Let us denote by $\mathcal{N}'_t(\mathcal{X}) = |\mathcal{X}'_t|.$

Lemma 5.2. For any $\mathcal{X} \subset\subset \mathbb{Z}^d$, $\alpha \in [0,1]$, $R \in \mathbb{N}$ and $t \geq 0$ we have

$$\mathbb{E}\left[\alpha^{\mathcal{N}_t(\mathcal{X})}\right] \le \mathbb{E}\left[\alpha^{\mathcal{N}_t'(\mathcal{X})}\right]. \tag{5.8}$$

Proof. As soon as we show $\mathcal{X}'_t \subseteq \mathcal{X}_t$, the inequality (5.8) will immediately follow. Let us assume that $D_t^{(x,y)} - D_{t-}^{(x,y)} = 1$ for some $x \in \mathcal{X}_{t-}$ and that $\mathcal{X}'_{t-} \subseteq \mathcal{X}_{t-}$ holds. One can readily check using (5.6) and (5.7) that we also have $\mathcal{X}'_t \subseteq \mathcal{X}_t$ by considering the cases

(a)
$$x \in \mathcal{X}_{t-} \setminus \mathcal{X}'_{t-}$$
, (b) $x \in \mathcal{X}'_{t-}$, $y \notin \mathcal{X}'_{t-}$, (c) $x \in \mathcal{X}'_{t-}$, $y \in \mathcal{X}'_{t-}$

separately. Since $\mathcal{X}_0 = \mathcal{X}_0' = \mathcal{X}$, the inclusion $\mathcal{X}_t' \subseteq \mathcal{X}_t$ for all $t \in \mathbb{R}_+$ follows by induction.

Let us now give an alternative construction of \mathcal{X}'_t on a different probability space. Recall the notation $\mathcal{X} = \{x_1, \dots, x_{|\mathcal{X}|}\}$. Let X_t^i , $1 \leq i \leq |\mathcal{X}|$ denote independent Rspread-out random walks with $X_0^i = x_i$. For $1 \le i < j \le |\mathcal{X}|$, we denote

$$\tau(i,j) = \inf\{t : X_t^i = X_t^j\}. \tag{5.9}$$

We also define the set-valued stochastic process $\mathcal{I}_t \subseteq [1, \dots, |\mathcal{X}|]$ and the stopping times T_0, T_1, T_2, \ldots by letting $T_0 = 0$ and $\mathcal{I}_{T_0} = [1, \ldots, |\mathcal{X}|]$, and then inductively for $k \geq 1$ by

$$T_k := \inf\{ \tau(i,j) : i < j, i, j \in \mathcal{I}_{T_{k-1}} \}, T_k = \tau(i^*, j^*), \mathcal{I}_{T_k} = \mathcal{I}_{T_{k-1}} \setminus \{i^*, j^*\}.$$

In words, T_k is the time of the k'th annihilation and \mathcal{I}_{T_k} is the set of indices of those walkers that are still alive after the k'th annihilation. Of course if $T_k = +\infty$ for some $k \ge 1$ then we stop our inductive definition. We define $\mathcal{I}_t = \mathcal{I}_{T_{k-1}}$ for any $T_{k-1} \le t < T_k$.

Claim 5.3. The set-valued process $\mathcal{X}'_t = \{X^i_t : i \in \mathcal{I}_t\}$ has the same law as the annihilating walks described in (5.7).

The proof of this claim is straightforward and we omit it. From now on we will use this new definition of annihilating walks. For $1 \le i < j \le |\mathcal{X}|$ we also define the indicators

$$\eta_{i,j} = \mathbb{1}\left[\tau(i,j) < +\infty, \ \tau(i,j) = T_k \text{ for some } k\right],\tag{5.10}$$

thus $\eta_{i,j}$ is the indicator that the walkers indexed by i and j annihilate each other before any other walker annihilates either of them. Let us define

$$\mathcal{A}_{\infty}(\mathcal{X}) = \sum_{i=1}^{|\mathcal{X}|} \sum_{i=i+1}^{|\mathcal{X}|} \eta_{i,j}$$
(5.11)

the total number of annihilations that ever occurred. Now we have

$$\lim_{t \to \infty} \mathcal{N}'_t(\mathcal{X}) =: \mathcal{N}'_{\infty}(\mathcal{X}) = |\mathcal{X}| - 2\mathcal{A}_{\infty}(\mathcal{X}), \tag{5.12}$$

since each annihilation event kills two walkers. By (3.5), Lemma 5.2 and (5.12) we only need to prove

$$\mathbb{E}\left(\alpha^{-2\mathcal{A}_{\infty}(\mathcal{X})}\right) \leq \prod_{1 \leq i < j \leq |\mathcal{X}|} \left(1 + h_R(x_i, x_j)\left(\alpha^{-2} - 1\right)\right) \tag{5.13}$$

in order to complete the proof of Lemma 5.1. Let us introduce auxiliary Bernoulli random variables $\eta_{i,j}^*$, $1 \le i < j \le |\mathcal{X}|$ such that they are independent and

$$\mathbb{P}[\eta_{i,j}^* = 1] = 1 - \mathbb{P}[\eta_{i,j}^* = 0] \stackrel{(2.9)}{=} h_R(x_i, x_j) \stackrel{(5.9)}{=} \mathbb{P}[\tau(i,j) < +\infty]. \tag{5.14}$$

Similarly to (5.11), let us define

$$\mathcal{A}_{\infty}^{*}(\mathcal{X}) = \sum_{i=1}^{|\mathcal{X}|} \sum_{j=i+1}^{|\mathcal{X}|} \eta_{i,j}^{*}.$$
 (5.15)

Now the right-hand side of (5.13) is equal to $\mathbb{E}\left(\alpha^{-2\mathcal{A}_{\infty}^{*}(\mathcal{X})}\right)$, thus in order to prove (5.13) we only need to show that for any $\lambda \geq 0$ we have

$$\mathbb{E}\left[e^{\lambda \mathcal{A}_{\infty}(\mathcal{X})}\right] \le \mathbb{E}\left[e^{\lambda \mathcal{A}_{\infty}^{*}(\mathcal{X})}\right]. \tag{5.16}$$

By taking the Taylor expansion of the above exponential functions about $\lambda = 0$, we see that we only need to prove

$$\mathbb{E}\left[\left(\mathcal{A}_{\infty}(\mathcal{X})\right)^{k}\right] \leq \mathbb{E}\left[\left(\mathcal{A}_{\infty}^{*}(\mathcal{X})\right)^{k}\right], \qquad k \geq 0.$$

in order to achieve (5.16). By expanding the k'th power of the sums in the definitions of $\mathcal{A}_{\infty}(\mathcal{X})$ (see (5.11)) and $\mathcal{A}_{\infty}^{*}(\mathcal{X})$ (see (5.15)), we see that we only need to prove

$$\mathbb{P}\left[\eta_{i_1,j_1} = \dots = \eta_{i_k,j_k} = 1\right] \le \mathbb{P}\left[\eta_{i_1,j_1}^* = \dots = \eta_{i_k,j_k}^* = 1\right] \tag{5.17}$$

for any $k \geq 1$ and any $1 \leq i_l < j_l \leq |\mathcal{X}|$, $1 \leq l \leq k$. First, we may assume that the the list of pairs $\{i_1, j_1\}, \ldots, \{i_k, j_k\}$ does not contain the same pair more than once, because we can throw out such duplicates and reduce the value of k without changing the probabilities on either side of (5.17). Second, we may also assume that the sets $\{i_1, j_1\}, \ldots, \{i_k, j_k\}$ are disjoint, because if some of these sets have non-empty intersection, then the left-hand side of (5.17) is equal to zero by the definition of the indicators $\eta_{i,j}$ (see (5.10)): a walker can only be annihilated once. Now if the sets $\{i_1, j_1\}, \ldots, \{i_k, j_k\}$ are disjoint, then

$$\mathbb{P}\left[\eta_{i_{1},j_{1}} = \dots = \eta_{i_{k},j_{k}} = 1\right] \overset{(5.10)}{\leq} \mathbb{P}\left[\tau(i_{1},j_{1}) < +\infty, \dots, \tau(i_{k},j_{k}) < +\infty\right] \overset{(*)}{=} \prod_{l=1}^{k} \mathbb{P}\left[\tau(i_{l},j_{l}) < +\infty\right] \overset{(5.14)}{=} \prod_{l=1}^{k} \mathbb{P}\left[\eta_{i_{l},j_{l}}^{*} = 1\right] \overset{(**)}{=} \mathbb{P}\left[\eta_{i_{1},j_{1}}^{*} = \dots = \eta_{i_{k},j_{k}}^{*} = 1\right],$$

where (*) holds because the walkers X_t^i , $1 \le i \le |\mathcal{X}|$ are independent and the sets $\{i_1, j_1\}, \ldots, \{i_k, j_k\}$ are disjoint, and (**) holds because $\eta_{i,j}^*$, $1 \le i < j \le |\mathcal{X}|$ are independent. The proof of (5.17) and Lemma 5.1 is complete.

6 Nearest-neighbour model, $d \geq 5$

The goal of this section is to prove Theorem 1.1. More specifically, we will show that (4.1) holds for any $d \ge 5$ and R = 1 and some $\alpha_0 = \alpha_0(d) > 0$. Note that the same proof would work for any $R \ge 1$; the only reason we stick to the classical nearest-neighbour case is to ease notation. Also note that a slight generalization of the method presented in this section would yield a proof of Theorem 1.1 as well as Theorem 1.2, however we chose to also present in Section 5 a relatively short argument which only proves Theorem 1.2.

We use the graphical construction of $\xi^{(\alpha)}$ distributed as μ_{α} (see (3.8)), however we will often drop the dependence on α from our notation, especially if a particular calculation works for any $\alpha \in (0, 1)$.

We will use the renormalization scheme of Section 4.1. In order to specify the value of ℓ in (4.3) we fix the exponents

$$\varepsilon = \frac{1}{4d}, \qquad \delta = \frac{\varepsilon}{d}.$$
 (6.1)

The following choice of ℓ in (4.3) will be suitable for our purposes:

$$\ell = 3^{1/\delta}.\tag{6.2}$$

The choice of a large enough L in (4.3) will be specified later in Section 6.4. Choosing ℓ as in (6.2) we have

$$|\Lambda_N| \stackrel{(4.5)}{=} \le C^{2^N}$$
 for some $C = C(d)$. (6.3)

Combining (6.3) and Lemma 4.3 in a union bound, we get, for any N,

$$\mathbb{P}[B(L_N) \overset{*\xi}{\longleftrightarrow} B(2L_N)^c] \le C^{2^N} \max_{\mathcal{T} \in \Lambda_N} \mathbb{P}\left[\cap_{m \in T_{(N)}} \{B(\mathcal{T}(m), L) \overset{*\xi}{\longleftrightarrow} B(\mathcal{T}(m), 2L)^c \} \right]$$
(6.4)

We now fix N and a proper embedding $\mathcal{T} \in \Lambda_N$ with the aim of bounding the probability on the right-hand side in (6.4) (see (6.14) below). We recall the definition of ε from (6.1) and let

$$T = L^{2-\varepsilon} \tag{6.5}$$

and, for $x, y \in \mathbb{Z}^d$, we define the events

$$E_x = \left\{ \max_{0 \le t \le T} |Y_t^x - x| > \frac{1}{4}L \right\},\tag{6.6}$$

$$E_{x,y} = E_x^c \cap E_y^c \cap \{Y_t^x \neq Y_t^y, \ 0 \le t \le T\},$$
(6.7)

$$F_{x,y} = E_{x,y} \cap \{\xi(x) = \xi(y) = 1\}. \tag{6.8}$$

Remark 6.1. We defined $T \ll L^2$ because we want $\mathbb{P}[E_x] \ll 1$, see (6.16) and (6.40).

Claim 6.2. For any $z \in L\mathbb{Z}^d \stackrel{(4.4)}{=} \mathcal{L}_0$, the following inclusion holds:

$$\{B(z,L) \stackrel{*\xi}{\longleftrightarrow} B(z,2L)^c\} \subseteq \left(\bigcup_{x \in B(z,2L)} E_x\right) \cup \left(\bigcup_{\substack{x,y \in B(z,2L)\\|x-y|=1}} F_{x,y}\right)$$
(6.9)

Proof. Assume that the event on the left-hand side occurs. Then there exists a *-connected path (z_1, \ldots, z_k) with $|z_1| = L$, $|z_k| = 2L$ and $\xi(z_i) = 1$ for each i. For one such path, define

$$i^* = \max\{i \le k : Y_t^{z_i} = Y_t^{z_1} \text{ for some } t \le T\}.$$

If $i^* = k$, then $E_{z_1} \cup E_{z_k}$ occurs, since $|z_1 - z_k| \ge L$. If $i^* < k$, then the walks $(Y_t^{z_{i^*}})$ and $(Y_t^{z_{i^*+1}})$ do not meet before time T, so either $E_{z_{i^*}} \cup E_{z_{i^*+1}}$ or $F_{z_{i^*}, z_{i^*+1}}$ occurs.

With (6.9) in mind, given $\mathcal{T} \in \Lambda_N$ we choose two sets $\mathcal{X}, \mathcal{Y} \subset \mathbb{Z}^d$.

Definition 6.3. The pair $(\mathcal{X}, \mathcal{Y})$, $\mathcal{X}, \mathcal{Y} \subset \bigcup_{m \in T_{(N)}} B(\mathcal{T}(m), 2L)$ is called admissible if

- (i) for any $m \in T_{(N)}$, $(|B(\mathcal{T}(m), 2L) \cap \mathcal{X}|, |B(\mathcal{T}(m), 2L) \cap \mathcal{Y}|)$ is either (2,0) or (0,1);
- (ii) if $B(\mathcal{T}(m), 2L) \cap \mathcal{X} = \{x, y\}$, then |x y| = 1.

The set of all admissible pairs $(\mathcal{X}, \mathcal{Y})$ associated to \mathcal{T} is denoted $\mathcal{P}_{\mathcal{T}}$.

Lemma 6.4. Given $T \in \Lambda_N$,

1. For any $(\mathcal{X}, \mathcal{Y}) \in \mathcal{P}_{\mathcal{T}}$ we have

$$\frac{1}{2}|\mathcal{X}| + |\mathcal{Y}| = 2^{N}. (6.10)$$

2. There exists C = C(d) such that the number of admissible pairs can be bounded by

$$|\mathcal{P}_{\mathcal{T}}| \le (CL^d)^{2^N}. \tag{6.11}$$

3. We have

$$\bigcap_{m \in T_{(m)}} \{B(\mathcal{T}(m), L) \stackrel{*\xi}{\longleftrightarrow} B(\mathcal{T}(m), 2L)^c\} \subset \bigcup_{\substack{(\mathcal{X}, \mathcal{Y}) \in \mathcal{P}_{\mathcal{T}} \\ |x-z|=1}} \left(\bigcap_{\substack{\{x, z\} \in \mathcal{X} \\ |x-z|=1}} F_{x, z}\right) \cap \left(\bigcap_{y \in \mathcal{Y}} E_y\right).$$
(6.12)

4. For every $x \in \mathcal{X}$, we have

$$\left| \mathcal{X} \cap B(x, \ell^k L/2) \right| \le 2^k, \qquad k \ge 1.$$
 (6.13)

Proof. Given an admissible pair $(\mathcal{X}, \mathcal{Y})$ associated to \mathcal{T} , define

$$\mathcal{A}_{(\mathcal{X},\mathcal{Y})} = \{ m \in T_{(N)} : (|B(\mathcal{T}(m), 2L) \cap \mathcal{X}|, |B(\mathcal{T}(m), 2L) \cap \mathcal{Y}|) = (2, 0) \},$$

so that, by Definition 6.3 (i), we have

$$T_{(N)} \setminus \mathcal{A}_{(\mathcal{X},\mathcal{Y})} = \{ m \in T_{(N)} : (|B(\mathcal{T}(m), 2L) \cap \mathcal{X}|, |B(\mathcal{T}(m), 2L) \cap \mathcal{Y}|) = (0, 1) \}$$

and thus,

$$2^{N} = |T_{(N)}| = |\mathcal{A}_{(\mathcal{X},\mathcal{Y})}| + |T_{(N)} \setminus \mathcal{A}_{(\mathcal{X},\mathcal{Y})}| = \frac{1}{2}|\mathcal{X}| + |\mathcal{Y}|.$$

Additionally, by Definition 6.3, the pair $(\mathcal{X}, \mathcal{Y})$ is determined when we choose $\mathcal{A}_{(\mathcal{X}, \mathcal{Y})}$ and then, for each $m \in \mathcal{A}_{(\mathcal{X}, \mathcal{Y})}$, we choose two *-connected vertices in $B(\mathcal{T}(m), 2L)$ and for each $m \in \mathcal{T}_{(N)} \setminus \mathcal{A}_{(\mathcal{X}, \mathcal{Y})}$, we choose one vertex in $B(\mathcal{T}(m), 2L)$. Thus (6.11) indeed holds:

$$|\mathcal{P}_{\mathcal{T}}| \le \sum_{A \subseteq T(N)} (|B(2L)| \cdot 3^d)^{|A|} \cdot |B(2L)|^{2^N - |A|} \le (CL^d)^{2^N}.$$

The inclusion (6.12) is a consequence of (6.9).

The bound (6.13) follows from Lemma 4.4 and the fact that for each $m \in T_{(N)}$ we have $|\mathcal{X} \cap B(\mathcal{T}(m), 2L)| \leq 2$ by Definition 6.3 (i).

Putting together (6.4), (6.11) and (6.12), we obtain

$$\mathbb{P}[B(L_N) \overset{*\xi}{\longleftrightarrow} B(2L_N)^c] \\
\leq \left(CL^d\right)^{2^N} \max_{\substack{\mathcal{T} \in \Lambda_N \\ (\mathcal{X}, \mathcal{Y}) \in \mathcal{P}_{\mathcal{T}}}} \mathbb{P}\left[\left(\bigcap_{\substack{\{x, z\} \in \mathcal{X} \\ |x-z|=1}} F_{x, z}\right) \cap \left(\bigcap_{y \in \mathcal{Y}} E_y\right)\right] \tag{6.14}$$

for some constant C = C(d).

The main ingredient in the proof of (4.1) is the following proposition.

Proposition 6.5. For every $d \ge 5$, there exist $L_{(0)} \ge 2$ and $C = C(d) < +\infty$ such that for any $L \ge L_{(0)}$, any $\alpha \le L^{2-d+1/4}$ and any $N \ge 1$ we have

$$\max_{\substack{\mathcal{T} \in \Lambda_N \\ (\mathcal{X}, \mathcal{Y}) \in \mathcal{P}_{\mathcal{T}}}} \mathbb{P} \left[\left(\bigcap_{\substack{\{x, z\} \in \mathcal{X} \\ |x-z|=1}} F_{x, z} \right) \cap \left(\bigcap_{y \in \mathcal{Y}} E_y \right) \right] \leq \left(\mathcal{C}L^{4-2d+1/2} \right)^{2^N}. \tag{6.15}$$

Together with (6.14) and the assumption $d \ge 5$, this proposition immediately yields the desired result (4.1). We will explain why our method fails to prove (4.1) if d = 3, 4 and R = 1 in Remark 7.1.

The rest of this section is devoted to the proof of Proposition 6.5.

6.1 Reduction to delayed coalescing walks using the graphical construction

From now on, we fix not only $\mathcal{T} \in \Lambda_N$ (see Definition 4.1), but also $(\mathcal{X}, \mathcal{Y}) \in \mathcal{P}_{\mathcal{T}}$ (see Definition 6.3). Let us define

$$\beta = \beta(L, d) \stackrel{(6.6)}{=} \mathbb{P}[E_x] = \mathbb{P}[E_0].$$
 (6.16)

Lemma 6.6.

$$\mathbb{P}\left[\left(\bigcap_{\substack{\{x,z\}\in\mathcal{X}\\|x-z|=1}} F_{x,z}\right) \cap \left(\bigcap_{y\in\mathcal{Y}} E_y\right)\right] \leq \beta^{|\mathcal{Y}|} \cdot \mathbb{E}\left[\alpha^{\mathcal{N}_{\infty}(\mathcal{X})} \cdot \mathbb{1}_{\{\mathcal{N}_{T}(\mathcal{X})=|\mathcal{X}|\}}\right]$$
(6.17)

Remark 6.7. Recall the definition of $\mathcal{N}_{\infty}(\cdot)$ in (3.4). The event in the indicator on the right-hand side is simply the event that the walks started from the vertices of \mathcal{X} do not coalesce with each other before time T.

Proof. We will use the joint graphical construction of the system of coalescing walks and the configuration $\xi = \xi^{(\alpha)}$ described by equation (3.8). Since our set \mathcal{X} is fixed, we can and will assume that, in the enumeration of \mathbb{Z}^d that was needed for (3.7), the vertices in \mathcal{X} come before all other vertices of \mathbb{Z}^d . We can thus write

$$\mathcal{X} = \{x_1, x_2, \dots, x_{|\mathcal{X}|}\}. \tag{6.18}$$

We will now define some sigma-algebras involving the Poisson processes $(D_t^{(x,y)})$ that were used in the graphical construction of the coalescing random walks (Y_t^x) in (3.2) and

the uniform random variables (\mathcal{U}_n) that were used in (3.8). We let

$$\mathcal{G}_{y} = \sigma\left(D_{t}^{(z,w)}: z \sim w, |z - y| \leq L/4, t \leq T\right), \quad y \in \mathcal{Y};$$

$$\mathcal{G}_{\mathcal{X}} = \sigma\left(\begin{cases} D_{t}^{(z,w)}: z \sim w, \operatorname{dist}(\{z\}, \mathcal{X}) \leq L/4, t \leq T \\ \cup \left\{D_{t}^{(u,v)} - D_{T}^{(u,v)}: u \sim v, t > T \right\} \cup \left\{\mathcal{U}_{n}: 1 \leq n \leq |\mathcal{X}| \right\} \end{cases}\right).$$

Recalling (6.6)-(6.8) we note that

$$\left(\bigcap_{\substack{x,z\in\mathcal{X}\\|x-z|=1}}F_{x,z}\right)\in\mathcal{G}_{\mathcal{X}}\quad\text{and}\quad E_y\in\mathcal{G}_y\text{ for all }y\in\mathcal{Y}.$$

Using (4.6), we see that the sets

$$\{z \in \mathbb{Z}^d : |z - y| \le L/4\} \times [0, T], \qquad y \in \mathcal{Y}$$

are all disjoint, and also disjoint from

$$\left(\left\{z \in \mathbb{Z}^d : \operatorname{dist}(\left\{z\right\}, \mathcal{X}) \le L/4\right\} \times [0, T]\right) \cup \left(\mathbb{Z}^d \times (T, \infty)\right).$$

This implies that the sigma-algebras $\mathcal{G}_{\mathcal{X}}$ and \mathcal{G}_y for $y \in \mathcal{Y}$ are all independent. Consequently,

$$\mathbb{P}\left[\left(\bigcap_{\substack{x,z\in\mathcal{X}\\|x-z|=1}}F_{x,z}\right)\cap\left(\bigcap_{y\in\mathcal{Y}}E_{y}\right)\right]=\mathbb{P}\left[\bigcap_{\substack{x,z\in\mathcal{X}\\|x-z|=1}}F_{x,z}\right]\cdot\prod_{y\in\mathcal{Y}}\mathbb{P}\left[E_{y}\right]\stackrel{(6.16)}{=}\beta^{|\mathcal{Y}|}\cdot\mathbb{P}\left[\bigcap_{\substack{x,z\in\mathcal{X}\\|x-z|=1}}F_{x,z}\right].$$

We now define $\mathcal{M}_{\mathcal{X}} = \{\eta(x_k) : 1 \leq k \leq |\mathcal{X}|\}$, where η is defined in (3.7). For every non-empty $A \subseteq \{1, \dots, |\mathcal{X}|\}$ we have

$$\mathbb{P}\left[\left\{\mathcal{M}_{\mathcal{X}} = A\right\} \cap \bigcap_{\substack{x,z \in \mathcal{X} \\ |x-z|=1}} F_{x,z}\right] \stackrel{(3.8),(6.8)}{=} \alpha^{|A|} \cdot \mathbb{P}\left[\left\{\mathcal{M}_{\mathcal{X}} = A\right\} \cap \bigcap_{\substack{x,z \in \mathcal{X} : \\ |x-z|=1}} E_{x,z}\right]. \tag{6.19}$$

Note that, by (3.4), (3.7) and (6.18), we have $|\mathcal{M}_{\mathcal{X}}| = \mathcal{N}_{\infty}(\mathcal{X})$. Therefore

$$\mathbb{P}\left[\bigcap_{\substack{\{x,z\}\in\mathcal{X}\\|x-z|=1}} F_{x,z}\right] \stackrel{(6.19)}{=} \sum_{k=1}^{|\mathcal{X}|} \alpha^k \cdot \mathbb{P}\left[\left\{|\mathcal{M}_{\mathcal{X}}| = m\right\} \cap \bigcap_{\substack{x,z\in\mathcal{X}:\\|x-z|=1}} E_{x,z}\right] \\
= \mathbb{E}\left[\alpha^{\mathcal{N}_{\infty}(\mathcal{X})} \cdot \mathbb{1}\left\{\bigcap_{\substack{x,z\in\mathcal{X}:\\|x-z|=1}} E_{x,z}\right\}\right] \stackrel{(4.6),(6.6),(6.7)}{\leq} \mathbb{E}\left[\alpha^{\mathcal{N}_{\infty}(\mathcal{X})} \cdot \mathbb{1}_{\left\{\mathcal{N}_{T}(\mathcal{X})=|\mathcal{X}|\right\}}\right].$$

The proof of Lemma 6.6 is complete.

6.2 Reduction to independent random walks

We now turn to the bound on the expectation on the right-hand side of (6.17). We will need to take a close look at the coalescing walks $\{(Y_t^x)_{t\geq 0}: x\in \mathcal{X}\}$. For this, it will no longer be convenient to work with the graphical construction of the coalescing walks using Poisson processes $(D_t^{(x,y)})$ that we described in Section 3. Rather, we will switch to a new probability space, in which we will give a different representation of the system of coalescing walks.

The following construction will depend on the set \mathcal{X} which has been fixed at the beginning of Section 6.1 and also on the enumeration of \mathcal{X} that was fixed in (6.18). Let P denote a probability measure under which one defines a collection of processes $\{(X_t^x)_{t\geq 0}: x\in \mathcal{X}\}$ satisfying:

- for each $x \in \mathcal{X}$, $(X_t^x)_{t \geq 0}$ is a continuous-time, nearest neighbor random walk on \mathbb{Z}^d with jump rate 1 and $X_0^x = x$;
- these walks are all independent.

(We emphasize that this is *not* a system of coalescing walks). The expectation operator associated to P is denoted by E. We then define the processes:

• $\{(W_t^x)_{t\geq 0}: x\in \mathcal{X}\}$. They are defined by induction. Put $W_t^{x_1}=X_t^{x_1}$ for all t. Assume W^{x_1},\ldots,W^{x_n} are defined and let

$$\sigma = \inf\{t : X_t^{x_{n+1}} = W_t^{x_k} \text{ for some } k \le n\}.$$

On $\{\sigma=\infty\}$, let $W_t^{x_{n+1}}=X_t^{x_{n+1}}$ for all t. On $\{\sigma<\infty\}$, let K be the smallest index such that $X_{\sigma}^{x_{n+1}}=W_{\sigma}^{x_K}$. Put

$$W_t^{x_{n+1}} = \left\{ \begin{array}{ll} X_t^{x_{n+1}} & \text{if } t \leq \sigma; \\ W_t^{x_K} & \text{if } t > \sigma. \end{array} \right.$$

• $\{(Z_t^x)_{t\geq 0}: x\in \mathcal{X}\}$. These are defined exactly as above, with the only difference that in the induction step, σ is defined by

$$\sigma = \inf\{t \ge T : X_t^{x_{n+1}} = Z_t^{x_k} \text{ for some } k \le n\}.$$

Claim 6.8. (i) $\{(W_t^x)_{t\geq 0}: x\in \mathcal{X}\}\$ is a system of coalescing walks started from \mathcal{X} ; in particular, its law under P is the same as that of $\{(Y_t^x)_{t\geq 0}: x\in \mathcal{X}\}\$ under \mathbb{P} .

(ii) $\{(Z^x)_{t\geq 0}: x\in \mathcal{X}\}$ is a system of random walks that move independently (with no coalescence) up to time T and after time T, behave as a system of coalescing walks.

The proof of this claim is straightforward and we omit it.

Similarly to (3.4) we also define

$$\begin{split} \mathcal{N}_t^W &= |\{W_t^x : x \in \mathcal{X}\}|, \qquad \mathcal{N}_\infty^W = \lim_{t \to \infty} \mathcal{N}_t^W \\ \mathcal{N}_t^Z &= |\{Z_t^x : x \in \mathcal{X}\}|, \qquad \mathcal{N}_\infty^Z = \lim_{t \to \infty} \mathcal{N}_t^Z. \end{split}$$

We now have

$$\mathbb{E}\left[\alpha^{\mathcal{N}_{\infty}(\mathcal{X})} \cdot \mathbb{1}_{\{\mathcal{N}_{T}(\mathcal{X})=|\mathcal{X}|\}}\right] = E\left[\alpha^{\mathcal{N}_{\infty}^{W}} \cdot \mathbb{1}_{\{X_{t}^{x} \neq X_{t}^{y} \text{ for all } x, y \in \mathcal{X}, x \neq y \text{ and } t \leq T\}}\right]$$

$$= E\left[\alpha^{\mathcal{N}_{\infty}^{Z}} \cdot \mathbb{1}_{\{X_{t}^{x} \neq X_{t}^{y} \text{ for all } x, y \in \mathcal{X}, x \neq y \text{ and } t \leq T\}}\right] \leq E\left[\alpha^{\mathcal{N}_{\infty}^{Z}}\right]. \tag{6.20}$$

At this point one might be tempted to apply Lemma 5.2, i.e., to switch from coalescing to annihilating walks. In Remark 7.2 we explain why the method of Section 5 cannot be used to prove Theorem 1.1.

A stochastic domination result

In this subsection we state the ingredients of the proof of Proposition 6.5. Given these ingredients, we will prove Proposition 6.5 in Section 6.4. In Sections 6.5, 6.6 and 6.7 we prove the ingredient results.

Recall the notion of the enumeration $\mathcal{X} = \{x_1, x_2, \dots, x_{|\mathcal{X}|}\}$ from (6.18). Let us define

$$U_n := \mathbb{1}\{\exists \ k < n, \ t \ge T : \ X_t^{x_n} = X_t^{x_k}\}, \quad 1 \le n \le |\mathcal{X}|, \qquad U = \sum_{n=1}^{|\mathcal{X}|} U_n. \tag{6.21}$$

In words: U_n is the indicator of the event that the n'th walker hits any of the previous walkers after T. Recalling the construction of Section 6.2 we have

$$\mathcal{N}_{\infty}^{Z} = |\mathcal{X}| - \sum_{n=1}^{|\mathcal{X}|} \mathbb{1}\{\exists \ k < n, \ t \ge T \ : \ Z_{t}^{x_{n}} = Z_{t}^{x_{k}}\} \ge |\mathcal{X}| - U$$

and we can thus bound

$$E\left[\alpha^{\mathcal{N}_{\infty}^{Z}}\right] \le \alpha^{|\mathcal{X}|} \cdot E\left[\alpha^{-U}\right] \tag{6.22}$$

Let us now describe the main ideas of this subsection. The indicator variables U_n , $1 \le n \le |\mathcal{X}|$ are not independent; however, in Proposition 6.13 we will argue that their sum can be dominated by a sum of independent variables. Let us explain now the heuristics for this domination. Suppose we reveal the paths $(X_t^{x_n})_{t>0}, 1 \leq n \leq |\mathcal{X}|$ one by one, starting with $(X_t^{x_1})_{t\geq 0}$. We think of each path n as a trial: a success if it avoids all the previously revealed paths after time T (that is, if $U_n = 0$), and a failure otherwise. At the time of revealing path n, it should have a high probability of being a success (since the set $\{X_T^x: x \in \mathcal{X}\}\$ is very sparse), unless some path of index k < n behaved in an atypical manner that makes it exceptionally likely that $(X_t^{x_n})_{t\geq T}$ meets $(X_t^{x_k})_{t\geq T}$. In (6.27) below we will introduce the variable $V_{k,n}$ as the indicator of this event that path k endangers trial n. We then rely on two fundamental observations. First (see Lemma 6.11): since the random set $\{X_T^x: x \in \mathcal{X}\}$ is very sparse (as suggested by (6.13)), it is very unlikely that a path endangers a trial, so that the random variables $V_k = \sum_{n>k} V_{k,n}$, which represent the number of trials endangered by each path k, are equal to zero with high probability. Second (see Lemma 6.12): if trial n is not endangered by any path of index k < n, then it is very likely to be successful.

For any $x, y \in \mathbb{Z}^d$ let us define the random variable

$$M_{\infty}^{x,y,T} = \mathbb{P}[\exists s \ge T : X_s^y = X_s^x \mid X_u^x : 0 \le u < \infty]. \tag{6.23}$$

(the reason for the ∞ symbol in $M_{\infty}^{x,y,T}$ will become clear in Section 6.7). Recall the definition of $\varepsilon = \varepsilon(d)$ and $\delta = \delta(d)$ from (6.1).

Proposition 6.9. There exists $T_0 = T_0(d) < +\infty$ and $D_0 = D_0(d) < +\infty$ such that

$$P[M_{\infty}^{x,y,T} > T^{1-\frac{d}{2}+\varepsilon}] \le e^{-T^{\delta}}, \qquad x, y \in \mathbb{Z}^{d}, \qquad T \ge T_{0}, \quad (6.24)$$

$$P[M_{\infty}^{x,y,T} > |x-y|^{2-d+\varepsilon}] \le e^{-|x-y|^{\delta}}, \qquad x, y \in \mathbb{Z}^{d}, |x-y| \ge D_{0}, \qquad T \ge 0. \quad (6.25)$$

$$P[M_{\infty}^{x,y,T} > |x-y|^{2-d+\varepsilon}] \le e^{-|x-y|^{\delta}}, \qquad x,y \in \mathbb{Z}^d, |x-y| \ge D_0, \qquad T \ge 0.$$
 (6.25)

Remark 6.10. By (2.6) we have $E\left[M_{\infty}^{x,y,T}\right] \simeq T^{1-d/2} \wedge |x-y|^{2-d}$, thus (6.24) are (6.25) are bounds on the probability that the random variable $M_{\infty}^{x,y,T}$ deviates too much from its expectation.

The proof of Proposition 6.9 is postponed to Section 6.7.

We now fix T_0 and D_0 as in Proposition 6.9. Given these choices, we may then assume that the renormalization constant L satisfies

$$T \stackrel{(6.5)}{=} L^{2-\varepsilon} \ge T_0, \qquad L \ge D_0.$$
 (6.26)

We define for $1 \le k < n \le |\mathcal{X}|$ the random variables

$$V_{k,n} = \begin{cases} \mathbb{1}\left\{M_{\infty}^{x_k, x_n, T} > T^{1 - \frac{d}{2} + \varepsilon}\right\}, & \text{if } |x_k - x_n| = 1, \\ \mathbb{1}\left\{M_{\infty}^{x_k, x_n, T} > |x_n - x_k|^{2 - d + \varepsilon}\right\}, & \text{otherwise.} \end{cases}$$
(6.27)

In words: $V_{k,n}$ is the indicator of the event that $(X_t^{x_k})_{t\geq 0}$ endangers $(X_t^{x_n})_{t\geq 0}$. We also define

$$\underline{V}_k = (V_{k,k+1}, \dots, V_{k,|\mathcal{X}|}), \qquad V_k = \sum_{n=k+1}^{|\mathcal{X}|} V_{k,n}. \tag{6.28}$$

Now by (6.23) and (6.27), for any $1 \le k < n \le |\mathcal{X}|$

$$V_{k,n}$$
 is measurable with respect to $\sigma(X_t^{x_k}: t \ge 0),$ (6.29)

therefore

$$\underline{V}_1, \dots, \underline{V}_{|\mathcal{X}|}$$
 are independent, (6.30)

$$V_1, \dots, V_{|\mathcal{X}|}$$
 are independent. (6.31)

Now by (6.28) for any $1 \leq n \leq |\mathcal{X}|$ the random variable V_n is the number of trajectories that the trajectory $(X_t^{x_n})$ endangers. We will "throw away" those trajectories that are endangered by another trajectory, and it is going to be convenient to also throw away $(X_t^{x_n})$ itself if it endangers any other trajectory. Thus the number of trajectories that we will throw away because of the behaviour of $(X_t^{x_n})$ is equal to $V_n + \mathbb{1}\{V_n > 0\}$.

Lemma 6.11. If $e^{-T^{\delta}} + \sum_{k=1}^{\infty} 2^k \cdot e^{-(\frac{1}{2}\ell^k L)^{\delta}} \leq 1$ then for any $n \in \{1, \dots, |\mathcal{X}|\}$ the random variable $V_n + \mathbb{1}\{V_n > 0\}$ is stochastically dominated by a random variable V_0^* with probability mass function $\mathsf{p}_{V_0^*}$ supported on the set of integers $\{0, 2, 3, 5, \dots, 2^k + 1, \dots\}$ and given by

$$\mathsf{p}_{V_0^*}(2) = e^{-T^\delta}, \quad \mathsf{p}_{V_0^*}(2^k + 1) = 2^k \cdot e^{-(\frac{1}{2}\ell^k L)^\delta}, \ k \ge 1, \quad \mathsf{p}_{V_0^*}(0) = 1 - \sum_{k > 0} \mathsf{p}_{V_0^*}(k). \ \ (6.32)$$

In particular,

$$P[V_n > 0] \le P[V_0^* > 0] = e^{-T^{\delta}} + \sum_{k=1}^{\infty} 2^k \cdot e^{-(\frac{1}{2}\ell^k L)^{\delta}}.$$
 (6.33)

The proof of Lemma 6.11 is postponed until Section 6.5.

Recall the definition of U_n from (6.21). In words, the next lemma states that if a path is not endangered by any of the previous paths, then it is very likely to avoid all of them.

Lemma 6.12. For any $n \in \{1, ..., |\mathcal{X}|\}$,

$$P[U_{n} = 1 \mid X_{t}^{x_{k}} : 1 \leq k < n, \ t \geq 0] \cdot \mathbb{1} \left\{ \sum_{k=1}^{n-1} V_{k,n} = 0 \right\}$$

$$\leq T^{1-\frac{d}{2}+\varepsilon} + \sum_{k=1}^{\infty} 2^{k} \left(\frac{1}{2} \ell^{k} L \right)^{2-d+\varepsilon}. \tag{6.34}$$

The proof of Lemma 6.12 is postponed until Section 6.5.

In order to state the following proposition, and for the sake of clarity, we recapitulate some relevant definitions:

- U_n (for $1 \le n \le |\mathcal{X}|$) and U in (6.21);
- $V_{k,n}$ (for $1 \le k < n \le |\mathcal{X}|$), \underline{V}_n and V_n (for $1 \le n \le |\mathcal{X}|$) in (6.27) and (6.28);
- V_0^* in Lemma 6.11.

We add to this list one more definition; let

$$p := p(L, d) = \frac{T^{1 - \frac{d}{2} + \varepsilon} + \sum_{k=1}^{\infty} 2^k \left(\frac{1}{2} \ell^k L\right)^{2 - d + \varepsilon}}{P[V_0^* = 0]}.$$
 (6.35)

Proposition 6.13. Let $U^* \sim \text{Bin}(|\mathcal{X}|, p)$ and let V^* be independent from U^* , where V^* is the sum of $|\mathcal{X}|$ i.i.d. copies of V_0^* . Then

$$U$$
 is stochastically dominated by $U^* + V^*$. (6.36)

Remark 6.14. If the n'th path is not endangered by previous paths then the parameter of the Bernoulli variable U_n is bounded by the right-hand side of (6.34). The indicators U_n , $1 \le n \le |\mathcal{X}|$ are not independent, but we can dominate their correlations by slightly increasing the parameters of these indicators and by adding V^* .

We postpone the proof of Proposition 6.13 until Section 6.6.

6.4 Proof of Proposition 6.5

By (3.8) the left-hand side of (6.15) decreases with α , so it is enough to prove (6.15) for

$$\alpha = L^{2-d+\frac{1}{4}}. (6.37)$$

Remark 6.15. Let us comment about the choice of α . For the sake of this heuristic argument let us assume that $\mathcal{Y} = \emptyset$ in (6.39) below, so that $|\mathcal{X}| = 2 \cdot 2^N$, see (6.10). Comparing the combinatorial complexity term $(CL^d)^{2^N}$ of (6.14) with the terms

$$\alpha^{|\mathcal{X}|} \cdot e^{|\mathcal{X}|p/\alpha} = \left(\alpha^2 e^{2p/\alpha}\right)^{2^N}$$

in (6.43) below, we see that if we want $\mathbb{P}[B(L_N) \stackrel{*\xi}{\longleftrightarrow} B(2L_N)^c] \ll 1$ then it is a good idea to choose α so that

$$\alpha^2 L^d \ll 1, \qquad p/\alpha = \mathcal{O}(1).$$
 (6.38)

Now p is not much bigger than L^{2-d} (see (6.47) below), so if $d \ge 5$, then (6.37) is a good choice if we want α to satisfy the bounds (6.38).

Let us fix $\mathcal{T} \in \Lambda_N$ (see Definition 4.1) and $(\mathcal{X}, \mathcal{Y}) \in \mathcal{P}_{\mathcal{T}}$ (see Definition 6.3). We have

$$\mathbb{P}\left[\left(\bigcap_{\substack{\{x,z\}\in\mathcal{X}\\|x-z|=1}}F_{x,z}\right)\cap\left(\bigcap_{y\in\mathcal{Y}}E_{y}\right)\right]\stackrel{(6.17),(6.20)}{\leq}\beta^{|\mathcal{Y}|}E\left[\alpha^{\mathcal{N}_{\infty}^{Z}}\right]\stackrel{(6.22),(6.36)}{\leq}$$

$$\beta^{|\mathcal{Y}|}\cdot\alpha^{|\mathcal{X}|}\cdot E\left[\left(\frac{1}{\alpha}\right)^{U^{*}}\right]\cdot E\left[\left(\frac{1}{\alpha}\right)^{V^{*}}\right]. \quad (6.39)$$

Now we bound the terms on the right-hand side of (6.39).

$$\beta \stackrel{(6.6),(6.16)}{=} \mathbb{P}\left[\max_{0 \le t \le T} |Y_t^0| > \frac{1}{4}L\right] \stackrel{(2.3),(6.5)}{\le} 2d \exp\left(-\frac{1}{8}L\ln\left(1 + \frac{d}{4}L^{-1+\varepsilon}\right)\right). \tag{6.40}$$

Recall from Proposition 6.13 that $U^* \sim \text{Bin}(|\mathcal{X}|, p)$, where p = p(L, d) was defined in (6.35). For a random variable $Z \sim \text{Bin}(m, r)$ and $\theta \geq 0$, we have $E[\theta^Z] \leq e^{mr\theta}$, thus

$$E\left[\left(\frac{1}{\alpha}\right)^{U^*}\right] \le e^{|\mathcal{X}|p/\alpha}.\tag{6.41}$$

Recall from Proposition 6.13 that V^* is the sum of $|\mathcal{X}|$ independent copies of V_0^* .

$$E\left[\left(\frac{1}{\alpha}\right)^{V_0^*}\right] \stackrel{(6.32)}{=} \mathsf{p}_{V_0^*}(0) + \frac{e^{-T^{\delta}}}{\alpha^2} + \sum_{k=1}^{\infty} \left(\frac{1}{\alpha}\right)^{2^k + 1} 2^k e^{-(\frac{1}{2}\ell^k L)^{\delta}} \stackrel{(6.5),(6.37)}{=}$$

$$\mathsf{p}_{V_0^*}(0) + L^{2d - \frac{9}{2}} \cdot e^{-L^{(2-\varepsilon)\delta}} + \sum_{k=1}^{\infty} \exp\left((2^k + 1)(d - \frac{9}{4})\ln(L) + k\ln(2) - \frac{1}{2^{\delta}}(\ell^{\delta})^k L^{\delta}\right)$$

$$=: q \stackrel{(*)}{=} q(L, d), \quad (6.42)$$

where in (*) the parameter q is indeed only a function of L and d, because of the definition of ε and δ in (6.1) and ℓ in (6.2). We can thus bound

$$\mathbb{P}\left[\left(\bigcap_{\substack{\{x,z\}\in\mathcal{X}\\|x-z|=1}}F_{x,z}\right)\cap\left(\bigcap_{y\in\mathcal{Y}}E_{y}\right)\right] \stackrel{(6.39),(6.41),(6.42)}{\leq} \beta^{|\mathcal{Y}|}\cdot\alpha^{|\mathcal{X}|}\cdot e^{|\mathcal{X}|p/\alpha}\cdot q^{|\mathcal{X}|} = \exp\left\{|\mathcal{Y}|\ln\beta + \frac{1}{2}|\mathcal{X}|(2\ln q + 2\ln\alpha + 2p/\alpha)\right\}. \quad (6.43)$$

Recall our definition of ℓ from (6.2). We will choose L big enough so that it satisfies multiple criteria, as we now discuss. By (6.26) we need

$$L > L_{(1)} := T_0^{\frac{1}{2-\varepsilon}} \vee D_0.$$

Having already fixed ε , δ and ℓ , we assume that L satisfies

$$L \ge L_{(2)}$$
, so that $\exp\left(-L^{(2-\varepsilon)\delta}\right) + \sum_{k=1}^{\infty} 2^k \cdot \exp\left(-\left(\frac{1}{2}\ell^k L\right)^{\delta}\right) \le \frac{1}{2}$, (6.44)

so that the condition of Lemma 6.11 is satisfied for L. We will also assume

$$L \ge L_{(3)}, \text{ so that } q(L, d) \stackrel{(**)}{\le} 2.$$
 (6.45)

The inequality (**) can be achieved because $p_{V_0^*}(0) \leq 1$ (see (6.32)) and by our choice of ℓ in (6.2) we have $(\ell^{\delta})^k \geq 3^k$, thus the sum of the other terms in the definition (6.42) of q can be made arbitrarily small by making L large. Next we observe that

$$p(L,d) \le 4L^{2-d+\frac{1}{4}} \stackrel{(6.37)}{=} 4\alpha \quad \text{if} \quad L \ge L_{(2)}.$$
 (6.46)

To show that this inequality indeed holds, we estimate

$$p = p(L,d) \stackrel{(6.5),(6.35)}{=} \frac{L^{(2-\varepsilon)(1-\frac{d}{2}+\varepsilon)} + \sum_{k=1}^{\infty} 2^k \left(\frac{1}{2}\ell^k L\right)^{2-d+\varepsilon}}{P[V_0^* = 0]} \stackrel{(6.1)}{\leq} \frac{L^{2-d+\frac{1}{4}} + L^{2-d+\varepsilon} \sum_{k=1}^{\infty} 2^k \left(\frac{1}{2}\ell^k\right)^{2-d+\varepsilon}}{P[V_0^* = 0]} \stackrel{(6.2),(6.33),(6.44)}{\leq} \frac{2\left(L^{2-d+\frac{1}{4}} + L^{2-d+\varepsilon}\right) \stackrel{(6.1)}{\leq} 4L^{2-d+\frac{1}{4}}. \quad (6.47)$$

We can now bound the expression in the exponential in the right-hand side of (6.43):

$$\begin{split} |\mathcal{Y}|\ln(\beta) + \frac{|\mathcal{X}|}{2} (2\ln q + \frac{p}{\alpha} + 2\ln \alpha) &\overset{(6.45),(6.46),}{\leq} |\mathcal{Y}|\ln(\beta) + \frac{|\mathcal{X}|}{2} (2\ln 2 + 4 + 2\ln \alpha) &\overset{(6.37),(6.40)}{\leq} \\ |\mathcal{Y}| \left(\ln(2d) - \frac{1}{8}L\ln\left(1 + \frac{d}{4}L^{-1+\varepsilon}\right)\right) + \frac{|\mathcal{X}|}{2} \left(\widehat{\mathcal{C}} + \left(4 - 2d + \frac{1}{2}\right)\ln L\right) &\overset{(*)}{\leq} \\ \left(|\mathcal{Y}| + \frac{|\mathcal{X}|}{2}\right) \left(\widehat{\mathcal{C}} + \left(4 - 2d + \frac{1}{2}\right)\ln L\right) &\overset{(6.10)}{=} 2^{N} \cdot \left(\widehat{\mathcal{C}} + \left(4 - 2d + \frac{1}{2}\right)\ln L\right), \quad (6.48) \end{split}$$

where (*) holds for $L \ge L_{(4)}$. Plugging (6.48) back in (6.43), we obtain that the statement of Proposition 6.5 holds with ℓ as in (6.2) and $L_{(0)} := L_{(1)} \lor \cdots \lor L_{(4)}$.

6.5 Proof of Lemmas 6.11 and 6.12

We now prove the two lemmas of Section 6.3 bounding the probability that random walk paths endanger (Lemma 6.11) and intersect (Lemma 6.12) each other. These proofs simply put together results that have already been established. For Lemma 6.11, we combine Proposition 6.9 – which bounds the probability that a path endangers another path that starts at a given distance from it – with (6.13) – which bounds the number of points of \mathcal{X} that are within a given distance from a fixed point $x \in \mathcal{X}$. Lemma 6.12 is even simpler and follows from a combination of (6.13) with the definition of "endangering" in (6.27).

Proof of Lemma 6.11. Fix $n \in \{1, ..., |\mathcal{X}|\}$. We take a bijection

$$\theta: \{0, 1, \dots, |\mathcal{X}| - n\} \to \{n, n + 1, \dots, |\mathcal{X}|\}$$

with the property that

$$0 = |x_{\theta(0)} - x_n| \le |x_{\theta(1)} - x_n| \le \dots \le |x_{\theta(|\mathcal{X}| - n)} - x_n|.$$

We have

$$|\{i \ge n : |x_i - x_n| \le \ell^k L/2\}| \le |\mathcal{X} \cap B(x_n, \ell^k L/2)| \stackrel{(6.13)}{\le} 2^k, \quad k \ge 1,$$

so that

$$|x_{\theta(i)} - x_n| > \ell^k L/2$$
 for all $i \ge 2^k$, $k \ge 1$.

By our definition of ℓ (see (6.1),(6.2)) and L (see (6.26)) we have $\ell^k L/2 \geq D_0$, for any $k \geq 1$, moreover $T \geq T_0$ (see (6.26)), therefore we can use Proposition 6.9 to bound the probability of the event in the indicator $V_{n,\theta(i)}$ (see (6.27)) that trajectory n endangers trajectory $\theta(i)$:

$$P\left[V_{n,\theta(i)} = 1\right] \le \begin{cases} e^{-T^{\delta}} & \text{if } i = 1; \\ e^{-(\ell^{k}L/2)^{\delta}} & \text{if } i \ge 2^{k}, \ k \ge 1. \end{cases}$$
 (6.49)

Now, if $i \geq 2$, we have

$$P[V_n \ge i] \le P[V_n \ge 2^{\lfloor \log_2 i \rfloor}] \le \sum_{j \ge 2^{\lfloor \log_2 i \rfloor}} P[V_{n,\theta(j)} = 1] \stackrel{(6.49)}{\le} \sum_{k = \lfloor \log_2 i \rfloor}^{\infty} 2^k \cdot e^{-(\ell^k L/2)^{\delta}}$$

and similarly,

$$P[V_n \ge 1] \le e^{-T^{\delta}} + \sum_{k=1}^{\infty} 2^k \cdot e^{-(\ell^k L/2)^{\delta}}.$$

We then obtain

$$P[V_n + 1 \{V_n > 0\} \ge 1] = P[V_n + 1 \{V_n > 0\} \ge 2] \le e^{-T^{\delta}} + \sum_{k=1}^{\infty} 2^k \cdot e^{-(\ell^k L/2)^{\delta}}$$

and, for i > 2,

$$P[V_n + 1\{V_n > 0\} \ge i] \le P[V_n \ge i - 1] \le \sum_{k = \lfloor \log_2(i-1) \rfloor}^{\infty} 2^k \cdot e^{-(\ell^k L/2)^{\delta}}.$$

The statement of the lemma now follows from comparing these inequalities with the definition of the law of V_0^* in (6.32).

Proof of Lemma 6.12. We have

$$\begin{split} P\left[U_{n} = 1 | \, X_{t}^{x_{k}} : 1 \leq k < n, \, \, t \geq 0 \right] \\ & \leq \sum_{m=1}^{(6.21)} P\left[\exists s \geq T \, : \, X_{s}^{x_{n}} = X_{s}^{x_{m}} | \, X_{t}^{x_{k}} : 1 \leq k < n, \, \, t \geq 0 \right] \\ & = \sum_{m=1}^{n-1} P\left[\exists s \geq T \, : \, X_{s}^{x_{n}} = X_{s}^{x_{m}} | \, X_{t}^{x_{m}} : \, \, t \geq 0 \right] \stackrel{(6.23)}{=} \sum_{m=1}^{n-1} M_{\infty}^{x_{m}, x_{n}, T}, \end{split}$$

so that

$$P\left[U_{n} = 1 \mid X_{t}^{x_{k}} : 1 \leq k < n, \ t \geq 0\right] \cdot \mathbb{1}\left\{\sum_{k=1}^{n-1} V_{k,n} = 0\right\} \leq \sum_{m=1}^{n-1} M_{\infty}^{x_{m}, x_{n}, T} \cdot \mathbb{1}\left\{V_{m,n} = 0\right\}.$$

Now, by (6.27),

$$M_{\infty}^{x_m, x_n, T} \cdot \mathbb{1}\left\{V_{m,n} = 0\right\} \le \begin{cases} T^{1 - \frac{d}{2} + \varepsilon} & \text{if } |x_m - x_n| = 1; \\ |x_m - x_n|^{2 - d + \varepsilon} & \text{otherwise.} \end{cases}$$

The proof of (6.34) can now be completed by applying Definition 6.3 and (6.13) as we did in the proof of Lemma 6.11; we omit the details.

6.6 Proof of Proposition 6.13

In this section we will prove our stochastic domination result using a coupling argument. The key idea lies in the definition of some auxiliary random variables U_n^* , $1 \le n \le |\mathcal{X}|$, so let us start by explaining this informally (the precise definition is given in (6.64)). Define the events

$$A_n = \left\{ \sum_{k=1}^{n-1} V_{k,n} = 0, \ V_n = 0 \right\}, \quad 1 \le n \le |\mathcal{X}|. \tag{6.50}$$

In words: A_n is the event that the n'th random walk path is not endangered by previous paths and does not endanger upcoming paths. We will specify the key properties of $U_n^*, 1 \le n \le |\mathcal{X}|$ using the events A_n in (6.51) and (6.52) below. Suppose we fix n and we reveal all the paths $\{X_t^{x_i}: i < n, t \ge 0\}$, and moreover we reveal the vector \underline{V}_n (defined in (6.28)). Given all this information, we are able to determine whether or not A_n has occurred. Now,

- (a) assume A_n has occurred. At this point, we have full knowledge of all the paths with index smaller than n, and also some partial knowledge of the n'th path: we know \underline{V}_n , in fact we know that A_n occurred, which implies $V_n = \sum_{k=n+1}^{|\mathcal{X}|} V_{n,k} = 0$. In Lemma 6.17, we argue that the conditional probability of $\{U_n = 1\}$ given all this information is at most p (defined in (6.35)), a number that is not much larger than the bound we had given in (6.34) (which did not include the conditioning on $\{V_n = 0\}$). We are thus able to define U_n^* so that $U_n \leq U_n^*$ and $U_n^* \sim \text{Bernoulli}(p)$.
- (b) if A_n has not occurred, we simply prescribe (using extra, auxiliary randomness) that U_n^* is Bernoulli(p).

The sum $\sum_{n=1}^{|\mathcal{X}|} U_n \cdot \mathbb{1}_{A_n}$ is then dominated by $\sum_{n=1}^{|\mathcal{X}|} U_n^* \cdot \mathbb{1}_{A_n} \leq \sum_{n=1}^{|\mathcal{X}|} U_n^*$, and the sum $\sum_{n=1}^{|\mathcal{X}|} U_n \cdot \mathbb{1}_{A_n^c}$ is dominated by $\sum_{n=1}^{|\mathcal{X}|} (V_n + \mathbb{1}\{V_n > 0\})$ (see (6.53) below), which in turn is dominated by $\sum_{n=1}^{|\mathcal{X}|} V_n^*$, a sum of i.i.d. random variables distributed as V_0^* from Lemma 6.11. Finally, the desired independence properties of our construction follow from the fact that the distribution of U_n^* is the same regardless of the conditioning; this is formalized in Lemma 6.18.

Proof of Proposition 6.13. In a series of lemmas we will construct, by extending the probability space of the random walks $(X_t^{x_n}), 1 \leq n \leq |\mathcal{X}|$, random variables $U_1^*, \ldots, U_{|\mathcal{X}|}^*$ satisfying

$$U_n \cdot \mathbb{1}_{A_n} \le U_n^* \cdot \mathbb{1}_{A_n}; \tag{6.51}$$

$$U_n^* \sim \text{Ber}(p)$$
 and is independent of $((U_k^*)_{1 \le k \le n-1}, (\underline{V}_k)_{1 \le k \le |\mathcal{X}|})$. (6.52)

Here we show how this construction implies (6.36). We let $U^* = \sum_{n=1}^{|\mathcal{X}|} U_n^*$. We have

$$U \stackrel{(6.21)}{=} \sum_{n=1}^{|\mathcal{X}|} U_n \stackrel{(6.51)}{\leq} \sum_{n=1}^{|\mathcal{X}|} (U_n^* + \mathbb{1}_{A_n^c}) \stackrel{(6.50)}{\leq} U^* + \sum_{n=1}^{|\mathcal{X}|} \left(\mathbb{1}_{\{V_n > 0\}} + \sum_{k=1}^{n-1} V_{k,n} \right)$$

$$= U^* + \sum_{n=1}^{|\mathcal{X}|} \mathbb{1}_{\{V_n > 0\}} + \sum_{k=1}^{|\mathcal{X}|} \sum_{n=k+1}^{|\mathcal{X}|} V_{k,n} \stackrel{(6.28)}{=} U^* + \sum_{k=1}^{|\mathcal{X}|} (V_k + \mathbb{1}_{\{V_k > 0\}}). \quad (6.53)$$

Now (6.52) implies that $U^* \sim \text{Bin}(|\mathcal{X}|, p)$ and is independent of $V_1, \ldots, V_{|\mathcal{X}|}$, which are also independent by (6.31). Putting this together with Lemma 6.11 we obtain that

$$U^* + \sum_{n=1}^{|\mathcal{X}|} (\mathbb{1}_{\{V_n > 0\}} + V_n)$$
 is stochastically dominated by $U^* + V^*$,

where V^* is a sum of $|\mathcal{X}|$ independent copies of V_0^* . This completes the proof of Proposition 6.13 given (6.51) and (6.52).

The rest of this subsection is devoted to the construction of random variables $U_1^*, \ldots, U_{|\mathcal{X}|}^*$ satisfying (6.51) and (6.52). We start recalling a few standard facts about conditional expectations.

Lemma 6.16. Let $(\Omega, \mathcal{F}, \mathsf{P})$ be a probability space.

1. [Wi91, Section 9.7, Property (k)] If X is an \mathcal{F} -measurable and bounded random variable (r.v.), $\mathcal{G}, \mathcal{G}' \subset \mathcal{F}$ are sigma-algebras and \mathcal{G}' is independent of $\sigma(\mathcal{G} \cup \sigma(X))$, then

$$\mathsf{E}[X \mid \mathcal{G}, \mathcal{G}'] = \mathsf{E}[X \mid \mathcal{G}]. \tag{6.54}$$

2. [Kl05, Theorem 2.24] Let $\mathcal{H} \subseteq \mathcal{F}$ be a sigma-algebra, Z be an \mathcal{F} -measurable r.v. independent of \mathcal{H} , Y be an \mathcal{H} -measurable r.v., and $f: \mathbb{R}^2 \to \mathbb{R}$ be Borel-measurable and bounded. If we define $g(y) := \mathsf{E}[f(y,Z)]$ for $y \in \mathbb{R}$ then

$$\mathsf{E}\left[f(Y,Z)\,|\,\mathcal{H}\right] = g(Y). \tag{6.55}$$

We now extend the probability space of the walks $X_t^x, t \geq 0, x \in \mathcal{X}$ with an independent collection of auxiliary random variables

$$\zeta_k$$
, $1 \le k \le |\mathcal{X}|$, i.i.d. and unifolly distributed on [0,1]. (6.56)

For $1 \leq n \leq |\mathcal{X}|$, we introduce the sigma-field

$$\sigma_{n} := \sigma\left(\left(\zeta_{k} : 1 \leq k \leq n\right), \ \left(\underline{V}_{k} : 1 \leq k \leq |\mathcal{X}|\right), \ \left(X_{t}^{x_{k}} : t \geq 0, \ 1 \leq k \leq n\right)\right)
\stackrel{(6.29)}{=} \sigma\left(\left(\zeta_{k} : 1 \leq k \leq n\right), \ \left(\underline{V}_{k} : n < k \leq |\mathcal{X}|\right), \ \left(X_{t}^{x_{k}} : t \geq 0, \ 1 \leq k \leq n\right)\right).$$
(6.57)

Recalling (6.21) we also define the random variable

$$p_{n} := P\left[U_{n} = 1 \mid \sigma_{n-1}\right] = P\left[U_{n} = 1 \mid \frac{(\zeta_{k})_{1 \le k \le n-1}, \ (\underline{V}_{k})_{n \le k \le |\mathcal{X}|},}{X_{t}^{x_{k}}, t \ge 0, 1 \le k \le n-1}\right].$$
(6.58)

Lemma 6.17. The number p defined in (6.35), the event A_n , $1 \le n \le |\mathcal{X}|$ defined in (6.50) and the random variable p_n defined in (6.58) satisfy

$$p_n \cdot \mathbb{1}_{A_n} \le p. \tag{6.59}$$

Proof. By (6.29), (6.54) and the fact that $(X_t^{x_k})$, $1 \le k \le |\mathcal{X}|$ are independent random walks, we have

$$p_n = P[U_n = 1 \mid \underline{V}_n, X_t^{x_k}, t \ge 0, k \le n - 1].$$

Recall from (6.28) that V_n is the indicator of the event that the path $(X_t^{x_n})$ does not endanger any upcoming paths. We now claim that

$$p_n \cdot \mathbb{1} \left\{ V_n = 0 \right\} = \frac{P\left[U_n = 1, \ V_n = 0 \mid X_t^{x_k}, t \ge 0, \ k \le n - 1 \right]}{P\left[V_n = 0 \mid X_t^{x_k}, t \ge 0, \ k \le n - 1 \right]} \cdot \mathbb{1} \left\{ V_n = 0 \right\}. \tag{6.60}$$

Before we prove this, let us see how it allows us to conclude. Noting that

$$P[V_n = 0 \mid X_t^{x_k}, t \ge 0, \ k \le n-1] \stackrel{(6.29),(6.54)}{=} P[V_n = 0],$$
 (6.61)

we have

$$\begin{split} p_n \cdot \mathbbm{1}_{A_n} & \stackrel{(6.50)}{=} p_n \cdot \mathbbm{1} \left\{ V_n = 0 \right\} \mathbbm{1} \left\{ \sum_{k=1}^{n-1} V_{k,n} = 0 \right\} \\ & \stackrel{(6.60),(6.61)}{\leq} \frac{P\left[U_n = 1 \mid X_t^{x_k}, t \geq 0, \ k \leq n-1 \right] \cdot \mathbbm{1} \left\{ \sum_{k=1}^{n-1} V_{k,n} = 0 \right\}}{P[V_n = 0]} \cdot \mathbbm{1} \left\{ V_n = 0 \right\}. \end{split}$$

By applying Lemma 6.12 to the numerator and Lemma 6.11 to the denominator, we conclude that the right-hand side is smaller than p, thus (6.59) holds.

It remains to prove (6.60). To this end, we abbreviate

$$\mathcal{V} = \sigma(V_n), \qquad \mathcal{G} = \sigma(X_t^{x_k}: t \ge 0, \ 1 \le k \le n-1).$$

We must then prove that

$$\mathbb{1}_{\{V_n=0\}} \cdot P[V_n=0|\mathcal{G}] \cdot P[U_n=1|\mathcal{G},\mathcal{V}] = \mathbb{1}_{\{V_n=0\}} \cdot P[U_n=1,\ V_n=0|\mathcal{G}];$$

since V_n is \mathcal{V} -measurable and $P[V_n = 0|\mathcal{G}]$ is \mathcal{G} -measurable, this is the same as

$$E\left[\mathbb{1}_{\{U_n=1, V_n=0\}} \cdot P[V_n=0|\mathcal{G}] \mid \mathcal{G}, \mathcal{V}\right] = \mathbb{1}_{\{V_n=0\}} \cdot P[U_n=1, V_n=0|\mathcal{G}]. \tag{6.62}$$

We now check that the right-hand side of (6.62) satisfies the definition of the left-hand side. First, note that $\mathbb{1}_{\{V_n=0\}} \cdot P[U_n=1, V_n=0|\mathcal{G}]$ is measurable with respect to $\sigma(\mathcal{G}, \mathcal{V})$. Second, for any event $C \in \sigma(\mathcal{G}, \mathcal{V})$, we must check that

$$E\left[\mathbb{1}_{C} \cdot \mathbb{1}_{\{U_{n}=1, V_{n}=0\}} \cdot P[V_{n}=0|\mathcal{G}]\right] = E\left[\mathbb{1}_{C} \cdot \mathbb{1}_{\{V_{n}=0\}} \cdot P[U_{n}=1, V_{n}=0|\mathcal{G}]\right].$$
(6.63)

Now \mathcal{V} is an atomic sigma-algebra (since it is generated by finitely many events, see (6.28)) and $\{V_n = 0\}$ is an atom of \mathcal{V} , therefore the event $C \cap \{V_n = 0\}$ is equal to $G \cap \{V_n = 0\}$ for some $G \in \mathcal{G}$. Using this, (6.63) is equivalent to

$$E\left[\mathbb{1}_{G} \cdot \mathbb{1}_{\{U_{n}=1, V_{n}=0\}} \cdot P[V_{n}=0|\mathcal{G}]\right] = E[\mathbb{1}_{\{V_{n}=0\}} \cdot E[\mathbb{1}_{G} \cdot \mathbb{1}_{\{U_{n}=1, V_{n}=0\}}|\mathcal{G}]].$$

By taking $E[\cdot|\mathcal{G}]$ inside the expectation, we see that both sides are equal to

$$E\left[E[\mathbb{1}_G \cdot \mathbb{1}_{\{U_n=1, V_n=0\}} | \mathcal{G}] \cdot P[V_n=0 | \mathcal{G}]\right].$$

The proof of Lemma 6.17 is complete.

We are now ready to define

$$U_n^* := \mathbb{1}_{A_n} \cdot \left(U_n + (1 - U_n) \cdot \mathbb{1} \left\{ \zeta_n \le \frac{p - p_n}{1 - p_n} \right\} \right) + \mathbb{1}_{A_n^c} \cdot \mathbb{1} \{ \zeta_n \le p \}.$$
 (6.64)

Lemma 6.18. U_n^* satisfies (6.51) and (6.52).

Proof. That (6.51) is satisfied is obvious, so we turn to (6.52).

Recalling the definitions of U_n from (6.21), \underline{V}_k from (6.28), A_n from (6.50), ζ_k from (6.56), σ_n from (6.57) and p_n from (6.58) we note that

$$A_n$$
, U_n , ζ_n and p_n are all σ_n -measurable, $1 \leq n \leq |\mathcal{X}|$.

Consequently, U_1^*, \ldots, U_{n-1}^* are all σ_{n-1} -measurable. Since $(\underline{V}_k)_{n \leq k \leq |\mathcal{X}|}$ are also σ_{n-1} -measurable, we see that (6.52) will follow once we show that

$$\mathbb{E}[U_n^* \mid \sigma_{n-1}] = p. \tag{6.65}$$

We start with

$$\begin{split} E\left[U_{n}^{*}\mid\sigma_{n-1}\right] &\overset{(6.64)}{=} E\left[U_{n}\cdot\mathbbm{1}_{A_{n}}\mid\sigma_{n-1}\right] + E\left[\left(1-U_{n}\right)\cdot\mathbbm{1}_{A_{n}}\cdot\mathbbm{1}_{\left\{\zeta_{n}\leq\frac{p-p_{n}}{1-p_{n}}\right\}}\mid\sigma_{n-1}\right] \\ &\quad + P\left[A_{n}^{c}\cap\left\{\zeta_{n}\leq p\right\}\mid\sigma_{n-1}\right] \\ &\overset{(*)}{=} \mathbbm{1}_{A_{n}}\cdot E\left[U_{n}\mid\sigma_{n-1}\right] + \mathbbm{1}_{A_{n}}\cdot E\left[\left(1-U_{n}\right)\cdot\mathbbm{1}_{\left\{\zeta_{n}\leq\frac{p-p_{n}}{1-p_{n}}\right\}}\mid\sigma_{n-1}\right] + \mathbbm{1}_{A_{n}^{c}}\cdot P\left[\zeta_{n}\leq p\mid\sigma_{n-1}\right] \\ &\overset{(6.58)}{=} \mathbbm{1}_{A_{n}}\cdot p_{n} + \mathbbm{1}_{A_{n}}\cdot E\left[\left(1-U_{n}\right)\cdot\mathbbm{1}_{\left\{\zeta_{n}\leq\frac{p-p_{n}}{1-p_{n}}\right\}}\mid\sigma_{n-1}\right] + \mathbbm{1}_{A_{n}^{c}}\cdot p, \end{split}$$

where in (*) we used that $A_n \in \sigma_{n-1}$. The proof of (6.65) will be complete once we show

$$\mathbb{1}_{A_n} \cdot E\left[(1 - U_n) \cdot \mathbb{1}_{\left\{ \zeta_n \le \frac{p - p_n}{1 - p_n} \right\}} \middle| \sigma_{n-1} \right] = \mathbb{1}_{A_n} \cdot (p - p_n). \tag{6.66}$$

To this end, we first calculate

$$\mathbb{1}_{A_n} \cdot E\left[(1 - U_n) \cdot \mathbb{1}_{\{\zeta_n \le \frac{p - p_n}{1 - p_n}\}} \middle| \sigma_{n-1}, (X_t^{x_n})_{t \ge 0} \right]$$
(6.67)

$$= \mathbb{1}_{A_n} \cdot (1 - U_n) \cdot E \left[\mathbb{1}_{\{\zeta_n \le \frac{p - p_n}{1 - p_n}\}} \middle| \sigma_{n-1}, (X_t^{x_n})_{t \ge 0} \right]$$

$$\stackrel{(**)}{=} \mathbb{1}_{A_n} \cdot (1 - U_n) \cdot \frac{p - p_n}{1 - p_n}, \tag{6.68}$$

where (**) follows from (6.59) and (6.55), which can be applied because p_n is σ_{n-1} -measurable and ζ_n is independent of $\sigma(\sigma_{n-1}, (X_t^{x_n})_{t\geq 0})$.

To conclude the proof of (6.66), note that taking $E[\cdot | \sigma_{n-1}]$ on (6.67) (and again using the fact that $A_n \in \sigma_{n-1}$) gives the left-hand side of (6.66), whereas taking $E[\cdot | \sigma_{n-1}]$ on (6.68) (and using (6.58)) gives the right-hand side of (6.66). The proof of (6.65) and Lemma 6.18 is complete.

6.7 Proof of Proposition 6.9

The goal of this section is to prove Proposition 6.9. Recall the definition of $M_{\infty}^{x,y,T}$ from (6.23). We generalize this definition by setting, for any $t \in [T, \infty)$,

$$M_t^{x,y,T} = P[\exists u \ge T : X_u^y = X_u^x \mid \mathcal{F}_t^x], \qquad \mathcal{F}_t^x = \sigma(X_u^x : 0 \le u \le t).$$
 (6.69)

This defines a martingale indexed by $t \in [T, \infty]$. In order to simplify notation, we will omit the superscripts that indicate dependence on x, y and T.

Let us now describe the strategy of proof of Proposition 6.9. As suggested in Remark 6.10, we have $M_0 = E[M_\infty] \times T^{1-d/2} \wedge |x-y|^{2-d}$, so the event $\{M_\infty > T^{1-\frac{d}{2}+\varepsilon}\}$ (and $\{M_\infty > |x-y|^{2-d+\varepsilon}\}$ for the case of |x-y| large) can only occur if the terminal value of the martingale deviates too much from its initial value. This is where Theorem 2.3 comes into play. In order to apply this theorem, we will obtain estimates on the size of the jumps of (M_t) and on its predictable quadratic variation $\langle M \rangle_\infty$; these estimates are given in (6.71) and (6.72). We derive these estimates by comparing M_t with $M_t^{(e)}$, which arises from M_t by artificially forcing the walk $(X_s^x)_{s\geq 0}$ to jump at time t in the direction of the unit vector $e \sim 0$, see Definition 6.20. Specifically, in Lemma 6.23 we show that the jumps of M can be bounded in terms of $|M_t - M_t^{(e)}|$ and the predictable quadratic variation $\langle M \rangle_\infty$ can be expressed as an integral of $(M_t - M_t^{(e)})^2$. The difference $|M_t - M_t^{(e)}|$ is bounded in Lemma 6.24 using the random walk facts of Section 2.3.

Recall that (M_t) is càdlàg. Denote by

$$\Delta M_T = \sup_{t > T} |M_t - M_{t-}|$$

the maximal jump size of M_t after time T. Recall the notion of $\langle M \rangle_t$ from Definition 2.2.

Lemma 6.19. There exist dimension-dependent constants C_0, C_1, C_2 such that the following bounds almost surely hold:

$$M_T \le C_0 T^{1 - \frac{d}{2}} \tag{6.70}$$

$$\Delta M_T \le C_1 T^{\frac{1}{2} - \frac{d}{2}} \tag{6.71}$$

$$\langle M \rangle_{\infty} - \langle M \rangle_T \le C_2 T^{2-d}$$
 (6.72)

Before we prove Lemma 6.19 we use it to prove Proposition 6.9.

Proof of Proposition 6.9. We first prove (6.24):

$$P\left[M_{\infty} > T^{1-\frac{d}{2}+\varepsilon}\right] \overset{(6.70)}{\leq} P\left[M_{\infty} - M_{T} > T^{1-\frac{d}{2}+\varepsilon} - C_{0}T^{1-\frac{d}{2}}\right] \overset{(*)}{\leq}$$

$$P\left[M_{\infty} - M_{T} > \frac{1}{2}T^{1-\frac{d}{2}+\varepsilon}\right] \overset{(2.2),(6.71),(6.72)}{\leq}$$

$$\exp\left(-\frac{1}{2}\frac{\frac{1}{2}T^{1-\frac{d}{2}+\varepsilon}}{C_{1}T^{\frac{1}{2}-\frac{d}{2}}}\ln\left(1 + \frac{\frac{1}{2}T^{1-\frac{d}{2}+\varepsilon}C_{1}T^{\frac{1}{2}-\frac{d}{2}}}{C_{2}T^{2-d}}\right)\right) =$$

$$\exp\left(-\frac{1}{4C_{1}}T^{\frac{1}{2}+\varepsilon}\ln\left(1 + \frac{C_{1}}{2C_{2}}T^{-\frac{1}{2}+\varepsilon}\right)\right) \overset{(*)}{\leq} \exp\left(-T^{\varepsilon}\right), \quad (6.73)$$

where the inequalities marked by (*) hold if T is large enough. We have proved that (6.24) holds with $\delta = \varepsilon$.

We now turn to (6.25). Given our $x, y \in \mathbb{Z}^d$, we bound

$$P[M_{\infty} > |x - y|^{2 - d + \varepsilon}] \overset{(6.69)}{\leq} P\left[P[\exists u \geq \widehat{T} : X_u^y = X_u^x \mid \mathcal{F}_{\infty}^x] > \frac{1}{2}|x - y|^{2 - d + \varepsilon}\right] + P\left[P[\exists u \leq \widehat{T} : X_u^y = X_u^x \mid \mathcal{F}_{\infty}^x] > \frac{1}{2}|x - y|^{2 - d + \varepsilon}\right], \quad (6.74)$$

where \widehat{T} is defined by

$$\widehat{T} = C|x - y|^{2 - \frac{\varepsilon}{d - 2 - \varepsilon/2}}, \quad C = 2^{\frac{1}{d/2 - 1 - \varepsilon/4}},$$
so that
$$\frac{1}{2}|x - y|^{2 - d + \varepsilon} = \widehat{T}^{1 - \frac{d}{2} + \varepsilon/4}.$$
(6.75)

Assuming that |x-y| is large enough, we can bound the first term on the right-hand side of (6.74) analogously to (6.73):

$$P\left[P[\exists u \ge \widehat{T} : X_u^y = X_u^x \mid \mathcal{F}_{\infty}^x] > \frac{1}{2}|x - y|^{2 - d + \varepsilon}\right] \stackrel{(6.75)}{\le} \exp\left(-\widehat{T}^{\frac{\varepsilon}{4}}\right) \stackrel{(6.1)}{\le} e^{-|x - y|^{\varepsilon/4}}.$$
(6.76)

Now we bound the second term on the right-hand side of (6.74) using Markov's inequality:

$$P\left[P[\exists u \le \widehat{T} : X_u^y = X_u^x \mid \mathcal{F}_{\infty}^x] > \frac{1}{2}|x - y|^{2 - d + \varepsilon}\right] \le \frac{P[\exists u \le \widehat{T} : X_u^y = X_u^x]}{\frac{1}{2}|x - y|^{2 - d + \varepsilon}}, \quad (6.77)$$

and

$$P[\exists u \le \widehat{T} : X_u^y = X_u^x] \stackrel{(2.8)}{=} P[\exists u \le 2\widehat{T} : X_u^{x-y} = 0] \le P\left[\max_{u \le 2\widehat{T}} |X_u^0| \ge |x-y|\right] \stackrel{(2.3)}{\le} 2d \exp\left(-\frac{1}{2}|x-y|\ln\left(1 + \frac{d \cdot |x-y|}{2\widehat{T}}\right)\right) \stackrel{(**)}{\le} \exp\left(-|x-y|^{\frac{\varepsilon}{d-2}}\right), \quad (6.78)$$

where (**) holds by (6.75) if |x-y| is large enough. Thus

$$\begin{split} P[M_{\infty} > |x-y|^{2-d+\varepsilon}] &\overset{(6.74),(6.76),(6.77),(6.78)}{\leq} \\ &e^{-|x-y|^{\varepsilon/4}} + \frac{\exp\left(-|x-y|^{\frac{\varepsilon}{d-2}}\right)}{\frac{1}{2}|x-y|^{2-d+\varepsilon}} \overset{(*)}{\leq} \exp\left(-|x-y|^{\varepsilon/d}\right), \end{split}$$

where (*) holds if |x-y| is large enough. This completes the proof of (6.25) with $\delta = \varepsilon/d$, as required by (6.1). The proof of Proposition 6.9 is complete, given Lemma 6.19.

Now we prepare the ground for the proof of Lemma 6.19.

Definition 6.20. For any $t \in [T, +\infty)$ let us define for $e \in \mathbb{Z}^d$, $e \sim 0$, the random variable

$$M_t^{(e)} = P[\{\exists T \le s \le t : X_s^y = X_s^x\} \cup \{\exists s \ge t : X_s^y = X_s^x + e\} \mid \mathcal{F}_t^x]$$
 (6.79)

Recall that we assume that our random walks and martingales are càdlàg.

Definition 6.21. Denote by $\tau_1 < \tau_2 < \dots$ the jump times of the random walk (X_t^x) and let $\tau_0 = 0$. For any $n \ge 1$ let $e_n = X_{\tau_n}^x - X_{\tau_{n-}}^x$ denote the direction of the jump of (X_t^x) at time τ_n .

Note that

$$(\tau_n - \tau_{n-1})_{n \ge 1}$$
 are i.i.d. with Exp(1) distribution. (6.80)

The next claim states that M_t only jumps when X_t^x jumps and in between jumps M_t is Lipschitz-continuous. The proof is straightforward and we omit it.

Claim 6.22. For any n = 1, 2, ... we have

$$M_{\tau_n} = M_{\tau_n -}^{(e_n)}, \tag{6.81}$$

$$\frac{\mathrm{d}}{\mathrm{d}t}M_t = -\frac{1}{2d}\sum_{e \sim 0} (M_t^{(e)} - M_t), \quad \tau_{n-1} < t < \tau_n, \tag{6.82}$$

and in particular,

$$|M_t - M_s| \le t - s \quad \text{for any} \quad \tau_{n-1} < s \le t < \tau_n. \tag{6.83}$$

Lemma 6.23. We have

$$\Delta M_T \le \sup_{t>T} \max_{e \sim 0} |M_t^{(e)} - M_t| \tag{6.84}$$

$$\langle M \rangle_t - \langle M \rangle_T = \frac{1}{2d} \sum_{e \ge 0} \int_T^t (M_s^{(e)} - M_s)^2 \,\mathrm{d}s \tag{6.85}$$

Proof. The inequality (6.84) immediately follows from (6.81).

Now we prove (6.85). Recall Definition 2.2. The right-hand side of (6.85) is adapted to (\mathcal{F}_t^x) and continuous in t, hence it is predictable (see Definition 2.1), thus we only need to check that that for any $T \leq s \leq t$ we have

$$E[M_t^2 - M_s^2 \mid \mathcal{F}_s^x] = E\left[\frac{1}{2d} \sum_{e \sim 0} \int_s^t (M_u^{(e)} - M_u)^2 \,\mathrm{d}u \mid \mathcal{F}_s^x\right]. \tag{6.86}$$

Let us define for $\delta > 0$ and $u \geq T$ the random variable

$$\psi_u^{\delta} := \frac{1}{\delta} E[(M_{u+\delta} - M_u)^2 \mid \mathcal{F}_u^x] \stackrel{(*)}{=} \frac{1}{\delta} E[M_{u+\delta}^2 - M_u^2 \mid \mathcal{F}_u^x], \tag{6.87}$$

where (*) follows from the fact that M_t is a martingale. Using (6.81), (6.83) and that (X_t) is a continuous-time simple random walk on \mathbb{Z}^d we obtain

$$\lim_{\delta \to 0_+} \psi_u^{\delta} = \frac{1}{2d} \sum_{e \sim 0} (M_u^{(e)} - M_u)^2, \qquad \mathbb{P} - \text{a.s.}$$
 (6.88)

It follows from the definition (6.87) that for any $\delta > 0$ we have

$$E\left[\int_{s}^{t} \psi_{u}^{\delta} du \,|\, \mathcal{F}_{s}^{x}\right] = E\left[\frac{1}{\delta} \int_{t}^{t+\delta} M_{u}^{2} du - \frac{1}{\delta} \int_{s}^{s+\delta} M_{u}^{2} du \,|\, \mathcal{F}_{s}^{x}\right].$$

From this, (6.80) and Claim 6.22 it follows that

$$\lim_{\delta \to 0_{+}} E\left[\int_{s}^{t} \psi_{u}^{\delta} du \,|\, \mathcal{F}_{s}^{x}\right] = E\left[M_{t}^{2} - M_{s}^{2} \,|\, \mathcal{F}_{s}^{x}\right], \qquad \mathbb{P} - \text{a.s.}$$

$$(6.89)$$

Now (6.86) will follow from (6.88) and (6.89) by dominated convergence as soon as we prove that for any $T \leq u$ and $0 < \delta \leq 1$ we have $\psi_u^{\delta} \leq 2$. This bound follows from (6.80) and Claim 6.22.

Lemma 6.24. There exists C > 0 such that for any $t \ge T \ge 1$ and $e \sim 0$,

$$|M_t^{(e)} - M_t| \le Ct^{\frac{1}{2} - \frac{d}{2}}. (6.90)$$

Before we prove Lemma 6.24, let us deduce Lemma 6.19 from it.

Proof of Lemma 6.19. We begin with (6.70). We first observe that, for any $y, z \in \mathbb{Z}^d$,

$$\sum_{w \in \mathbb{Z}^d} p_T(y, w) \cdot P\left[\exists t \ge 0 : X_t^w = X_t^z\right] \stackrel{(2.10), (2.5)}{\le} \int_T^\infty p_t(y, z) dt \stackrel{(2.6)}{\le} C_0 T^{1 - \frac{d}{2}}. \tag{6.91}$$

With this at hand, we derive (6.70):

$$M_T \stackrel{(6.69)}{=} \sum_{z,w} \mathbb{1}\{X_T^x = z\} \cdot p_T(y,w) \cdot P\left[\exists t \ge T : X_t^y = X_t^x \mid X_T^x = z, \ X_T^y = w\right]$$
$$= \sum_z \mathbb{1}\{X_T^x = z\} \cdot \sum_w p_T(y,w) \cdot P\left[\exists t \ge 0 : X_t^w = X_t^z\right] \stackrel{(6.91)}{\le} C_0 T^{1-\frac{d}{2}}.$$

The bound (6.71) follows from (6.84) and (6.90). Now we prove (6.72):

$$\langle M \rangle_{\infty} - \langle M \rangle_T \stackrel{(6.85)}{=} \frac{1}{2d} \sum_{e \sim 0} \int_T^{\infty} (M_s^{(e)} - M_s)^2 \, \mathrm{d}s \stackrel{(6.90)}{\leq} \int_T^{\infty} C s^{1-d} \, \mathrm{d}s = C T^{2-d}.$$

Proof of Lemma 6.24. Given $t \ge T$ let us define $A = \{\exists T \le s \le t : X_s^y = X_s^x\}$. We have

$$M_{t} \stackrel{(6.69),(2.10)}{=} P[A \mid \mathcal{F}_{t}^{x}] + \sum_{v,w \in \mathbb{Z}^{d}} P[A^{c} \cap \{X_{t}^{y} = w\} \mid \mathcal{F}_{t}^{x}] \mathbb{1}[X_{t}^{x} = v] \cdot \frac{g(v,w)}{g(0,0)},$$

$$M_{t}^{(e)} \stackrel{(6.79),(2.10)}{=} P[A \mid \mathcal{F}_{t}^{x}] + \sum_{v,w \in \mathbb{Z}^{d}} P[A^{c} \cap \{X_{t}^{y} = w\} \mid \mathcal{F}_{t}^{x}] \mathbb{1}[X_{t}^{x} = v] \cdot \frac{g(v + e, w)}{g(0,0)},$$

thus we obtain (6.90):

$$|M_t^{(e)} - M_t| \leq \sum_{v,w \in \mathbb{Z}^d} P[A^c \cap \{X_t^y = w\} \mid \mathcal{F}_t^x] \cdot \mathbb{1}[X_t^x = v] \cdot \frac{|g(v + e, w) - g(v, w)|}{g(0, 0)} \stackrel{(2.4)}{\leq}$$
$$\sum_{v,w \in \mathbb{Z}^d} P[X_t^y = w] \cdot \mathbb{1}[X_t^x = v] \cdot |g(v + e, w) - g(v, w)| \stackrel{(2.14)}{\leq} Ct^{\frac{1}{2} - \frac{d}{2}}.$$

7 Concluding remarks

Remark 7.1. Let us briefly explain why the method of Section 6 fails to prove (4.1) if d=3,4 and R=1 by arguing that the right-hand side of (6.14) does not go to zero. Rather than fixing the value of ℓ as in (6.2), in this heuristic argument we will keep track of the dependence on ℓ as well as on L of the terms on the right-hand side of (6.14). If we assume $\mathcal{Y} = \emptyset$, then by (6.10) we have $|\mathcal{X}| = 2 \cdot 2^N$. We will bound the probability of the event the right-hand side of (6.14) from below. For any fixed $\alpha > 0$ we can bound

$$\mathbb{P}\left[\bigcap_{\substack{\{x,z\}\in\mathcal{X}\\|x-z|=1}} F_{x,z}\right] \stackrel{(3.4),(6.8)}{\geq} \alpha \mathbb{P}\left[\mathcal{N}_{\infty}(\mathcal{X}) = 1, \bigcap_{\substack{\{x,z\}\in\mathcal{X}\\|x-z|=1}} E_{x,z}\right]. \tag{7.1}$$

Now the probability that $E_{x,z}$ occurs and yet $Y_t^x = Y_t^z$ for some t > T is roughly $\sqrt{T}^{2-d} = L^{(1-\varepsilon/2)(2-d)}$ by (2.6), moreover we can use the binary tree structure of \mathcal{X} to construct a scenario where $\mathcal{N}_{\infty}(\mathcal{X}) = 1$ and give a (heuristic) lower bound the probability on the right-hand side of (7.1) by

$$\left(L^{(1-\varepsilon/2)(2-d)} \right)^{2^N} \cdot \prod_{k=1}^N \left(L\ell^k \right)^{(2-d)2^{N-k}} \asymp \left(L^{(2-d)(2-\varepsilon/2)} \cdot \ell^{(2-d)\cdot 2} \right)^{2^N}.$$

If we multiply this with the combinatorial complexity term $(L^d\ell^{2d-2})^{2^N}$ that appears on the right-hand side of (6.14) then the resulting product goes to infinity as $N \to \infty$.

Remark 7.2. Let us explain why the method of Section 5 cannot be used to prove Theorem 1.1. Let us assume $\mathcal{Y} = \emptyset$ (so that by (6.10) we have $|\mathcal{X}| = 2 \cdot 2^N$) and bound the probability of the event of the right-hand side of (6.14):

$$\mathbb{P}\left[\bigcap_{\substack{\{x,z\}\in\mathcal{X}\\|x-z|=1}} F_{x,z}\right] \overset{(6.17),(6.20)}{\leq} E\left[\alpha^{\mathcal{N}_{\infty}^{Z}}\right].$$

Now we try to bound this using the idea of Lemma 5.2, i.e., we let random walks starting from the vertices of \mathcal{X} run independently until time T and then we let them annihilate each other. Let us denote by $\mathcal{N}_{\infty}^{Z'}$ the number of walkers that do not get annihilated. Then we have $E\left[\alpha^{\mathcal{N}_{\infty}^{Z'}}\right] \leq E\left[\alpha^{\mathcal{N}_{\infty}^{Z'}}\right]$, but using an argument similar to the one used in Remark 7.1 we can (non-rigorously) bound

$$E\left[\alpha^{\mathcal{N}_{\infty}^{Z'}}\right] \ge P[\mathcal{N}_{\infty}^{Z'} = 0] \gtrsim \left(L^{(1-\varepsilon/2)(2-d)}\right)^{2^N},$$

and this term is not small enough to beat the combinatorial complexity term $(CL^d)^{2^N}$ on the right-hand side of (6.14).

Acknowledgements: We would like to thank Alain-Sol Sznitman for bringing the problem under study here to our attention and hence initiating this project. We would also like to thank Roberto Imbuzeiro Oliveira for suggesting useful references on martingale concentration inequalities, Tim Hulshof for directing us to the reference [HvdHS03], and Ed Perkins for inspiring discussions.

The work of Balázs Ráth is partially supported by OTKA (Hungarian National Research Fund) grant K100473, the Postdoctoral Fellowship of the Hungarian Academy of Sciences and the Bolyai Research Scholarship of the Hungarian Academy of Sciences.

References

- [BLM87] J. Bricmont, J. Lebowitz, C. Maes, *Percolation in strongly correlated systems:* the massless Gaussian Field. Journal of Statistical Physics, Vol. 48, Nos. 5/6 (1987).
- [CS73] P. Clifford, A. Sudbury, A model for spatial conflict. Biometrika 60, 581-588 (1973).
- [HW83] B.I. Halperin, A. Weinrib, Critical phenomena in systems with long-range-correlated quenched disorder. Phys. Rev. B, 27(1), 413-427, (1983).
- [HvdHS03] T. Hara, G. Slade, R. van der Hofstad, Critical two-point functions and the lace expansion for spread-out high-dimensional percolation and related models. Annals of Probability 31(1), 349-408, (2003).
- [HL75] R. Holley, T.M. Liggett, Ergodic theorems for weakly interacting systems and the voter model. Annals of Probability 3, 643-663, (1975).
- [HMN14] M. Holmes, Y. Mohylevskyy, C. Newman, The voter model chordal interface in two dimensions. http://arxiv.org/abs/1409.0136 (2014).
- [Kal] O. Kallenberg, Foundations of Modern Probability. New York: Springer (1997).
- [K86] H. Kesten. Aspects of first-passage percolation. École d'été de probabilité de Saint-Flour XIV, Lecture Notes in Math 1180, Springer-Verlag, 125-264 (1986).
- [Kl05] F. Klebaner, *Introduction To Stochastic Calculus With Applications*. Second edition, World Scientific, (2005).
- [L96] G.F. Lawler, *Intersections of random walks*. Probability and its Applications, Birkhäuser Boston Inc. (1996).
- [LS86] J. Lebowitz, H. Saleur, Percolation in strongly correlated systems. Physica 138 A, 194-205 (1986).
- [LM06] J. Lebowitz, V. Marinov, Percolation in the harmonic crystal and voter model in three dimensions. Physical Review E 74, 031120 (2006).
- [Li85] T. Liggett, *Interacting particle systems*. Grundlehren der mathematischen Wissenschaften 276, Springer (1985).
- [Ma07] V. Marinov, *Percolation in correlated systems*. PhD thesis, Rutgers University (2007).
- [Ra05] B. Ráth, A short proof of the phase transition for the vacant set of random interlacements. Electronic Communications in Probability 20(3), 1-11 (2015).
- [RS13] P.-F. Rodriguez, A.-S. Sznitman, *Phase transition and level set percolation for the Gaussian free field.* Communications in Mathematics Physics 320 (2), 571–601 (2013).
- [S10] V. Sidoravicius and A.-S. Sznitman, Percolation for the vacant set of random interlacements. Communications on Pure and Applied Mathematics 62 (6), 831–858 (2009).
- [Sz10] A.-S. Sznitman, Vacant set of random interlacements and percolation. Annals of Mathematics 171(2), 2039–2087 (2010).

- [Sz12] A.-S. Sznitman, Decoupling inequalities and interlacement percolation on $G \times \mathbb{Z}$. Inventiones mathematicae, 187, 3, 645-706 (2012).
- [W84] A. Weinrib, Long-range correlated percolation. Phys. Rev. B, **29(1)**, 387–395, (1984).
- [Wi91] D. Williams, *Probability with martingales*, Cambridge University Press (1991).