

# PRELIMINARY STRUCTURAL OPTIMIZATION OF SOME FUMONISIN METABOLITES BY DENSITY FUNCTIONAL THEORY CALCULATION

Bors, I.<sup>(1)</sup>, Szabó-Fodor, J.<sup>(1)</sup>, Kovács, M.<sup>(1,2)</sup>

Original scientific paper

## SUMMARY

**Maize (*Zea mays* L.) is often contaminated with *Fusarium verticillioides*. This harmful fungus produces fumonisins as secondary metabolites. These fumonisins can appear both free and hidden form in planta. The hidden form is usually bound covalently to cereal starch. From the hidden fumonisins, during enzymatic degradation, glycosides are formed, and the fumonisin is further decomposed during a de-esterification step. In this short communication some preliminary DFT calculated structural results which could be useful in the future to help to understand the van der Waals force controlled molecular interactions between these kinds of mycotoxin molecules and enzymes are demonstrated.**

**Key-words: Fumonisin, fumonisin metabolites, DFT, electron density surfaces, structural optimization**

## INTRODUCTION

The term "mycotoxin" is usually reserved for the secondary metabolites produced by *fungi* that readily colonize crops. The sphingosine-like fumonisins are produced by several species of *Fusarium* molds, such as *Fusarium verticillioides*. The most commonly contaminated crop is maize (*Zea mays*). Fumonisin is a common mycotoxin in maize, although these toxins can occur in a few other crops as well (Placinta et al., 1999). The primary health concerns associated with fumonisins are carcinogenic properties and acute toxic effects (Voss et al., 2007). The disruption of sphingolipid metabolism by inhibition of ceramide synthase has been proposed to be responsible for the carcinogenicity and toxicity (Wang et al., 1991). Four groups of fumonisins (FA, FB, FC and FP) were classified based on different structure of the carbon backbone and the location of the nitrogen functional group (Musser and Plattner, 1997). Fumonisin (FB<sub>1</sub>) is the most common and economically important form, followed by B<sub>2</sub> and B<sub>3</sub>, where the index numbers refer to the different location of the hydroxyl groups on the carbon chain. Therefore, the metabolic investigations are focused on FB<sub>1</sub> *in vivo* (Fodor et al., 2008) and *in vitro* (Cirlini et al. 2015; Falavigna et al. 2012).

Mycotoxins undetectable by conventional, extraction-based analytical methods are termed as masked or hidden mycotoxins (Berthiller et al., 2013). Extractable mycotoxins can be easily detected but bound and/or hidden mycotoxins cannot be directly analysed. They have to be liberated from the matrix by chemical or enzymatic pre-treatment prior to chemical analysis. Dall'Asta et al. (2010) reported that with an *in vitro* digestion model method - after an enzymatic pre-treatment -, significantly more (30-40%) fumonisin was detected as compared to the measured moiety with conventional extraction method. In case of naturally contaminated (field derived) crop samples the hidden proportion of fumonisin is identical (35.6±22.3%, Dall'Asta et al., 2010). In case of inoculated crop cultures produced at laboratory-scale this value is 38.6±18.5% (Szabó-Fodor et al., 2015). Chemically modified mycotoxins are currently the largest group of modified mycotoxins and can be classified as "thermally formed" and "non-thermally formed" (Rychlik et al., 2014). Thermal degradation products have been described for several mycotoxins. A prominent example is fumonisin FB<sub>1</sub> which can react

(1) Ph.D. István Bors (bors.istvan@ke.hu), Ph.D. Judit Szabó-Fodor, Prof. Dr. Melinda Kovács - MTA-KE Mycotoxins in the Food Chain Research Group, 7400 Kaposvár, Guba S. u. 40. Hungary (2) Prof. Dr. Melinda Kovács, Kaposvár University, Faculty of Agricultural and Environmental Sciences, Guba S. u. 40., Kaposvár, H-7400, Hungary

in a Maillard-type reaction with reducing sugars leading to *N*-(1-deoxy-D-fructos-1-yl) fumonisin B<sub>1</sub> (NDF). The formation of hydrolyzed fumonisins (HFB<sub>1</sub>), on one hand, are biologically modified and formed by the intestinal

microbiota (Fodor et al., 2008) and, on the other, are formed under alkaline conditions during food processing (Humpf and Voss, 2004).




Figure 1. Structures of mentioned fumonisins

FB<sub>1</sub> is an inhibitor of sphinganine *N*-acyltransferase and increases the ratio of sphinganine/ sphingosine (Sa/So). Thus, elevation in the Sa/So ratio in different tissues (e.g. serum, urine, liver, kidney) is the exclusive biomarker of fumonisin exposure in exposed animals or human. However, best to our knowledge, no additional computation study was made to clarify the type of connection between sphinganine *N*-acyltransferase and fumonisin's monogastic metabolites. Density Functional Theory (DFT) calculation is an efficient method to determine the hydrophilic and hydrophobic site of molecules and it provides feasible input structures for the profound enzyme-substrate docking studies, in order to support the toxicological investigations.

From structural point of view, fumonisins provide a special nature: the long hydrophobic carbon chain bears electronegative (hydroxyl, amino- and carboxyl) functional groups. Due to free rotation around the carbon chain's single bonds, there could be several conformations – with different spatial location of functional groups –, being convenient for the above mentioned enzymes as “key”, according to the lock and key enzyme model.

Several toxin molecules were investigated by theoretical DFT studies (e.g. Song et al., 2011; Rahmani brothers, 2014) including also mycotoxins (e.g. Türker and Gümüş, 2009). The first step of these computational studies is always the structure optimization followed by enzyme docking studies (e.g. Kumar and Garg, 2014). Docking studies needs huge computing power, usually executed in a computer cluster, which was not available in this case. The aim of this work was to highlight the lack amidst fumonisin metabolites and computa-

tional chemistry, as well as to provide some preliminary results for further, enzyme docking studies.

## MATERIAL AND METHODS

All calculations were performed by Gaussian 03W. The structures were built in Advanced Chemistry Development, Inc. (ACD/Labs-ACD/3D) as .mol files and then converted to GaussView 3.09. The geometry of fumonisin metabolites were first pre-optimized with semi-empirical PM3 method and these structures were re-optimized in terms of DFT Gaussian 03 calculations for structural parameters using B3LYP (Becke's three-parameter functional with exact HF exchange and Lee-Yang-Parr exchange-correlation)/6-311G basis set. The FMO analysis (HOMO - highest occupied molecular orbital, LUMO - lowest unoccupied molecular orbital), are acronyms for and, involving hybridizations of selected bonds are also calculated at B3LYP methods and 6-311G level of the theory. Although most of all biologically relevant DFT studies set water as model solvent to investigate the solvation effect, in this case solvent interactions were neglected because of the long computation time; therefore all model calculations were executed in vacuum. GaussView 3.09's cube generator was used to calculate and visualize the electrostatic potential maps from the DFT results.


## RESULTS AND DISCUSSION

Although the calculations are performed in vacuum – which is obviously not the medium of a living cell –, some ascertainment should be noted. The optimized structure of FB<sub>1</sub> is more spheroidal than expected which

is be explained with the zwitterion-like force between the Lewis base amine and acidic carboxyl group. It is in agreement with the more linear structure of NDF, where amine form *N*-glycosidic bond which is unable to show the above mentioned intermolecular interaction. HFB<sub>1</sub> became also a semicircle instead of a line, but with a smaller curve than FB<sub>1</sub>, representing the ground state torsional angle between bonds (Figure 2.). Computed energy gap values ( $\Delta\varepsilon$ ) are shown in Table 1. The results are in agreement with the awaited stability (the first and second hydrolyzation step of NDF): HFB<sub>1</sub> > FB<sub>1</sub> > NDF.

**Table 1. B3LYP/6-311G calculated LUMO, HOMO and  $\Delta\varepsilon$  (energies, are in eV) ( $\Delta\varepsilon = \varepsilon_{\text{LUMO}} - \varepsilon_{\text{HOMO}}$ )**

Energy – (eV)	FB <sub>1</sub>	HFB <sub>1</sub>	NDF
LUMO	-4.08	-2.48	-5.01
HOMO	-9.82	-9.88	-8.98
$\Delta\varepsilon$	5.74	7.40	3.97



**Figure 2. Optimized geometry of the investigated fumonisin molecules at 37°C *in vacuo* – tube model**

## CONCLUSION

The structures of FB<sub>1</sub>, NDF and HFB<sub>1</sub> have been optimized by the Gaussian 03W at B3LYP/6-311G level. With the help of the results – and within the framework of the lock and key model –, researchers can better picture how the active site of the sphinganine *N*-acyltransferase approaches and joins the investigated fumonisin metabolites inhibiting the function of the enzyme in question.

By all means, these calculations should be handled as preliminary results, since solvation effects haven't been taken into consideration. Moreover, pH has momentous influence on the structure, especially in the case of FB<sub>1</sub>. It should be calculated at isoelectric point as a zwitterion, with  $-\text{NH}_3^+$  and  $-\text{COO}^-$  functional groups. Due to the limited computational sources, we have to decide in the future to use smaller basis set instead of B3LYP/6311G and recomputing with solvent effect or to use a computer cluster to assess exact calculations.

## ACKNOWLEDGEMENT

The research was supported by the Hungarian Academy of Sciences (within the frame of the MTA-KE "Mycotoxins in the food chain" Research Group), and by the Bolyai János research grant (BO\_499\_13 to J. Sz-F.)

## REFERENCES

1. ACD/Structure Elucidator, version 12.01, Advanced Chemistry Development, Inc., Toronto, ON, Canada, www.acdlabs.com, 2014.
2. Berthiller, F., Crews, C., Dall'Asta, C., De Saeger, S., Haesaert, G., Karlovsky, P., Oswald, I.P., Seefelder, W., Speijers, G. and Stroka, J. (2013): Masked Mycotoxins: A Review. *Mol. Nutr. Food Res.*, 57: 165-186. doi: <http://dx.doi.org/10.1002/mnfr.201100764>
3. Cirlini, M., Hahn, I., Varga, E., Dall'Asta, M., Falavigna, C., Calani, L., Berthiller, F., Del Rio, D., Dall'Asta, C. (2015): Hydrolysed fumonisin B<sub>1</sub> and *N*-(deoxy-D-fructos-1-yl)-fumonisin B<sub>1</sub>: stability and catabolic fate under simulated human gastrointestinal conditions. *Int. J. Food Sci. Nutr.* 4, 66(1): 98-103. doi: <http://dx.doi.org/10.3109/09637486.2014.979316>

4. Dall'Asta, C., Falavigna, C., Galaverna, G., Dossena, A. and Marchelli, R. (2010): *In vitro* digestion assay for determination of hidden fumonisins in maize. *J. Agr. Food Chem.*, 58: 12042-12047. doi: <http://dx.doi.org/10.1021/jf103799q>
5. Falavigna, C., Cirlini, M., Galaverna, G., Dall'Asta, C. (2012): Masked fumonisins in processed food: Co-occurrence of hidden and bound forms and their stability under digestive conditions. *World Mycotoxin J.*, 5(3): 325-334. doi: <http://dx.doi.org/10.3390/toxins6020523>
6. Fodor, J., Balogh, K., Weber, M., Mézes, M., Kametler, L., Pósa, R., Mamet, R., Bauer, J., Horn, P., Kovács, F., Kovács, M. (2008): Absorption, distribution and elimination of fumonisin B<sub>1</sub> metabolites in weaned piglets. *Food Addit. Contam.: Part A.* 25(1): 88-96. doi: <http://dx.doi.org/10.1080/02652030701546180>
7. Gaussian 03, Revision C.02, Gaussian, Inc., Wallingford CT, 2004 ([http://www.gaussian.com/g\\_misc/g03/citation\\_g03.htm](http://www.gaussian.com/g_misc/g03/citation_g03.htm))
8. Hahn, I., Nagl, V., H. Elisabeth, S-Z., Varga, E., Schwarz, C., Slavik, V., Reisinger, N., A Malachová, A., Cirlini, M., Generotti, S., Dall'Asta, C., Krska, R., Moll, W-D., Berthiller, F. (2015): Effects of orally administered fumonisin B<sub>1</sub> (FB<sub>1</sub>), partially hydrolysed FB<sub>1</sub>, hydrolysed FB<sub>1</sub> and *N*-(1-deoxy-D-fructos-1-yl) FB<sub>1</sub> on the sphingolipid metabolism in rats. *Food Chem. Toxicol.*, 76: 11–18. doi: <http://dx.doi.org/10.1016/j.fct.2014.11.020>
9. Humpf, H., Voss, K.A. (2004): Effects of thermal food-processing on the chemical structure and toxicity of fumonisin mycotoxins. *Mol. Nutr. Food Res.*, 48: 255-269. doi: <http://dx.doi.org/10.1002/mnfr.200400033>
10. Kumar, N., Garg, A. (2014): Structural optimization and docking studies of anatoxin-a: A potent neurotoxin. *Afr. J. Biotechnol.*, 13(30): 3092-3100. doi: <http://dx.doi.org/10.5897/AJB2014.13671>
11. Merrill, AH. Jr., Wang, E., Gilchrist, DG., Riley, RT. (1993): Fumonisin and other inhibitors of *de novo* sphingolipid biosynthesis. *Adv. Lipid Res.* 26: 215-234. link
12. Musser, S.M., Plattner, R.D. (1997): Fumonisin composition in cultures of *Fusarium moniliforme*, *Fusarium proliferatum* and *Fusarium nygami*. *J. Agric. Food Chem.*, 45: 1169-1173. doi: <http://dx.doi.org/10.1021/jf960663t>
13. Placinta, C.M., D'Mello, J.P.F., Macdonald, A.M.C. (1999): A review of worldwide contamination of cereal grains and animal feed with *Fusarium* mycotoxins. *Anim. Feed Sci. Technol.*, 78: 21-37. doi: [http://dx.doi.org/10.1016/S0377-8401\(98\)00278-8](http://dx.doi.org/10.1016/S0377-8401(98)00278-8)
14. Rahmani, A., Rahmani, S., Rahmani, M., Hosseini, S. J. (2014): Theory Functional Calculations (DFT), Structural Properties, Natural Bond and Energies for the Haloryanic Compound. *J. Appl. Sci. & Agric.*, 9(7): 2745-2748. link
15. Rychlik, M., Humpf, H-U., Marko, D., Dänicke, S., Mally, A., Berthiller, F., Klaffke, H., Lorenz, N. (2014): Proposal of a comprehensive definition of modified and other forms of mycotoxins including "masked" mycotoxins. *Mycotoxin Res.*, 30: 197–205. doi: <http://dx.doi.org/10.1007/s12550-014-0203-5>
16. Song, Y., Ambati, J., Parkin, S., Rankin, S. E., Robertson, L. W., Lehmler, H-J. (2011): Crystal Structure and Density Functional Theory Studies of Toxic Quinone Metabolites of Polychlorinated Biphenyls. *Chemosphere*, 85(3): 386-392. doi: <http://dx.doi.org/10.1016/j.chemosphere.2011.07.004>
17. Szabó-Fodor, J., C. Dall'Asta, C., Falavigna, C., Kachlek, M., Szécsi, Á., Szabó, A., Kovács, M. (2015): Determination of the amount of bioaccessible fumonisin B<sub>1</sub> in different matrices after *in vitro* digestion. *World Mycotoxin J.*, 8(3): 261-267. doi: <http://dx.doi.org/10.3920/WMJ2014.1771>
18. Türker, L., Gümüş, S. (2009): A theoretical study on vomitoxin and its tautomers. *J. Hazard. Mater.*, 163: 285-294. doi: <http://dx.doi.org/10.1016/j.jhazmat.2008.06.087>
19. Voss, K.A., Smith, G.W., Haschek, W.M. (2007): Fumonisin: toxicokinetics, mechanism of action and toxicity. *Anim. Feed Sci. Technol.*, 137: 299-325. doi: <http://dx.doi.org/10.1016/j.anifeedsci.2007.06.007>
20. Wang, E., Norred, W.P., Bacon, C.W., Riley, R.T., Merrill, A.H. Jr. (1991): Inhibition of sphingolipid biosynthesis by fumonisins. Implications for diseases associated with *Fusarium moniliforme*. *J. Biol. Chem.*, 266(22): 14486-14490.

(Received on 29 April 2015; accepted on 7 July 2015)