


AUTHOR QUERY FORM

| | | |
|---|---|--|
|  | <p>Journal: NSC</p> <p>Article Number: 14033</p> | <p>Please e-mail or fax your responses and any corrections to:</p> <p>E-mail: corrections.essd@elsevier.sps.co.in</p> <p>Fax: +31 2048 52799</p> |
|---|---|--|

Dear Author,

Please check your proof carefully and mark all corrections at the appropriate place in the proof (e.g., by using on-screen annotation in the PDF file) or compile them in a separate list. Note: if you opt to annotate the file with software other than Adobe Reader then please also highlight the appropriate place in the PDF file. To ensure fast publication of your paper please return your corrections within 48 hours.

For correction or revision of any artwork, please consult <http://www.elsevier.com/artworkinstructions>.

Any queries or remarks that have arisen during the processing of your manuscript are listed below and highlighted by flags in the proof. Click on the 'Q' link to go to the location in the proof.

| Location in article | Query / Remark: click on the Q link to go Please insert your reply or correction at the corresponding line in the proof |
|---------------------------|--|
| <u>Q1</u> | Please confirm that given names and surnames have been identified correctly. |
| <u>Q2</u> | Please select six key words from list. |
| <u>Q3</u> | Please provide city, state and country for manufacturers 'Cell Signaling, Santa Cruz, Sigma-Aldrich, etc.' |
| <u>Q4</u> | The references 'Um and coworkers (2011), Steiner et al. (2011), Koltai et al. (2012), Yu and co-workers (2010), Gomez-Pinilla et al. (2008), and Bayod et al. (2011)' are cited in the text but not listed. Please check. |
| | <div style="border: 1px solid black; padding: 5px; display: inline-block;"> <p style="color: red; margin: 0;">Please check this box if you have no corrections to make to the PDF file</p> <input style="width: 30px; height: 20px; margin-left: 10px;" type="checkbox"/> </div> |

Thank you for your assistance.

Neuroscience xxx (2012) xxx–xxx

Highlights

Neuroscience xx (2012) xxx

► Long term physical exercise reduced the levels of ROS and protein carbonyls in the hippocampus of aging rats. ► Levels of antioxidant enzymes /Gpx and SOD-1/ were increased by exercise. ► AMPK and PGC-1 α are important in mediating the beneficial effects of exercise in the hippocampus. © 2012 Published by Elsevier Ltd. on behalf of IBRO.

Please cite this article in press as: Marosi K et al. Long-term exercise treatment reduces oxidative stress in the hippocampus of aging rats. *Neuroscience* (2012), <http://dx.doi.org/10.1016/j.neuroscience.2012.09.001>

Neuroscience xxx (2012) xxx–xxx

LONG-TERM EXERCISE TREATMENT REDUCES OXIDATIVE STRESS IN THE HIPPOCAMPUS OF AGING RATS

K. MAROSI, Z. BORI, N. HART, L. SÁRGA, E. KOLTAI, Z. RADÁK AND C. NYÁKAS*

Semmelweis University, Institute of Sport Science, Budapest, Hungary

Abstract—Exercise can exert beneficial effects on cognitive functions of older subjects and it can also play an important role in the prevention of neurodegenerative diseases. At the same time it is perceivable that limited information is available on the nature of molecular pathways supporting the antioxidant effects of exercise in the brain. In this study 12-month old, middle-aged female Wistar rats were subjected to daily moderate intensity exercise on a rodent treadmill for a period of 15 weeks which covered the early aging period unmasking already some aging-related molecular disturbances. The levels of reactive oxygen species (ROS), the amount of protein carbonyls, the levels of antioxidant intracellular enzymes superoxide dismutases (SOD-1, SOD-2) and glutathione peroxidase (GPx) were determined in the hippocampus. In addition, to identify the molecular pathways that may be involved in ROS metabolism and mitochondrial biogenesis, the activation of 5'-AMP-activated protein kinase (AMPK), the protein level of peroxisome proliferator-activated receptor gamma coactivator 1-alpha (PGC-1 α), nuclear respiratory factor 1 (NRF-1) and mitochondrial transcription factor A (mtTFA) were measured. Our results revealed a lower level of ROS associated with a reduced amount of protein carbonyls in the hippocampus of physically trained rats compared to sedentary controls. Furthermore, exercise induced an up-regulation of SOD-1 and GPx enzymes, p-AMPK and PGC-1 α , that can be related to an improved redox balance in the hippocampus. These results suggest that long-term physical exercise can comprises antioxidant properties and by this way protect neurons against oxidative stress at the early stage of aging. © 2012 Published by Elsevier Ltd. on behalf of IBRO.

Key words: exercise, hippocampus, aging, ROS, SOD, GPx, AMPK, PGC-1 α .

*Corresponding author. Address: Semmelweis University, Institute of Sport Science, Laboratory of Brain Physiology, Alkotás utca 44, 1123 Budapest, Hungary. Tel: +36-30-8467149; fax: +36-1-3566337.

E-mail address: nyakas@tf.hu (C. Nyakas).

Abbreviations: AMPK, 5'-AMP-activated protein kinase; CaMKK, calmodulin-dependent kinase; Gpx, GSH peroxidase; mtTFA, mitochondrial transcription factor A; NRF-1, nuclear respiratory factor-1; PGC-1 α , peroxisome proliferator-activated receptor gamma coactivator 1-alpha; ROS, reactive oxygen species; RT, room temperature; SOD, superoxide dismutase; TBS-T, Tris-buffered saline Tween-20.

0306-4522/12 \$36.00 © 2012 Published by Elsevier Ltd. on behalf of IBRO. <http://dx.doi.org/10.1016/j.neuroscience.2012.09.001>

INTRODUCTION

Reactive oxygen species (ROS) are the products of cellular aerobic metabolism. When ROS are presented in physiological concentrations they play an important role in the modulation of gene expression and signal transduction pathways (Sen and Packer, 1996; Finkel, 2003; Esposito et al., 2004). However, when ROS are produced in excess for a considerable time, they can attack cellular macromolecules like proteins, membrane lipids and DNA. Extensive damage to these biomolecules in the brain can cause neuronal dysfunction and trigger apoptosis (Emerit et al., 2004).

A number of studies have demonstrated that increased levels of ROS are involved in the aging process (Gerschman et al., 1954; Harman, 1956; Droge, 2003) and contribute to pathological changes in neurodegenerative disorders (Floyd and Hensley, 2002). The major targets of ROS are the amino acid residues of proteins. The oxidation of lysine, arginine, proline and threonine residues results in carbonyl derivatives that are used as markers of oxidative stress on proteins (Levine and Stadtman, 2001). ROS-induced carbonylation leads to dysfunctional proteins and enzymes with reduced catalytic activity. Enhanced levels of protein carbonyls have been shown to impair cognitive processes and correlate with the progression of several neuronal pathologies (Stadtman, 2001). The accumulation of carbonyl derivatives have been dramatically raised in vulnerable neurons in Alzheimer disease (Aksenov et al., 2001), and Parkinson disease (Offen et al., 1999).

The hippocampus is highly vulnerable to oxidative damage during aging due to the reduced capacity of neurons to maintain redox homeostasis (Serrano and Klann, 2004). Since the hippocampus is involved in certain forms of learning and memory consolidation (Douglas, 1967; Meissner, 1967; Morris, 2006) oxidative damage to this brain area can cause impairment in cognitive functions (Serrano and Klann, 2004). The maintenance of a normal redox state in hippocampal neurons therefore, is important in the prevention of cognitive decline during aging.

The antioxidant defense at first line is constituted by the antioxidant enzymatic actions. Superoxide radicals are converted to hydrogen peroxide by superoxide dismutase (SOD), and the hydrogen peroxide is eliminated by glutathione peroxidase (GPx) and/or catalase. Catalase activity has been found to be low in the brain (Gaunt and de Duve, 1976), the enzyme GPx,

therefore, is primarily responsible for destruction of excess hydrogen peroxide formed in the nervous tissue (Sinét et al., 1980). The activity of antioxidant enzymes in the brain is modulated by various factors including aging (Venkateshappa et al., 2012) and physical activity (Radak et al., 2001).

Numerous studies indicate that exercise can reduce the risk of oxidative stress-related diseases and play an important role against age-associated cognitive decline (Bergersen and Storm-Mathisen, 2006; Ferrari, 2007; Hollmann et al., 2007; Shin, 2009). It has been proposed that regular physical training induces an adaptation process in ROS-detoxifying systems, resulting in increased resistance of cells to oxidative challenges (Radak et al., 1999a). Most of the beneficial effects of exercise on aged rodents (Devi and Kiran, 2004) and elderlies (Asha Devi, 2009) were related to long-term exercise of moderate intensity, while the acute training was proved to induce oxidative insults in the nervous system (Aydin et al., 2009). The adaptive response to long-term exercise is rather complex and not fully elucidated; it may involve the modulation of redox-sensitive transcriptional factors (Toldy et al., 2005). Peroxisome proliferator-activated receptor gamma coactivator 1-alpha (PGC-1 α) may play an important role in the control of ROS metabolism, since it regulates the expression of ROS-detoxifying enzymes (Valle et al., 2005). In addition, PGC-1 α interacts with a broad range of transcriptional factors involved in the regulation of mitochondrial electron transport activity and mitochondrial biogenesis (Ventura-Clapier et al., 2008). PGC-1 α has been reported to induce the transcription of nuclear respiratory factor 1 (NRF-1) leading to the increased expression of mitochondrial transcription factor A (mtTFA), a key regulator of mitochondrial DNA replication (Ventura-Clapier et al., 2008).

In the skeletal muscle, phosphorylation of PGC-1 α is mediated by AMPK-activated protein kinase (AMPK) (Jager et al., 2007), a molecule emerging as a central regulator of energy balance principally as fuel sensor. The exercise-induced activation of AMPK in skeletal muscle has been confirmed by many studies (Lee-Young et al., 2009; Palacios et al., 2009; O'Neill et al., 2011). Nonetheless, the effects of physical activity on AMPK, PGC-1 α and the related molecular processes in the hippocampus are not yet completely elucidated.

The purpose of the present study was to investigate how long-term moderate intensity physical exercise influences the oxidative status in the hippocampus at the early stage of aging, when the neuronal functions have been acknowledged to decline slowly but gradually. To study the impacts of forced physical activity on the hippocampal redox state 12-month-old female rats were subjected to long-term exercise intervention in our experiment. It is known that the normal cyclic female rats show a gradual decrease in serum estradiol level starting at the age of 12 months (Lapolt et al., 1988; Moorthy et al., 2005b). Thus 12 months can be considered as the starting period of "menopause" and the consequent months as the early

stage of "postmenopausal" period by attempting to draw reference to human condition. Estrogens, otherwise, can regulate the expression and activity of antioxidant enzymes itself (Moorthy et al., 2005a) and affect mitochondrial functions (Klinge, 2008). Whatever is the initial deteriorating factor during early aging on the redox balance physical activity could be a significant non-pharmacological tool to attenuate the dysregulation of antioxidant system as it has been proposed in the present study.

EXPERIMENTAL PROCEDURES

Animals and treatments

Middle-aged (12 months old) female Wistar rats were selected for the study. Animals were housed in a room maintained 22 ± 1 °C with a 12:12-h light/dark cycle starting the light period at 7:00. Food and tap water were available *ad libitum*. The animals were divided into two experimental groups: one group served as a sedentary control group ($n = 6$), while the other group was subjected to exercise treatment ($n = 6$). The exercise protocol included moderate intensity running on a rodent treadmill. Exercise trained rats were first introduced to treadmill running for 2 days on a 0% incline with 10 and 15 m/min, respectively. Afterwards for the next 4 days the running speed of the daily exercise sessions were gradually increased to 60% of the animals' VO₂ max assayed earlier in this particular age group. From the second week of the training program, the animals ran daily at 18 m/min, on a 0% incline, for 30 min.

Twenty-four hours after the last exercise treatment session the animals were sacrificed by decapitation under light CO₂ anesthesia and the brains were quickly removed. The two hemispheres were rapidly separated along the midline on an ice-cooled glass plate. The hippocampus was quickly excised bilaterally and immediately frozen on dry ice. The hippocampal samples were stored at -80 °C until processing. All experimental procedures which were carried out on the animals had been approved by the Animal Examination Ethics Council of the Animal Protection Advisory Board at Semmelweis University, Budapest.

Western blots

The hippocampi of each animal were homogenized in lysis buffer containing 137 mM NaCl, 20 mM Tris-HCl pH 8.0, 2% Nonidet P-40, 10% glycerol and protease inhibitors. The homogenate was sonicated for 30 s in a cold pack. Lysates were centrifuged for 15 min at 15,300g at 4 °C and the supernatants were collected and stored at -20 °C until use. The concentration of protein was determined using the Bradford assay (Bradford and Williams, 1976). Twenty micrograms of protein were electrophoresed on 8–12% (v/v) polyacrylamide SDS-PAGE gels. Proteins were electro transferred onto PVDF membranes (Amersham, Piscataway, NJ, USA). The nonspecific binding of immunoproteins was blocked with 5% non-fat dry powdered milk dissolved in Tris-buffered saline Tween-20 (TBS-T) for 2 h at room temperature (RT). After blocking, the membranes were incubated with primary antibodies overnight at 4 °C. Antibodies were dissolved in TBS-T containing 5% non-fat powdered milk. The primary antibodies were: AMPK: 1:2000, #2532 Cell Signaling; p-AMPK: 1:1000, #2531 Cell Signaling; PGC-1 α : 1:1000, #sc13067 Santa Cruz; NRF-1: 1:1000, #sc33771 Santa Cruz; mtTFA: 1:750, #sc30963 Santa Cruz; SOD-1: 1:1000, #AV45752 Sigma-Aldrich; GPx-1 (ISO1): 1:1000, #7283P1 Sigma-Aldrich; SOD-2: 1:1000, #SAB1406465 Sigma-Aldrich). The membranes were rinsed in TBS-T followed by 1-h

182 incubation with HRP-conjugated secondary antibody at RT. After
 183 incubation the membranes were repeatedly washed in TBS-T
 184 and incubated with an enhanced chemiluminescence reagent
 185 (ECL plus, RPN 2132, Amersham). The protein bands were
 186 visualized on X-ray films. The bands were quantified by Image
 187 J software, and standardized to β -actin (1:2000, #sc-47778
 188 Santa Cruz). With this software the optical density of the
 189 protein bands was measured. Results were expressed in
 190 relative density units. The phosphorylation of the AMPK was
 191 evaluated by dividing the phospho-specific form by the
 192 dephospho form of AMPK.

193 Detection of reactive oxygen species (ROS)

194 The overall ROS was determined by using modifications of
 195 the dichlorodihydrofluorescein diacetate (H₂DCF-DA) staining
 196 method (Kim et al., 2000). This assay approximates levels of
 197 reactive species, such as superoxide radical, hydroxyl radical,
 198 and hydrogen peroxide. H₂DCF-DA (2',7'-
 199 dichlorodihydrofluorescein diacetate, #D-399 Invitrogen) was
 200 dissolved at a concentration of 12.5 mM in ethanol before use.
 201 For fluorescence reactions, 96-well black microplates were
 202 loaded with phosphate buffer (pH 7.4) to a final concentration
 203 of 152 μ M/well. Then 8 μ l diluted freshly prepared
 204 hippocampus homogenate and 40 μ l of 125 μ M dye were
 205 added to achieve a final dye concentration of 25 μ M. The
 206 changes in the fluorescent signal of the oxidized H₂DCF-DA
 207 were recorded at three time points (0, 1 and 30 min), using a
 208 micro plate fluorescence reader (excitation/emission
 209 wavelengths of 485–538 nm, Fluoroskan Ascent FL). The
 210 fluorescence intensity unit was normalized with the protein
 211 content and expressed in relative unit production per minute.

212 Detection of protein carbonyls

213 The levels of oxidized proteins were determined using an Oxyblot
 214 kit (S7150, Chemicon/Millipore, Temecula, CA, USA). Proteins
 215 were derivatized with 4-dinitrophenylhydrazine (DNPH) for
 216 15 min followed by incubation at room temperature with a
 217 neutralization buffer. Derivatized proteins were electrophoresed
 218 on a 10% SDS-PAGE and blotted on PVDF membranes. Blots
 219 were blocked with 5% non-fat dry milk in Dulbecco's PBS
 220 containing 0.05% Tween 20 (PBS-T) for 3 h at 4 °C followed by
 221 incubation with anti-DNP primary antibody (1:150, #S7150
 222 Chemicon/Millipore) overnight at 4 °C. After three washes with
 223 PBS-T, membranes were incubated for 1 h at room
 224 temperature with HRP-secondary antibodies (1:300, #S7150
 225 Chemicon/Millipore). Immunocomplexes were visualized using
 226 ECL plus reagent. The bands were quantified by Image J
 227 software, and standardized to β -actin (1:2000, #sc-47778 Santa
 228 Cruz). Results on the figures were expressed in density units.

229 Statistical analyses

230 The results from experimental groups were compared by paired
 231 *t*-test. Statistical significance was set at $p < 0.05$. Means
 232 and standard errors of means (SEM) were presented to
 233 demonstrate the results. All statistical analyses were done
 234 applying the Statistica 8.8 program.

235 RESULTS

236 Physical exercise decreased the level of ROS by 28% in
 237 the hippocampus measured at the age of 15 months
 238 after the 15-week-long training period ($t = 3.472$,
 239 $p < 0.01$) as it is shown in Fig. 1. In addition, Fig. 2
 240 shows that at the same time the amount of protein
 241 carbonyls in the hippocampus decreased in the trained




Fig. 1. Reduction of reactive oxygen species (ROS) in the hippocampus by physical exercise as compared to the control (sedentary) group is shown (** $p < 0.01$ vs. control). Columns represent means \pm SEMs for six animals per group.

group compared to sedentary controls ($t = 3.179$,
 $p < 0.01$). In this figure (Fig. 2) the carbonylated bands
 are also shown. Exercise decreased the densities in
 most of the proteins visualized.

In order to evaluate the protein levels of antioxidant
 enzymes in the hippocampi of 15-month-old rats, the
 immunoreactivity of SOD-1, SOD-2 and GPx were
 measured by Western blot (Figs. 3 and 4). Exercise was
 effective in up-regulating the expression of SOD-1 in the
 hippocampus of aging animals ($t = 2.330$, $p < 0.05$;
 Fig. 3, left side panel; 52% increase). Fig. 3 also
 presents that there was a trend toward an increased
 expression of SOD-2 in the trained group ($t = 1.897$,
 $p = 0.087$; 25% increment). Student *t*-test revealed
 a significant increment in the GPx protein levels (Fig. 4)
 in the physically trained group compared to the control
 group ($t = 2.374$, $p < 0.05$; the increment after exercise
 was 34%). Upper parts of the figures show the densities
 of immunoreactive Western blot spots of all animals
 investigated and the calculated densities were
 normalized to β -actin and used for statistical processing
 which are represented by the columns and SEMs.

The phosphorylation of the AMPK was evaluated by
 dividing its phospho-specific form by its dephospho-
 form and was normalized to β -actin (Fig. 5). The
 phosphorylation of AMPK molecules was increased
 significantly by the exercise treatment ($t = 3.285$, $p <$
 0.01 ; the increment compared to controls was 40%).

PGC-1 α , NRF-1 and mtTFA are transcriptional factors
 involved in the regulation of cell metabolisms and
 mitochondrial functions. Western blot analysis showed
 that exercise enhanced the protein level of PGC-1 α
 compared to the control group in the hippocampus
 ($t = 3.523$, $p < 0.01$; Fig. 6). The increment was 55%
 which may be considered as a marked enhancement.
 Fig. 7 shows that the protein levels of NRF-1 is tended
 to be increased by exercise, however the increment just
 approached significance level ($t = 2.182$, $p = 0.054$;
 left side of the figure). Physical activity did not influence
 the level of mtTFA expression (Fig. 7, right side) shown
 by Student *t*-test.




Fig. 2. Protein carbonyl content in the hippocampus decreased by long-term exercise compared to the sedentary control group ($*p < 0.05$ vs. control). Columns represent means \pm SEMs for six animals per group. The representative Western blots show the immunoreactivities of protein carbonyls and β -actin (used as a loading control) at the right side of figure.




Fig. 3. Intracellular SOD-1 protein levels (left side) and mitochondrial SOD-2 protein content (right side) are shown in the hippocampus of 15-month-old aging female rats. SOD-1 protein levels are increased as a result of exercise ($**p < 0.01$ vs. control). The right side histogram represents that SOD-2 protein levels tended to be increased in exercised rats compared to the sedentary control group ($#p = 0.087$ vs. control). Columns represent means \pm SEMs, $n = 6$. The representative western blots taken from each individual rats show the immunoreactivities of SOD-1 and SOD-2 (upper lines) and below that of β -actin (used as a loading control).

DISCUSSION

283

284 The aim of present study was to examine the effects of
 285 long-term exercise on the hippocampal oxidative state
 286 and the related molecular processes in the course of
 287 early brain aging period in female rats. It was found that
 288 the long-term physical exercise resulted in reduced
 289 levels of reactive oxygen radicals (ROS) in the
 290 hippocampus of physically trained group, measured 24 h
 291 after the last exercise session under resting state. Due
 292 to the lower levels of free radicals the oxidative damage
 293 to proteins was significantly reduced in the trained rats.
 294 An important finding of this study is that exercise
 295 resulted in elevated levels of intracellular antioxidant
 296 enzymes in the early phase of aging. The activation of
 297 AMPK and the expression of PGC-1 α were also
 298 promoted by exercise, which can serve for possible

molecular pathway regulating the antioxidant gene
 transcription in the hippocampus. 299

In this study, the aged animals were subjected to long-
 term, moderate intensity exercise paradigm, which was
 effective in reducing the free radical levels in the
 hippocampus. Previous studies confirmed the role of
 training intensities in the modulation of redox state in
 different tissues. A single bout of exercise and
 exhaustive training have been reported to increase the
 ROS content and oxidative damage in the muscle and
 liver (Radak et al., 1999b; Nakamoto et al., 2007)
 assayed after a 20-h resting state. In contrast, long term
 training on moderate intensity has been found to reduce
 oxidative stress markers in the rat brain (Ogonovszky
 et al., 2005) measured 24 h after the last exercise
 session, which is in agreement with our observations.
 As a consequence of reduced ROS levels, lower 315




Fig. 4. Physical exercise was effective to increase the levels of GPx protein levels in the hippocampus shown by the columns below (* $p < 0.05$ vs. control, means \pm SEMs, $n = 6$). The representative Western blots show the immunoreactivities of GPx and β -actin in all individual rats.




Fig. 6. Exercise up-regulated the PGC-1 α protein levels in the hippocampus compared to the control group (** $p < 0.01$). Columns represent means \pm SEMs for six animals per group. The representative western blots of PGC-1 α and β -actin are shown above.




Fig. 5. The activation of AMPK is enhanced by exercise in the hippocampus. The histogram represents that the phosphorylation of AMPK is increased in the physically trained group (exercise) compared to the sedentary control group (** $p < 0.01$ vs. control). The phosphorylation of the AMPK was evaluated by dividing the phospho-specific form by the dephospho-form. Columns represent means \pm SEMs for six animals per group. The representative Western blots show the immunoreactivities of p-AMPK, AMPK and β -actin in all animals.

316 amount of protein **carbonyls** was detected in the trained
 317 animals. Similarly, Radak and co-workers (2001) also
 318 found decreased protein carbonyl levels in the brain of
 319 **14-month-old** male rats exposed to a long-term
 320 swimming regime. Together with previous studies our
 321 data show that regular exercise can exert beneficial
 322 impacts on redox state suggesting that long-term
 323 training on moderate intensity could increase the ability
 324 of neurons to cope with oxidative stress during the
 325 **aging** process in females as well.

The beneficial effects of exercise on the oxidative 326
 status could be related to the enzymatic adaptation 327
 processes. GPx and two types of SOD were assayed, 328
 namely copper–zinc SOD (SOD-1) located in the cytosol 329
 and mitochondrial manganese SOD (SOD-2). The 330
 protein levels of SOD-1 and GPx have been elevated in 331
 response to long-term exercise intervention in this study 332
 and the level of SOD-2 was only slightly increased. Our 333
 results are in agreement with Um and coworkers' data 334
 (2011) demonstrating significant increment in SOD-1 335
 expression after a **3-month-long** training period. 336

In the present experiment, long-term exercise 337
 enhanced the protein levels of PGC-1 α in the 338
 hippocampus. Recently, Steiner et al. (2011) also 339
 demonstrated an up-regulation of PGC-1 α mRNA 340
 expression in specific brain regions of male rodents 341
 after **8** weeks of training. PGC-1 α could be an important 342
 modulator of redox balance in neuronal cells, because it 343
 is required for the induction of the gene expression of 344
 several antioxidant enzymes including SOD-1, SOD-2 345
 and GPx (St-Pierre et al., 2006). Furthermore, PGC-1 α 346
 can rescue neurons from **oxidative-stress-mediated** cell 347
 death and plays a role in the prevention of 348
 neurodegenerative processes (St-Pierre et al., 2006). 349
 The induced expression of PGC-1 α thus could serve an 350
 explanation for the elevated levels of SOD-1 and GPx 351
 found in the hippocampi of trained animals. 352

Besides its regulatory role on intracellular antioxidant 353
 gene transcription, PGC-1 α can also influence the 354
 mitochondrial functions through increasing the 355
 transcriptional activity of NRF-1 resulting in a 356
 downstream activation of mtTFA. In the present study 357
 no significant increment in the protein levels of NRF-1 358
 and mtTFA were observed, although NRF-1 expression 359
 exposed a near significant elevation. Koltai et al. (2012) 360
 also showed that exercise increased PGC-1 α 361
 expression, but the NRF-1 and mtTFA protein remained 362
 to be unchanged in the muscle of aged rats. To date, no 363




Fig. 7. Elevated (close to significant, $^{\#}p = 0.054$ vs. control) NRF-1 protein level in response to physical exercise is shown in the left panel. The mtTFA protein levels remained unchanged in the hippocampus following exercise (right panel). Means \pm SEMs are shown. The representative Western blots show the immunoreactivities of both transcription factors.

364 studies have examined the effects of physical activity on
 365 NRF-1, and mtTFA in the hippocampus. These
 366 molecules can share the common components of
 367 mitochondrial biogenesis and respiration, however it
 368 may be added that the regulation of mitochondrial
 369 protein formation might involve multiple, probably
 370 cooperative regulatory mechanisms in tissue- and in a
 371 cell-specific manner. Therefore, the exact individual
 372 contribution of each of the factors supporting
 373 mitochondrial functions and mitochondrial biogenesis is
 374 rather difficult to dissect (Lopez-Lluch et al., 2008).
 375 Moreover deacetylases, such as SIRT-1 exert the
 376 posttranslational modification of PGC-1 α promoter,
 377 which can affect its regulatory role on NRF-1 and
 378 mtTFA transcription (Canto and Auwerx, 2009). The
 379 complexity of this regulation is further demonstrated by
 380 the exercise-intensity-dependent regulation of PGC-1 α
 381 mRNA induction in the skeletal muscle (Egan et al.,
 382 2010). Further research is needed to establish the
 383 exercise-mediated molecular pathways involved in
 384 mitochondrial functions in cell specific manner.

385 Our results demonstrated that p-AMPK levels have
 386 been increased in the hippocampus following the 15-
 387 week-long physical training. The activation of PGC-1 α
 388 may be linked to the upstream activation of AMPK in
 389 the brain. Yu and co-workers (2010) found that the
 390 pharmacological stimulation of AMPK promoted the
 391 transcriptional activation of PGC-1 α in visual cortical
 392 neurons. The exercise-induced phosphorylation of
 393 AMPK has been well established in skeletal muscle
 394 (Canto et al., 2010). The regulation of AMPK activity by
 395 physical activity and its effect on hippocampal plasticity
 396 and metabolism has been proposed by Gomez-Pinilla
 397 et al. (2008). Recently, Bayod et al. (2011) also showed
 398 enhanced p-AMPK levels after 36 weeks of physical
 399 training in different brain regions including the
 400 hippocampus. Based on these observations, it is likely
 401 that the activation of AMPK could be a key molecular
 402 target of exercise in the hippocampus as well.

The major impact on AMPK signaling is the
 relationship between neuronal activity and energy
 demand (Yu and Yang, 2010). Calmodulin-dependent
 kinases (CaMKKs) have also been found to serve as
 upstream regulators of AMPK signaling in neurons
 (Hawley et al., 2005). Physical training elicits neuronal
 activation and modulates CaMKK (Hescham et al.,
 2009; Nishijima et al., 2012) which could underlay the
 stimulation of AMPK by physical activity in the
 hippocampus. On the other hand, moderate hypoxia
 and oxidative agents have been also shown to trigger
 CaMKK and AMPK (Mungai et al., 2011). It is possible
 that the exercise generates modest amount of ROS,
 which actually results in the activation of CaMKK/AMPK/
 PGC-1 α pathway and the subsequent activation of
 antioxidant enzymes. Thus exercise can increase the
 resistance toward further oxidative insults in long-term.
 As a consequence of adaptation, the vulnerability of the
 hippocampus to oxidative stress and diseases could be
 attenuated by physically active lifestyle.

CONCLUSIONS

In summary, the data from this study demonstrate that
 long term physical exercise results in lower levels of
 ROS and protein carbonyls as well as elevated levels of
 antioxidant enzymes in the hippocampus of aging rats.
 Physical training, thus, can be an effective option to
 regulate the oxidative balance and thus delay the onset
 of oxidative stress-related neurodegenerative processes.
 This study also provides further evidence that AMPK
 and PGC-1 α could be important molecules in mediating
 the beneficial effects of exercise in the hippocampus.

Acknowledgment—National Science and Research Foundation
 (K75702) awarded to Z. Radák. C. Nyakas was supported by
 Neuropeptide Foundation from The Netherlands.

REFERENCES

- 437
- 438 Aksenov MY, Aksenova MV, Butterfield DA, Geddes JW, Markesbery
439 WR (2001) Protein oxidation in the brain in Alzheimer's disease.
440 Neuroscience 103:373–383. 507
- 441 Asha Devi S (2009) Aging brain: prevention of oxidative stress by
442 vitamin E and exercise. Sci World J 9:366–372. 508
- 443 Aydin C, Sonat F, Sahin SK, Cangul IT, Ozkaya G (2009) Long term
444 dietary restriction ameliorates swimming exercise-induced
445 oxidative stress in brain and lung of middle-aged rat. Indian J
446 Exp Biol 47:24–31. 509
- 447 Bergersen LH, Storm-Mathisen J (2006) Training and brain health.
448 Tidsskr Nor Laegeforen 126:3253. 510
- 449 Bradford MM, Williams WL (1976) New, rapid, sensitive method for
450 protein determination. Fed Proc 35:274. 511
- 451 Canto C, Auwerx J (2009) PGC-1alpha, SIRT1 and AMPK, an energy
452 sensing network that controls energy expenditure. Curr Opin
453 Lipidol 20:98–105. 512
- 454 Canto C, Jiang LQ, Deshmukh AS, Matakis C, Coste A, Lagouge M,
455 Zierath JR, Auwerx J (2010) Interdependence of AMPK and
456 SIRT1 for metabolic adaptation to fasting and exercise in skeletal
457 muscle. Cell Metab 11:213–219. 513
- 458 Devi SA, Kiran TR (2004) Regional responses in antioxidant system
459 to exercise training and dietary vitamin E in aging rat brain.
460 Neurobiol Aging 25:501–508. 514
- 461 Douglas RJ (1967) The hippocampus and behavior. Psychol Bull
462 67:416–422. 515
- 463 Droge W (2003) Oxidative stress and aging. Adv Exp Med Biol
464 543:191–200. 516
- 465 Egan B, Carson BP, Garcia-Roves PM, Chibalin AV, Sarsfield FM,
466 Barron N, McCaffrey N, Moyna NM, Zierath JR, O'Gorman DJ
467 (2010) Exercise intensity-dependent regulation of peroxisome
468 proliferator-activated receptor coactivator-1 mRNA abundance is
469 associated with differential activation of upstream signalling
470 kinases in human skeletal muscle. J Physiol 588:1779–1790. 517
- 471 Emerit J, Edeas M, Bricaire F (2004) Neurodegenerative diseases
472 and oxidative stress. Biomed Pharmacother 58:39–46. 518
- 473 Esposito F, Ammendola R, Faraonio R, Russo T, Cimino F (2004)
474 Redox control of signal transduction, gene expression and cellular
475 senescence. Neurochem Res 29:617–628. 519
- 476 Ferrari CK (2007) Functional foods and physical activities in health
477 promotion of aging people. Maturitas 58:327–339. 520
- 478 Finkel T (2003) Oxidant signals and oxidative stress. Curr Opin Cell
479 Biol 15:247–254. 521
- 480 Floyd RA, Hensley K (2002) Oxidative stress in brain aging.
481 Implications for therapeutics of neurodegenerative diseases.
482 Neurobiol Aging 23:795–807. 522
- 483 Gaunt GL, de Duve C (1976) Subcellular distribution of D-amino
484 acid oxidase and catalase in rat brain. J Neurochem
485 26:749–759. 523
- 486 Gerschman R, Gilbert DL, Nye SW, Dwyer P, Fenn WO (1954)
487 Oxygen poisoning and X-irradiation: a mechanism in common.
488 Science 119:623–626. 524
- 489 Harman D (1956) Aging: a theory based on free radical and radiation
490 chemistry. J Gerontol 11:298–300. 525
- 491 Hawley SA, Pan DA, Mustard KJ, Ross L, Bain J, Edelman AM,
492 Frenguelli BG, Hardie DG (2005) Calmodulin-dependent protein
493 kinase kinase-beta is an alternative upstream kinase for AMP-
494 activated protein kinase. Cell Metab 2:9–19. 526
- 495 Heschem S, Grace L, Kellaway LA, Bugarith K, Russell VA (2009)
496 Effect of exercise on synaptophysin and calcium/calmodulin-
497 dependent protein kinase levels in prefrontal cortex and
498 hippocampus of a rat model of developmental stress. Metab
499 Brain Dis 24:701–709. 527
- 500 Hollmann W, Struder HK, Tagarakis CV, King G (2007) Physical
501 activity and the elderly. Eur J Cardiovasc Prev Rehabil
502 14:730–739. 528
- 503 Jager S, Handschin C, St-Pierre J, Spiegelman BM (2007) AMP-
504 activated protein kinase (AMPK) action in skeletal muscle via
505 direct phosphorylation of PGC-1alpha. Proc Natl Acad Sci USA
506 104:12017–12022. 529
- 519 Kim HJ, Kim KW, Yu BP, Chung HY (2000) The effect of age on
520 cyclooxygenase-2 gene expression: NF-kappaB activation and
521 IkappaBalpha degradation. Free Radic Biol Med 28:683–692. 530
- 522 Klinge CM (2008) Estrogenic control of mitochondrial function and
523 biogenesis. J Cell Biochem 105:1342–1351. 531
- 524 Lapolt PS, Yu SM, Lu JK (1988) Early treatment of young female rats
525 with progesterone delays the aging-associated reproductive
526 decline: a counteraction by estradiol. Biol Reprod 38:987–995. 532
- 527 Lee-Young RS, Griffiee SR, Lynes SE, Bracy DP, Ayala JE,
528 McGuinness OP, Wasserman DH (2009) Skeletal muscle AMP-
529 activated protein kinase is essential for the metabolic response to
530 exercise in vivo. J Biol Chem 284:23925–23934. 533
- 531 Levine RL, Stadtman ER (2001) Oxidative modification of proteins
532 during aging. Exp Gerontol 36:1495–1502. 534
- 533 Lopez-Lluch G, Irujo PM, Navas P, de Cabo R (2008) Mitochondrial
534 biogenesis and healthy aging. Exp Gerontol 43:813–819. 535
- 535 Meissner WW (1967) Hippocampus and learning. Int J
536 Neuropsychiatry 3:298–310. 536
- 537 Moorthy K, Sharma D, Basir SF, Baquer NZ (2005a) Administration of
538 estradiol and progesterone modulate the activities of antioxidant
539 enzyme and aminotransferases in naturally menopausal rats. Exp
540 Gerontol 40:295–302. 537
- 541 Moorthy K, Yadav UC, Siddiqui MR, Mantha AK, Basir SF, Sharma D,
542 Cowsik SM, Baquer NZ (2005b) Effect of hormone replacement
543 therapy in normalizing age related neuronal markers in different
544 age groups of naturally menopausal rats. Biogerontology
545 6:345–356. 538
- 546 Morris RG (2006) Elements of a neurobiological theory of
547 hippocampal function: the role of synaptic plasticity, synaptic
548 tagging and schemas. Eur J Neurosci 23:2829–2846. 539
- 549 Mungai PT, Waypa GB, Jairaman A, Prakriya M, Dokic D, Ball MK,
550 Schumacker PT (2011) Hypoxia triggers AMPK activation through
551 reactive oxygen species-mediated activation of calcium release-
552 activated calcium channels. Mol Cell Biol 31:3531–3545. 540
- 553 Nakamoto H, Kaneko T, Tahara S, Hayashi E, Naito H, Radak Z,
554 Goto S (2007) Regular exercise reduces 8-oxodG in the nuclear
555 and mitochondrial DNA and modulates the DNA repair activity in
556 the liver of old rats. Exp Gerontol 42:287–295. 541
- 557 Nishijima T, Okamoto M, Matsui T, Kita I, Soya H (2012)
558 Hippocampal functional hyperemia mediated by NMDA receptor/
559 NO signaling in rats during mild exercise. J Appl Physiol
560 112:197–203. 542
- 561 O'Neill HM, Maarbjerg SJ, Crane JD, Jeppesen J, Jorgensen SB,
562 Schertzer JD, Shyroka O, Kiens B, van Denderen BJ,
563 Tarnopolski MA, Kemp BE, Richter EA, Steinberg GR (2011)
564 AMP-activated protein kinase (AMPK) beta1beta2 muscle null
565 mice reveal an essential role for AMPK in maintaining
566 mitochondrial content and glucose uptake during exercise. Proc
567 Natl Acad Sci USA 108:16092–16097. 543
- 568 Offen D, Hochman A, Gorodin S, Ziv I, Shirvan A, Barzilai A,
569 Melamed E (1999) Oxidative stress and neuroprotection in
570 Parkinson's disease: implications from studies on dopamine-
571 induced apoptosis. Adv Neurol 80:265–269. 544
- 572 Ogonovszky H, Berkes I, Kumagai S, Kaneko T, Tahara S, Goto S,
573 Radak Z (2005) The effects of moderate-, strenuous- and over-
574 training on oxidative stress markers, DNA repair, and memory, in
575 rat brain. Neurochem Int 46:635–640. 545
- 576 Palacios OM, Carmona JJ, Michan S, Chen KY, Manabe Y, Ward 3rd
577 JL, Goodyear LJ, Tong Q (2009) Diet and exercise signals
578 regulate SIRT3 and activate AMPK and PGC-1alpha in skeletal
579 muscle. Aging (Albany, NY) 1:771–783. 546
- 580 Radak Z, Kaneko T, Tahara S, Nakamoto H, Ohno H, Sasvari M,
581 Nyakas C, Goto S (1999a) The effect of exercise training on
582 oxidative damage of lipids, proteins, and DNA in rat skeletal
583 muscle: evidence for beneficial outcomes. Free Radic Biol Med
584 27:69–74. 547
- 585 Radak Z, Pucsek J, Mecsek S, Csont T, Ferdinandy P (1999b)
586 Muscle soreness-induced reduction in force generation is
587 accompanied by increased nitric oxide content and DNA
588 damage in human skeletal muscle. Free Radic Biol Med
589 26:1059–1063. 548

- 578 Radak Z, Taylor AW, Ohno H, Goto S (2001) Adaptation to exercise-
579 induced oxidative stress: from muscle to brain. *Exerc Immunol*
580 *Rev* 7:90–107.
- 581 Sen CK, Packer L (1996) Antioxidant and redox regulation of gene
582 transcription. *FASEB J* 10:709–720.
- 583 Serrano F, Klann E (2004) Reactive oxygen species and synaptic
584 plasticity in the aging hippocampus. *Ageing Res Rev* 3:431–443.
- 585 Shin MK (2009) Effects of an exercise program on frontal lobe
586 cognitive function in elders. *J Korean Acad Nurs* 39:107–115.
- 587 Sinet PM, Heikkila RE, Cohen G (1980) Hydrogen peroxide
588 production by rat brain in vivo. *J Neurochem* 34:1421–1428.
- 589 St-Pierre J, Drori S, Uldry M, Silvaggi JM, Rhee J, Jager S,
590 Handschin C, Zheng K, Lin J, Yang W, Simon DK, Bachoo R,
591 Spiegelman BM (2006) Suppression of reactive oxygen species
592 and neurodegeneration by the PGC-1 transcriptional coactivators.
593 *Cell* 127:397–408.
- 594 Stadtman ER (2001) Protein oxidation in aging and age-related
595 diseases. *Ann N Y Acad Sci* 928:22–38.
- 596 Toldy A, Stadler K, Sasvari M, Jakus J, Jung KJ, Chung HY, Berkes I,
597 Nyakas C, Radak Z (2005) The effect of exercise and nettle
supplementation on oxidative stress markers in the rat brain. *Brain Res Bull* 65:487–493.
- 598 Valle I, Alvarez-Barrientos A, Arza E, Lamas S, Monsalve M (2005)
599 PGC-1alpha regulates the mitochondrial antioxidant defense
600 system in vascular endothelial cells. *Cardiovasc Res* 66:
601 562–573.
- 602 Venkateshappa C, Harish G, Mahadevan A, Srinivas Bharath MM,
603 Shankar SK (2012) Elevated oxidative stress and decreased
604 antioxidant function in the human hippocampus and frontal cortex
605 with increasing age: implications for neurodegeneration in
606 Alzheimer's disease. *Neurochem Res* 37:1601–1614.
- 607 Ventura-Clapier R, Garnier A, Veksler V (2008) Transcriptional
608 control of mitochondrial biogenesis: the central role of PGC-
609 1alpha. *Cardiovasc Res* 79:208–217.
- 610 Yu L, Yang SJ (2010) AMP-activated protein kinase mediates activity-
611 dependent regulation of peroxisome proliferator-activated
612 receptor gamma coactivator-1alpha and nuclear respiratory
613 factor 1 expression in rat visual cortical neurons. *Neuroscience*
614 169:23–38.
- 615
616

(Accepted 1 September 2012)
(Available online xxxx)

UNCORRECTED PROOF