

This is the author's pre-print version.

The final publication is available at Springer via DOI 10.1007/s11738-014-1753-z

Citation: Ayaydin et al. *Acta Physiol Plant* (2015) 37:3

1

2

3

4

5

6

7

8

9

10

11

12 ***Arabidopsis* NAP-related proteins (NRPs) are soluble nuclear proteins**

13 **immobilized by heat**

14

15 Ferhan Ayaydin, Judit Bíró, Mónika Domoki, Györgyi Ferenc, Attila Fehér

16 *Institute of Plant Biology, Biological Research Center, Hungarian Academy of Sciences,*

17 *Temesvári krt. 62., H-6726 Szeged, Hungary*

18

19

20

21

22

23

24

25

26

27

28

29

30

31 Corresponding author:

32 Attila Fehér, Institute of Plant Biology, Biological Research Centre, Hungarian Academy of

33 Sciences, Temesvári krt. 62., H-6726 Szeged, Hungary; P.O.Box. 521., H-6701 Szeged,

34 Hungary

35 Phone: 36 62599701; Fax: 36 62433434; e-mail: feher.attila@brc.mta.hu

This is the author's pre-print version.

The final publication is available at Springer via DOI 10.1007/s11738-014-1753-z

Citation: Ayaydin et al. *Acta Physiol Plant* (2015) 37:3

## 1 **Abstract**

2 Nucleosome assembly protein-related proteins (NRPs) are multifunctional proteins having histone  
3 chaperone and phosphatase inhibitor properties. Although it is believed that these proteins are nuclear  
4 and bind the chromatin, they can be detected in the cytoplasmic but not in the nuclear protein fraction  
5 by immunoblotting analysis. It is shown here that under normal conditions NRPs are nuclear but soluble  
6 and leak out of the nuclei during their purification. However, under elevated temperatures (above 42°C),  
7 NRPs display significantly reduced mobility and are retained in the nuclei during purification probably  
8 due to binding other immobile macromolecules in the nucleus. Our observations highlight the necessity  
9 to use different techniques in parallel to unambiguously determine the intracellular localization of  
10 proteins. As heat adapted (38°C 2 h followed by 2 h recovery) and heat shocked (45°C, 1 h) *Arabidopsis*  
11 seedlings were found to have phenotypes similar to those observed in the NRP loss-of-function mutants  
12 *nrp1-1 nrp2-1* (short, branching roots, increased bleomycin sensitivity), it was also investigated whether  
13 the immobilization of NRPs by heat results in disturbed NRP functions. The results indicated, however,  
14 that heat affected the investigated traits independent of the presence of NRPs.

15

16

17

18 **Keywords:** *Arabidopsis thaliana* (L.), cellular localization, heat shock, nuclear protein,  
19 nucleosome assembly protein-related protein, protein mobility

20

21

22

23

24

25

26

27

28

29

30

31

32

33

This is the author's pre-print version.

The final publication is available at Springer via DOI 10.1007/s11738-014-1753-z

Citation: Ayaydin et al. *Acta Physiol Plant* (2015) 37:3

1

2

3 **Author Contribution Statement**

4 Ferhan Ayaydin: cell biology, microscopy, manuscript revision; Judit Bíró: protein purification,

5 immunoblotting; Mónika Domoki: *Arabidopsis* culture, phenotype analysis; Györgyi Ferenc:

6 RT-qPCR; Attila Fehér: research concept, experimental design, data analysis and evaluation,

7 manuscript writing.

## 1 **1. Introduction**

2 Even small environmental changes can induce gene expression or repression of hundreds of  
3 genes in plants. Regulation of such synchronized events requires chromatin remodeling – a  
4 process that involves histones, the core protein molecules of nucleosomes (Vermaak 2003).  
5 One group of proteins implicated in the regulation of histone modification patterns are the NAP  
6 (Nucleosome Assembly Protein) family of histone chaperones, which possess a central  
7 conserved NAP domain, and a highly acidic C-terminal domain (Park and Luger 2006). Histone  
8 chaperones play a crucial role in nucleosome assembly and are thought to be necessary for  
9 prevention of nonproductive aggregation between the highly positive charged histones and the  
10 highly negative charged DNA.

11 Plant NAP-related proteins (NRPs), form a distinct phylogenetic group within the NAP family  
12 and are closely related to the animal SET/I<sub>2</sub><sup>PP2A</sup> (patient SE translocation/inhibitor 2 of protein  
13 phosphatase 2A) protein (Dong et al. 2003). SET was first identified as the product of a  
14 translocated gene in acute undifferentiated leukemia (Adachi et al. 1994). In biochemical  
15 assays, SET/I<sub>2</sub><sup>PP2A</sup> proteins stimulate replication of the adenovirus genome and inhibit protein  
16 phosphatase 2A (PP2A) (Haruki et al. 2006; Li et al. 1996). The SET/I<sub>2</sub><sup>PP2A</sup> protein was  
17 identified in protein complexes with histone acetylation or methylation enzymes (Cervoni et al.  
18 2002), B-type cyclins (Kellogg and Kikuchi 1995), a granzyme A-activated DNase (Fan et al.  
19 2003) or transcription coactivators (Karetsou et al. 2005). SET/I<sub>2</sub><sup>PP2A</sup> was also implicated in the  
20 heat shock response of fruit fly blocking histone H3 dephosphorylation at heat-shock loci  
21 (Nowak and Corces 2000). Concerning the plant NRPs, it has been shown that *Arabidopsis*  
22 NRP1 (UniProt accession: Q9CA59) and NRP2 (UniProt accession: Q8LC68) have important  
23 roles in cell cycle regulation and root meristem maintenance in addition to their potential  
24 histone chaperone and histone phosphatase inhibitor functions (Bíró et al. 2012; Zhu et al.

This is the author's pre-print version.

The final publication is available at Springer via DOI 10.1007/s11738-014-1753-z

Citation: Ayaydin et al. *Acta Physiol Plant* (2015) 37:3

1 2006). Although the *Arabidopsis* single *nrp1-1* or *nrp2-1* mutants do not exhibit any observable  
2 phenotype under *in vitro* or greenhouse conditions (Zhu et al. 2006), *nrp1-1 nrp2-1* double  
3 mutants have short branched roots, increased sensitivity against genotoxic stress, and impaired  
4 somatic homologous recombination (Gao et al. 2012; Zhu et al. 2006).  
5 In contrast to their above described nuclear functions, plant NRPs could be detected in  
6 cytoplasmic cell fractions by immunoblotting (Bíró et al. 2013). Here we report a series of  
7 experiments that was carried out in order to clarify the intracellular localization of NRPs. It was  
8 established that *Arabidopsis* NRPs are indeed nuclear proteins and their accumulation in the  
9 supernatant during the isolation of purified nuclei for immunoblotting analysis is only an  
10 experimental artefact due to their small size and high mobility. The obtained results underline  
11 the necessity to use various techniques to validate the intracellular localization of proteins since  
12 the isolation of nuclei, fixation, or the attachment of a large GFP-tag to small proteins all can  
13 strongly influence the obtained results as exemplified here by NRPs. The experimental results  
14 also indicate that the NRP proteins despite their known histone-binding capability are not  
15 strongly bound to the chromatin under normal conditions. However, NRPs become cross-linked  
16 to other proteins in plant nuclei subjected to heat shock (above 42°C for 30-60 min). The  
17 potential contribution of this phenomenon to the phenotype of heat-treated *Arabidopsis* plants  
18 similar to that of NRP-null mutants was also investigated. It was found that heat affected the  
19 phenotype of *Arabidopsis* plantlets independent of the presence of NRPs. The dissimilarity  
20 between plant and fruit fly NRP-type proteins in respect to their role in heat responses is  
21 discussed.

22

## 1 **2. Materials and methods**

### 2 **2.1. Plant cultures and heat treatment**

3 Experiments were carried out with the wild-type Columbia ecotype of *Arabidopsis thaliana*  
4 (L.) Heynh or its double mutant *nrp1-1 nrp2-1*. Seeds of the *nrp1-1 nrp2-1* double mutant  
5 *Arabidopsis* plants deficient in At NRP1 and At NRP2 expression were kindly provided by Dr.  
6 Yan Zhu (State Key Laboratory of Genetic Engineering, Institute of Plant Biology, School of  
7 Sciences, Fudan University, Shanghai, China). Neither the transcripts (Zhu et al. 2006) nor the  
8 proteins (Bíró et al. 2012) of NRP1 or NRP2 could be detected in the mutant seedlings.

9 Seeds of *Arabidopsis thaliana* were surface-sterilized by soaking in 70% ethanol (v/v) for 1  
10 min then in 30% (v/v) commercial bleach (with 4.8% w/w sodium hypochloride) for 10 min  
11 and then rinsed five times with sterile distilled water and plated on solid Murashige and Skoog  
12 (MS) medium (Duchefa Biochemie, B.V., Haarlem, The Netherlands) containing 3% (w/v)  
13 sucrose, 0.6% (w/v) agar, with pH adjusted to 5.8. The seeds were germinated at 24°C using an  
14 8 h light/16 h dark regime. Experiments were carried out at least in triplicates with 7-8-days-  
15 old seedlings. Heat shock treatment was applied in a water bath at 45°C for 1 h with or without  
16 preceding heat acclimation (38°C for two hours followed by two hours recovery at 22°C).  
17 Survival rates were determined by scoring the percentage of healthy seedlings with green leaves  
18 that showed continuous growth during six weeks after the treatment.

### 19 **2.2. Protein purification and detection**

20 Total plant proteins were extracted by grinding samples with quartz sand in the homogenization  
21 buffer containing 25 mM Tris-HCl pH 7.6, 15 mM MgCl<sub>2</sub>, 15 mM EGTA, 75 mM NaCl, 1 mM  
22 NaF, 60 mM β-glycero-phosphate, 1 mM PMSF, 2 mM DTT and 10 μl/ml plant-specific  
23 protease inhibitor cocktail (Sigma, St-Louis, USA). Nuclei and the nuclear protein fraction were  
24 purified according to (Bowler et al. 2004). The supernatant following the precipitation of nuclei

This is the author's pre-print version.

The final publication is available at Springer via DOI 10.1007/s11738-014-1753-z

Citation: Ayaydin et al. *Acta Physiol Plant* (2015) 37:3

1 by centrifugation was further cleared by an additional centrifugation at 10000 g for 10 min, at  
2 4°C, and was used for the isolation of the cytosolic protein fraction. Detection of NRP proteins  
3 by immunoblotting was performed according to standard procedures as described elsewhere in  
4 more details (Bíró et al. 2012).

### 5 **2.3. Immunohistochemistry on protoplasts and isolated nuclei**

6 Protoplasts were prepared from *Arabidopsis* cell suspension with the enzyme solution of 3%  
7 Cellulase RS, 1% Cellulase R-10, 1% Macerozyme R-10, 0.5 M mannitol, 5 mM CaCl<sub>2</sub>, pH  
8 5.3. They were then fixed with 4% formaldehyde (Sigma). In experiments where isolated nuclei  
9 were used for immunofluorescence (Fig 2d and 2e), nuclei were isolated by chopping unfixed  
10 (heat-treated or control) cells on ice with a razor blade (Kotogány et al. 2010) then fixed with  
11 1% formaldehyde on ice for 30 min. Following fixation, cells or nuclei were immobilized on  
12 poly-L-lysine coated slides and rehydrated with 0.05% Triton X-100. The polyclonal anti-NRP  
13 antibody (see Bíró et al. 2012) was diluted 1:200 in phosphate-buffered saline (PBS). After 1 h  
14 incubation with the diluted primary antibody, the slides were washed three times with PBS.  
15 Samples were then incubated with the secondary anti-rabbit antibody conjugated to fluorescein  
16 isothiocyanate (FITC) (Sigma). Following final washes with PBS, slides were counterstained  
17 with 4',6-diamidino-2-phenylindole (DAPI). Fluorescence was recorded using an Olympus  
18 FV1000 confocal laser scanning microscope.

### 19 **2.4. Transient gene expression and the fluorescence loss in photobleaching** 20 **(FLIP) assay**

21 For the transient expression assay, the pK7FW62 vector (Karimi et al. 2002) was digested with  
22 *Hind*III and *Xba*I and the resulted fragment was subcloned into pBSK vector cut with the same  
23 restriction enzymes resulting in pBSK-GFP-GW. A PCR fragment carrying the coding

This is the author's pre-print version.

The final publication is available at Springer via DOI 10.1007/s11738-014-1753-z

Citation: Ayaydin et al. *Acta Physiol Plant* (2015) 37:3

1 sequence of At NRP1 was inserted into the vector by GATEWAY recombination using LR  
2 recombinase (Invitrogen, Carlsbad, USA). The constructs were sequence verified.  
3 Isolation and transfection of *Arabidopsis* cell culture protoplasts were made according to  
4 (Mathur et al. 1995).  
5 Fluorescence loss in photobleaching (FLIP) assay was performed using an Olympus FV1000  
6 confocal laser scanning microscope as described in Ayaydin and Dasso (2004). After heat-  
7 treatment (1h at 45°C or 25°C) of one-day-old *Arabidopsis* protoplasts expressing the  
8 GFP:NRP1 protein, an intranuclear spot (1  $\mu\text{m}$  x 1  $\mu\text{m}$  square) was continuously photobleached  
9 using high intensity 488nm Argon laser excitation. During spot bleaching, images of the whole  
10 nucleus were also captured. On captured images, intranuclear GFP:NRP1 intensity changes  
11 were recorded at a fixed distance (5  $\mu\text{m}$ ) away from the bleaching spot. In case of immobile  
12 proteins only the tight bleaching spot is expected to be photobleached and depleted, while for  
13 highly mobile nuclear proteins whole nuclear fluorescence intensity is expected to drop  
14 considerably. Recorded values were plotted to compare the intranuclear mobility of GFP:NRP1  
15 on heat-treated and untreated cells.

## 16 **2.5. Gene expression analysis by real-time quantitative PCR (RT-qPCR)**

17 RNA isolation, cDNA synthesis, and RT-qPCR analysis by an ABI Prism 7700 sequence  
18 detection system have been described earlier in details (Domoki et al. 2006). DNA damage and  
19 repair-associated (*PARP2*, At4g02390; *RAD51*, At5g20850; *AtMYB*, At5g03780; *RNR2*,  
20 At3g27060) as well as the control unresponsive (*UBC18*, At5g42990) genes were selected  
21 based on the work of Chen et al. (2003). Primer sequences are listed as supplementary material  
22 in Supplementary Table 1.

1

## 2 **3. Results and Discussion**

3

### 4 **3.1. NRPs are soluble nuclear proteins**

5 The animal homologue of plant NRPs, the SET protein, has various functions ranging from the  
6 inhibition of different enzyme complexes (protein phosphatase 2A (PP2A), histone deacetylase,  
7 granzyme A-activated DNase) to the regulation of transcription and nucleosome assembly (Park  
8 and Luger 2006). The information regarding the function of plant NRPs is limited, but these  
9 proteins are most likely multifunctional, as well. They have been reported to inhibit PP2A  
10 activity *in vitro* (Bíró et al. 2012) and to possess histone chaperone-like properties (Zhu et al.  
11 2006). Their possible roles in cell division control, gene silencing, homologous recombination,  
12 and sensitivity against DNA damaging agents have also been shown (Gao et al. 2012, Zhu et  
13 al. 2006) Most of the above mentioned functions require the nuclear localization of the NRP  
14 proteins. Fluorescent protein-tagged NRPs localize to the nuclei of *Arabidopsis* cells (Fig 1a)  
15 in accord with their possible functions (Zhu et al. 2006). However, the detection of NRP  
16 proteins in cytoplasmic cell fractions by immunoblotting contradicts these results (Fig. 1c; Bíró  
17 et al. 2013).

18 In order to clarify this discrepancy regarding the localization of NRPs, we carried out further  
19 experiments. In transfected protoplasts, the ectopically expressed GFP-fused At NRP1 could  
20 exclusively be detected by immunoblotting in the nuclear fraction, while the endogenous  
21 untagged NRPs were accumulating in the cytoplasmic (supernatant) fraction (Fig. 1d). This  
22 result suggested that the increased size due to the GFP-tag may be responsible for the nuclear  
23 retention of the protein fusion. We tested this hypothesis by determining the localization of  
24 endogenous NRP proteins in intact cells by indirect immunofluorescence using the same At

This is the author's pre-print version.

*The final publication is available at Springer via DOI 10.1007/s11738-014-1753-z*

*Citation: Ayaydin et al. Acta Physiol Plant (2015) 37:3*

1 NRP1 antibody as in immunoblotting experiments. The strong nuclear fluorescence indicated  
2 that the NRP proteins are predominantly nuclear in formaldehyde-fixed intact cells (Fig. 1b,  
3 asterisks indicate two NRP-containing nuclei). The GFP-tag, therefore, is not required for the  
4 nuclear localization of NRPs. In a subsequent experiment, we investigated the effect of cell  
5 fixation on NRP detection in nuclear and cytoplasmic fractions. Formaldehyde-fixed and  
6 unfixed cells were fractionated, and the fractions were tested for the presence of NRPs by  
7 immunoblotting. The NRP proteins could only be detected in the nuclei of fixed but not that of  
8 the unfixed cells (Fig. 1e). Moreover, the anti-NRP antibodies in immunolabeling experiments  
9 did not label the mitotic chromosomes in dividing cells, indicating that the protein is not  
10 chromatin-bound in these cells, either (Fig. 1b, arrow indicates a mitotic cell devoid of NRP  
11 labeling).

12 These data altogether can be explained if we suppose that the relatively small (<30 kD), nuclear  
13 NRP proteins leak out from the nuclei into the supernatant during cell fractionation. GFP-tagged  
14 NRP proteins having a size of approximately 60 kD are most likely large enough to be retained  
15 within the isolated nuclei, while formaldehyde fixation may crosslink NRPs to other  
16 macromolecules and prevents leakage. YFP-NRP fusions were reported to be bound to the  
17 chromatin (Zhu et al. 2006). Our observations using an anti-NRP1 antibody and cell  
18 fractionation are against a strong chromatin binding and support a view that NRPs are soluble  
19 and mobile within unfixed nuclei. Further experimental data obtained during the investigation  
20 of the effect of heat on NRP solubility supported this hypothesis.

21

### 22 **3.2 Heat shock immobilizes NRPs within the nucleus**

23 In previous publications, we have reported the analysis of the potential role of NRPs in heat  
24 stress response of plants (Bíró et al. 2012, 2013). During our experiments we have realized that

This is the author's pre-print version.

The final publication is available at Springer via DOI 10.1007/s11738-014-1753-z

Citation: Ayaydin et al. *Acta Physiol Plant* (2015) 37:3

1 strong heat-treatment (45°C, 1 h) of *Arabidopsis* cells resulted in the retention of NRPs in  
2 isolated nuclei in contrast to the untreated cells (Bíró et al. 2013). In our follow up experiments  
3 presented here, it was established that lower temperature (37°C) was not but shorter time (half  
4 an hour) at 45°C heat treatment was sufficient to achieve this retention (Fig. 2a,b). Moreover,  
5 it was found that the retention was sustained. Nuclei which were isolated 42 h after the one-  
6 hour heat treatment still contained NRPs (Fig. 2b). Nuclear retention of NRPs in response to  
7 heat was also verified in heat-treated *Arabidopsis* seedlings (Fig. 2c). Preconditioning of the  
8 plantlets at 37°C prior to heat-treatment (45°C, 1 h) did not prevent this nuclear retention  
9 phenomenon (Fig. 2c). In order to verify the results obtained by immunoblotting,  
10 immunofluorescence-based microscopy assays were performed using nuclei isolated from  
11 unfixed control and heat-treated cells (Fig. 2d,e). Only the nuclei derived from heat-treated cells  
12 exhibited immunofluorescence signal indicating the presence of NRPs. Other stress treatments,  
13 including, heavy metal, salt, genotoxic and UV stress, did not result in the retention of NRPs  
14 during the isolation of nuclei (Supplementary Fig. 1) indicating that the phenomenon is heat  
15 specific.

16 Fluorescence loss in photobleaching (FLIP) assay using laser scanning confocal microscope on  
17 live cells provided further proof that NRPs are mobile soluble nuclear proteins, but are  
18 immobilized in response to heat stress. In this experiment, *Arabidopsis* protoplasts were  
19 transfected with a GFP-At NRP construct. GFP fluorescence was bleached repetitively only in  
20 a restricted, 1 square micron area within selected fluorescent nuclei, and the depletion of  
21 emission was measured at a fixed position away from the depletion zone (Ayaydin and Dasso  
22 2004). In control protoplasts (no heat treatment), successive spot-bleaching resulted in the  
23 gradual loss of fluorescence in the whole nucleus due to the mobility (solubility) of NRPs (Fig.  
24 3a). In contrast, when the protoplasts were treated by heat (45°C, 1 h), only the excited area lost  
25 the fluorescence signal indicating that the NRP proteins got immobilized and could not diffuse

This is the author's pre-print version.

The final publication is available at Springer via DOI 10.1007/s11738-014-1753-z

Citation: Ayaydin et al. *Acta Physiol Plant* (2015) 37:3

1 to the site of bleaching. Quantitative analyses corroborated these microscopy observations in  
2 that the kinetics of loss of fluorescence displayed a dramatic drop of nuclear fluorescence  
3 intensity in untreated samples as compared to heat-treated samples (Fig. 3b).  
4 DNase or RNase treatment of the heat-treated nuclei prior to immunoblotting, did not affect  
5 nuclear retention (Fig. 3c), suggesting that the heat treatment resulted in crosslinking of NRPs  
6 to other nuclear proteins rather than nucleic acid polymers.  
7 The animal SET/I<sub>2</sub><sup>PP2A</sup> protein, the homologue of plant NRPs, is considered to be a scaffold  
8 protein being present in several large protein complexes. The same may hold true for plant  
9 NRPs. However, until now, *Arabidopsis* NRPs were shown to interact only with histones and  
10 the catalytic subunit of the plant PP2A phosphatase (Bíró et al. 2012; Zhu et al. 2006). It has  
11 long been known that heat causes protein aggregation also in the nuclei of animal cells (Borrelli  
12 et al. 1996; Laszlo et al. 1992; Littlewood et al. 1987; Roti Roti and Turkel 1994 a,b; Warters  
13 et al. 1986). Although the degree of heat-induced protein aggregation could be correlated with  
14 the thermotolerance of HeLa cells (Roti Roti and Turkel 1994 a,b) and several oncogenes have  
15 also been found to be associated with these complexes (Littlewood et al. 1987), the significance  
16 of the phenomenon is unclear.

### 17 **3.3. NRP immobilization does not result in an NRP loss-of-function phenotype**

18 Although acquired thermotolerance can ensure survival, extreme heat alters the development  
19 of even acclimatized plants. *Arabidopsis* seedlings exposed to a short but strong heat treatment  
20 (45°C, 1 h) following an acclimation period (38°C, 2 h) exhibit an almost 100% survival rate  
21 but show reduced hypocotyl and root length as compared to untreated controls (see e.g.  
22 Larkindale and Knight 2002). This indicates that the thermotolerance mechanisms do not  
23 prevent certain sublethal developmental changes caused by heat.

This is the author's pre-print version.

The final publication is available at Springer via DOI 10.1007/s11738-014-1753-z

Citation: Ayaydin et al. *Acta Physiol Plant* (2015) 37:3

1 In order to determine whether the heat-mediated crosslinking of NRPs to the chromatin  
2 contributes to the phenotype of heat-treated seedlings, we compared the NRP loss-of-function  
3 *nrp1-1 nrp2-1* double mutants (Zhu et al. 2006) to seedlings acclimatized at 38°C for two hours  
4 followed by one hour heat shock (45°C) (see Materials and methods for details). Hundred  
5 percent of acclimatized plants could survive the applied, otherwise lethal, heat shock treatment  
6 that resulted in the crosslinking of NRPs to the chromatin (Fig 4a, see also Fig 2c). Therefore,  
7 these plants were suitable for comparison to the *nrp1-1 nrp2-1* mutants.

8 The *nrp1-1 nrp2-1* mutants have been characterized by Zhu et al. (2006). One of the  
9 characteristic phenotypes of *nrp1-1 nrp2-1* seedlings is altered root growth resulting in a small,  
10 branched root system, very similar to that of heat-treated *Arabidopsis* plantlets (Fig. 4a).  
11 However, heat-treatment of the *nrp1-1 nrp2-1* plantlets resulted in further root growth  
12 retardation (Fig. 4b) indicating that the effects of heat and NRP loss-of-function are additive  
13 and therefore it is unlikely that they affect the same pathway.

14 Another distinctive phenotype of the *nrp1-1 nrp2-1* mutants is their sensitivity towards  
15 bleomycin, a DNA damaging agent (Zhu et al. 2006). The survival rate of bleomycin-treated  
16 *nrp1-1 nrp2-1* mutant seedlings (~50%) was similar to wild type ones subjected to combined  
17 genotoxic and heat stresses (~39%) (Fig. 5a). Heat and *nrp1-1 nrp2-1* mutation together  
18 decreased the survival rate of seedlings on bleomycin plates down to ~23% indicating again  
19 that heat and NRP loss-of-function do not affect the same pathway. This sensitivity was not due  
20 to the altered expression of genes implicated in DNA repair as these genes exhibited similar  
21 induction under the investigated conditions (Fig. 5b).

22

### 1 **3.4. NRPs and the heat response of *Arabidopsis***

2 The NRP homolog animal SET/I<sub>2</sub><sup>PP2A</sup> protein regulates the expression of heat shock protein  
3 (HSP) genes through the inhibition of histone H3 dephosphorylation at heat-shock gene loci  
4 (Nowak et al. 2003). We have previously shown that NRPs are dispensable for the heat-induced  
5 expression of heat shock protein genes in *Arabidopsis*, despite the fact that they act as potent  
6 histone H3 phosphatase inhibitors *in vitro* (Bíró et al. 2012). Here, we provide further evidence  
7 that NRPs are not responsible for heat-induced phenotypes of high temperature acclimated  
8 *Arabidopsis* plants; although there are remarkable phenotypic similarities between NRP-  
9 deficient mutants and heat-treated acclimated wild types (Figs. 4-5). Therefore, NRPs seem not  
10 to be involved in acquired thermotolerance in a way described for the orthologous *Drosophila*  
11 SET protein. Nevertheless, the overexpression of the At NRP1 protein slightly increased the  
12 survival of non-acclimated *Arabidopsis* cells/seedlings exposed to high temperature (45°C for  
13 1 h; Bíró et al. 2013). In contrast, the *nrp1-1 nrp2-1 Arabidopsis* mutants did not exhibit  
14 increased heat sensitivity according to our investigations (data not shown). We hypothesize that  
15 the improved survival of NRP-overexpressing *Arabidopsis* cells exposed to heat might be  
16 associated with the crosslinking of NRPs to other nuclear proteins in response to heat (see 3.2).  
17 This hypothesis, however, still needs to be experimentally validated.

18

19

### 20 **Acknowledgements**

21 The presented work was supported by the Hungarian Scientific Research Fund (grant# K68051)  
22 and the National Innovation Office (contract# OTKA\_A\_07-2-2008-0022).

23

This is the author's pre-print version.

*The final publication is available at Springer via DOI 10.1007/s11738-014-1753-z*

*Citation: Ayaydin et al. Acta Physiol Plant (2015) 37:3*

1 **Conflict of interest**

2 The authors declare that they have no conflict of interest.

3

1

## 2 **References**

- 3 Adachi Y, Pavlakis GN, Copeland TD (1994) Identification and characterization of SET, a  
4 nuclear phosphoprotein encoded by the translocation break point in acute  
5 undifferentiated leukemia. *Mol Cell* 269:2258–2262.
- 6 Ayaydin F, Dasso M (2004) Distinct *in vivo* dynamics of vertebrate SUMO paralogues. *Mol*  
7 *Biol Cell* 15:5208–5218. doi: 10.1091/mbc.E04-07-0589
- 8 Bíró J, Domoki M, Fehér A (2013) NAP-Related Protein 1 (Atnrp1) overexpression increases  
9 the heat tolerance of *Arabidopsis* cells/plantlets. *J Plant Biochem Physiol* 1:1–3. doi:  
10 10.4172/2329-9029.1000115
- 11 Bíró J, Farkas I, Domoki M, Ötvös K, Bottka S, Dombrádi V, Fehér A (2012) The histone  
12 phosphatase inhibitory property of plant nucleosome assembly protein-related proteins  
13 (NRPs). *Plant Physiol Biochem* 52:162–168. doi: 10.1016/j.plaphy.2011.12.010
- 14 Borrelli MJ, Lepock JR, Frey HE, Lee YJ, Corry PM (1996) Excess protein in nuclei isolated  
15 from heat-shocked cells results from a reduced extractability of nuclear proteins. *J Cell*  
16 *Physiol* 167:369–379. doi: 10.1002/(SICI)1097-4652(199606)167:3<369::AID-  
17 JCP1>3.0.CO;2-N
- 18 Bowler C, Benvenuto G, Laflamme P, Molino D, Probst AV, Tariq M, Paszkowski J (2004)  
19 Chromatin techniques for plant cells. *Plant J* 39:776–789. doi: 10.1111/j.1365-  
20 313X.2004.02169.x
- 21 Cervoni N, Detich N, Seo S-B, Chakravarti D, Szyf M (2002) The oncoprotein Set/TAF-  
22 1beta, an inhibitor of histone acetyltransferase, inhibits active demethylation of DNA,  
23 integrating DNA methylation and transcriptional silencing. *J Biol Chem* 277:25026–  
24 25031. doi: 10.1074/jbc.M202256200
- 25 Chen I-P, Haehnel U, Altschmied L, Schubert I, Puchta H (2003) The transcriptional response  
26 of *Arabidopsis* to genotoxic stress - a high-density colony array study (HDCA). *Plant J*  
27 35:771–786. doi: 10.1046/j.1365-313X.2003.01847.x
- 28 Domoki M, Györgyey J, Bíró J, Pasternak TP, Zvara A, Bottka S, Puskás LG, Dudits D,  
29 Fehér A (2006) Identification and characterization of genes associated with the induction  
30 of embryogenic competence in leaf-protoplast-derived alfalfa cells. *Biochim Biophys*  
31 *Acta* 1759:543–551. doi: 10.1016/j.bbaexp.2006.11.005
- 32 Dong A, Zhu Y, Yu Y, Cao K, Sun C, Shen WH (2003) Regulation of biosynthesis and  
33 intracellular localization of rice and tobacco homologues of nucleosome assembly  
34 protein 1. *Planta* 216:561–570. doi: 10.1007/s00425-002-0910-6
- 35 Fan Z, Beresford P, Oh D, Zhang D, Lieberman J (2003) Tumor suppressor NM23-H1 is a  
36 granzyme A-activated DNase during CTL-mediated apoptosis, and the nucleosome

This is the author's pre-print version.

The final publication is available at Springer via DOI 10.1007/s11738-014-1753-z

Citation: Ayaydin et al. *Acta Physiol Plant* (2015) 37:3

- 1 assembly protein SET is its inhibitor. *Cell* 112:659–672. doi: 10.1016/S0092-  
2 8674(03)00150-8
- 3 Gao J, Zhu Y, Zhou W, Molinier J, Dong A, Shen, W-H (2012) NAP1 family histone  
4 chaperones are required for somatic homologous recombination in *Arabidopsis*. *Plant*  
5 *Cell* 24:1437–1447. doi: 10.1105/tpc.112.096792
- 6 Haruki H, Okuwaki M, Miyagishi M, Taira K, Nagata K (2006) Involvement of template-  
7 activating factor I/SET in transcription of adenovirus early genes as a positive-acting  
8 factor. *J Virol* 80:794–801. doi: 10.1128/JVI.80.2.794-801.2006
- 9 Karetsov Z, Martic G, Sflomos G, Papamarcaki T (2005) The histone chaperone SET/TAF-  
10 Ibeta interacts functionally with the CREB-binding protein. *Biochem Biophys Res*  
11 *Commun* 335:322–327. doi: 10.1016/j.bbrc.2005.06.210
- 12 Karimi M, Inzé D, Depicker A (2002) GATEWAY vectors for *Agrobacterium*-mediated plant  
13 transformation. *Trends Plant Sci* 7:193–195. doi: 10.1016/S1360-1385(02)02251-3
- 14 Kellogg D, Kikuchi A (1995) Members of the NAP/SET family of proteins interact  
15 specifically with B-type cyclins. *J Cell Biol* 130:661–673.
- 16 Kotogány E, Dudits D, Horváth GV, Ayaydin F. (2010) A rapid and robust assay for detection  
17 of S-phase cell cycle progression in plant cells and tissues by using ethynyl  
18 deoxyuridine. *Plant Methods*. 6(1):5 doi: 10.1186/1746-4811-6-5.
- 19 Larkindale J, Knight MR (2002) Protection against heat stress-induced oxidative damage in  
20 *Arabidopsis* involves calcium, abscisic acid, ethylene, and salicylic acid. *Plant Physiol*  
21 128:682–695. doi: 10.1104/pp.010320
- 22 Laszlo A, Wright W, Roti Roti JL (1992) Initial characterization of heat-induced excess  
23 nuclear proteins in HeLa cells. *J Cell Physiol* 151:519–532. doi:  
24 10.1002/jcp.1041510311
- 25 Li M, Makkinje A, Damuni Z (1996) Thymeloid leukemia-associated protein SET is a potent  
26 inhibitor of protein phosphatase 2A. *J Biol Chem* 271:11059–11062. doi:  
27 10.1074/jbc.271.19.11059
- 28 Littlewood TD, Hancock DC, Evan GI (1987) Characterization of a heat shock-induced  
29 insoluble complex in the nuclei of cells. *J Cell Sci* 88:65–72.
- 30 Mathur J, Koncz C, Szabados L (1995) A simple method for isolation, liquid culture,  
31 transformation and regeneration of *Arabidopsis thaliana* protoplasts. *Plant Cell Rep*  
32 14:221–226. doi: 10.1007/BF00233637
- 33 Nowak SJ, Corces VG (2000) Phosphorylation of histone H3 correlates with transcriptionally  
34 active loci. *Genes Dev* 14:3003–3013. doi: 10.1101/gad.848800

This is the author's pre-print version.

The final publication is available at Springer via DOI 10.1007/s11738-014-1753-z

Citation: Ayaydin et al. *Acta Physiol Plant* (2015) 37:3

- 1 Nowak SJ, Pai CY, Corces VG (2003) Protein phosphatase 2A activity affects histone H3  
2 phosphorylation and transcription in *Drosophila melanogaster*. *Mol Cell Biol* 23:6129–  
3 6138. doi: 10.1128/MCB.23.17.6129
- 4 Park Y, Luger K (2006) Structure and function of nucleosome assembly proteins 1. *Biochem*  
5 *Cell Biol* 84:549–558. doi: 10.1139/O06-088
- 6 Roti Roti JL, Turkel N (1994a) Heat-induced changes in nuclear-associated proteins in normal  
7 and thermotolerant HeLa cells. *Radiat Res* 139:73–81.
- 8 Roti Roti JL, Turkel N (1994b) Heat-shock-induced changes in nuclear protein and cell  
9 killing in thermotolerant HeLa cells. *Radiat Res* 138:286–290.
- 10 Vermaak D (2003) Maintenance of chromatin states: an open-and-shut case. *Curr Opin Cell*  
11 *Biol* 15:266–274. doi: 10.1016/S0955-0674(03)00043-7
- 12 Warters RL, Brizgys LH, Sharma R, Roti Roti JL (1986) Heat shock (45°C) results in an  
13 increase of nuclear matrix protein mass in HeLa cells. *Int J Radiat Biol* 50:253-268. doi:  
14 10.1080/09553008614550641
- 15 Zhu Y, Dong A, Meyer D, Pichon O, Renou J-P, Cao K, Shen W-H (2006) *Arabidopsis* NRP1  
16 and NRP2 encode histone chaperones and are required for maintaining postembryonic  
17 root growth. *Plant Cell* 18:2879–2892. doi: 10.1105/tpc.106.046490

18

## 1 **Figure legends**

2 **Fig. 1** Fractionation and microscopy analyses of intracellular compartmentalization of

3 *Arabidopsis* NRP (At NRP) proteins

4 (a) Intracellular localization analysis on protoplasts shows nuclear accumulation of

5 GFP:NRP1. Nuclei and nucleoli (arrowheads) of GFP:NRP1-transfected protoplasts are

6 located using differential interference contrast (DIC) imaging. (b) In formaldehyde-fixed

7 cultured *Arabidopsis* cells, NRP immunodetection (green) shows nuclear accumulation of

8 NRP in interphase cells (asterisks) but not in mitotic cells (arrow). Nuclei are counterstained

9 using DAPI (pseudocolored red). Scale bars are 5  $\mu$ m. (c) Immunoblotting and detection of At

10 NRP after fractionation of nuclei (N) and supernatant (S) of unfixed cells. NRP is detected in

11 supernatant fraction. Immunodetection of a nucleocytoplasmic protein PP2Ac (Protein

12 phosphatase 2A catalytic subunit) is used as loading control on the lower panel. (d)

13 Immunodetection of NRP after fractionation of nuclei (N) and supernatant (S) of GFP:NRP1

14 expressing unfixed cells. Note that GFP-tagged NRP1 is retained in the nuclear fraction. (e)

15 Nuclear and cytoplasmic fractionation and immunodetection of NRP using fixed and unfixed

16 cells. Note that fixed cells' nuclear fractions and unfixed cells' cytoplasmic fractions contain

17 NRP.

18

19 **Fig. 2** The effect of heat shock on nuclear retention of *Arabidopsis* NRP (At NRP) proteins.

20 (a) Immunoblotting of At NRP on nuclear (N) and supernatant (S) fractions after 1 h heat

21 treatment at various temperatures. NRP of heat-treated cells (45°C and 50°C) remains nuclear

22 during fractionation procedure. (b) Immunodetection analysis of induction and longevity of

23 temperature-induced nuclear NRP retention using suspension cells. Left panel shows the

24 nuclear detection of NRP in the nuclei after half an hour of heat treatment at 45°C. Right

This is the author's pre-print version.

The final publication is available at Springer via DOI 10.1007/s11738-014-1753-z

Citation: Ayaydin et al. *Acta Physiol Plant* (2015) 37:3

1 panel shows persistence of nuclear retention after 30 h and 42 h following a 1 h, 45°C  
2 treatment. (c) Immunodetection of At NRP in nuclear protein fractions isolated from heat-  
3 treated Arabidopsis seedlings. Treatment temperature and durations are indicated below the  
4 image. (d), (e) Immunolocalization of NRP (green) in isolated nuclei of control (25°C) and  
5 heat treated (45°C) cells. Unfixed cells were used for isolation of nuclei which were then  
6 fixed with formaldehyde (see Methods). DAPI as nuclear counterstain is shown in red  
7 pseudocolor. Merged images with differential interference contrast (DIC) overlay are shown  
8 at the last panel. Scale bar is 5 µm.

9

10 **Fig. 3** Immobilization of soluble NRPs in heat-shocked nuclei.

11 (a) Fluorescence loss in photobleaching (FLIP) assay. Live cell images of heat-shocked  
12 (45°C, upper panel) and control (25°C, lower panel) GFP:NRP1-transfected Arabidopsis  
13 protoplasts were collected during successive laser bleaching of a localized intranuclear spot.  
14 Black squares of 1 µm width indicate bleached region. Yellow triangles represent repetitive  
15 activation of laser bleaching. White circles are 5 µm away from the bleaching spot and were  
16 used to measure the depletion of fluorescence to assess the mobility of intranuclear  
17 GFP:NRP1. Prebleach images indicate the initial fluorescence intensity of nuclei and the  
18 position of nucleoli (dark circular regions). Representative timepoints are shown. Scale bar is  
19 5 µm. (b) Kinetics of loss of fluorescence during FLIP assay of heat-treated (45°C) and  
20 control (25°C) GFP:NRP1-transfected samples. Intensity values were recorded at the  
21 intranuclear region which was 5 µm away from the bleached spot as shown on part (a).  
22 Averages and standard deviations of five measurements on different nuclei are shown. (c)  
23 Heat-treatment-induced nuclear retention (immobilization) of NRPs in isolated nuclei cannot  
24 be reverted by DNase and/or RNase treatment. Electrophoretic analysis of nucleic acids in an  
25 ethidium bromide-stained agarose gel (left) and the detection of NRPs in isolated nuclei of

1 heat-treated cells by an anti-NRP antibody (right). M: molecular weight marker; NT: Not  
2 treated, D: DNase-treated, D+R: DNase and RNase-treated.

3

4 **Fig. 4** Effect of heat shock on root growth of wild type and NRP-deficient *nrp1-1 nrp 2-1*  
5 mutant seedlings.

6 (a) Two-weeks-old wild type (WT) and NRP-deficient (-NRP) seedlings have been subjected  
7 to heat adaptation and heat shock (38°C, 2h▶22°C, 2h ▶45°C, 1h) or left untreated as  
8 control. Root images were taken on the 10th day post-treatment. (b) The length of the main  
9 roots was measured daily during the investigated period. Average main root length is shown  
10 for each treatment with standard deviation (n>25).

11

12 **Fig. 5** Effect of heat shock on bleomycin sensitivity of wild type and *nrp1-1 nrp 2-1* mutant  
13 seedlings.

14 (a) One week old wild type (WT) and NRP-deficient *nrp1-1 nrp 2-1* mutant (-NRP) seedlings,  
15 untreated control (CTRL) or heat adapted then heat shocked (HS) (38°C, 2h▶22°C, 2h  
16 ▶45°C, 1h) have been transferred to fresh media supplemented with 1 mg/L bleomycin and  
17 their survival was scored after six weeks. Averages and standard deviations from three  
18 experiments are shown. (b) The expression of the indicated genes, known to have a role in  
19 genotoxic stress response (Chen et al. 2003), was determined by real-time quantitative PCR  
20 using whole seedlings one hour after being transferred to bleomycin (BLEO) plates. The  
21 expression of the UBC18 (At5g42990) gene was used as reference for normalization and  
22 expression of untreated seedlings was used as a reference value (value = 1; a.u.: arbitrary  
23 unit).

24




Fig. 1




Fig. 2






Fig. 3





Fig. 4





Fig. 5



**Supplementary Figure 1. NRP retainment in the nucleus is not a general stress response.**

Heavy metal stress (CdCl<sub>2</sub>, **a**), salt stress (NaCl, **b**) genotoxic stress (bleomycin, **c**) or UV-C irradiation (**d**) did not result in nuclear NRP retention in cultured *Arabidopsis* cells. N: nuclear pellet, S: supernatant

<b>AGI code</b>	<b>Primer name</b>	<b>Sequence (5'-3')</b>
At5g42990	UBC18 FW	ACAGCAATGGACATATTTGTTTAGA
	UBC18 REV	TGATGCAGACTGAACTCACTGTC
At4g02390	PARP2 FW	GCGAACTATTGCTATGCCAAC
	PARP2 REV	ATGTCTCCCAAAGCAACCTC
At5g20850	RAD51 FW	GATCACGGGAGCTCGATAAA
	RAD51 REV	GCGGAACTCACCATATAACTCTG
At3g27060	RNR2 FW	GAATCATCGCTTTTCGCTTG
	RNR2 REV	TTCAGCCAGAAGATTGAACAAA
At5g03780	Myb FW	TGTGGCTAAGTCAAAGACAGTGA
	Myb REV	TCCACTCCACCTTTAGCAT
At2g26760	ATCYCB1FW	CCTTAAGAGCTCGAAGCAAGG
	ATCYCB1REV	AGCATCCACAGCATCAATGT

**Supplementary Table 1. Primers used in the study**