

## Title page

**Title:** Neural mechanisms of predatory aggression in rats - implications for abnormal intraspecific aggression

**Running title:** Neural background of predatory aggression

**Authors:** Aron Tulogdi, Laszlo Biro, Beata Barsvari, Mona Stankovic, Jozsef Haller, Mate Toth

**Affiliation:** Institute of Experimental Medicine, Department of Behavioral Neuroscience, P.O. Box 67, H-1450 Budapest, Hungary

**Corresponding author:**

József Haller

Institute of Experimental Medicine

Department of Behavioral Neuroscience

H-1450 Budapest

P.O. Box 67

Hungary

*E-mail:* [haller.jozsef@koki.mta.hu](mailto:haller.jozsef@koki.mta.hu); *Phone:* +36 12109406; *Fax:* +36 12109951

Number of words in the abstract: 250

Number of words in text: 3376

Number of figures: 5

Number of tables: 2

## Abstract

1  
2 Our recent studies showed that brain areas that are activated in a model of escalated  
3  
4 aggression overlap with those that promote predatory aggression in cats. This finding raised  
5  
6 the interesting possibility that the brain mechanisms that control certain types of abnormal  
7  
8 aggression include those involved in predation. However, the mechanisms of predatory  
9  
10 aggression are poorly known in rats, a species that is in many respects different from cats. To  
11  
12 get more insights into such mechanisms, here we studied the brain activation patterns  
13  
14 associated with spontaneous muricide in rats. Subjects not exposed to mice, and those which  
15  
16 did not show muricide were used as controls. We found that muricide increased the activation  
17  
18 of the central and basolateral amygdala, and lateral hypothalamus as compared to both  
19  
20 controls; in addition, a ventral shift in periaqueductal gray activation was observed.  
21  
22

23  
24 Interestingly, these are the brain regions from where predatory aggression can be elicited, or  
25  
26 enhanced by electrical stimulation in cats. The analysis of more than 10 other brain regions  
27  
28 showed that brain areas that inhibited (or were neutral to) cat predatory aggression were not  
29  
30 affected by muricide. Brain activation patterns partly overlapped with those seen earlier in the  
31  
32 cockroach hunting model of rat predatory aggression, and were highly similar with those  
33  
34 observed in the glucocorticoid dysfunction model of escalated aggression. These findings  
35  
36 show that the brain mechanisms underlying predation are evolutionarily conservative, and  
37  
38 indirectly support our earlier assumption regarding the involvement of predation-related brain  
39  
40 mechanisms in certain forms of escalated social aggression in rats.  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50

51 **Keywords:** muricide; aggression; c-Fos immunohistochemistry; hypothalamus; amygdala;  
52  
53 periaqueductal grey  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## 1. Introduction

The general functions of aggression fall into two main categories, particularly social competition and predation [1-3]. The first entails conspecifics, which fight for access to resources in a broad meaning (e.g. food, territory, social rank, etc.). This form of aggression is associated with high physiological arousal, and covers social communication. E.g. threat signals convey information on 'attack intentions'; moreover, threats may replace actual fights by the process of ritualization. In contrast, predation aims at killing an individual that belongs to a different species. This behavior is associated with minimal arousal, and does not involve social communication. These two forms of aggression are controlled by distinct neural circuits as shown by feline stimulation studies [4-6]. Based on phenomenological and physiological similarities, these types of aggressive behavior were proposed to be analogous with particular forms of psychopathological human aggression [7-10]. E.g. exacerbated affective aggression is seen in intermittent explosive disorder, which is a violent response to a perceived threat. Other forms of pathological aggression, e.g. those seen in antisocial personality disorder, have different characteristics: such aggression is often gain-oriented, and is associated with limited emotional arousal and low intention signaling [7-9]. The phrase 'predatory aggression' is frequently used to emphasize these characteristics [11-13].

In recent years, the differentiation of the two types of aggression and the idea that they are governed by distinct neural mechanisms gained attention in both human and animal research. In humans, a psychiatric inventory was developed to differentiate reactive and proactive aggression [14], and current theories deal with their neural underpinnings differentially [15, 16]. We recently developed two laboratory models that mimic important characteristics of affective/reactive and instrumental/proactive forms of aggression, and proposed behavioral methods to differentiate species-typical and abnormal forms of aggression [17-20]. Importantly for the present study, we found that rats submitted to one of

1 these models –the glucocorticoid dysfunction model of abnormal aggression– deliver bites to  
2 vulnerable body parts of conspecifics (head, throat and belly), which is accompanied by low  
3  
4 intention signaling by threats, disturbed social behavior, and reduced autonomic arousal,  
5  
6 features that are in many respects to similar to the symptoms of aggressive antisocial  
7  
8 personality disordered subjects [17-20]. We found that in this model, aggressive encounters  
9  
10 increase the activation of the lateral hypothalamus, central amygdala and ventral  
11  
12 periaqueductal grey (PAG) above the levels seen in controls submitted to fights (i.e. these  
13  
14 regions were overactivated) [4, 18, 19, 21, 22]. Moreover, the activation of the central  
15  
16 amygdala and lateral hypothalamus correlated significantly with the share of abnormal,  
17  
18 predatory-like attacks in this model [22]. As the very same brain regions were shown to  
19  
20 control predatory attacks in cats [1, 4-6], we proposed that antisocial-like aggressiveness in  
21  
22 rats has a 'predatory dimension' as it regards both behavior and brain function. Unfortunately,  
23  
24 however, the brain mechanisms of predatory aggression are less well known in rats than in  
25  
26 cats. Early electrophysiological studies identified the hypothalamic locus of control of frog  
27  
28 and mouse killing in rats but in contrast to cats, such studies provided limited information on  
29  
30 other modulatory brain regions [23-26]. More recently, neural mechanisms were evaluated in  
31  
32 rats by using c-Fos immunohistochemistry to investigate brain activation patterns of  
33  
34 cockroach hunting as a model of predatory aggression [27, 28]. In these studies, food intake  
35  
36 inherently associated with cockroach hunting was carefully controlled. While the activated  
37  
38 brain areas overlapped in many respects with the circuitry that controls predation in cats,  
39  
40 important differences were also observed. E.g. the lateral hypothalamus showed similar levels  
41  
42 of activation in insect hunting rats and their feeding controls, despite the fact that this brain  
43  
44 area is considered the most important control region of rat killing in cats and frog and mouse  
45  
46 killing in rats [1, 4, 29]. According to our own observation, muricide and insect hunting are  
47  
48 behaviorally different, which may explain these discrepancies. As such, studies using  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 muricide as a model seem necessary to fully understand the brain mechanisms of predatory  
2 aggression in rats, but such studies are unavailable at present.  
3

4 Here we investigated c-Fos activation in 15 aggression-related brain regions in adult  
5 male Wistar rats that spontaneously killed a mouse in their home-cage. Rats which did not  
6 attack the mice and rats without mouse exposure were used as controls. This study was  
7 motivated by multiple goals. Firstly, we aimed at describing brain activation patterns  
8 associated with muricide, a work that has not been performed so far. Secondly, we aimed at  
9 comparing these findings with those obtained earlier in cat stimulation studies to establish the  
10 cross-species stability of predation-related brain mechanisms. We also aimed at comparing  
11 findings with those obtained in the cockroach-hunting model, to investigate the impact of the  
12 prey on brain mechanisms. Finally, we aimed at providing a more proper comparison for the  
13 recently described "predatory-like aggression network" activated in the aforementioned model  
14 of violent social aggression [22].  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30

## 31 **2. Material and methods**

### 32 *2.1. Animals*

33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
Subjects were adult male Wistar rats raised in the breeding facility of our Institute. Parents were obtained from Charles River Laboratories (Germany). Rats were housed in macrolon cages in groups of 4-6. Food and water were available *ad libitum* throughout, temperature and relative humidity was kept at 22±2 °C and 60±10 %, respectively. Rats were maintained in a light cycle of 12:12 hours with lights off at 1000h. The weight of subjects was 350-450 g when behaviorally tested. Behavioral tests were conducted in the early phase of the dark period, under dim red illumination. 50-70 days-old male CD1 mice from the same source were used as stimulus animals. Mice were housed in a different room, but otherwise were maintained under similar conditions. The experiments were carried out in accordance

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

with the European Communities Council Directive of 24 November 1986 (86/609/EEC) and were reviewed and approved by the Animal Welfare Committee of the Institute of Experimental Medicine.

## 2.2. *Experimental procedure*

Subjects were housed individually for one week before behavioral testing but otherwise were maintained under the same conditions as earlier. The experiment was started by placing a mouse in the home-cage of the rat. Subjects have never encountered a mouse before. If the rat killed the mouse, the latency to kill was recorded, the killed mouse was removed immediately, and at the same time, another, uninjured mouse was removed from the home cage of a randomly chosen rat to assure that the time of stimulation/interaction was equal between groups (“muricide” vs. “no muricide” control). The cut-off time for mouse-killing was 20 min. On each experimental day, control rats unexposed to mice („no mouse” control) were also randomly chosen from rats that were not exposed to mice to assess baseline c-Fos activity. The experiment was continued until sample sizes reached 7 per group.

## 2.3. *Brain processing and immunohistochemistry*

Rats were left undisturbed for 120 minutes after the encounters to allow c-Fos signal to develop. Afterwards, they were deeply anesthetized by an i.p. injection of a mixture of ketamine, xylazine and pipolphen (5, 10 and 5 mg/kg, respectively) and perfused through the ascending aorta with 100 ml ice-cold 0.1M phosphate-buffered saline followed by approximately 200 ml 4% paraformaldehyde dissolved in 0.1M phosphate-buffered saline. The brains were removed, post-fixed in the same solution for 3h and cryoprotected overnight by 20% sucrose in phosphate-buffered saline at 4 °C. 30µm frozen sections were cut in the frontal plane on a sliding microtome. The c-Fos protein was labeled with a rabbit polyclonal

1 antibody raised against the amino terminus of c-Fos p62 (Santa Cruz Biotechnology, USA, sc-  
2 52) as described earlier [20, 22]. The primary antibody (1:5000) was detected by biotinylated  
3 anti-rabbit goat serum (1:1000; Jackson Laboratories, USA) and avidin-biotin complex (ABC,  
4 1:1000; Vector Laboratories, Burlingame, CA, USA). The peroxidase reaction was developed  
5 in the presence of diaminobenzidine tetrahydrochloride (0.2 mg/ml), nickel–ammonium  
6 sulphate (0.1%) and hydrogen peroxide (0.003%) dissolved in Tris buffer.  
7  
8  
9  
10  
11  
12  
13

14 Table 1 shows the brain structures investigated in the present study; anteroposterior  
15 levels and frame sizes are also shown. The number of investigated levels depended on the  
16 length of the particular brain region. At each level, the c-Fos signal was counted bilaterally,  
17 and the average of counts was considered. Section planes were standardized according to the  
18 atlas of Paxinos and Watson [30]. Microscopic images were digitized by an OLYMPUS CCD  
19 camera using a 10x magnification lens and stained particles were counted by means of the  
20 ImageJ v1.41o software (<http://rsbweb.nih.gov/ij/>). Uniform thresholds and minimum sizes  
21 of stained particles were considered.  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35

#### 36 *2.4. Statistical analyses*

37 Data are expressed as means  $\pm$  SEM. C-Fos counts were square-root transformed if  
38 necessary and were analyzed by one or two-factor ANOVA as shown below. The Duncan test  
39 was used for post-hoc comparisons. Correlation between latency of muricide and neuronal  
40 activation was analyzed by using the Spearman test. Significance level was set at  $p < 0.05$   
41 throughout; trends are indicated between  $0.05 < p < 0.10$ .  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52

### 53 **3. Results**

#### 54 *3.1. Muricide*

55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Muricide occurred suddenly without threats or other social-like signals, and was very short, almost a "point-like" behavioral act. The latency of muricidal behavior was  $6.0 \pm 1.2$  min; its range was 1 to 10 min. 'No muricide' rats were rather passive except for occasional trials of sniffing at the mouse that showed obvious trials to avoid this contact.

### 3.2. Brain activation patterns

As shown above, each brain area was investigated at 2 or 3 rostro-caudal levels depending on its length. While c-Fos counts varied rostro-caudally in some areas, no interaction with group assignment was observed; therefore, the average of values obtained at these levels was considered in statistical calculi.

C-fos counts were different in the three areas of the orbitofrontal cortex ( $F_{\text{area}}(2, 34) = 10.85$ ;  $p < 0.001$ ). A significant group difference was also observed ( $F_{\text{group}}(2, 17) = 3.34$ ;  $p = 0.05$ ) but the two factors did not interact ( $F_{\text{interaction}}(4, 34) < 1$ ; n.s.) (Fig. 1a). In the medial prefrontal cortex, the three areas were also different ( $F_{\text{area}}(2, 38) = 68.5$ ;  $p < 0.0001$ ), but no group effects were observed ( $F_{\text{group}}(2, 18) = 1.72$ ;  $p > 0.2$ ;  $F_{\text{interaction}}(4, 36) < 1$ ; n.s.) (Fig. 1b). No effects were observed in the lateral septum ( $F_{\text{group}}(2, 18) = 2.07$ ;  $p = 0.15$ ; Fig. 1c).

There was a significant interaction between factors in the amygdala ( $F_{\text{area*group}}(4, 36) = 8.28$ ;  $p < 0.0001$ ) (Fig. 2). The central and basolateral amygdala were not activated by mouse exposure, but a strong activation was observed after muricide. The activation of the medial amygdala increased in both 'no muricide' controls and muricidal rats but the activation was stronger in the latter. By contrast, the bed nucleus of stria terminalis (BNST) was activated neither by mouse exposure nor by muricide.

A similar phenomenon was seen in the hypothalamus, where the factors interacted significantly ( $F_{\text{area*group}}(4, 38) = 5.46$ ;  $p < 0.01$ ). C-Fos counts changed neither in the paraventricular nucleus of the hypothalamus (PVN) nor in the hypothalamic attack area



(HAA). By contrast, a strong increase in activation was observed in the lateral hypothalamus, specifically in muricidal rats (Fig. 3).

Overall, no significant group differences were observed in the activation of the periaqueductal grey (PAG) ( $F_{\text{group}}(2, 18) = 0.26$ ;  $p = 0.76$ ;  $F_{\text{group*area}}(4, 36) = 0.73$ ;  $p = 0.57$ ) (Fig. 4a). However, there was an activity shift between ventral and dorsal columns at the rostral and intermediate levels, resulting in increased ventral/dorsal ratio in 'muricide' rats (rostral part:  $F_{\text{group}}(2,18) = 4.86$ ,  $p < 0.05$ ; middle part:  $F_{\text{group}}(2,18) = 3.34$ ,  $p = 0.059$ ; Fig. 4b). Monoaminergic nuclei showed no activation in either group (all regions:  $F_{\text{group}}(2,18) < 1.06$ , ns; Table 2).

C-Fos activity in the medial amygdala, but none of the other regions, showed a positive correlation with the latency of muricide ( $R = 0.793$ ,  $p < 0.05$ ; Fig. 2).

#### 4. Discussion

##### *Main findings*

Muricidal attacks induced strong c-Fos activation in brain areas regulating aggressive behavior which was markedly distinct from activation exhibited by rats showing no muricide during the interaction. Muricide increased the activation of the medial, central, and basolateral amygdala, and of the lateral hypothalamus. In addition, muricide shifted PAG activations from dorsal to ventral columns. This pattern of brain activation was highly congruent with findings obtained in cat stimulation studies, was in several respects similar to c-Fos findings obtained with the rat cockroach-hunting model, and was highly similar with the findings obtained in rats submitted to the glucocorticoid dysfunction-model of abnormal aggression (Fig. 5). In the rat models, overactivations were seen in brain areas that promoted predatory attacks in cats. In brain areas where electrical stimulation inhibited predatory aggression in cats, no activation was seen in any of the rat models, except for the medial amygdala, which was activated by muricide. However, this brain response correlated positively with the latency

1 of muricide, suggesting that similar to cats, the medial amygdala is involved in the negative  
2 modulation of predatory aggression. Taken together, our findings show that brain mechanisms  
3 underlying predatory aggression are highly conservative, and strongly support our earlier  
4 suggestions on the similarities between the brain mechanisms of predatory aggression and  
5 those underlying certain forms of abnormal social aggression.  
6  
7  
8  
9  
10

### 11 *Comparisons with earlier findings*

12  
13  
14 The immediate-early gene product c-Fos has been widely used as an indicator of acute  
15 neuronal activation to describe brain activation patterns during different aggressive  
16 interactions in several laboratory species, including mice, hamsters and rats [31-34]. It was  
17 also used to study the brain mechanisms of cockroach hunting, another model of rat predatory  
18 aggression [27, 28]. The study presented here was performed to serve as a comparison for  
19 other models of predatory aggression (stimulation-induced aggression in cats [1, 4], and  
20 cockroach hunting in rats [27, 28]), as well as for certain models of abnormal aggression in  
21 rats (glucocorticoid dysfunction model, [18, 21, 35]). These models will be compared by  
22 starting with the hypothalamus that plays crucial roles in the elicitation of biting attacks in  
23 both cats and rats [1].  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38

39 Cholinergic drugs injected into the lateral hypothalamus elicited muricide in rats [23,  
40 26], whereas the electrical stimulation of this brain area induced quiet biting in cats, a  
41 predatory form of aggression in this species [1, 6, 24, 36, 37]. In our muricidal rats, strong and  
42 highly muricide-specific activation was seen in the lateral hypothalamus along its whole  
43 rostro-caudal extent, which supports the notion that this region is a primary center for  
44 predatory aggression. In the cockroach-hunting model, this region was also activated;  
45 however, its activity was not specific to hunting as the effects of feeding (the control  
46 treatment in this model) were similarly strong [28]. Indeed, the lateral hypothalamus is  
47 crucially involved in feeding [38]; hence, one can assume that the effects of cockroach killing  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 were masked by the effects of food intake. Interestingly, the lateral hypothalamus was also  
2 overactivated in the glucocorticoid dysfunction model of abnormal aggression, the subjects of  
3 which deliver bites to highly vulnerable targets of their opponents (head, throat and belly) on  
4 the background of diminished social signaling of attacks by threats, which make their  
5 behavior similar to that seen during predation [22]. In contrast to the lateral hypothalamic  
6 region, the mediobasal hypothalamus (often called the 'hypothalamic attack area' in rats)  
7 elicits affective aggression in both cats and rats, and is activated by resident/intruder  
8 interactions in the latter species [1, 4, 21, 37, 39, 40]. Cat studies revealed a reciprocal  
9 inhibition between the lateral and mediobasal regions, which involves a context-dependent  
10 control of attacks, making social and predatory aggressions mutually exclusive [4, 39].  
11 Although similar direct evidence is missing in rats, one cannot rule out a potentially similar  
12 inhibitory mechanism; for instance, mediobasal hypothalamic lesions result in 'uncontrolled'  
13 muricidal behavior which suggests cross-species similarities [41]. The mediobasal  
14 hypothalamic area was activated neither in our muricidal rats, nor in the cockroach-hunting  
15 model [28]. In the glucocorticoid dysfunction model, the mediobasal hypothalamus was  
16 activated by fights as compared to non-fighting controls, but its activation was similar to that  
17 seen in sham operated fighting controls. Taken together, these findings suggest that the  
18 mediobasal hypothalamus plays no particular roles in muricide, cockroach hunting, or in the  
19 abnormal attack features of rats submitted to the glucocorticoid-dysfunction model. The  
20 activation of the paraventricular nucleus of the hypothalamus –a brain region that orchestrates  
21 the glucocorticoid stress response– was activated in neither rat model discussed here, which is  
22 in line with the low emotionality of both predatory aggression and glucocorticoid  
23 dysfunction-induced abnormal aggression [18, 19, 28].

24 Hypothalamic attack areas receive inputs from the amygdala, which plays a subregion-  
25 specific role in aggression control. In cats, the medial amygdala and its major output region  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 BNST inhibit predatory attacks, whereas the central and basolateral nuclei facilitate this  
2 response [1, 42-46]. Our findings are in line with these reports by showing a muricide-specific  
3  
4 activation of the central and basolateral nuclei. Highly similar findings were obtained in the  
5  
6 cockroach-hunting model of predatory aggression [28]. The over-activation of the central  
7  
8 nucleus is also well-documented in the glucocorticoid-dysfunction model, where predatory-  
9  
10 like attacks are shown in a social context [19, 21, 47]. No medial amygdala activation  
11  
12 surpassing that seen in controls were observed in the rat models discussed here except for  
13  
14 muricide. Our correlation analysis, however, suggests that this region is involved in the  
15  
16 negative modulation of muricide, which is congruent with findings obtained in cats, where the  
17  
18 concomitant stimulation of the medial amygdala and lateral hypothalamus decreased the  
19  
20 efficacy of the latter in inducing predatory attacks [1].  
21  
22  
23  
24  
25

26 Hypothalamic attack centers send projections to the PAG, which controls behaviors and  
27  
28 autonomic functions during aggressive conflicts [1]. This brain area is organized in columns,  
29  
30 which are implicated in behavioral control differentially. In cats, the stimulation of ventral  
31  
32 columns (lateral and ventrolateral PAG) induces predatory attacks, whereas dorsal columns  
33  
34 (dorsomedial and dorsolateral PAG) are involved in defensive responses [1, 6, 48-50]. Here  
35  
36 we found that muricidal attacks were accompanied by a dominantly ventral activation of the  
37  
38 PAG. Congruent findings were obtained in the cockroach-hunting model [27, 28, 51]. A  
39  
40 comparison of cockroach hunting- and muricide-induced c-Fos activity also suggests a rostro-  
41  
42 caudal heterogeneity of the PAG and a potential role of rostral regions in the switch between  
43  
44 intraspecific and predatory attack patterns. Noteworthy, a ventral shift in PAG activation was  
45  
46 also seen in the abnormal aggression model [21, 22, 35, 47, 52].  
47  
48  
49  
50  
51  
52

53 Our findings are equivocal as it regards the involvement of prefrontal regions in  
54  
55 aggression control. Muricide increased the activation of the orbitofrontal cortex as compared  
56  
57 to 'no-mouse' controls but not as compared to 'no-muricide' controls. A similar situation  
58  
59  
60  
61  
62  
63  
64  
65

1 occurred in the glucocorticoid dysfunction model, where aggression did increase the  
2 activation of the orbitofrontal cortex as compared to non-fighting controls, but this activation  
3  
4 was not larger than that seen in sham operated fighting controls [35]. The orbitofrontal cortex  
5  
6 was not investigated in the cockroach model so far, but findings on the activation of the  
7  
8 medial prefrontal cortex were similarly equivocal. The activation of this area was larger in  
9  
10 cockroach-hunting rats as compared with home cage controls, but smaller than that seen in  
11  
12 feeding controls [28]. These findings suggest that the involvement of the prefrontal cortex in  
13  
14 aggression is more complex than expected, and invites further research.  
15  
16  
17

### 18 *Conclusions*

19  
20  
21 (i) The neural underpinnings of predatory aggression show remarkable cross-species  
22  
23 similarities. Muricide enhanced the activation of brain areas which promote, but did not affect  
24  
25 the activation of areas that inhibit predatory aggression in cats (Fig. 5). Although enhanced  
26  
27 activation was seen in the medial amygdala, which inhibits predatory attacks in cats, this brain  
28  
29 region appears to modulate muricide negatively as shown by correlation analysis.  
30  
31  
32

33  
34 (ii) The prey has an impact on brain activation patterns. The effects of cockroach  
35  
36 hunting and muricide were similar in some, but not in all brain regions. As the food intake  
37  
38 component of cockroach hunting was controlled in the study by Comoli et al. [28], differences  
39  
40 in brain activations are likely due to behavioral differences inherently associated with the  
41  
42 nature of the prey. According to our observations, explicit acts of killing are present in the  
43  
44 muricide, but mostly absent in cockroach hunting model. Still, the brain mechanisms  
45  
46 underlying the two forms of rat predation overlap to a certain extent showing that different  
47  
48 forms of predation are controlled by overlapping but not entirely similar networks.  
49  
50  
51

52  
53 (iii) Brain activation patterns elicited by muricide and glucocorticoid dysfunction-  
54  
55 induced abnormal aggression were highly similar, and support the notion that this type of  
56  
57 abnormal aggression activates brain mechanisms involved in predation. This parallels  
58  
59  
60  
61  
62  
63  
64  
65

1 behavioral and physiological similarities; in both models, attacks are targetted on vulnerable  
2 body parts, social communication is deficient, and physiological arousal is low. These  
3  
4 observations support earlier clinical descriptions that differentiate reactive/affective from  
5  
6 *predatory*/instrumental types of human aggression, and implicate that their neural  
7  
8 underpinnings are different [7-9].  
9

10  
11 The present study used a 'descriptive mapping' approach. Future studies are required to  
12  
13 elucidate the causality between brain activation patterns and aggressive behavior.  
14  
15  
16

### 17 18 19 **Acknowledgements** 20

21 This work was supported by Hungarian National Science and Research Fund (OTKA) grant  
22  
23 No 112907. The authors thank Éva Mikics for her excellent comments on the manuscript.  
24  
25

26 The authors declare no conflict of interest.  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**Table 1.** The investigated brain areas.

<b>Area</b>	<b>Level of analysis (mm from Bregma)</b>	<b>Number of slices</b>	<b>Frame size (mm<sup>2</sup>)</b>
Orbitofrontal cortex	4.20 to 3.70	2	0.564
Medial prefrontal cortex	3.20 to 2.70	2	0.469
Lateral septum	0.70 to 0.20	3	0.880
Mediobasal hypothalamus ("hypothalamic attack area")	-1.60 to -2.30	3	0.575
Lateral hypothalamic area	-1.88 to -3.60	3	0.673
Paraventricular nucleus of the hypothalamus (parvocellular region)	-1.40 to -1.80	3	0.161
Medial amygdala	-2.30 to -2.80	2	0.697
Central amygdala	-1.80 to -2.56	3	0.576
Basolateral amygdala	-2.30 to -3.14	3	0.748
Bed nucleus of the stria terminalis, medial part	-0.80 to -0.92	2	0.403
Ventral tegmental area	-5.20 to -6.04	2	0.733
Dorsal raphe	-7.64 to -8.00	2	0.368
Locus coeruleus	-9.80 to -10.04	2	0.149
Periaqueductal grey, dorsomedial part	-6.72 to -8.30	3	0.184
Periaqueductal grey, dorsolateral part	-6.72 to -7.80	2	0.207
Periaqueductal grey, lateral part	-6.72 to -8.30	3	0.282
Periaqueductal grey, ventrolateral part	-7.80 to -8.30	2	0.368

Note that the relatively broad spatial range of section planes is explained by the fact that 2-3 sections were investigated in the case of each area, to investigate potential rostro-caudal differences in activation patterns. Post-hoc analysis showed no such differences; therefore, c-Fos counts were averaged over levels. The periaqueductal gray was an exception in this respect, because the structure of this brain area changes substantially in the rostro-caudal direction. Section planes were standardized according to the Rat Brain Atlas of Paxinos and Watson [30]

**Table 2.** Muricide-related neuronal activity in monoaminergic nuclei as indicated by c-Fos expression.

<b>Area</b>	<b>No mouse</b>	<b>No muricide</b>	<b>Muricide</b>
Dorsal raphe	18.79±5.20	19.43±4.61	22.19±3.64
Locus coeruleus	4.93±0.91	4.83±1.06	5.91±1.17
Ventral tegmental area	16.79±4.39	15.36±3.28	21.44±6.32

Data (mean ± standard error of the mean) shows c-Fos counts in the frames specified in Table

1. No statistically significant differences were observed.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



## References

- [1] Siegel A, Roeling TA, Gregg TR, Kruk MR. Neuropharmacology of brain-stimulation-evoked aggression. *Neuroscience and biobehavioral reviews*. 1999;23:359-89.
- [2] Archer J. The nature of human aggression. *International journal of law and psychiatry*. 2009;32:202-8.
- [3] Darwin C. *The Descent of Man and Selection in Relation to Sex*. : Murray, London. ; 1871.
- [4] Gregg TR, Siegel A. Brain structures and neurotransmitters regulating aggression in cats: implications for human aggression. *Progress in neuro-psychopharmacology & biological psychiatry*. 2001;25:91-140.
- [5] Shaikh MB, Brutus M, Siegel HE, Siegel A. Topographically organized midbrain modulation of predatory and defensive aggression in the cat. *Brain research*. 1985;336:308-12.
- [6] Shaikh MB, Barrett JA, Siegel A. The pathways mediating affective defense and quiet biting attack behavior from the midbrain central gray of the cat: an autoradiographic study. *Brain research*. 1987;437:9-25.
- [7] Blair RJ. Neurocognitive models of aggression, the antisocial personality disorders, and psychopathy. *Journal of neurology, neurosurgery, and psychiatry*. 2001;71:727-31.
- [8] Meloy JR. Empirical basis and forensic application of affective and predatory violence. *The Australian and New Zealand journal of psychiatry*. 2006;40:539-47.
- [9] Vitiello B, Behar D, Hunt J, Stoff D, Ricciuti A. Subtyping aggression in children and adolescents. *The Journal of neuropsychiatry and clinical neurosciences*. 1990;2:189-92.
- [10] Siegel A, Victoroff J. Understanding human aggression: New insights from neuroscience. *International journal of law and psychiatry*. 2009;32:209-15.
- [11] Anderson PD, Bokor G. Forensic aspects of drug-induced violence. *Journal of pharmacy practice*. 2012;25:41-9.
- [12] Felson RB, Massoglia M. When is violence planned? *Journal of interpersonal violence*. 2012;27:753-74.
- [13] Masi G, Manfredi A, Milone A, Muratori P, Polidori L, Ruglioni L, et al. Predictors of nonresponse to psychosocial treatment in children and adolescents with disruptive behavior disorders. *Journal of child and adolescent psychopharmacology*. 2011;21:51-5.
- [14] Raine A, Dodge K, Loeber R, Gatzke-Kopp L, Lynam D, Reynolds C, et al. The Reactive-Proactive Aggression Questionnaire: Differential Correlates of Reactive and Proactive Aggression in Adolescent Boys. *Aggressive behavior*. 2006;32:159-71.
- [15] Blair RJ. Neuroimaging of psychopathy and antisocial behavior: a targeted review. *Current psychiatry reports*. 2010;12:76-82.
- [16] Blair RJ. Psychopathy, frustration, and reactive aggression: the role of ventromedial prefrontal cortex. *British journal of psychology*. 2010;101:383-99.
- [17] Miczek KA, de Boer SF, Haller J. Excessive aggression as model of violence: a critical evaluation of current preclinical methods. *Psychopharmacology*. 2013;226:445-58.
- [18] Haller J, Kruk MR. Normal and abnormal aggression: human disorders and novel laboratory models. *Neuroscience and biobehavioral reviews*. 2006;30:292-303.
- [19] Haller J, van de Schraaf J, Kruk MR. Deviant forms of aggression in glucocorticoid hyporeactive rats: a model for 'pathological' aggression? *Journal of neuroendocrinology*. 2001;13:102-7.
- [20] Toth M, Tulogdi A, Biro L, Soros P, Mikics E, Haller J. The neural background of hyper-emotional aggression induced by post-weaning social isolation. *Behavioural brain research*. 2012;233:120-9.

- 1 [21] Halasz J, Liposits Z, Kruk MR, Haller J. Neural background of glucocorticoid  
2 dysfunction-induced abnormal aggression in rats: involvement of fear- and stress-related  
3 structures. *The European journal of neuroscience*. 2002;15:561-9.
- 4 [22] Tulogdi A, Toth M, Halasz J, Mikics E, Fuzesi T, Haller J. Brain mechanisms involved  
5 in predatory aggression are activated in a laboratory model of violent intra-specific  
6 aggression. *The European journal of neuroscience*. 2010;32:1744-53.
- 7 [23] Bandler RJ, Jr. Cholinergic synapses in the lateral hypothalamus for the control of  
8 predatory aggression in the rat. *Brain research*. 1970;20:409-24.
- 9 [24] Smith DE, King MB, Hoebel BG. Lateral hypothalamic control of killing: evidence for a  
10 cholinceptive mechanism. *Science*. 1970;167:900-1.
- 11 [25] Vergnes M, Karli P. [Elicitation of aggressive behavior by electric stimulation of the  
12 medial hypothalamus in the rat]. *Physiology & behavior*. 1970;5:1427-30.
- 13 [26] Woodworth CH. Attack elicited in rats by electrical stimulation of the lateral  
14 hypothalamus. *Physiology & behavior*. 1971;6:345-53.
- 15 [27] Comoli E, Ribeiro-Barbosa ER, Canteras NS. Predatory hunting and exposure to a live  
16 predator induce opposite patterns of Fos immunoreactivity in the PAG. *Behavioural brain  
17 research*. 2003;138:17-28.
- 18 [28] Comoli E, Ribeiro-Barbosa ER, Negrao N, Goto M, Canteras NS. Functional mapping of  
19 the prosencephalic systems involved in organizing predatory behavior in rats. *Neuroscience*.  
20 2005;130:1055-67.
- 21 [29] Siegel A, Bhatt S, Bhatt R, Zalzman SS. The neurobiological bases for development of  
22 pharmacological treatments of aggressive disorders. *Current neuropharmacology*. 2007;5:135-  
23 47.
- 24 [30] Paxinos G, Watson C. *The rat brain in stereotaxic coordinates*. 4th ed. San Diego:  
25 Academic Press; 1998.
- 26 [31] Davis ES, Marler CA. c-fos Changes following an aggressive encounter in female  
27 California mice: a synthesis of behavior, hormone changes and neural activity. *Neuroscience*.  
28 2004;127:611-24.
- 29 [32] Lin D, Boyle MP, Dollar P, Lee H, Lein ES, Perona P, et al. Functional identification of  
30 an aggression locus in the mouse hypothalamus. *Nature*. 2011;470:221-6.
- 31 [33] Kollack-Walker S, Newman SW. Mating and agonistic behavior produce different  
32 patterns of Fos immunolabeling in the male Syrian hamster brain. *Neuroscience*.  
33 1995;66:721-36.
- 34 [34] Halasz J, Liposits Z, Meelis W, Kruk MR, Haller J. Hypothalamic attack area-mediated  
35 activation of the forebrain in aggression. *Neuroreport*. 2002;13:1267-70.
- 36 [35] Halasz J, Toth M, Kallo I, Liposits Z, Haller J. The activation of prefrontal cortical  
37 neurons in aggression--a double labeling study. *Behavioural brain research*. 2006;175:166-75.
- 38 [36] Levison PK, Flynn JP. The objects attacked by cats during stimulation of the  
39 hypothalamus. *Animal behaviour*. 1965;13:217-20.
- 40 [37] Wasman M, Flynn JP. Directed attack elicited from hypothalamus. *Archives of  
41 neurology*. 1962;6:220-7.
- 42 [38] Bernardis LL, Bellinger LL. The lateral hypothalamic area revisited: ingestive behavior.  
43 *Neuroscience and biobehavioral reviews*. 1996;20:189-287.
- 44 [39] Han Y, Shaikh MB, Siegel A. Medial amygdaloid suppression of predatory attack  
45 behavior in the cat: II. Role of a GABAergic pathway from the medial to the lateral  
46 hypothalamus. *Brain research*. 1996;716:72-83.
- 47 [40] Kruk MR, Van der Poel AM, Meelis W, Hermans J, Mostert PG, Mos J, et al.  
48 Discriminant analysis of the localization of aggression-inducing electrode placements in the  
49 hypothalamus of male rats. *Brain research*. 1983;260:61-79.
- 50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

- 1 [41] Albert DJ, Walsh ML, Ryan J, Siemens Y. Mouse killing in rats: a comparison of  
2 spontaneous killers and rats with lesions of the medial hypothalamus or the medial accumbens  
3 nucleus. *Physiology & behavior*. 1982;29:989-94.
- 4 [42] Egger MD, Flynn JP. Amygdaloid suppression of hypothalamically elicited attack  
5 behavior. *Science*. 1962;136:43-4.
- 6 [43] McIntyre DC. Amygdala kindling and muricide in rats. *Physiology & behavior*.  
7 1978;21:49-56.
- 8 [44] Han Y, Shaikh MB, Siegel A. Medial amygdaloid suppression of predatory attack  
9 behavior in the cat: I Role of a substance P pathway from the medial amygdala to the medial  
10 hypothalamus. *Brain research*. 1996;716:59-71.
- 11 [45] Shaikh MB, Brutus M, Siegel HE, Siegel A. Regulation of feline aggression by the bed  
12 nucleus of stria terminalis. *Brain research bulletin*. 1986;16:179-82.
- 13 [46] Shaikh MB, Steinberg A, Siegel A. Evidence that substance P is utilized in medial  
14 amygdaloid facilitation of defensive rage behavior in the cat. *Brain research*. 1993;625:283-  
15 94.
- 16 [47] Haller J, Toth M, Halasz J, De Boer SF. Patterns of violent aggression-induced brain c-  
17 fos expression in male mice selected for aggressiveness. *Physiology & behavior*.  
18 2006;88:173-82.
- 19 [48] Bandler R, Shipley MT. Columnar organization in the midbrain periaqueductal gray:  
20 modules for emotional expression? *Trends in neurosciences*. 1994;17:379-89.
- 21 [49] Behbehani MM. Functional characteristics of the midbrain periaqueductal gray. *Progress*  
22 *in neurobiology*. 1995;46:575-605.
- 23 [50] Brutus M, Shaikh MB, Siegel A. Differential control of hypothalamically elicited flight  
24 behavior by the midbrain periaqueductal gray in the cat. *Behavioural brain research*.  
25 1985;17:235-44.
- 26 [51] Sukikara MH, Mota-Ortiz SR, Baldo MV, Felicio LF, Canteras NS. A role for the  
27 periaqueductal gray in switching adaptive behavioral responses. *The Journal of neuroscience* :  
28 the official journal of the Society for Neuroscience. 2006;26:2583-9.
- 29 [52] Haller J, Halasz J, Mikics E, Kruk MR. Chronic glucocorticoid deficiency-induced  
30 abnormal aggression, autonomic hypoarousal, and social deficit in rats. *Journal of*  
31 *neuroendocrinology*. 2004;16:550-7.
- 32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## Figure captions and legends

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**Fig. 1.** Neuronal activity in the (a) orbitofrontal and (b) medial prefrontal cortex as well as (c) the lateral septum as indicated by c-Fos expression. The schematics illustrate the regions investigated (gray squares) based on Paxinos and Watson [30]. *ac*, anterior commissure; *cc*, corpus callosum; *Cg1*, anterior cingulate cortex; *fmi*, forceps minor of corpus callosum; *IL*, infralimbic cortex; *LS*, lateral septum; *MO*, medial orbitofrontal cortex; *mPFC*, medial prefrontal cortex; *LO*, lateral orbitofrontal cortex; *OB*, olfactory bulb; *PrL*, prelimbic cortex; *VO*, ventral orbitofrontal cortex; \*, significant difference from ‘no mouse’ controls ( $p < 0.05$ ).

**Fig. 2.** Neuronal activity in the extended amygdala. The schematics illustrate the regions investigated (gray ovals) based on Paxinos and Watson [30]. Panel ‘b’ shows correlation between medial amygdala activity and the latency of muricide. *MeA*, medial amygdala. *ac*, anterior commissure; *BLA*, basolateral amygdala; *BNST*, bed nucleus of stria terminalis, medial part; *CeA*, central amygdala; *ic*, internal capsule; *MeA*, medial amygdala; *ot*, optic tract; *st*, stria terminalis; \*, significant difference from ‘no mouse’ controls; #, significant difference from ‘no muricide’ controls ( $p < 0.05$ ).

**Fig. 3.** Hypothalamic neuronal activity following an interaction with a mouse as indicated by c-Fos expression. The schematics illustrate the regions investigated (gray ovals) based on Paxinos and Watson [30]. *f*, fornix; *HAA*, hypothalamic attack area (mediobasal hypothalamus); *LH*, lateral hypothalamus; *ot*, optic tract; *PVN*, paraventricular nucleus of the hypothalamus; \*, significant difference from ‘no mouse’ controls; #, significant difference from ‘no muricide’ controls ( $p < 0.05$ ).

**Fig. 4.** Overall c-Fos counts (a) and the ventral/dorsal ratio of activation (b) at different rostro-caudal levels of the periaqueductal grey. The schematics illustrate the regions investigated (gray ovals) based on Paxinos and Watson [30]. *DL*, dorsolateral part; *DM*, dorsomedial part; *L*, lateral part; *VL*, ventrolateral part; *V/D*; ventral/dorsal ratio of activation; \*, significant difference from ‘no mouse’ controls ( $p < 0.05$ ); +, marginally significant difference from ‘no mouse’ controls ( $0.1 > p < 0.05$ ); #, significant difference from ‘no muricide’ controls ( $p < 0.05$ ).



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

**Fig. 5.** A comparison of four aggression models. Data obtained in the muricide (based on present findings), cockroach hunting (based on [20, 21]), and glucocorticoid dysfunction induced abnormal aggression models (based on [11, 14, 52]) were grouped according to the effects of electrical stimulation on predatory aggression in cats (based on [1, 2]). *OFC*, orbitofrontal cortex; *mPFC*, medial prefrontal cortex; *LS*, lateral septum; *BNST*, bed nucleus striae terminalis; *HAA*, hypothalamic attack area (mediobasal hypothalamus); *MeA*, medial amygdala; *CeA*, central amygdala; *BLA*, basolateral amygdala; *LH*, lateral hypothalamus; *PAG*, periaqueductal gray; *horizontal gray arrows*, no change in c-Fos activation; *vertical black arrows*, increased c-Fos activation; \* and °, similar and discrepant, respectively, findings in muricide and cockroach-hunting models; *empty cells*, no data.



*Note.* The following contrasts were considered: no muricide vs. muricide (mouse killing); feeding vs. cockroach hunting (cockroach hunting); sham-operated fighting vs. adrenalectomized fighting (glucocorticoid dysfunction-induced abnormal aggression).

Figure(s)



**(a) Orbitofrontal cortex**



**(b) Medial prefrontal cortex**





**(c) Lateral septum**




- No mouse
- No muricide
- Muricide



**(a) Extended amygdala**



**(b) Correlation between MeA activation and muricide**




## Hypothalamus








**(a) Periaqueductal gray – activation levels**



**(b) Periaqueductal gray – ventral/dorsal activation ratio**



***A comparison of the models considered here***

**Areas that *inhibit*  
predatory attacks in cats**

**Mouse killing  
Rat**

**Abnormal aggression  
Rat**

OFC	→	→
mPFC	→ ○	→
LS	→ *	→
BNST	→ ○	→
HAA	→ *	→
MeA	↑ ○	→

**Areas that *stimulate*  
predatory attacks in cats**

CeA	↑ *	↑
BLA	↑ *	
LH	↑ ○	↑
ventral PAG	ventral shift in PAG activation *	ventral shift in PAG activation