



Light absorption properties of laboratory generated tar ball particles

A. Hoffer et al.

Light absorption properties of laboratory generated tar ball particles

A. Hoffer¹, A. Tóth², I. Nyirő-Kósa¹, M. Pósfai², and A. Gelencsér^{1,2}

¹MTA-PE Air Chemistry Research Group, Veszprém, P.O. Box 158, 8201, Hungary

²Department of Earth and Environmental Sciences, University of Pannonia, Veszprém, P.O. Box 158, 8201, Hungary

Received: 3 April 2015 – Accepted: 19 May 2015 – Published: 16 June 2015

Correspondence to: A. Gelencsér (gelencs@almos.uni-pannon.hu)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Tar balls (TBs) are a specific particle type which is abundant in the global troposphere, in particular in biomass smoke plumes. These particles belong to the family of atmospheric brown carbon (BrC) which can absorb light in the visible range of the solar spectrum. Albeit TBs are typically present as individual particles in biomass smoke plumes, their absorption properties have been only indirectly inferred from field observations or calculations based on their electron energy-loss spectra. This is because in biomass smoke TBs coexist with various other particle types (e.g. organic particles with inorganic inclusions and soot, the latter is emitted mainly during flaming conditions) from which they cannot be physically separated; thus, a direct experimental determination of their absorption properties is not feasible. Very recently we have demonstrated that TBs can be generated in the laboratory from droplets of wood tar that resemble atmospheric TBs in all of their observed properties. As a follow-up study we have installed on-line instruments to our laboratory set-up generating pure TB particles to measure the absorption and scattering, as well as size distribution of the particles. In addition, samples were collected for transmission electron microscopy (TEM) and total carbon (TC) analysis. The effects of experimental parameters were also studied. The mass absorption coefficients of the laboratory generated TBs were found to be in the range of $0.8\text{--}3.0\text{ m}^2\text{ g}^{-1}$ at 550 nm, with absorption Ångström exponents (AAE) between 2.7 and 3.4 (average 2.9) in the wavelength range 467–652 nm. The refractive index of TBs as derived from Mie calculations was about $1.84\text{--}0.21i$ at 550 nm. In the brown carbon continuum these values fall closer to those of soot than to other light-absorbing species such as humic-like substances (HULIS). Considering the abundance of TBs in biomass smoke and the global magnitude of biomass burning emissions, these findings may have substantial influence on the understanding of global radiative energy fluxes.

Light absorption properties of laboratory generated tar ball particles

A. Hoffer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1 Introduction

Tar balls (TBs) are abundant and represent a peculiar particle type emitted from biomass burning. They can be readily identified by transmission electron microscopy (TEM) by their morphology, chemical composition, and amorphous structure. TBs are homogeneous and spherical particles that can withstand the high-energy electron beam of the TEM. They are most often present in external mixture, i.e. as individual standalone particles. Their sizes range from 30 to 500 nm in optical diameter as determined by TEM (Pósfai et al., 2004; Cong et al., 2009; Adachi and Buseck, 2010; Fu et al., 2012; China et al., 2013). Very recently we have demonstrated that TBs can be generated in the laboratory from droplets of wood tar that resemble atmospheric TBs in all of their observed properties (Tóth et al., 2014). These particles belong to the family of atmospheric brown carbon (BrC) which can absorb light in the visible range of the solar spectrum (Andreae and Gelencsér, 2006). Chung et al. (2012) have estimated that the global contribution of BrC to light absorption may be as high as 20 % at 550 nm. Given that the estimated contribution of humic-like substances (HULIS) to solar absorption can be only few per cent at 500 nm (Hoffer et al., 2006), a substantial fraction of BrC absorption may be attributed to TBs. So far a direct experimental determination of absorption properties of TBs has not been feasible because in biomass smoke TBs coexist with various other particle types from which they cannot be separated. Thus, their absorption properties have been so far only indirectly inferred from field observations (Hand et al., 2004; Chakrabarty et al., 2010) or calculations based on their dielectric functions obtained from electron energy-loss spectrometry (Alexander et al., 2008).

Hand et al. (2004) were the first to estimate the optical properties of TB by measuring the optical properties of ambient particles emitted from biomass burning during the YACS (Yosemite Aerosol Characterization Study) conducted from July to September 2002 in the western United States. The derived (estimated from OC/EC and scattering data) ensemble complex index of refraction of TBs was found to be 1.56–0.02i

Light absorption properties of laboratory generated tar ball particles

A. Hoffer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Pella Inc., USA) fixed on 13.1 mm spots of quartz filters placed in the filter holder that were used for sampling for elemental analysis as well.

The morphologies of the particles were studied in bright-field TEM images obtained using a Philips CM20 TEM operated at 200 kV accelerating voltage. The possible presence of an internal structure was checked in high-resolution electron micrographs and in selected-area electron diffraction patterns. The electron microscope was equipped with an ultra-thin-window Bruker Quantax X-ray detector that allowed the energy-dispersive X-ray analysis (EDS) of the elemental compositions of individual particles. Spectra were acquired for 60 s, with the diameter of electron beam adjusted to include the entire individual TB particles.

3 Results

3.1 Morphology, elemental composition and structure of the generated particles

Two samples were collected for TEM analysis to investigate the morphology and elemental composition of the generated particles, one representing the particles generated from the aqueous phase of the tar, whereas the other was collected from the oily phase. In both cases the oven temperature was set to 650 °C, the flows and other experimental parameters were similar to those applied for samples collected for TC analysis.

As it can be observed in Fig. 1, the particles generated from the aqueous phase were spherical. From the oily phase more irregularly shaped particles with oval two-dimensional outlines were produced, indicating that in the latter case the particles were not perfectly solid at the time of collection. It was observed during the TEM analysis that all of the generated particles can withstand the high-energy electron beam of the instrument: they did not evaporate or shrink while exposed to the electron beam.

Light absorption properties of laboratory generated tar ball particles

A. Hoffer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Light absorption properties of laboratory generated tar ball particles

A. Hoffer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The observed sizes of the particles vary widely (up to ~ 360 nm in diameter), the number size distribution peaks at ~ 100 nm as determined from the TEM images. Bimodal number size distribution was obtained from the DMPS measurements for the particles produced from both the aqueous and oily phase of the tar (Fig. 2). For the particles aged at 650°C the two modes are centred around 20–40 and 100–140 nm. The number size distributions of nigrosin and the blank (pure methanol) are unimodal peaking at 117 and 41 nm, respectively. Nevertheless, in cases when the ageing temperature was higher than 500°C the mass and volume of the particles are dominated by the larger particles: at least 86 and 70 % of the total mass is represented by the larger particle mode in the case of the aqueous and oily samples, respectively. Considering that both the absorption and scattering efficiencies are very small for small particles, the optical properties are also determined by the particles of the larger mode. (Here we note that the mass absorption coefficient was calculated only for size distributions in which the relative contribution of the second mode to total volume was larger than 93 %.) The sizes of the particles of the second mode were similar to those determined for ambient TB particles observed in samples from K-pusztá and Southern Africa (Pósfai et al., 2004).

The EDS spectra of the particles generated from both the aqueous and oily phase indicated that the particles consist predominantly of carbon and oxygen. In the case of the particles formed from the aqueous phase the average carbon to oxygen molar ratio was 10 : 1, with 90 mol % C (RSD = 10 %), 9 mol % O (RSD = 16 %) and N, Na, Si, S, K only in trace amounts. The limitations of determining molar ratios by this method are described in detail elsewhere (Pósfai et al., 2003). It should be noted that the spectra were practically indistinguishable from those obtained from atmospheric TBs. Both HRTEM images and electron diffraction confirm that the particles in both samples are perfectly amorphous, lacking even the short-range order that is characteristic of ns-soot (Buseck et al., 2014).

3.2 Measurement uncertainties

In order to estimate the measurement uncertainties, nigrosin dye (Sigma-Aldrich, Acid black 2, water soluble) was measured with the same setup that was used for the measurements of TBs. The nigrosin was also dissolved in methanol and particles were generated with the process used for the TB samples. Oven temperature was set to 65 °C in order to evaporate methanol from the droplets without inducing compositional changes of nigrosin. According to Massoli et al. (2009), the scattering coefficient of absorbing particles with single scattering albedo (SSA) = 0.4 (at 532 nm) is overestimated by 25 % using the Anderson and Ogren correction (Anderson and Ogren, 1998) for the raw data measured by a TSI nephelometer. Since in our case the SSA of the generated nigrosin was ~ 0.4 at 550 nm, the scattering coefficient might be also overestimated by $\sim 25\%$. The uncertainty of the measurements of Particle Soot Absorption Photometer (PSAP) whose measurement principle is very similar to that of the CLAP is 20–30 % (Bond, 1999). It was demonstrated that the presence of organic compounds (secondary organic aerosol, SOA) causes positive bias and enhances the uncertainty of the PSAP (Cappa et al., 2008; Lack et al., 2008). This effect has to be considered in the case of particles generated from tar which contain condensable organic compounds as well. Based on the above, if we consider that the CLAP overestimated the absorption of nigrosin by 25 % and the scattering is also overestimated by 25 %, we obtain a refractive index of $1.65-0.29i$ and $1.77-0.27i$ for nigrosin at wavelengths of 550 and 652 nm, respectively. The index of refraction of nigrosin at 633 nm was reported to be $1.67-0.26i$ (Pinnick et al., 1973). By assuming that the absorption is similar at both 633 and 652 nm, Mie calculations using the refractive index of nigrosin ($1.67-0.26i$) and the measured size distribution yield scattering and absorption coefficients at 652 nm higher by $\sim 17\%$ and lower by $\sim 2\%$, respectively, as compared to those directly measured. These uncertainties are considered when interpreting the results.

spectra (Alexander et al., 2008). The real part of the index of refraction as measured in our experiment is higher by about 10 % than the one calculated for the carbon spheres.

7 Conclusions

Tar balls have been shown to be abundant in the troposphere impacted by biomass smoke which is now the main global source of anthropogenic aerosol particles. The contribution of TBs to the number concentration of particles could be as high as 80 % in the vicinity of biomass burning sources (Pósfai et al., 2003), while it was in the range of 6–14 % away from the sources (Adachi et al., 2011), as observed using TEM. At a site that represents regional background conditions (K-puszta) the abundance of TBs varied from 0 to 40 % depending on the season and time of sampling (Pósfai et al., 2004). Even over the Himalaya TB particles accounted for 3 % of all observed particles (Cong et al., 2009). Near the Arctic, in Hyytiälä during a pollution episode 1–4 % of the particles were identified as TBs (Niemi et al., 2006). Given the abundance of TBs in the global troposphere and their relatively high absorption efficiency over the entire solar spectrum, their contribution to column absorption can be clearly significant. This is particularly true for immense geographical regions impacted by Atmospheric Brown Clouds (ABCs) where TBs may make a contribution to solar absorption comparable to that of BC. The last question that remains is where TBs are positioned in the black-to-brown carbon continuum of atmospheric aerosols (Andreae and Gelencsér, 2006; Sun et al., 2007). Somewhat surprisingly, their optical properties suggest that they are not very far from BC or amorphous carbon, despite their markedly different formation mechanism and chemical composition. On the other hand, it is clear that TBs are very much different from faintly coloured species such as HULIS or SOA in their absorption properties. We suggest that TBs are on the dark side of brownness of aerosol carbon, but clearly out of the BC regime both in terms of their key absorption parameters (e.g. refractive index and AAE) and for lack of fundamental properties (Petzold et al., 2013).

Light absorption properties of laboratory generated tar ball particles

A. Hoffer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



tory measurements, *Aerosol Sci. Tech.*, 42, 1022–1032, doi:10.1080/02786820802389285, 2008.

Chakrabarty, R. K., Moosmüller, H., Chen, L.-W. A., Lewis, K., Arnott, W. P., Mazzoleni, C., Dubey, M. K., Wold, C. E., Hao, W. M., and Kreidenweis, S. M.: Brown carbon in tar balls from smoldering biomass combustion, *Atmos. Chem. Phys.*, 10, 6363–6370, doi:10.5194/acp-10-6363-2010, 2010.

China, S., Mazzoleni, C., Gorkowski, K., Aiken, A. C., and Dubey, M. K.: Morphology and mixing state of individual freshly emitted wildfire carbonaceous particles, *Nature Comm.*, 4, 2122, doi:10.1038/ncomms3122, 2013.

Chung, C. E., Ramanathan, V., and Decremier, D.: Observationally constrained estimates of carbonaceous aerosol radiative forcing, *P. Natl. Acad. Sci. USA*, 109, 11624–11629, doi:10.1073/pnas.1203707109, 2012.

Clarke, A. D., Shinozuka, Y., Kapustin, V. N., Howell, S., Huebert, B., Doherty, S., Anderson, T. Covert, D., Anderson, J., Hua, X., Moore II, K. G., McNaughton, C., Carmichael, G., and Weber, R.: Size distributions and mixtures of dust and black carbon aerosol in Asian outflow: physiochemistry and optical properties, *J. Geophys. Res.*, 109, D15S09, doi:10.1029/2003JD004378, 2004.

Cong, Z., Kang, S., Dong, S., and Zhang, Y.: Individual particle analysis of atmospheric aerosols at Nam Co, Tibetan Plateau, *Aerosol Air Qual. Res.*, 9, 323–331, doi:10.4209/aaqr.2008.12.0064, 2009.

Fu, H., Zhang, M., Li, W., Chen, J., Wang, L., Quan, X., and Wang, W.: Morphology, composition and mixing state of individual carbonaceous aerosol in urban Shanghai, *Atmos. Chem. Phys.*, 12, 693–707, doi:10.5194/acp-12-693-2012, 2012.

Guyon, P., Boucher, O., Graham, B., Beck, J., Mayol-Bracero, O. L., Roberts, G. C., Maenhaut, W., Artaxo, P., and Andreae, M. O.: Refractive index of aerosol particles over the Amazon tropical forest during LBA-EUSTACH 1999, *J. Aerosol. Sci.*, 34, 883–907, doi:10.1016/S0021-8502(03)00052-1, 2003.

Hand, J. L., Malm, W. C., Laskin, A., Day, D., Lee, T., Wang, C., Carrico, C., Carrillo, J., Cowin, J. P., Collet Jr., J., and Iedema, M. J.: Optical, physical and chemical properties of tar balls observed during the Yosemite Aerosol Characterisation Study, *J. Geophys. Res.-Atmos.*, 110, D21210, doi:10.1029/2004JD005728, 2005.

Light absorption properties of laboratory generated tar ball particles

A. Hoffer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Light absorption properties of laboratory generated tar ball particles

A. Hoffer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Hoffer, A., Gelencsér, A., Guyon, P., Kiss, G., Schmid, O., Frank, G. P., Artaxo, P., and Andreae, M. O.: Optical properties of humic-like substances (HULIS) in biomass-burning aerosols, *Atmos. Chem. Phys.*, 6, 3563–3570, doi:10.5194/acp-6-3563-2006, 2006.
- Lack, D. A., Cappa, C. D., Covert, D. S., Baynard, T., Massoli, P., Sierau, B., Bates, T. S., Quinn, P. K., Lovejoy, E. R., and Ravishankara, A. R.: Bias in filter-based aerosol light absorption measurements due to organic aerosol loading: evidence from ambient measurements, *Aerosol Sci. Tech.*, 42, 1033–1041, doi:10.1080/02786820802389277, 2008.
- Maschio, G., Koufopoulos, C., and Lucchesi, A.: Pyrolysis, a promising route for biomass utilization, *Bioresource Technol.*, 42, 219–231, 1992.
- Massoli, P., Murphy, D. M., Lack, D. A., Baynard, T., Brock, C. A., and Lovejoy, E. R.: Uncertainty in light scattering measurements by TSI nephelometer: results from laboratory studies and implications for ambient measurements, *Aerosol Sci. Tech.*, 43, 1064–1074, doi:10.1080/02786820903156542, 2009.
- Moosmüller, H., Chakrabarty, R. K., and Arnott, W. P.: Aerosol light absorption and its measurement: a review, *J. Quant. Spectrosc. Ra.*, 110, 844–878, 2009.
- Moosmüller, H., Chakrabarty, R. K., Ehlers, K. M., and Arnott, W. P.: Absorption Ångström coefficient, brown carbon, and aerosols: basic concepts, bulk matter, and spherical particles, *Atmos. Chem. Phys.*, 11, 1217–1225, doi:10.5194/acp-11-1217-2011, 2011.
- Niemi, J. V., Saarikoski, S., Tervahattu, H., Mäkelä, T., Hillamo, R., Vehkamäki, H., Sogacheva, L., and Kulmala, M.: Changes in background aerosol composition in Finland during polluted and clean periods studied by TEM/EDX individual particle analysis, *Atmos. Chem. Phys.*, 6, 5049–5066, doi:10.5194/acp-6-5049-2006, 2006.
- Okuyama, K. and Lenggoro, I. W.: Preparation of nanoparticles via spray route, *Chem. Eng. Sci.*, 58, 537–547, doi:10.1016/S0009-2509(02)00578-X, 2003.
- Petzold, A., Ogren, J. A., Fiebig, M., Laj, P., Li, S.-M., Baltensperger, U., Holzer-Popp, T., Kinne, S., Pappalardo, G., Sugimoto, N., Wehrli, C., Wiedensohler, A., and Zhang, X.-Y.: Recommendations for reporting "black carbon" measurements, *Atmos. Chem. Phys.*, 13, 8365–8379, doi:10.5194/acp-13-8365-2013, 2013.
- Pinnick, R. G., Rosen, J. M., and Hofmann, D. J.: Measured light-scattering properties of individual aerosol particles compared to Mie scattering theory, *Appl. Optics*, 12, 37–43, 1973.
- Pósfai, M., Simonics, R., Li, J., Hobbs, P. V., and Buseck, P. R.: Individual aerosol particles from biomass burning in southern Africa: 1. Compositions and size distributions of carbonaceous particles, *J. Geophys. Res.-Atmos.*, 108, 8483, doi:10.1029/2002JD002291, 2003.

**Light absorption
properties of
laboratory generated
tar ball particles**

A. Hoffer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Pósfai, M., Gelencsér, A., Simonics, R., Arató, K., Li, J., Hobbs, P. V., and Buseck, P. R.: Atmospheric tar balls: particles from biomass and biofuel burning, *J. Geophys. Res.-Atmos.*, 109, D06213, doi:10.1029/2003JD004169, 2004.
- 5 Schnaiter, M., Horvath, H., Mohler, O., Naumann, K. H., Saathoff, H., and Schock, O. W.: UV-VIS-NIR spectral optical properties of soot and soot-containing aerosols, *J. Aerosol. Sci.*, 34, 1421–1444, 2003.
- Sun, H., Biedermann, L., and Bond, T.: Color of brown carbon: a model for ultraviolet and visible light absorption by organic carbon aerosol, *Geophys. Res. Lett.*, 34, L17813, doi:10.1029/2007GL029797, 2007.
- 10 Taha, G., Box, P. G., Cohen, D. D., and Stelcer, E.: Black carbon measurement using laser integrating plate method, *Aerosol Sci. Tech.*, 41, 266–276, doi:10.1080/02786820601156224, 2007.
- 15 Tóth, A., Hoffer, A., Nyiró-Kósa, I., Pósfai, M., and Gelencsér, A.: Atmospheric tar balls: aged primary droplets from biomass burning?, *Atmos. Chem. Phys.*, 14, 6669–6675, doi:10.5194/acp-14-6669-2014, 2014.

Light absorption properties of laboratory generated tar ball particles

A. Hoffer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1. Optical parameters of tar ball particles generated from the aqueous phase.

Sample name	Oven temperature (°C)	AAE	MAE (m ² g ⁻¹)	<i>Re</i>	<i>Im</i>	Volume of large particles (%)
18-d1	500	3.4				59
14-d1	650	2.8	0.8–2.5	1.88	0.27	98
15-d1	650	3.4		1.79	0.15	86
16-d2	650	2.8		1.87	0.27	99
17-d1	650	2.8	1.0–3.0	1.82	0.18	93
22-d2	650	3.0	0.8–2.3			93
25-d2	650	2.7	0.8–2.3	1.84	0.17	95
20-d1	800	3.0	1.0–3.1			97
20-d2	800	3.0				96




Figure 1. TEM images of tar balls generated from the aqueous (**a, b**) (sample 16-d2) and oily (**c, d**) phase of tar obtained from dry distillation of wood.

Light absorption properties of laboratory generated tar ball particles

A. Hoffer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Light absorption properties of laboratory generated tar ball particles

A. Hoffer et al.




Figure 2. Number and volume size distribution of particles generated from aqueous tar (sample 16-d2) measured with DMPS and TEM.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Light absorption properties of laboratory generated tar ball particles

A. Hoffer et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Figure 3. The effect of heat shock (oven temperature) on the Ångström exponent of TB particles generated from the aqueous and from the oily phase of wood tar.