

Coverings by few monochromatic pieces - a transition between two Ramsey problems

András Gyárfás

Computer and Automation Research Institute
Hungarian Academy of Sciences
Budapest, P.O. Box 63
Budapest, Hungary, H-1518
`gyarf@szta.hu`

Gábor N. Sárközy*

Computer Science Department
Worcester Polytechnic Institute
Worcester, MA, USA 01609
`gsarkozy@cs.wpi.edu`

and

Computer and Automation Research Institute
Hungarian Academy of Sciences
Budapest, P.O. Box 63
Budapest, Hungary, H-1518

Stanley Selkow

Computer Science Department
Worcester Polytechnic Institute
Worcester, MA, USA 01609
`sms@cs.wpi.edu`

April 4, 2013

Abstract

¹ The typical problem in (generalized) Ramsey theory is to find the order of the largest monochromatic member of a family \mathcal{F} (for example matchings, paths, cycles, connected subgraphs) that must be present in any edge coloring of a complete graph K_n with t colors. Another area is to find the minimum number of monochromatic members of \mathcal{F} that partition or cover the vertex set of every edge colored complete graph. Here we propose a problem that connects these areas: for a fixed positive integers $s \leq t$, at least how many vertices can be covered by the vertices of no more than s monochromatic members of \mathcal{F} in every edge coloring of

*Research supported in part by the National Science Foundation under Grant No. DMS-0968699.

¹Apart from a footnote referring to Pokrovskiy [24], this manuscript was submitted to Graphs and Combinatorics in May, 2011

K_n with t colors. Several problems and conjectures are presented, among them a possible extension of a well-known result of Cockayne and Lorimer on monochromatic matchings for which we prove an initial step: every t -coloring of K_n contains a $(t-1)$ -colored matching of size k provided that

$$n \geq 2k + \left\lfloor \frac{k-1}{2^{t-1}-1} \right\rfloor.$$

1 Introduction

The typical problem in (generalized) Ramsey theory is to find the order of the largest monochromatic member of a family \mathcal{F} (for example matchings, paths, cycles, connected subgraphs) that must be present in any edge coloring of a complete graph K_n with t colors. For easier reference these problems are called Ramsey problems in this paper. Another well studied area, we call them *cover problems*, is to find the minimum number of monochromatic members of \mathcal{F} that partition or cover the vertex set of every edge colored complete graph.

Here we propose a common generalization of Ramsey and cover problems. For a fixed positive integer s , at least how many vertices can be covered by the vertices of no more than s monochromatic members of \mathcal{F} in every edge coloring of K_n with t colors? A somewhat related problem was proposed by Chung and Liu [4]: for a given graph G and for fixed s, t , find the smallest n such that in every t -coloring of the edges of K_n there is a copy of G colored with at most s colors.

Several problems and conjectures are formulated, among them a possible extension of a well-known result of Cockayne and Lorimer on monochromatic matchings [5]. Our main result (Theorem 2.3) is that every t -coloring of K_n contains a $(t-1)$ -colored matching of size k provided that

$$n \geq 2k + \left\lfloor \frac{k-1}{2^{t-1}-1} \right\rfloor.$$

This result is sharp. A simple consequence (Corollary 2.4) is that every t -colored K_{2^t-2} has a perfect matching missing at least one color. This is a special case of a conjecture posed in [11].

1.1 Path and cycle covers

As far as path covers in infinite graphs are concerned, Rado [25] has a “perfect” result stated here in an abridged form with its simplified original proof.

Theorem 1.1. *The vertex set of any t -colored countable complete graph can be partitioned into finite or one-way infinite monochromatic paths, each of a different color.*

Proof. Call a set $C \subseteq \{1, \dots, t\}$ of k colors, $1 \leq k \leq t$, *perfect* if there exists a set $\mathcal{P} = \{P_1, \dots, P_k\}$ of k vertex disjoint finite paths $P_1 = \dots x_1, \dots, P_k = \dots, x_k$ with the following property: P_i is monochromatic in color c_i and there is an

infinite set Y of vertices such that Y is disjoint from the paths of \mathcal{P} and for each $i \in \{1, \dots, k\}$ and for all $y \in Y$, $x_i y$ is colored with c_i . A perfect color set exists since any color c_1 present on infinitely many edges of a star incident to vertex x forms such a (one-element) set. Select a perfect set C of k colors so that k is as large as possible ($k \leq t$), this is witnessed by \mathcal{P} and Y . Let u be an arbitrary vertex not covered by \mathcal{P} . Consider a color c such that uy has color c for every $y \in Y^*$ where $Y^* \subseteq Y$, is infinite. It follows from the choice of k that $c \in C$. Now u can be added to the end of the c -colored path of \mathcal{P} , either directly if $u \in Y$, or through a vertex $v \in Y^*$ if $u \notin Y$. The infinite set witnessing the extension is either Y^* or $Y^* \setminus \{v\}$. Clearly the extensions can be continued to place all vertices of the countable complete graph so that all paths of \mathcal{P} are finite or one-way infinite. \square

There are several possibilities to “finitize” Theorem 1.1. The 2-color version works perfectly as noted in a footnote in [8].

Proposition 1.2. *The vertex set of any 2-colored finite complete graph can be partitioned into monochromatic paths, each of a different color.*

Proof. If $P_1 = \dots, x_1, P_2 = \dots, x_2$ are red and blue paths and v is uncovered then either v can be placed as the last vertex of one of the paths P_i or one of the bypasses P_1, x_2, v or P_2, x_1, v extends one monochromatic path (and shortens the other). \square

Lehel conjectured that Proposition 1.2 remains true if paths are replaced by cycles (where the empty set, one vertex and one edge are accepted as a cycle). Although the existence of a ‘near partition’ (where the two monochromatic cycles intersect in at most one vertex) follows easily, see [10], it took a long time until this was proved for large n in [21], [1]. Recently an elementary proof was found by Bessy and Thomassé [3] that works for all n .

Theorem 1.3. ([3]) *The vertex set of any 2-colored complete graph can be partitioned into two monochromatic cycles of different colors.*

In [16] and [6] several possible extensions of Proposition 1.2 were suggested. It turned out that for 3 colors one can not except full partition by distinct colors, the first example of this phenomenon is from Heinrich [18]. Recently the asymptotic ratios of monochromatic path and cycle partitions with three distinct colors was obtained in [13].

Theorem 1.4. ([13]) *In every 3-colored K_n at least $(\frac{3}{4} - o(1))n$ vertices can be partitioned into monochromatic cycles of distinct colors.*

We note that here the asymptotic ratio $\frac{3}{4}$ is best possible. Nevertheless it was conjectured in [6] that Proposition 1.2 and Theorem 1.3 carries over to any number of colors if repetitions of colors are possible.

Conjecture 1.5. ([6]) *The vertex set of every t -colored complete graph can be partitioned into t monochromatic cycles.*

² The case of three colors was recently solved in asymptotic sense.

²Pokrovskiy [24] proved that Conjecture is not quite true for $t = 3$

Theorem 1.6. ([13]) *In every 3-colored K_n at least $(1 - o(1))n$ vertices can be partitioned into three monochromatic cycles.*

The proofs of Theorems 1.4, 1.6 rely on the Regularity Lemma and relaxations of cycles to connected matchings (see Subsection 1.2). The new problem we propose here is the following.

Problem 1.7. *Suppose $1 \leq s \leq t$. What is the maximum number of vertices that can be covered by s monochromatic cycles (paths) in every t -coloring of the edges of K_n ?*

We do not have a general conjecture here, not even for the asymptotics (for fixed s, t and large n). The case $s = 1$ is the path Ramsey number where the case $t = 3$ ([12]) is the only evidence that perhaps $\frac{n}{t-1}$ is the true asymptotic value ($\frac{n}{t}$ is an easy lower bound). The case $s = t$ is in Conjecture 2.

The first interesting special case is $t = 3, s = 2$.

Conjecture 1.8. *In any 3-colored K_n there are two vertex disjoint monochromatic paths (cycles) covering at least $\frac{6n}{7}$ vertices.*

A weaker form (for matchings instead of paths) of Conjecture 1.8 follows from Theorem 2.3 below (when $t = 3$).

1.2 Connected matchings.

One technique used recently in many papers (for example [7], [12],[15]) in Ramsey and in covering problems related to paths or cycles is to replace the paths or cycles by a simpler structure, *monochromatic connected matchings*, and rely on the Regularity and Blow-up Lemmas to create paths or cycles from them. A connected monochromatic matching means that all edges of the matching are in the same component of the subgraph induced by the edges in the color of the matching.

Thus a natural step towards proving an asymptotic (or sharp for large enough n) version of Conjecture 1.8 would be to prove it for connected matchings. However, in the problem mentioned above we are a step behind, we could only prove Conjecture 1.8 for matchings (without the connectivity condition). A logical plan is to treat connected pieces and matchings separately, this is done in Subsection 1.3 and in Section 2.

1.3 Covers by connected components

Since every connected component contains a spanning tree of the component, we use here the somewhat simpler tree language. A special case of a conjecture attributed to Ryser, (appearing in his student, Henderson's thesis [19]) states that every intersecting t -partite hypergraph has a transversal of at most $t - 1$ elements. Using the dual of the hypergraph of monochromatic components in a t -coloring of complete graphs, one can easily see that the following form of the conjecture (introduced in [9]) is equivalent.

Conjecture 1.9. *In every t -coloring of the edges of a complete graph, the vertex set can be covered by the vertices of at most $t - 1$ monochromatic trees.*

If Conjecture 1.9 is true then an easy averaging argument would easily extend it as follows.

Conjecture 1.10. *For every $1 \leq s \leq t - 1$ and for every t -coloring of K_n at least $\frac{ns}{t-1}$ vertices can be covered by the vertices of at most s monochromatic trees.*

Since Conjecture 1.9 is known to be true for $2 \leq t \leq 5$, Conjecture 1.10 is true for $1 \leq s \leq t \leq 5$. Also, the case $s = 1$ is known for arbitrary t (originally in [9], [17] is a recent survey). Perhaps a good test case is to try to prove Conjecture 1.10 for $s = 2$ (and for general t).

Since Ryser's conjecture is extended further in [6] by changing cover to partition in Conjecture 1.9, one may perhaps even require partition in Conjecture 1.10 as well.

1.4 Covers by copies of a fixed graph

It seems that to find the percentage of vertices that can be covered by monochromatic copies of a fixed graph H having at most s colors is a difficult problem. Indeed, even the case when H is a single edge seems difficult. However, somewhat surprisingly, for any fixed *connected non-bipartite graph* H and for any fixed $t \geq 3$ and fixed $s \leq t$, the percentage of vertices of K_n that can be covered by vertex disjoint monochromatic copies having at most s colors can be rather well approximated. In fact, the following theorem can be easily obtained from the results of [22]. Let $R_t(H)$ denote the smallest integer m such that in every t -coloring of the edges of K_n there is a monochromatic copy of H .

Theorem 1.11. *Suppose that $t \geq 3, 1 \leq s \leq t$ and H is a connected non-bipartite graph. Then in every t -coloring of the edges of K_n , at least $\frac{s(n-R_t(H))}{t}$ vertices can be covered by vertex disjoint monochromatic copies of H using at most s colors. On the other hand, for any n that is divisible by t , the edges of K_n can be t -colored so that at most $\frac{sn}{t}$ vertices can be covered by vertex disjoint monochromatic copies of H having at most s colors.*

Proof. The first part follows by selecting successively monochromatic copies of H , removing after each step the part covered so far. Clearly, the process stops only when less than $R_t(H)$ vertices remain. Then, an obvious averaging argument gives that the copies in s suitable colors cover the claimed quantity.

The second part follows from the following construction. Partition $V(K_n)$ into t equal parts and color the edges within the parts with t different colors, say within part i every edge gets color i . The crossing edges (going from one part to another) are all colored with the same color between any fixed pair of parts. There are two rules. On one hand, crossing edges of color i cannot be incident to part i . On the other hand, the union of crossing edges of color i should span a bipartite graph. It is easy to see that these rules can be easily guaranteed for

$t \geq 3$ (and impossible to meet for $t = 2$). Because H is connected, not bipartite and crossing edges of color i are not adjacent to part i , each monochromatic copy of H must be completely within a part. Therefore copies of H having at most s colors are covered by the at most s parts, proving the second statement of the theorem. \square

2 Covers by matchings - how to generalize Cockayne - Lorimer theorem?

Here we return to the most basic case, when we want to cover by copies of an edge, i.e. by matchings. A matching in a t -colored complete graph is called an s -colored matching if at most s colors are used on its edges. To describe easily certain t -colorings of K_n we need, consider partition vectors with t positive integer coordinates whose sum is equal to n . Assume that $V(K_n) = \{1, 2, \dots, n\}$. Then $[p_1, p_2, \dots, p_t]$ represents the coloring obtained by partitioning $V(K_n)$ into parts A_i so that $|A_i| = p_i$ for $i = 1, 2, \dots, t$ and the color of any edge $e = (x, y)$ is the minimum j for which $\{x, y\}$ has non-empty intersection with A_j .

Problem 2.1. *Suppose $1 \leq s \leq t$. What is the size of the largest s -colored matching that can be found in every t -coloring of the edges of K_n ?*

The Ramsey problem, the case $s = 1$ in Problem 2.1, was completely answered by Cockayne and Lorimer [5]. Here we state its diagonal case only.

Theorem 2.2. ([5]) *Assume $n \geq (t + 1)p + 2$ and K_n is arbitrarily t -colored. Then there is a monochromatic matching of size $p + 1$.*

Observe that Theorem 2.2 is sharp, the coloring $[p, p, \dots, p, 2p + 1]$ of $K_{(t+1)p+1}$ does not contain a monochromatic matching with $p + 1$ edges.

Notice that the case $s = t$ of Problem 2.1 is trivial, any perfect (or near-perfect if n is odd) matching is obviously optimal. In this paper we settle the case $s = t - 1$, by showing that the extremal coloring is close to the coloring $[p, 2p, 4p, \dots, 2^{t-1}p]$. More precisely we prove the following.

Theorem 2.3. *Every t -coloring of K_n contains a $(t - 1)$ -colored matching of size k provided that*

$$n \geq 2k + \left\lfloor \frac{(k - 1)}{2^{t-1} - 1} \right\rfloor.$$

This is sharp for every $t \geq 2, k \geq 1$.

In case of $t = 3$ Theorem 2.3 gives Conjecture 1.8 in a weaker form. Noting that for $k < 2^{t-1}$ the second term is zero in Theorem 2.3, we get the following.

Corollary 2.4. *Every t -colored K_{2^t-2} has a perfect matching missing at least one color.*

We note here that for $t = 2, 3, 4$ there are results stronger than Theorem 2.3. Namely, not only a $(t - 1)$ -colored matching of size k can be guaranteed, but a $(t - 1)$ -colored path on $2k$ vertices. For $t = 2$ this is a well-known result [8], for $t = 3$ it was proved in [23] and for $t = 4$ in [20]. In fact, it was conjectured in [20] that Theorem 2.3 holds also if the matching of size k is replaced by P_{2k} .

For the case $t = 4, s = 2$ we suspect that the extremal coloring is essentially $[p, p, 2p, 4p]$. That leads to

Conjecture 2.5. *If $n \geq \lfloor \frac{8k-2}{3} \rfloor$ then every 4-coloring of K_n contains a 2-colored matching of size k .*

For the case $s = 2, t = 5$, the coloring $[p, p, p, 2p, 4p]$ and the coloring $[p, p, p, p, 2p]$ that belongs to Theorem 2.2 give essentially the same parameters so we do not risk a conjecture here. Moreover, for $s = 2, t = 6$ the latter coloring $[p, p, p, p, p, 2p]$ is better than $[p, p, p, p, 2p, 4p]$. This leads to the dilemma whether there are better coloring in this case or $[p, p, p, p, p, 2p]$ is the extremal one? The latter possibility would be similar to the phenomenon discussed in Subsection 1.4, saying vaguely that in a 6-colored complete graph the size of the largest 2-matching is twice the size of the largest monochromatic matching.

3 Large $(t - 1)$ -colored matchings in t -colored complete graphs.

Here we prove Theorem 2.3. To show that it is sharp, set $N = 2k - 1 + \lfloor \frac{(k-1)}{2^{t-1}-1} \rfloor = 2k - 1 + p$, where $p = \lfloor \frac{N}{2^t-1} \rfloor = \lfloor \frac{k-1}{2^{t-1}-1} \rfloor$. Consider the coloring $[p, 2p, 4p, \dots, 2^{t-2}p, q]$ of K_N with $q = N - (2^{t-1} - 1)p$.

If a matching in this coloring misses color $j \neq t$ then it misses at least $2^{j-1}p - \sum_{i < j} 2^{i-1}p = p$ vertices from the vertex set to which color class j is incident to. Thus at most $N - p = 2k - 1$ vertices are covered by this matching so its size is smaller than k . A matching that misses color t has at most $\sum_{i < t} 2^{i-1}p = (2^{t-1} - 1)p \leq k - 1$ edges.

To prove the upper bound, consider a t -colored K_n where $n = 2k + \lfloor \frac{(k-1)}{2^{t-1}-1} \rfloor = 2k + p$. Set $V = V(K_n)$, let G_i denote the subgraph of K_n with vertex set V and containing edges of colors different from color i , $1 \leq i \leq t$. We are going to show that for at least one i , G_i has a matching of size k . The proof is indirect: if the maximum matching of G_i , $\nu(G_i)$, is at most $k - 1$ for each i , then for the deficiency of G_i , $def(G_i)$, defined as the the number of vertices uncovered by any maximum matching of G_i , we have

$$def(G_i) \geq n - 2\nu(G_i) \geq 2k + p - 2(k - 1) = p + 2.$$

We apply the following well-known result, where $c_o(G)$ is the number of odd components of G .

Theorem 3.1. (Berge formula) $def(G) = \max\{c_o(V(G) \setminus X) - |X| : X \subset V(G)\}$.

Thus, for each i , there is a set $X_i \subset V$ such that

$$c_o(V \setminus X_i) \geq |X_i| + p + 2. \quad (1)$$

Assume w.l.o.g that $|X_1| \leq \dots \leq |X_t|$ and observe that the edges between connected components of G_i in $V \setminus X_i$ are all colored with color i . Let $C_1^1, \dots, C_{m_1}^1$ be the vertex sets of the connected components of G_1 in $V \setminus X_1$, from (1) we have $m_1 \geq |X_1| + p + 2$.

Lemma 3.2. *There is an index $l \in \{1, 2, \dots, m_1\}$, say $l = 1$, such that for every $j > 1$, $\cup_{i \neq 1} C_i^1 \subset X_j$.*

Proof. Suppose that $v, w \notin X_2$ where $v \in C_q^1, w \in C_r^1$ and $q \neq r$. This implies that the edges of color 1 form a complete multipartite graph M on $V \setminus (X_1 \cup X_2)$ with at least two partite classes. Therefore all vertices of M must be in the same connected component of G_2 in $V \setminus X_2$. Thus G_2 has at most $1 + |X_1| \leq 1 + |X_2| < |X_2| + p + 2$ odd components in $V \setminus X_2$, contradicting (1). Thus X_2 must cover all but at most one among the C_i^1 -s, say C_1^1 can be uncovered.

Next we show that for all $j \geq 2$, we have $\cup_{i \neq 1} C_i^1 \subset X_j$. We have seen this for $j = 2$, so assume $j > 2$. The argument of the previous paragraph gives that X_j covers all but one C_i^1 , say the exceptional one is C_l^1 . Suppose $l \neq 1$, say $l = 2$. The inequality $|X_2| \leq |X_j|$ implies $|X_2 \setminus X_j| \leq |X_j \setminus X_2|$ and from this

$$|X_1 \cap (X_2 \setminus X_j)| + |C_2^1 \setminus X_j| \leq |X_1 \cap (X_j \setminus X_2)| + |C_1^1 \setminus X_2|. \quad (2)$$

On the other hand, using (1),

$$\begin{aligned} |V \setminus X_j| &= |X_1 \cap (X_2 \setminus X_j)| + |X_1 \setminus (X_2 \cup X_j)| + |C_2^1 \setminus X_j| \geq c_o(V \setminus X_j) \geq |X_j| + p + 2 \\ &\geq |X_1 \cap (X_j \setminus X_2)| + |C_1^1 \setminus X_2| + |X_j \cap C_2^1| + |\cup_{i \geq 3} C_i^1| + p + 2 \end{aligned}$$

which can be rearranged as

$$\begin{aligned} (|X_1 \cap (X_2 \setminus X_j)| + |C_2^1 \setminus X_j|) - (|X_1 \cap (X_j \setminus X_2)| + |C_1^1 \setminus X_2|) + |X_1 \setminus (X_2 \cup X_j)| &\geq \\ &\geq |X_j \cap C_2^1| + |\cup_{i \geq 3} C_i^1| + p + 2. \end{aligned} \quad (3)$$

Note that from (2) the left hand side of (3) is at most $|X_2 \cup X_j|$. Thus from (3) we get

$$|X_1| \geq |X_1 \setminus (X_2 \cup X_j)| \geq |X_j \cap C_2^1| + |\cup_{i \geq 3} C_i^1| + p + 2 \geq m_1 - 2 + p + 2 = m_1 + p$$

and this contradicts $m_1 \geq |X_1| + p + 2$ and finishes the proof of the lemma. \square

Call $Ker_1 = C_2^1 \cup \dots \cup C_{m_1}^1$ the first kernel. With this notation Lemma 3.2 claims that each X_j with $j > 1$ contains Ker_1 . We may iterate Lemma 3.2 to define the set $Ker_i = C_2^i \cup \dots \cup C_{m_i}^i$, the i -th kernel, so that each X_j with $j > i$ contains Ker_i . Furthermore, these kernels are disjoint, since X_{i+1} contains Ker_i , but Ker_{i+1} is contained in $V \setminus X_{i+1}$. This implies that we have the following recursion on the sizes of the X_i 's.

Claim 1. *For every $2 \leq i \leq t$ we have*

$$|X_i| \geq |X_1| + \dots + |X_{i-1}| + (i-1)(p+1).$$

Indeed, X_i contains all the disjoint kernels Ker_1, \dots, Ker_{i-1} and thus using (1) we get

$$|X_i| \geq \sum_{j=1}^{i-1} |Ker_j| \geq \sum_{j=1}^{i-1} (c_0(V \setminus X_j) - 1) \geq \sum_{j=1}^{i-1} (|X_j| + p + 1),$$

as desired.

Claim 1 implies easily by induction the following

$$|X_i| \geq (2^{i-1} - 1)(p + 1). \quad (4)$$

But then, since the kernels are disjoint, using (1) again we get the following contradiction

$$\begin{aligned} n &\geq \sum_{i=1}^t |Ker_i| \geq \sum_{i=1}^t (c_0(V \setminus X_i) - 1) \geq \sum_{i=1}^t (|X_i| + p + 1) = \\ &= \sum_{i=1}^t |X_i| + t(p + 1) \geq \sum_{i=1}^t 2^{i-1}(p + 1) = (2^t - 1)(p + 1) > n. \end{aligned}$$

Here for the last inequality we have to check

$$(2^t - 1)(p + 1) \geq n + 1 = 2k + p + 1.$$

This is equivalent to

$$p + 1 \geq \frac{k}{2^{t-1} - 1}, \quad (5)$$

which is always true for our choice $p = \lfloor \frac{k-1}{2^{t-1}-1} \rfloor$. Indeed,

$$\left\lfloor \frac{k-1}{2^{t-1}-1} \right\rfloor = \left\lfloor \frac{k}{2^{t-1}-1} \right\rfloor$$

(and so (5) is trivially true) for all cases except when $\frac{k}{2^{t-1}-1}$ is an integer, but (5) is true in this case as well, finishing the proof of Theorem 2.3. \square

References

- [1] P. Allen, Covering two-edge-coloured complete graphs with two disjoint monochromatic cycles, *Combinatorics, Probability and Computing*, **17** (2008), 471-486.
- [2] S. A. Burr, P. Erdős, J. H. Spencer, Ramsey theorems for multiple copies graphs, *Transactions of the American Mathematical Society* **209** (1975), pp. 87-99.

- [3] S. Bessy, S. Thomassé, Partitioning a graph into a cycle and an anticycle, a proof of Lehel's conjecture, *Journal of Combinatorial Theory B.*, **100** (2009), 176-180.
- [4] K.M. Chung, C.L. Liu, A generalization of Ramsey theory for graphs, *Discrete Mathematics* **2** (1978) 117-127.
- [5] E. J. Cockayne, P. J. Lorimer, The Ramsey number for stripes, *J. Austral. Math. Soc.* **19** (1975), pp. 252-256.
- [6] P. Erdős, A. Gyárfás, L. Pyber, Vertex coverings by monochromatic cycles and trees, *Journal of Combinatorial Theory B* **51** (1991) 90-95.
- [7] A. Figaj, T. Luczak, The Ramsey number for a triple of long even cycles, to appear in the *Journal of Combinatorial Theory, Ser. B* **97** (2007) 584-596.
- [8] L. Gerencsér, A. Gyárfás, On Ramsey type problems, *Ann. Univ. Sci. Eötvös, Budapest* **10** (1967) 167 - 170.
- [9] A. Gyárfás, Partition coverings and blocking sets in hypergraphs (in Hungarian) *Communications of the Computer and Automation Institute of the Hungarian Academy of Sciences* **71** (1977) 62 pp.
- [10] A. Gyárfás, Vertex coverings by monochromatic paths and cycles, *Journal of Graph Theory* **7**. (1983)131-135.
- [11] A. Gyárfás, Large matchings with few colors - a problem for 'Emléktábla' workshop, manuscript, 2010.
- [12] A. Gyárfás, M. Ruszinkó, G. N. Sárközy, E. Szemerédi, Three-color Ramsey numbers for paths, *Combinatorica*, **27** (2007), pp. 35-69.
- [13] A. Gyárfás, M. Ruszinkó, G. N. Sárközy, E. Szemerédi, Partitioning 3-colored complete graphs into three monochromatic cycles, *Electronic J. of Combinatorics* **18**(2011) N.53.
- [14] A. Gyárfás, G. N. Sárközy, E. Szemerédi, The Ramsey number of diamond matchings and loose cycles in hypergraphs, *Electronic Journal of Combinatorics*, **15** (2008), R126.
- [15] A. Gyárfás, M. Ruszinkó, G. N. Sárközy, E. Szemerédi, One-sided coverings of complete bipartite graphs, in Algorithms and Combinatorics 26, Topics in Discrete Mathematics, 133-144.
- [16] A. Gyárfás, Monochromatic path covers, *Congressus Numerantium* **109** (1995), 201-202.
- [17] A. Gyárfás, Large monochromatic components in edge colorings of graphs - a survey, '*Ramsey Theory Yesterday, Today and Tomorrow, DIMACS workshop 2008* To appear in 'Progress in Mathematics' series.

- [18] Kathy Heinrich, personal communication, 1994.
- [19] J. R. Henderson, Permutation Decomposition of (0-1)-Matrices and Decomposition Transversals, Ph.D. thesis, Caltech, 1971.
- [20] A. Khamseh, G. R. Omid, A generalization of Ramsey theory for linear forests, manuscript submitted in 2010.
- [21] T. Łuczak, V. Rödl, E. Szemerédi, Partitioning two-colored complete graphs into two monochromatic cycles, *Combinatorics, Probability and Computing*, **7** (1998), 423-436.
- [22] P.J. Lorimer, R.J. Segegin, Ramsey numbers for multiple copies of complete graphs, *Journal of Graph Theory*, **2** (1978) 89-91.
- [23] R.Meenakshi, P.S. Sundararaghavan, Generalized Ramsey numbers for paths in 2-chromatic graphs, *Internat. J. Math. and Math. Sci.* **9** (1986) 273-276.
- [24] A. Pokrovskiy, Partitioning edge-coloured complete graphs into monochromatic cycles and paths, arXiv:12.05.5492v1
- [25] R. Rado, Monochromatic paths in graphs, *Annals of Discrete Mathematics* **3** (1987) 89-91.