

EVALUATION OF CELL DISRUPTION FOR PARTIAL ISOLATION OF INTRACELLULAR PYRUVATE DECARBOXYLASE ENZYME BY SILVER NANOPARTICLES METHOD

J. TANGTUA^a, C. TECHAPUN^b, R. PRATANAPHON^b, A. KUNTIYA^b, V. SANGUANCHAIWONG^c, T. CHAIYASO^b, P. HANMUANGJAI^b, P. SEESURIYACHAN^b, N. LEKSAWASDI^d, and N. LEKSAWASDI^{d*}

^aDivision of Food Science and Technology, Faculty of Science and Technology,
Kamphaeng Phet Rajabhat University, Kamphaeng Phet, 62000. Thailand

^bDivision of Biotechnology & Bioprocess Research Cluster, School of Agro-Industry, Faculty of Agro-Industry,
Chiang Mai University, Chiang Mai, 50100. Thailand

^cDepartment of Applied Biology & Bioprocess Research Cluster, Faculty of Science, King Mongkut's Institute of
Technology Ladkrabang, Bangkok, 10520. Thailand

^dDivision of Food Engineering, School of Agro-Industry, Faculty of Agro-Industry, Chiang Mai University,
Chiang Mai, 50100. Thailand

(Received: 11 February 2014; accepted: 14 April 2014)

Candida tropicalis TISTR 5350 was used in the comparison of seven concentration levels of silver nanoparticles (0, 5, 10, 15, 20, 25, and 30 $\mu\text{g mL}^{-1}$) for cell disruption methods. The optimized cell disruption strategy was selected based on the optimal protein yield and biological activity. The maximum volumetric and specific pyruvate decarboxylase (PDC, EC 4.1.1.1) activities ($0.53 \pm 0.05 \text{ U mL}^{-1}$ and $0.17 \pm 0.02 \text{ U mg}^{-1} \text{ protein}$, respectively) were observed at 15 $\mu\text{g mL}^{-1}$ silver nanoparticles. The silver nanoparticle concentration level of 15 $\mu\text{g mL}^{-1}$ was investigated further by comparing the reaction mixtures at different time intervals of 0, 1, 2, 3, 4, 5, and 6 min. The result showed that the highest specific PDC activity of $0.39 \pm 0.01 \text{ U mg}^{-1} \text{ protein}$ was obtained from mixing for 3 min. This was not significantly different ($P \leq 0.05$) from other mixing time intervals.

Keywords: silver nanoparticles, cell disruption, pyruvate decarboxylase, protein concentration, enzyme activity

The disruption of cell walls is an important step to isolate and extract substances from inside the cells and for preparation of intracellular products for biopharmaceutical usage. The overall goal in cell disruption is to obtain the desired product from within the cells, thus cell walls must be disrupted to release the contents of the cells. In essence, the objective of cell disruption is to achieve high protein recovery, while maintaining biological activity and high reproducibility (GERDEA et al., 2012).

Current commercial phenylacetylcarbinol (PAC) production processes involve the use of fermenting yeast to produce sufficient biomass for the associated accumulation of the intracellular pyruvate and the induction of PDC synthesis in a fed-batch process (TANGTUA et al., 2013). An enzymatic process based on extracted pyruvate decarboxylase (PDC, EC 4.1.1.1) overcomes the problem of by-product benzyl alcohol formation, since there is no regeneration of electron donors (e.g. NADH) in the cell-free system. Up to 28.6 g l⁻¹ PAC with a yield on initial benzaldehyde of 95.3% has been reported for the cell-free process using partially purified *Candida utilis* PDC (SHIN & ROGERS, 1996). In addition, a possible disadvantage of using whole cells as biotransformation catalysts was putative mass transfer

* To whom correspondence should be addressed.

Phone: +66-87-6985528; fax: +66-53-94-8206; e-mail: noppol@hotmail.com

limitation of substrates and products across the cell envelope (SATIANEGARA, 2006; SAMBERG et al., 2010). Silver nanoparticles are applied with various technologies and its application for antimicrobial coatings is increasingly common (SAMBERG et al., 2010). Silver ions and silver salts have been used for decades as antimicrobial agents in various fields because of the growth inhibiting property against microorganisms. Silver ions may bind non-specifically to cell surfaces, causing some disruption to the cellular membrane function and allowing the silver ions to penetrate the microbial structure. The electrostatic interaction between positively charged ions, such as silver (Ag^+), and the negatively charged cell wall of microorganisms leads to the built up of stress and eventual to the breakdown of cell walls (VARNER & SANFORD, 2010). The mechanisms of such deactivation support the hypothesis that silver nanoparticles are suitable for new cell disruption methods.

The objective of this research was to consider the best concentration levels of silver nanoparticles and mixing time interval for partial isolation of intracellular PDC enzyme.

1. Materials and methods

1.1. Microorganism

C. tropicalis TISTR 5350 from Thailand Institute of Scientific and Technological Research (TISTR) was cultivated in 10 ml inoculation medium for later propagation at a 100 ml scale. The cultivation was carried out at 30 °C on a rotary shaker at 200 r.p.m. for 24 h (LALUCE et al., 2009). The cell pellet was separated by centrifugation at $2822\times g$ for 15 min using a centrifuge machine. Cell pellet was recovered from 10 ml culture broth and washed twice with 10 ml distilled water. This was performed prior to resuspending in citrate buffer (pH 6.0).

1.2. Silver nanoparticles

Silver nanoparticles were obtained from the courtesy of Sensor Research Unit, Department of Chemistry, Faculty of Science, Chulalongkorn University. Colloidal suspension of silver nanoparticles was stabilized in 2% (w/v) soluble starch. The transmission electron microscopy (TEM) image of the silver synthesized is presented in Figure 1 and indicates well dispersed particles. These were spherical in shape with an average diameter of 6.03 ± 2.60 nm. The appearance of the solution was dark yellow liquid.




Fig. 1. TEM image of spherical silver nanoparticles

1.3. Estimation of silver nanoparticle concentration for cell disruption

Various concentration levels of silver nanoparticles at 0, 5, 10, 15, 20, 25, and 30 $\mu\text{g ml}^{-1}$ were evaluated for the microbial cell disruption based on the protein yield and biological activity (LEE et al., 2010). Silver nanoparticle solution was added in the cell suspension, which was recovered from 10 ml culture broth prior to resuspension in citrate buffer (pH 6.0). The mixture was vortexed at a maximum speed ($2350\times g$) for 1 min. After the sample was cooled down for 2 min in ice–water mixture. The supernatant was separated by centrifugation for subsequent enzyme activity and protein concentration analyses. The best concentration level of silver nanoparticles was selected based on PDC activity for comparison of time dependent interaction of silver nanoparticles for cell disruption.

1.4. Time dependent interaction of silver nanoparticles

The most appropriate concentration level of silver nanoparticles was suspended in cell suspension to perform the time dependent cell disruption study of silver nanoparticles. The mixture was vortexed at a maximum speed for 1 min with varying time interval of 0, 1, 2, 3, 4, 5, and 6 min. After each time interval treatment, cell debris was removed by centrifugation at $2822\times g$ for 15 min and the supernatant was collected and stored at $-20\text{ }^{\circ}\text{C}$ for enzyme assays and protein concentration estimation.

1.5. Analytical methods

PDC carboligase activity was measured as a formation of PAC in 20 min at $25\text{ }^{\circ}\text{C}$ from 80 mM benzaldehyde and 200 mM pyruvate in carboligase buffer. One unit (U) carboligase activity was defined as the amount of enzyme that produced 1 μmol PAC from pyruvate and benzaldehyde per min at pH 6.4 and $25\text{ }^{\circ}\text{C}$ in a carboligase assay as specified by ROSCHE and co-workers (2002). The detection of protein concentration was performed according to BRADFORD (1976) using bovine serum albumin for construction of a standard curve. Specific carboligase activity was determined based on protein concentrations in the sample and expressed in units of enzyme per milligram protein (U mg^{-1}).

2. Results and discussion

2.1. Estimation of silver nanoparticle concentration for cell disruption

Seven concentration levels of silver nanoparticles, namely, 0, 5, 10, 15, 20, 25, and 30 $\mu\text{g ml}^{-1}$ were evaluated for the microbial cell disruption based on the biological activity and product yield. The highest volumetric and specific PDC activities ($0.53\pm 0.05\text{ U ml}^{-1}$ and $0.17\pm 0.02\text{ U mg}^{-1}$ protein, respectively) were observed at 15 $\mu\text{g ml}^{-1}$ of silver nanoparticle concentration as indicated in Table 1. This was different statistically ($P\leq 0.05$) from the lower concentration level of 0, 5, and 10 $\mu\text{g ml}^{-1}$ at 0.00 ± 0.00 , 0.14 ± 0.01 , and $0.12\pm 0.01\text{ U mg}^{-1}$ protein, respectively. SONDI and SALOPEK-SONDI (2004) suggested that the antimicrobial activity was dependent on the concentration of silver nanoparticles, and was closely associated with the formation of pits in the cell wall. The accumulation of silver nanoparticles in the bacterial membrane caused increasing permeability, releasing desired biomolecules from within the cells. The mechanism of silver ions inhibitory effects included the degradation effect of the membrane structure through the electrostatic attraction between negatively

charged cell membranes and positively charged nanoparticles (KIM et al., 2007). PANACEK and co-workers (2009) revealed that *Candida* spp. was inhibited by silver nanoparticles at very low concentration levels. The minimum inhibitory concentration (MIC) was found at 1.69 $\mu\text{g ml}^{-1}$ for the growth of *C. parapsilosis*. However, they also suggested that silver nanoparticle inhibition was dependent on the yeast species tested. Ionic silver caused death of *C. tropicalis* immediately even at such low concentrations as 0.84 mg l^{-1} (KVITEK et al., 2011). PRASAD and co-workers (2011) stated that silver nanoparticles showed good antimicrobial activity against *Escherichia coli*, *Bacillus cereus*, and *Candida tropicalis*.

Table 1. Comparative effect of different silver nanoparticle concentration levels on PDC activity and protein concentration

Silver nanoparticle concentration levels ($\mu\text{g ml}^{-1}$)	Volumetric PDC activity (U ml^{-1})		Specific PDC activity (U mg^{-1} protein)		Protein concentration (mg ml^{-1})	
0	0.00 \pm 0.00	D	0.00 \pm 0.00	C	0.00 \pm 0.00	E
5	0.34 \pm 0.03	B	0.14 \pm 0.01	AB	2.51 \pm 0.06	D
10	0.36 \pm 0.02	B	0.12 \pm 0.01	B	3.03 \pm 0.01	C
15	0.53 \pm 0.05	A	0.17 \pm 0.02	A	3.19 \pm 0.07	B
20	0.20 \pm 0.03	C	0.05 \pm 0.01	C	3.63 \pm 0.10	A
25	0.11 \pm 0.02	C	0.03 \pm 0.01	C	3.68 \pm 0.07	A
30	0.10 \pm 0.02	C	0.03 \pm 0.01	C	3.49 \pm 0.12	B

The number with the same alphabet (A–E) indicated no significant difference ($P>0.05$) for comparison between different rows of the same columns.

The higher concentration levels (20, 25, and 30 $\mu\text{g ml}^{-1}$) of silver nanoparticles were inversely proportional to the PDC activity with the corresponding levels of 0.20 \pm 0.03, 0.11 \pm 0.02, and 0.10 \pm 0.02 U ml^{-1} . This was compared to PAC production levels of 1.78 \pm 0.31, 1.02 \pm 0.15, and 0.88 \pm 0.18 mM, respectively. This might be due to the high reactivity of silver ions, which readily bounded to electron donor groups. The prime targets were thiol groups (-SH), which are commonly found in enzymes within the microbe (MARIA et al., 2010). This caused the enzyme to denature with certain degree of microbial cell disruption at the higher silver nanoparticle concentration level of 15 $\mu\text{g ml}^{-1}$.

The obtained results showed that silver nanoparticles at 15 $\mu\text{g ml}^{-1}$ proved to be more effective for *C. tropicalis* TISTR 5350 cell disruption than at other concentration levels.

2.2. Time dependent interaction of silver nanoparticles

Silver nanoparticles at 15 $\mu\text{g ml}^{-1}$ concentration were mixed into the microbial cell suspension. The highest PDC activity of 0.39 \pm 0.01 U mg protein^{-1} was observed when treating cells with silver nanoparticles for 3 min (Table 2). The protein concentration, although increased with time, was inversely proportional to PDC activity when the treatment was carried out for longer than 3 min. It is possible that the membrane of microorganism contains a number of sulphate-containing functional groups that can interact with silver ions and results in the

changes of the membrane morphology and membrane vulnerability that allowed penetration for silver ions. Sulphate-groups in residue 221 of yeast PDC, which are located in the vicinity of the active site and are involved in the activation by substrate (BABURINA et al., 1994), could also interact with silver ions. This might contribute to the partial loss of PDC activity (TOLAYMAT et al., 2010). The morphological differences between control (no treatment) and treated ($15 \mu\text{g ml}^{-1}$ silver nanoparticles for 3 min) *C. tropicalis* TISTR 5350 cells were observed under LV-SEM (Figs 2–3). The direct exposure to silver nanoparticles resulted in some distortion and collapsing of some cells as silver ions could easily interact with membrane and were able to penetrate into the cells.




Fig. 2. LV-SEM photomicrograph of *C. tropicalis* TISTR 5350 before treatment (control)




Fig. 3. LV-SEM photomicrographs of *C. tropicalis* TISTR 5350 cells after treatment with $1.5 \mu\text{g ml}^{-1}$ silver nanoparticles for 3 min

Table 2. Comparative effect of different time interval on PDC activity and protein concentration

Time mixing interval (min)	Volumetric PDC activity (U ml ⁻¹)		Specific PDC activity (U mg ⁻¹ protein)		Protein concentration (mg ml ⁻¹)	
0	0.49±0.01	C	0.27±0.01	C	1.80±0.01	BC
1	0.55±0.02	B	0.27±0.01	C	0.14±0.01	D
2	0.57±0.01	B	0.33±0.01	B	1.73±0.02	BC
3	0.66±0.01	A	0.39±0.01	A	1.70±0.03	BC
4	0.63±0.02	A	0.32±0.01	B	1.94±0.01	B
5	0.51±0.02	BC	0.31±0.01	B	1.67±0.01	C
6	0.37±0.02	D	0.17±0.01	D	2.15±0.03	A

The number with the same alphabet (A–D) indicated no significant difference ($P>0.05$) for comparison between different rows of the same columns.

3. Conclusions

The effect of silver nanoparticles to the observed PDC activity was both dose and time dependent. The maximum level of PDC activity (0.66 ± 0.01 U ml⁻¹ and 0.39 ± 0.01 U mg⁻¹ protein) was observed at $15\text{ }\mu\text{g ml}^{-1}$ silver nanoparticle concentration for 3 min. Silver nanoparticles could thus also be used as an alternative to be used to disrupt cells by increasing the permeability of the cell membrane, which contributes to the leakage of cell contents. Moreover, silver nanoparticles method could reduce time investment and renders unavailable equipment unnecessary for cell disruption method. Silver nanoparticles method is also suitable for application to commercial processes, and it is an uncomplicated process with low costs. To improve the current research, further investigation, such as elimination of silver contaminants from PDC enzyme, should be done for the preservation of activity.

*

The authors gratefully acknowledge the financial support of the National Research University (NRU), the Office of Higher Education Commission (OHEC), the Ministry of Education, Thailand, the National Research Council of Thailand (NRCT). The authors thank the Thailand Institute of Scientific and Technological Research (TISTR) for the supplying of microbial strains for this project, as well as Bioprocess Research Cluster (BRC), School of Agro-Industry, Faculty of Agro-Industry, Chiang Mai University. We would also like to thank Associate Professor SANONG EKGASIT the courtesy of Sensor Research Unit, Department of Chemistry, Faculty of Science, Chulalongkorn University, Thailand to provide the silver nanoparticles used in the current study.

References

- BABURINA, I., GAO, Y., HU, Z., JORDAN, F., HOHMANN, S. & FUREY, W. (1994): Substrate activation of brewer's yeast pyruvate decarboxylase is abolished by mutation of cysteine 221 to serine. *Biochem.*, **33**, 5630–5635.
- BRADFORD, M.M. (1976): A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.*, **72**, 248–254.
- GERDEA, J.A., MONTALBO-LOMBOYA, M., YAOA, L., GREWELL, D. & WANGA, T. (2012): Evaluation of microalgae cell disruption by ultrasonic treatment. *Bioresource Technol.*, **125**, 175–181.
- KIM, J.S., KUK, E., YU, K.N., KIM, J.H., PARK, S.J., LEE, H.J., KIM, S.H., PARK, Y.K., PARK, Y.H., HWANG, C.Y., KIM, Y.K., LEE, Y.S., JEONG, D.H. & CHO, M.H. (2007): Antimicrobial effects of silver nanoparticles. *Nanomed.-Nanotechnol.*, **3**, 95–101.

- KVITEK, L., PANACEK, A., PRUCEK, R., SOUKUPOVA, J., VANICKOVA, M., KOLAR, M. & BORIL, R.Z. (2011): Antibacterial activity and toxicity of silver – nanosilver versus ionic silver. *J. Phys. Conf. Ser.*, **304**, 12–29.
- LALUCE, C., TOGNOLLI, J.O., DE OLIVEIRA, K.F., SOUZA, C.S. & MORAIS, M.R. (2009): Optimization of temperature, sugar concentration, and inoculum size to maximize ethanol production without significant decrease in yeast cell viability. *Appl. Microbiol. Biot.*, **83**, 627–637.
- LEE, J., KIM, K.J., SUNG, W.S., KIM, J.K. & LEE, D.G. (2010): The silver nanoparticle (nano-Ag): A new model for antifungal agents. Available at: <http://www.intechopen.com/books/silver-nanoparticles/the-silver-nanoparticle-nano-ag-a-new-model-for-antifungal-agents>. Last accessed 5 Oct 2010.
- MARIA, L.C.S., SANTOS, A.L.C., OLIVEIRA, P.C., VALLE, A.S.S., BARUD, H.S., MESSADDEQ, Y. & RIBEIRO, S.J.L. (2010): Preparation and antibacterial activity of silver nanoparticles impregnated in bacterial cellulose. *Polimeros*, **20**, 72–77.
- PANACEK, A., KOLÁR, M., VECEROVÁ, R., PRUCEK, R., SOUKUPOVÁ, J., KRYSOF, V., HAMAL, P., ZBORIL, R. & KVÍTEK, L. (2009): Antifungal activity of silver nanoparticles against *Candida* spp. *Biomaterials*, **30**, 6333–6340.
- PRASAD, T.N.V.K.V., ELUMALAI, E.K. & KHATEEJA, S. (2011): Evaluation of the antimicrobial efficacy of phytogenic silver nanoparticles. *Asian Pac. J. Trop. Biomed.*, **1**, S82–S85.
- ROSCHKE, B., LEKSAWASDI, N., SANDFORD, V., BREUER, M., HAUER, B. & ROGERS, P.L. (2002): Enzymatic (R)-phenylacetylcarbinol production in benzaldehyde emulsions. *Appl. Microbiol. Biot.*, **60**, 94–100.
- SAMBERG, M.E., OLDENBURG, S.J. & MONTEIRO-RIVIERE, N.A. (2010): Evaluation of silver nanoparticle toxicity in skin *in vivo* and keratinocytes *in vitro*. *Environ. Health Persp.*, **118**, 407–413.
- SATIANEGARA, G. (2006): *Comparative studies on different enzyme preparations for (R)-phenylacetylcarbinol production*. Dissertation, University of New South Wales, Sydney, Australia. 162 pages.
- SHIN, H.S. & ROGERS, P.L. (1996): Production of L-phenylacetylcarbinol (L-PAC) from benzaldehyde using partially purified pyruvate decarboxylase (PDC). *Biotechnol. Bioeng.*, **49**, 52–62.
- SONDI, I. & SALOPEK-SONDI, B. (2004): Silver nanoparticles as antimicrobial agent: A case study on *E. coli* as a model for Gram-negative bacteria. *J. Colloid Interf. Sci.*, **275**, 177–182.
- TANGTUA, J., TECHAPUN, C., PRATANAPHON, R., KUNTIYA, A., CHAIYASO, T., HANMUANGJAI, P., SEESURIYACHAN, P. & LEKSAWASDI, N. (2013): Screening of 50 microbial strains for production of ethanol and (R)-phenylacetylcarbinol. *Chiang Mai J. Sci.*, **40**, 299–304.
- TOLAYMAT, T.M., AMRO, M., EL BADAWY, A.M., ASH GENAIDY, A., SCHECKEL, K.G., LUXTON, T.P. & SUIDAN, M. (2010): An evidence-based environmental perspective of manufactured silver nanoparticle in syntheses and applications: A systematic review and critical appraisal of peer-reviewed scientific papers. *Sci. Total Environ.*, **408**, 999–1006.
- VARNER, K. & SANFORD, J. (2010): State of the science literature review: everything nanosilver and more. Available from: <http://www.epa.gov/nanoscience/files/NanoPaper1.pdf>. Last accessed 3 Aug 2010.