

COMPARING THE EFFECTS OF EXCESS COPPER IN THE LEAVES OF *BRASSICA JUNCEA* (L. CZERN) AND *BRASSICA NAPUS* (L.) SEEDLINGS: GROWTH INHIBITION, OXIDATIVE STRESS AND PHOTOSYNTHETIC DAMAGE

GÁBOR FEIGL,^{2*} DEVANAND KUMAR,¹⁺ NÓRA LEHOTAI,² ANDREA PETŐ,²
ÁRPÁD MOLNÁR,² ÉVA RÁCZ,² ATTILA ÖRDÖG,² LÁSZLÓ ERDEI,²
ZSUZSANNA KOLBERT² and GÁBOR LASKAY²

¹Department of Life Sciences and Bioinformatics, Assam University, Silchar-788 011, Assam, India

²Department of Plant Biology, University of Szeged, Szeged, Hungary

(Received: May 13, 2014; accepted: July 18, 2014)

Hydroponic experiments were conducted to compare the effects of excess copper (Cu) on growth and photosynthesis in young Indian mustard (*Brassica juncea*) and oilseed rape (*Brassica napus*). We compared the effects of excess Cu on the two *Brassica* species at different physiological levels from antioxidant levels to photosynthetic activity. Nine-day-old plants were treated with Cu (10, 25 and 50 μ M CuSO₄) for 7 and 14 days. Both species took up Cu from the external solution to a similar degree but showed slight root-to-shoot translocation. Furthermore, after seven days of treatment, excess Cu significantly decreased other microelement content, such as iron (Fe) and manganese (Mn), especially in the shoots of *B. napus*. As a consequence, the leaves of young *Brassica napus* plants showed decreased concentrations of photosynthetic pigments and more intense growth inhibition; however, accumulation of highly reactive oxygen species (hROS) were not detected. After 14 days of Cu exposure the reduction of Fe and Mn contents and shoot growth proved to be comparable in the two species. Moreover, a significant Cu-induced hROS accumulation was observed in both *Brassica* species. The diminution in pigment contents and photosynthetic efficiency were more pronounced in *B. napus* during prolonged Cu exposure. Based on all the parameters, *B. juncea* appears to be more resistant to excess Cu than *B. napus*, rendering it a species with higher potential for phytoremediation.

Keywords: *Brassica juncea* – *Brassica napus* – copper – oxidative stress – phytoremediation – photosynthesis

INTRODUCTION

Among heavy metals, copper in trace amounts is essential for plant life but it becomes toxic at higher concentrations. In excess, this heavy metal seriously inhibits leaf expansion at the early growth stage and it causes changes in physiological processes such as transpiration, photosynthetic electron transport and biosynthesis of chlorophyll [30]. Indeed, photosynthesis is the most heavy metal-sensitive process [1] since

*Corresponding author; e-mail address: fglgbr@gmail.com

+These authors contributed equally to this work.

excess Cu can, *inter alia*, affect photosynthetic electron transport on the reducing side of PSI at the level of ferredoxin. In addition, it can alter the PSII on the oxidising side by inhibiting the electron transport at P680 as well as by inactivating some PSII reaction centres [41]. A few studies indicate that excess Cu can impair the PSII electron transport on its reducing side by affecting the rate of electron transfer from tyrosine (Y_Z) to the oxidized primary donor $P680^+$ [16].

At the molecular level, excess Cu is able to induce the formation of reactive oxygen species (ROS) *via* the Fenton or Haber–Weiss reactions which subsequently damage proteins, nucleic acids and lipids [14]. This effect of excess Cu was supported by the positive correlation between Cu-treatment and the production of hydroxyl radicals in *Arabidopsis* [9]. Besides, excess Cu can indirectly cause oxidative stress by disrupting the balance between ROS generation and detoxification [34]. The activities of several ROS scavenging enzymes (such as superoxide dismutase, SOD; catalase, CAT and peroxidases, POD) are modulated by excess Cu but the effect depends on the plant species, the concentration and the duration of exposure [41].

Brassica species are economically important as food and oilseed plants. Indian mustard (*Brassica juncea*) is primary source of the anticarcinogenic 3-butenyl glucosynolate [19]. Oilseed rape (*Brassica napus*) is also known as an oilseed plant offering a high yield potential due to the high chloroplast number per unit leaf area [8]. Moreover, these species were chosen for the experiments because they possess rapid growth, high biomass, and a remarkable capacity to take up toxic metals [13, 18].

Accumulation of heavy metals in the environment and their toxic effects through the food chain can lead to serious ecological and health problems. Phytoremediation is a suitable way for clean-up the metal-rich soils; although soil pollution is always a combination of inorganic elements and/or organic compounds [32], but there are only few hyperaccumulator plants having the ability for accumulating more metals. To resolve this problem, it is needed to widen the field of application for phytoremediation and the potential of crop-related species (such as *Brassica* species) should be taken into account [31]. Therefore, it is important to determine and compare the susceptibility of these promising crop-related species for phytoremediation to Cu exposure.

For that reason, the aim of the present study was to compare the effects of excess Cu at different concentrations and treatment durations on the morphological changes of young *B. juncea* and *B. napus* leaves and the background of this response such as changes in the element content and the levels of highly reactive oxygen species, furthermore the behaviour of the antioxidant defence and photosynthetic systems.

MATERIALS AND METHODS

Plant material and growth conditions

The study was conducted on *B. juncea* L. Czern. and *B. napus* L. The seeds were surface sterilised by 5% (v/v) sodium hypochlorite for 10 min, then rinsed and imbibed for 30 minutes in running water and germinated and grown in perlite supported hydroponically for 9 days under controlled condition in a greenhouse (12/12 day/night period, 150 $\mu\text{mol}/\text{m}^2/\text{s}$ light intensity, relative humidity of 55–60% and 25 ± 2 °C). Nine-day-old *Brassica* plants were grown in full-strength Hoagland solution supplemented with 10, 25 and 50 μM CuSO_4 . As controls, untreated plants were used. Shoots were sampled for experimental analysis on the 7th and 14th days of the treatment.

Measurement of the shoot size

The plants treated with different Cu concentrations were separated carefully and washed thoroughly. Fresh and dry weights (g) of the shoots were measured on the 7th and 14th days of the treatments. Leaf area was also determined on the 7th and 14th days, on at least 10 specimens from every concentration in every case by using a grid and ImageJ (National Institute of Mental Health, Bethesda, Maryland, USA) image processing software.

Measurement of Cu concentration by ICP-MS

For determination of Cu concentrations in the plant tissues, inductively-coupled plasma mass spectrometer (ICP-MS, Thermo Scientific XSeries II, Asheville, USA) was applied. Shoots of control, 10, 25 and 50 μM Cu-treated *B. juncea* and *B. napus* plants were rinsed with distilled water. After drying at 70 °C for 72 hours, nitric acid (65%, w/v) and H_2O_2 (30%, w/v) was added to the samples, which were destructed at 200 °C and 1600 W for 15 minutes. Values of Cu concentrations are given in $\mu\text{g}/\text{g}$ dry weight (DW) and the bioaccumulation factor (BAF) was calculated as follows:

$\text{BAF} = \text{Cu concentration in plant tissues } (\mu\text{g}/\text{g}) / \text{Initial Cu concentration in the nutrient solution } (\mu\text{g}/\text{g})$.

In situ detection of highly reactive ROS and measurement of antioxidants

Highly reactive ROS, such as peroxynitrite (ONOO^-), hydroxyl radical (OH^\bullet) and hypochlorite anion (OCl^-) were visualized by 3'-(p-aminophenyl) fluorescein (APF) [17]. Leaf discs with a diameter of 7.5 mm were cut representatively from the fully developed leaves. They were incubated in 10 μM APF solution for 1 hour at

room temperature in darkness, and were washed twice with TRIS-HCl buffer (10 mM, pH 7.4). Experiments were carried out using a Zeiss Axiovert 200M inverted-fluorescence microscope (Carl Zeiss, Jena, Germany) equipped with a high resolution digital camera (Axiocam HR, HQ CCD) with filter set 10 (excitation: 450–490 nm, emission: 515–565 nm). Fluorescence intensities (pixel intensities) were measured on digital images within circular areas of 600 μm radii using Axiovision Rel. 4.8 software. The radii of circles were not modified during the experiments.

SOD (EC 1.15.1.1) activity was determined by measuring the ability of the enzyme to inhibit the photochemical reduction of nitro blue tetrazolium (NBT) in the presence of riboflavin in light [6]. The enzyme activity was expressed in terms of U/mg fresh weight; one unit (U) of SOD corresponds to the amount of enzyme causing 50% inhibition of NBT reduction in light.

CAT (EC 1.11.1.6) activity was determined by the decomposition of H_2O_2 and was measured spectrophotometrically by following the decrease in absorbance at 240 nm [45]. One U = the amount of H_2O_2 (in μmol) decomposed in 1 min. For measuring SOD and CAT the same crude extract was used. 250 mg plant material was ground with 10 mg polyvinyl polyvinylpyrrolidone (PVPP) and 1 ml 50 mM phosphate buffer (pH 7.0, with 1 mM EDTA added).

Activity of ascorbate peroxidase (APX) (EC 1.11.1.11) was measured according to a modified method by Nakano and Asada [34] by monitoring the decrease in ascorbate content at 265 nm ($E = 14 \text{ mM cm}^{-1}$). For the enzyme extract, 250 mg plant material was ground with 1.5 ml extraction buffer containing 1 mM EDTA, 50 mM NaCl and 900 μM ascorbate. Data are expressed as specific activity (Unit/g fresh weight).

Determination of ascorbate (ASA)/dehydroascorbate (DHA) contents was carried out by the method of Law et al. [23]. Plant material (250 mg) was ground in 1 ml 5% (w/v) trichloroacetic acid (TCA). The measurement is based on the reduction of Fe^{3+} to Fe^{2+} by ascorbate and then Fe^{2+} forms a complex with bipyridyl resulting in pink colour with an absorption maximum at 525 nm. The amount of total ASA was determined by the reduction of dehydroascorbate to ascorbate by dithiothreitol (DTT). ASA/DHA contents were expressed in $\mu\text{mol/g}$ fresh weight.

Measurement of pigment composition and chlorophyll fluorescence

The amount of chlorophyll a, b and total carotenoids were determined using the method described by Lichtenthaler [25]. The pigments were extracted with 80% acetone and then the extract's absorbance was measured at 663, 646 and 470 nm. The calculated amounts of the pigments are expressed as μg pigment/g fresh weight.

Chlorophyll fluorescence parameters were measured using a Pulse Amplitude-Modulated Fluorometer (Program "Run 8", PAM 200 Chlorophyll Fluorometer, Heinz Walz GmbH, Effeltrich, Germany). Leaves of treated and control plants were first dark adapted for 30 minutes and F_m , F_m' , F_t and F_o' parameters were measured in the function of increasing light intensity (PAR = Photosynthetic Active Radiation)

from 60 to 850 $\mu\text{mol photons/m}^2/\text{s}$. From these parameters the effective quantum yield of PSII [$\text{Yield} = (F_m' - F_t)/F_m'$], electron transport rate ($\text{ETR} = \text{Yield} \times \text{PAR} \times 0.5 \times 0.84$), photochemical quenching [$qP = (F_m' - F_t)/(F_m' - F_o')$] and non-photochemical quenching [$\text{NPQ} = (F_m - F_m')/F_m'$] were calculated and recorded. All measurements were carried out on leaves from five different plants in three parallel experiments.

Statistical analysis

The results are expressed as $\text{mean} \pm \text{SE}$. Multiple comparison analyses were performed with SigmaStat 12 software using analysis of variance (ANOVA, $P < 0.05$) and Duncan's test. In some cases, Microsoft Excel 2010 and Student's *t*-test were used ($*P \leq 0.05$, $**P \leq 0.01$, $***P \leq 0.001$). All experiments were carried out at least two times. In each treatment at least 10 samples were measured.

RESULTS AND DISCUSSION

Morphological changes, Cu accumulation capability and tolerance

Regarding all the three examined shoot growth parameters, a remarkable and concentration-dependent reduction was found in the *Brassica* plants. In the short-term treatments (up to 7 days) both fresh and dry weight was reduced more significantly in *B. napus*. However, by the 14th day of the treatment this difference disappeared and both species exhibited a significant reduction in the shoot growth. The highest doses of Cu treatment resulted in 80% decrease in both fresh and dry mass (Fig. 1a, b). On the contrary, the biggest reduction in leaf area was observed in *B. juncea* and this was caused by 25 and 50 μM Cu at both short- and long-term treatments (Fig. 1c, Fig. 2). According to Maksymiec [30], the serious shoot growth inhibition induced by copper is characteristic in the early developmental stage of the plants. Despite the growth inhibition, no visible symptoms of cell death (e.g. necrotic spots, lesions) were observed on the leaves during the treatment period.

Both *Brassica* species possessed similar basal Cu contents ($\sim 12\text{--}18 \mu\text{g/g DW}$) in their shoots, which is in agreement with the results published by Russo et al. [39], where the Cu content in the control leaves of *B. napus* was determined as $18 \mu\text{g/g DW}$. Both species showed similar Cu accumulation rates and the enhancement of Cu contents in the shoots depends on the metal concentration in the nutrient solution. A notable increment in Cu content was measured already in *Brassica* shoots treated with the lowest Cu concentration. However, in cases of 25 and 50 μM Cu treatments the enhancement of Cu content in the shoot tissues is lower, compared to the lowest external Cu concentration (Table 1). The root-to-shoot Cu transport can be conceived by the movement of nicotianine complexes via the xylem vessels [6].

Similarly to the results of Ebbs and Kochian [11], excess Cu induced deficiency in other microelements such as Fe and Mn in the shoots. In the short term, Fe and Mn




Fig. 1. Shoot fresh weight (in control %, a), shoot dry weight (in control %, b) and leaf area (c) of untreated and Cu-treated *B. juncea* and *B. napus* on the 7th and the 14th day. Significant differences according to Student's *t*-test ($n = 10$, * $P \leq 0.05$, ** $P \leq 0.01$, *** $P \leq 0.001$) are indicated

Table 1
Concentrations ($\mu\text{g/g DW}$) of Cu, Fe, Mn and bioaccumulation factor for Cu in the shoots of *B. juncea* and *B. napus* after 7 and 14 days of Cu treatment

		7th day			
		Control	10 $\mu\text{M Cu}$	25 $\mu\text{M Cu}$	50 $\mu\text{M Cu}$
<i>B. juncea</i>	Cu	12.93 \pm 0.66	45.35 \pm 1.52***	65.22 \pm 1.28***	162.2 \pm 4.15***
	Cu BAF	391.81 \pm 7.25	68.50 \pm 2.56***	40.83 \pm 1.97***	40.46 \pm 1.88***
	Fe	90.99 \pm 0.75	61.95 \pm 1.45***	37.04 \pm 0.42***	73.03 \pm 1.22***
	Mn	48.66 \pm 0.36	25.86 \pm 1.33***	17.79 \pm 0.20***	19.85 \pm 0.33***
<i>B. napus</i>	Cu	18.52 \pm 0.33	40.50 \pm 1.75***	66.60 \pm 1.01***	85.34 \pm 2.32***
	Cu BAF	561.21 \pm 9.58	61.17 \pm 2.33***	41.70 \pm 2.01***	21.29 \pm 0.32***
	Fe	104.20 \pm 1.03	37.02 \pm 1.50***	85.69 \pm 0.50	48.63 \pm 1.12***
	Mn	82.52 \pm 0.42	21.86 \pm 0.17***	20.08 \pm 0.13***	19.88 \pm 0.05***
		14th day			
		Control	10 $\mu\text{M Cu}$	25 $\mu\text{M Cu}$	50 $\mu\text{M Cu}$
<i>B. juncea</i>	Cu	12.41 \pm 0.42	49.79 \pm 1.33***	79.56 \pm 3.30***	88.29 \pm 2.87***
	Cu BAF	376.1 \pm 8.11	75.21 \pm 3.02***	49.81 \pm 2.18***	22.28 \pm 0.56***
	Fe	83.66 \pm 0.99	30.43 \pm 0.71***	32.73 \pm 0.88***	48.44 \pm 0.44***
	Mn	60.25 \pm 0.41	27.78 \pm 1.02***	15.40 \pm 0.14***	13.81 \pm 0.08***
<i>B. napus</i>	Cu	12.47 \pm 0.15	57.66 \pm 0.36***	74.74 \pm 4.66***	82.01 \pm 5.15***
	Cu BAF	377.87 \pm 7.98	87.09 \pm 3.35***	46.80 \pm 2.04***	20.46 \pm 0.78***
	Fe	74.26 \pm 0.75	43.03 \pm 0.84***	37.62 \pm 1.35***	39.44 \pm 0.79***
	Mn	60.01 \pm 0.16	35.36 \pm 0.10***	15.77 \pm 0.33***	15.96 \pm 0.04***

Significant differences according to Student's *t*-test ($n = 10$, *** $P \leq 0.001$) are indicated.

contents were lower in Cu-treated *B. napus* than in *B. juncea*. As the effect of 14 days of Cu exposure, the difference between Fe and Mn contents between the species disappeared. Iron depletion in the shoot derives from the competition between Fe and Cu during the uptake [24]. Intervenal chlorosis as the major symptom of iron deficiency [44] was very conspicuous and visible on the leaves of both *Brassica* species treated with Cu (Fig. 2). According to Lidon and Henriques [26], excess Cu changes the uptake rate of manganese, which can explain the Cu-induced reduction in its content.

The bioaccumulation factor (BAF) gives information about the metal accumulation potential, as well as the phytoremediation ability of plants [46]. The BAF values of the shoot system significantly decreased in both species, since the rate of Cu accumulation was lower in the shoot system than the enhancement in Cu content in the external solu-

tion (Table 1). On the contrary, BAF of the root enhanced in the function of increasing external Cu concentrations [12]. These suggest the strong ability of these plants to extract Cu from the medium and to effectively accumulate it in their root system; however, they possess a slighter root-to-shoot translocation of Cu. Similarly, Cd was shown to accumulate preferentially in the roots and part of it was translocated to the shoot of *B. juncea* [40]. In other papers, *B. napus* and *B. juncea* was shown to accumulate Cd and Zn but not Cu in the shoot [11]. Therefore, the application of these *Brassica* species in rhizofiltration purposes can be suggestible [10, 12].




Fig. 2. Representative photographs of *Brassica* shoots in case of control and 10, 25 and 50 μM Cu treatment

Oxidative damage and activation of the antioxidant defence system

Oxidative processes taking place in the leaves can be inferred by measuring the levels of highly reactive ROS (hydroxyl radical, peroxynitrite and hypochlorite radical) and the activities of antioxidants. In *B. juncea* 7-day exposure to excess Cu caused a significant accumulation of ROS, but not in *B. napus*. In the long term (14 days) both species exhibited significant ROS production which was elicited by 25 and 50 μM Cu in *B. juncea*, whereas in *B. napus* 10 μM was effective, too (Fig. 3a).

Being a transition metal, Cu in excess induces the formation of reactive oxygen species (ROS) based on the Fenton or Haber–Weiss reactions [14] and it changes the activities or transcription levels of antioxidants [20, 27, 28]. In the short term, the activity of the H_2O_2 decomposing enzyme CAT decreased by ~40% as the effect of excess Cu in *B. juncea* shoots. In contrast, a significant and concentration-dependent increase in CAT activity was observed in *B. napus*. In long term, no significant alteration of CAT activity was observed (Fig 3b). In the case of APX enzyme, the 7-day-long Cu exposure did not lead to activity changes, while in long term the activity of this antioxidant enzyme decreased in both species (Fig. 3c). Similarly, in the work of Luna et al. [29] reduced APX activity was detected in Cu-exposed oat plants, which can contribute to ROS accumulation. However, in our previous work a significant Cu-induced induction of SOD activity was observed in the root system of *Brassica* [12], in the shoot the activity of this enzyme did not show any significant




Fig. 3. Intensity of hROS-dependent fluorescence (a), activity of CAT (b), APX (c) and SOD (d) in the leaves of *B. juncea* and *B. napus*. Different letters indicate significant differences according to Duncan-test ($n = 10$, $P \leq 0.05$), n.s. = not significant




Fig. 4. Ascorbate contents in the shoot system of *B. juncea* and *B. napus*. Significant differences according to Student's *t*-test ($n = 10$, $*P \leq 0.05$, $***P \leq 0.001$) are indicated

Table 2
Ratios of ASA and DHA in the shoots of control and Cu-treated *B. juncea* and *B. napus*

	ASA/DHA ratios			
	<i>B. juncea</i> 7 th day	<i>B. napus</i> 7 th day	<i>B. juncea</i> 14 th day	<i>B. napus</i> 14 th day
Control	3.55	5.42	1.39	4.09
10 μ M Cu	1.51	2.71	1.22	1.35
25 μ M Cu	1.57	2.29	1.09	1.41
50 μ M Cu	1.58	1.76	1.13	1.32

change during the experimental period (Fig. 3d). Similarly, Singh et al. [43] observed an enhancement of SOD activities in *Brassica* shoots only caused by 50 μ M or higher Cu concentration. The amount of ascorbate and dehydroascorbate (Fig. 4) and their ratios (ASA/DHA, Table 2) were also determined in the shoot system. The concentration of the reduced ascorbate form (ASA) did not show significant changes in none of the examined species during the experimental period. On the 7th day, Cu treatments caused a concentration- and time-dependent accumulation of oxidized ascorbate (DHA) in both species; however, this effect of excess Cu proved to be slighter in the shoot system of *B. juncea*. In case of long Cu exposure, the enhancement of DHA content was significant only in *B. juncea* treated with 50 μ M Cu. Under control conditions, *B. napus* showed more reduced ascorbate pool (lower ASA/DHA ratios) in the shoot system on both sampling days than *B. juncea*. As the effect of the increasing applied Cu concentrations, the reduced ASA ratio remarkably decreased in both species after a 7-day-long exposure. In long term, the diminution of the ASA/DHA ratio was more pronounced in *B. napus*, which can be explained by the higher APX activity. Based on these observations we can conclude that highly reactive ROS (e.g. hydroxyl radical, peroxynitrite) are formed in both *Brassica* species in the long-term treatment and furthermore in *B. juncea* after seven days as well. Parallel with the accumulation of ROS, the activity of antioxidants decreased, which can contribute to the increased efficiency of toxic oxygen radicals [29].

Pigment composition and photosynthetic capacity

Copper treatment decreased both chlorophyll and carotenoid contents significantly in these two *Brassica* species, more or less in a dose-dependent manner, but in *B. juncea* the effects were less pronounced (Table 3). The rate of the decrease was greater in case of chl b compared to chl a, which resulted in the increment of chl a/b ratios especially in longer term (14 days). The enhancement of chl a/b ratios were Cu concentration-dependent in *B. juncea* leaves, while *B. napus* showed the most intense increase of it in the case of 25 μ M Cu (Table 3). The reason for copper-triggered pigment loss is that Cu in excess inhibits chlorophyll and carotenoid biosynthesis and

Table 3
Concentration of photosynthetic pigments ($\mu\text{g/g}$ FW) and the Chl a/b ratios in the leaves of control and Cu-treated *B. juncea* and *B. napus* after 7 and 14 days

		Chl a	Chl b	Chl a/b	Total chl	Carotenoids
<i>B. juncea</i> 7th day	Control	11.1907 \pm 0.0046	3.1180 \pm 0.0049	3.589	14.3081 \pm 0.0093	3.0664 \pm 0.0045
	10 μM Cu	5.8517 \pm 0.0048***	1.3959 \pm 0.0072***	4.192	7.2484 \pm 0.0119***	1.9772 \pm 0.0013***
	25 μM Cu	4.3665 \pm 0.0050	1.0617 \pm 0.0054	4.113	5.4274 \pm 0.0104	1.5175 \pm 0.0004
	50 μM Cu	3.0658 \pm 0.1221	0.0886 \pm 0.8970	34.588	3.2384 \pm 0.7748	1.3802 \pm 0.3867
<i>B. juncea</i> 14th day	Control	10.3395 \pm 0.0027	4.1937 \pm 0.0056	2.465	14.5343 \pm 0.0083	3.3623 \pm 0.0047
	10 μM Cu	5.0749 \pm 0.0029***	1.8118 \pm 0.0080***	2.801	6.8857 \pm 0.0107***	1.6174 \pm 0.0010***
	25 μM Cu	2.9784 \pm 0.0128	0.9924 \pm 0.0067	3.001	3.9753 \pm 0.0099	1.0397 \pm 0.0126
	50 μM Cu	2.4494 \pm 0.0040	0.4693 \pm 0.0093	5.219	2.9180 \pm 0.0132	1.0900 \pm 0.0030
<i>B. napus</i> 7th day	Control	10.8251 \pm 0.0028	3.1109 \pm 0.0080	3.480	13.9370 \pm 0.0107	3.2593 \pm 0.0014
	10 μM Cu	8.8388 \pm 0.0051***	2.6375 \pm 0.0047***	3.351	11.4774 \pm 0.0096***	2.7681 \pm 0.0006***
	25 μM Cu	4.3885 \pm 0.0528	1.2287 \pm 0.0175	3.572	5.6215 \pm 0.0361	1.4429 \pm 0.0084
	50 μM Cu	5.4949 \pm 0.0028	1.6088 \pm 0.0080	3.416	7.1030 \pm 0.0108	1.8564 \pm 0.0010
<i>B. napus</i> 14th day	Control	14.0376 \pm 0.0057	4.0549 \pm 0.0036	3.462	18.0916 \pm 0.0093	4.2415 \pm 0.0015
	10 μM Cu	7.4806 \pm 0.0021***	1.6289 \pm 0.0100***	4.592	9.1086 \pm 0.0121***	2.5135 \pm 0.0031***
	25 μM Cu	3.5589 \pm 0.0058	0.6393 \pm 0.0031	5.567	4.1974 \pm 0.0087	1.2139 \pm 0.0016
	50 μM Cu	7.1903 \pm 0.0029	1.8830 \pm 0.0106	3.819	9.0732 \pm 0.0135	2.3191 \pm 0.0030

Significant differences according to Student's *t*-test ($n = 10$, *** $P \leq 0.001$) are indicated.




Fig. 5. Chlorophyll fluorescence parameters (Yield, ETR, qP, NPQ) of *B. juncea* (a) and *B. napus* (b) leaves after a 14-day-long Cu exposure

retards the incorporation of these pigments in photosystems [4, 5, 27]. In addition, the substitution of the central magnesium ion in the porphyrin ring of chlorophyll by excess Cu may also damage the chlorophyll synthesizing system [21]. Furthermore, the synthesis of chlorophyll requires iron, thus the Cu-induced Fe deficiency (see Table 1) may contribute to the diminution of chlorophyll concentration and the increment of chl a/b ratio [36].

Chlorophyll fluorescence parameters are recognized as powerful tools to study physiological responses of plants to metal-induced stress and offer a reliable method for assessing photosynthetic activity [22, 37]. Exposure to excess Cu induced inhibition of photosynthesis in both species after 14 days, but the pattern of inhibition was different. Interestingly, the Yield, ETR and qP parameters of *B. juncea* were not affected by 25 μ M Cu treatment significantly, but they were equally inhibited by 10 and 50 μ M Cu (Fig. 5a). On the other hand, in *B. napus* clear dose-response relationships were seen in all these three parameters (Yield, ETR, qP; Fig. 5b), indicating that excess Cu is an effective blocker of PSII function. Indeed, through binding to both donor and acceptor sides of PSII, excess Cu inhibits electron transfer processes of PSII *in vitro* [15, 43]. Also, PSII showed greater sensitivity to Cu excess because of degradation and leakage of chloroplasts membranes, inducing a decrease in yield of O₂ evolution [35]. Moreover, Cu in excess is more efficient at inactivating the ferredoxin-dependent reactions of the photosynthetic electron transport (such as NADP⁺ photoreduction) than the light reactions in isolated chloroplasts of spinach [42]. No significant alteration in NPQ was seen in either species (Fig. 5a, b), indicating that Cu treatment did not increase the probability of dissipating the excess excitation energy via this alternative route. Based on these, the photosynthesis of *B. juncea* was more sensitive to low applied Cu concentration, while *B. napus* showed susceptibility to more serious excess Cu (25 and 50 μ M Cu). Considering the remarkable ROS accumulation detected in the leaves we can assume that chloroplast membranes may suffer Cu-induced peroxidation, which can partially explain the diminution of photosynthetic efficiency [3]. Furthermore, free radicals can damage the photosynthetic apparatus [2] and may also catalyse protein degradation through oxidative modification and increased proteolytic activity [38], which may also lead to the decrease in photosynthesis. Another possible model of Cu effects on photosynthesis was suggested by Pätsikkä et al. [36], where Cu-induced Fe deficiency leads to the reduction of Chl content in the leaves, resulting increased sensitivity of PSII to photoinhibition.

CONCLUSIONS

Copper in excess became toxic and notably inhibited the growth of young *Brassica* shoots. This hindrance was more pronounced in *B. napus* in short treatment period. Excess Cu reduced the iron and manganese contents of *B. napus* shoots to a larger extent than those of *B. juncea* to levels associated with Fe and Mn deficiencies, which definitely contributed to growth inhibition. Highly reactive ROS (e.g. hydroxyl radical, peroxynitrite) were formed in both species with a parallel decrease in the activity

of antioxidants, which is indicative of oxidative stress induced by Cu excess. Summarizing, the extent of Cu-induced oxidative stress was greater in *B. napus* compared to *B. juncea*. In addition, decreases in the amounts of photosynthetic pigments along with a decrease in photosynthetic activity were more pronounced in *B. napus*. The decrease in pigment concentrations (or increase in their degradation), the consequent negative effect on photosynthetic electron transport and the oxidative damage to photosynthetic membranes could lead to a decrease in the photosynthetic activity of young *Brassica* leaves during Cu toxicity. Taken the parameters together, the growth and the photosynthesis of *B. juncea* plants were comparatively less affected by excess Cu compared to *B. napus*, indicating the species-specific nature of Cu tolerance in the early growth stage.

ACKNOWLEDGEMENTS

The experiments were conducted at the Department of Plant Biology, Faculty of Science and Informatics, University of Szeged, Szeged, Hungary, during a partial PhD programme for D. K. (C2 type Hungarian Balassi Fellowship). This work was financially supported by the Hungarian Scientific Research Fund (grant no. OTKA PD100504) and Hungary–Serbia IPA Cross-border Co-operation Program (PLANTTRAIN, HUSBR/1203/221/173).

REFERENCES

1. Ahmad, M. S. A., Ashraf, M., Tabassam, Q., Hussain, M., Firdous H. (2011) Lead (Pb)- induced regulation of growth, photosynthesis and mineral nutrition in maize (*Zea mays* L.) plants at early growth stages. *Biol. Trace Elem. Res.* 144, 1229–1239.
2. Arshad, M., Murtaza, G., Asada, K. (2006) Production and scavenging of reactive oxygen species in chloroplasts and their functions. *Plant Physiol.* 141, 391–396.
3. Babu, T. S., Marder, J. B., Tripuranthakam, S., Dixon, D. G., Greenberg, B. M. (2001) Synergistic effects of a photooxidized polycyclic aromatic hydrocarbon and copper on photosynthesis and plant growth: evidence that *in vivo* formation of reactive oxygen species is a mechanism of copper toxicity. *Environ. Toxicol. Chem.* 20, 1351–1358.
4. Boswell, C., Sharma, N. C., Sahi, S. V. (2002) Copper tolerance and accumulation potential of *Chlamidomonas reinhardtii*. *Bull. Environ. Contam. Toxicol.* 69, 546–553.
5. Böddi, B., Oravecz, A. R., Lehocski, É. (1995) Effect of cadmium on organization and photoreduction of protochlorophyllide in dark-grown leaves and etioplast inner membrane preparations of wheat. *Photosynth* 31, 411–420.
6. Burkhead, J. L., Reynolds, K. A. G., Abdel-Ghany, S. E. (2009) Copper homeostasis. *New Phytol.* 182, 799–816.
7. Dhindsa, R. S., Plumb-Dhindsa, P., Thorpe, T. A. (1981) Leaf senescence: correlated with increased levels of membrane permeability and lipid peroxidation, and decreased levels of superoxide dismutase and catalase. *J. Exp. Bot.* 32, 93–101.
8. Dixon, G. R. (2007) Vegetables *Brassicaceae* and related *Cruciferae*. *Crop production science in horticulture series:14*. Bibbles Ltd, King's Lynn.
9. Drązkiewicz, M., Skórzyńska-Polit, E., Krupa, Z. (2004) Copper-induced oxidative stress and antioxidant defence in *Arabidopsis thaliana*. *Biometals* 17, 379–387.
10. Dushenkov, V., Nanda Kumar, P. B. A., Motto, H., Raskin, I. (1995) Rhizofiltration: The use of plants to remove heavy metals from aqueous streams. *Environ. Sci. Technol.* 29, 1239–1245.

11. Ebbs, S. D., Kochian, L. V. (1997) Toxicity of zinc and copper to *Brassica* species: implications for phytoremediation. *J. Environ. Qual.* 26, 776–781.
12. Feigl, G., Kumar, D., Lehotai, N., Tugyi, N., Molnár, A., Ördög, A., Szepesi, A., Gémes, K., Laskay, G., Erdei, L., Kolbert, Z. (2013) Physiological and morphological responses of the root system of Indian mustard (*Brassica juncea* L. Czern.) and rapeseed (*Brassica napus* L.) to copper stress. *Ecotox. Environ. Safety* 94, 179–189.
13. Fellet, G., Marchiol, L., Zerbi, G. (2013) Potential for metal phytoextraction of *Brassica* oilseed species. In: Naser A. Anjum et al. (eds) *Phytotechnologies: Remediation of Environmental Contaminants*. CRC Press, Taylor & Francis Group, Boca Raton, pp. 180–201.
14. Halliwell, B., Gutteridge, J. M. C. (1984) Oxygen toxicity, oxygen radical, transition metals and disease. *Biochem. J.* 219, 1–14.
15. Jegerschöld, C., Arellano, J. B., Schröder, W. P., van Kan, P. J., Barón, M., Styring, S. (1995) Copper(II) inhibition of electron transfer through photosystem II studied by EPR spectroscopy. *Biochem. J.* 312, 12747–12754.
16. Jegerschöld, C., MacMillan, F., Lubitz, W., Rutherford, A. W. (1999) Effects of copper and zinc ions on photosystem II studied by EPR spectroscopy. *Biochem. J.* 342, 12439–12445.
17. Kolbert, Zs., Pető, A., Lehotai, N., Feigl, G., Ördög, A., Erdei, L. (2012) *In vivo* and *in vitro* studies on fluorophore-specificity. *Acta Biol. Szeged.* 56, 37–41.
18. Kumar, P. B. A. N., Dushenkov, V., Motto, H., Raskin, I. (1995) Phytoextraction: the use of plants to remove heavy metals from soils. *Environ. Sci. Tech.* 29, 1232–1238.
19. Kumar, S., Andy, A. (2012) Health promoting bioactive phytochemicals from *Brassica*. *Int. Food Res. J.* 19, 59–66.
20. Kurepa, J., Hérouart, D., van Montagu, M., Inzé, D. (1997) Differential expression of CuZn- and Fe-superoxide dismutase genes of tobacco during development, oxidative stress, and hormonal treatments. *Plant Cell Physiol.* 38, 463–470.
21. Küpper, H., Setlik, I., Spiller, M., Küpper, F., Prasil, O. (2002) Heavy metal-induced inhibition of photosynthesis: targets of *in vivo* heavy metal chlorophyll formation. *J. Phycol.* 38, 429–441.
22. Lahive, E., O'Halloran, J., Jansen, M. A. K. (2012) Frond development gradients are a determinant of the impact of zinc on photosynthesis in three species of *Lemnaceae*. *Aquat. Bot.* 101, 55–63.
23. Law, M. Y., Charles, S. A., Halliwell, B. (1983) Glutathione and ascorbic acid in spinach (*Spinacia oleracea*) chloroplasts. *Biochem. J.* 210, 899–903.
24. Lequeux, H., Hermans, C., Lutts, S., Verbruggen, N. (2010) Response to copper excess in *Arabidopsis thaliana*: Impact on the root system architecture, hormone distribution, lignin accumulation and mineral profile. *Plant Physiol. Biochem.* 48, 673–682.
25. Lichtenthaler, H. K. (1987) Chlorophylls and carotenoids: pigments of photosynthetic biomembranes. *Meth. Enzymol.* 148, 350–382.
26. Lidon, F. C., Henriques, F. S. (1993) Effects of copper toxicity on growth and the uptake and translocation of metals in rice plants. *J. Plant Nutr.* 16, 1449–1464.
27. Liu, J., Xiong, Z. T., Li, T. Y., Huang H. (2004) Bioaccumulation and ecophysiological responses to copper stress in two populations of *Rumex dentatus* L. from Cu contaminated and non-contaminated sites. *Environ. Exp. Bot.* 52, 43–51.
28. Lombardi, L., Sebastiani, L. (2005) Copper toxicity in *Prunus cerasifera*: growth and antioxidant enzymes responses of *in vitro*-grown plants. *Plant Sci.* 168, 797–802.
29. Luna, C. M., González, C. A., Trippi, V. S. (1994) Oxidative damage caused by excess of copper in oat leaves. *Plant Cell Physiol.* 35, 11–15.
30. Maksymiec, W. (1997) Effect of copper on cellular processes in higher plants. *Photosynth* 34, 321–342.
31. Marchiol, L., Sacco, P., Assolari, S., Zerbi, G. (2004) Reclamation of polluted soil: Phytoremediation potential of crop-related *Brassica* species. *Water, Air, and Soil Pollution* 158, 345–356.
32. Mattina, M. J. I., Lannucci-Berger, W., Musante, C., White, J. C. (2003) Concurrent plant uptake of heavy metals and persistent organic pollutants from soil. *Environ. Pollut.* 124, 375–378.

33. Møller, I. M., Jensen, P. E., Hansson, A. (2007) Oxidative modifications to cellular components in plants. *Annu. Rev. Plant Biol.* 58, 459–481.
34. Nakano, Y., Asada, K. (1981) Hydrogen peroxide is scavenged by ascorbate specific peroxidase in spinach chloroplasts. *Plant Cell Physiol.* 22, 867–880.
35. Ouzounidou, G., Lannoye, R., Karataglis, S. (1993) Photoacoustic measurements of photosynthetic activities in intact leaves under copper stress. *Plant Sci.* 89, 221–226.
36. Pätsikkä, E., Kairavuo, M., Šeršen, F., Tyystjärvi, E. A. E. (2002) Excess copper predisposes photosystem II to photoinhibition *in vivo* by outcompeting iron and causing decrease in leaf chlorophyll. *Plant Physiol.* 129, 1359–1367.
37. Roháček, K., Soukupová, J., Barták, M. (2008) Chlorophyll fluorescence: A wonderful tool to study plant physiology and plant stress. In: Benoît Schoefs (ed.) *Plant Cell Compartments – Selected Topics*. Research Signpost, Fort P.O., Trivandrum-695 023, Kerala, pp. 41–104.
38. Romero-Puertas, M. C., Palma, J. M., Gómez, M., Del Río, L. A., Sandalio, L. M. (2002) Cadmium causes the oxidative modification of proteins in pea plants. *Plant Cell Environ.* 25, 677–686.
39. Russo, M., Sgherri, C., Izzo, R., Navari-Izzo, F. (2008) *Brassica napus* subjected to copper excess: Phospholipases C and D and glutathione system in signalling. *Environ. Exp. Bot.* 62, 238–246.
40. Salt, D. E., Prince, R. C., Pickering, I. J., Raskin, I. (1995) Mechanisms of cadmium mobility and accumulation in Indian mustard. *Plant Physiol.* 109, 1427–1433.
41. Sharma, S. S., Dietz, K. J. (2008) The relationship between metal toxicity and cellular redox imbalance. *Trends Plant Sci.* 14, 43–50.
42. Shioi, Y., Tamai, H., Sasa, T. (1978) Effects of copper on photosynthetic electron transport systems in spinach chloroplasts. *Plant Cell Physiol.* 19, 203–209.
43. Singh, S., Singh, S., Ramachandran, V., Eapen, S. (2010) Copper tolerance and response of antioxidative enzymes in axenically grown *Brassica juncea* (L.) plants. *Ecotoxicol. Environ. Safety* 73, 1975–1981.
44. Taylor, G. J., Foy, C. D. (1985) Differential uptake and toxicity of ionic and chelated copper in *Triticum aestivum*. *Can. J. Bot.* 63, 1271–1275.
45. Upadhyaya, A., Sankhla, D., Davis, T. D., Sankhla, N., Smith, B. N. (1985) Effect of paclobutrazol on the activities of some enzymes of activated oxygen metabolism and lipid peroxidation in senescing soybean leaves. *J. Plant Physiol.* 121, 453–461.
46. Zhao, F. J., Lombi, E., McGrath, S. P. (2003) Assessing the potential for zinc and cadmium phytoremediation with the hyperaccumulator *Thlaspi caerulescens*. *Plant Soil* 249, 37–43.