

The effect of subthalamic nucleus deep brain stimulation on prospective memory functions in Parkinson's disease

Gyula Demeter (1)(2)*, Péter Pajkossy (1)(2), Ágnes Szöllősi (1), Ágnes Lukács (1), István Valálik (3), & Mihály Racsmány (1)(2)

- (1) Department of Cognitive Science, Budapest University of Technology and Economics
- (2) Frontostriatal System Research Group, Hungarian Academy of Sciences
- (3) Department of Neurosurgery, St. John's Hospital, Budapest

Abstract

The aim of the study was to assess the effect of bilateral deep brain stimulation (DBS) of subthalamic nucleus (STN) on prospective memory (PM) functions involving thirteen PD patients with DBS implantation (DBS group) and 15 PD wait-listed patients (control group). Tasks of executive functions and an event-based PM task were applied. Each task was administered twice: before and after surgery in the DBS group with the stimulators on and with a similar time interval between the two task-administration points in the control group. The DBS group showed a significant decline on semantic verbal fluency task and a decreased execution cost score of the PM task after surgery, suggesting possible effects of STN DBS on PM-related frontostriatal networks.

This research was supported by the research grant of the Hungarian Brain Research Program (KTIA NAP Grant ID: 13-2-2014-0020) awarded to Mihály Racsmány as PI.

* Gyula Demeter is a grantee of the János Bolyai Research Scholarship of the Hungarian Academy of Sciences.