

Making a C_6 -free graph C_4 -free and bipartite

Ervin Győri

Scott Kensell

Casey Tompkins

June 3, 2014

Abstract

We show that every C_6 -free graph G has a C_4 -free, bipartite subgraph with at least $3e(G)/8$ edges. Our proof is probabilistic and uses a theorem of Füredi, Naor and Verstraëte on C_6 -free graphs.

1 Introduction

For G a graph, let $e(G)$ denote the number of edges in G . We say G is H -free if it does not contain H as a subgraph. For a family of graphs \mathcal{F} , let $\text{ex}(n, \mathcal{F})$ denote the maximum number of edges an n -vertex graph G can have such that G is F -free for all $F \in \mathcal{F}$.

Győri [2] proved that every bipartite, C_6 -free graph contains a C_4 -free subgraph with at least half as many edges. Extending this result, Kühn and Osthus [3] showed that every bipartite, C_{2k} -free graph has a C_4 -free subgraph with at least $1/(k-1)$ of the original edges. In an extensive study of the Turán number $\text{ex}(n, C_6)$, Füredi, Naor and Verstraëte [1] gave another generalization of Győri's result by showing (Theorem 3.1) that a C_6 -free graph has a triangle-free, C_4 -free subgraph with at least half as many edges.

Using any of these results combined with the well-known fact that every graph has a bipartite subgraph with at least half as many edges, it is easy to show that any C_6 -free graph has a bipartite, C_4 -free subgraph with at least $1/4$ the original edges. Improving the constant $1/4$ is the main focus of this paper.

In general if we would like to make a C_6 -free graph C_4 -free and bipartite, we cannot hope to keep more than $2/5$ of its edges (consider many disjoint K_5 's). We show that if c is the maximum constant such that every C_6 -free graph G has a C_4 -free subgraph on $c \cdot e(G)$ edges then $3/8 \leq c \leq 2/5$.


Theorem 1. *Let G be a C_6 -free graph, then G contains a subgraph with at least $3e(G)/8$ edges which is both C_4 -free and bipartite.*

The result can also be phrased in the language of Turán theory: If \mathcal{C} denotes the set of all odd cycles, then $\text{ex}(n, C_6) \leq 8 \text{ex}(n, C_4, C_6, \mathcal{C})/3$.


Our proof is a probabilistic deletion procedure consisting of several steps. First we two-color the vertices, and then, focusing on specific edge-disjoint subgraphs, we delete certain edges given the outcome of the coloring. These edge-disjoint subgraphs are the maximal subgraphs obtained by pasting together edge-intersecting C_4 's and were characterized by Füredi, Naor and Verstraëte. We use the following slightly weaker formulation of their theorem.

Theorem 2. For a C_6 -free graph G , let H denote the graph whose vertex set is the collection of C_4 's in G and whose edge set represents edge-intersection. Each connected component of H corresponds to an induced subgraph of G of one of the following types:


- (0) the complete bipartite graph $K_{2,m}$ for some $m > 0$,
- (1) a triangle xyz with α additional vertices adjacent to x and y , and β more vertices adjacent to x and z ,
- (2) a K_4 with $\gamma \geq 0$ paths of length 2 (outside the K_4) between two of the vertices,
- (3) a K_5 , K_5 minus an edge, or a K_5 minus two non-adjacent edges.




Type 0



Type 1



Type 2



Type 3

2 Proof of Theorem 1

Independently at random, color all vertices in G red or blue with probability $1/2$ each. Deleting all monochromatic edges would yield a bipartite graph, but some C_4 's may remain. Thus, given the random coloring we will deterministically delete additional edges in such a way that, upon deletion of monochromatic edges, at least $3e(G)/8$ edges remain in expectation, but all C_4 's are deleted. Notice that after coloring, the C_4 's which require further edge deletion are exactly the properly colored C_4 's (those with no monochromatic edges).

For each component H of type 0, 1, 2, or 3 from Theorem 2 we will show that our vertex-coloring and subsequent edge-deletion procedure preserves at least $3e(H)/8$ edges in expectation. Since these components are edge-disjoint and cover all C_4 's, we are then done by linearity of expectation.

Case(H is of type 0): First, suppose H is a component of type 0. That is, H is a complete bipartite graph $K_{2,t}$. Let x and y be the vertices in the first class, and v_1, v_2, \dots, v_t be the vertices

in the second class. If x and y are opposite colors, then there are no properly colored C_4 's, and the expected number of remaining edges is exactly $e(H)/2$.

Now suppose that x and y are the same color, say red. If none of the v_i 's are colored blue then we lose all edges in H . If exactly s , $s \geq 1$, of the v_i 's are colored blue, then we must delete all but one of the edges emanating from x to the v_i 's for otherwise we would have a properly colored C_4 . Thus, exactly $s + 1$ edges will remain in H . The probability that s of the v_i are blue is $\binom{t}{s}/2^t$. Let N_0 be the random variable equal to the number of edges which remain in H , then

$$\begin{aligned} \mathbb{E}(N_0 \mid x \text{ and } y \text{ same color}) &= \frac{1}{2^t} 0 + \sum_{s=1}^t \frac{\binom{t}{s}}{2^t} (s + 1) \\ &= \frac{1}{2^t} \sum_{s=1}^t \binom{t}{s} s + \frac{1}{2^t} \sum_{s=1}^t \binom{t}{s} \\ &= \frac{1}{2^t} t 2^{t-1} + \frac{1}{2^t} (2^t - 1) \\ &\geq \frac{t}{2} + \frac{1}{2}. \end{aligned}$$

It follows that,

$$\begin{aligned} \mathbb{E}(N_0) &= \frac{1}{2} \mathbb{E}(N_0 \mid x \text{ and } y \text{ opposite color}) + \frac{1}{2} \mathbb{E}(N_0 \mid x \text{ and } y \text{ same color}) \\ &\geq \frac{e(H)}{4} + \frac{t}{4} + \frac{1}{4} \\ &= \frac{3e(H)}{8} + \frac{1}{4}. \end{aligned}$$

Case(H is of type 1): Now, assume that H is of type 1. Let x, y, z be as in the figure. Assume that there are α vertices adjacent to x and y (excluding z), and β vertices adjacent to x and z (excluding y). Notice that $2\alpha + 2\beta = e(H) - 3$.

First suppose x, y and z are the same color. This subcase occurs with probability $1/4$. The edges $\{x, y\}, \{x, z\}$ and $\{y, z\}$ are all monochromatic, so all properly colored C_4 's are contained in one of two bipartite graphs, a $K_{2,\alpha}$ or a $K_{2,\beta}$. By the reasoning in the previous case we can preserve,

$$\frac{\alpha}{2} + \frac{1}{2} + \frac{\beta}{2} + \frac{1}{2} = \frac{e(H)}{4} + \frac{3}{4}$$

edges in expectation.

Now, suppose x is one color and both y and z are the opposite color. This subcase also occurs with probability $1/4$. We have that exactly two of the edges in the triangle formed by x, y and z are preserved as are half of the remaining edges. Thus, in total we save $(e(H) - 3)/2 + 2 = e(H)/2 + 1/2$ edges in expectation.

Next, assume that x and y are one color and z is the opposite color. This again happens with probability $1/4$. In this subcase we must also consider C_4 's through x, y, z and one of the α vertices other than z adjacent to x and y . To this end, we immediately delete the edge $\{y, z\}$. Now, only one edge remains on the triangle through x, y and z which is not monochromatic. Each of the β vertices is on one monochromatic edge and one properly colored edge. The vertices x, y and their α common neighbors again form a $K_{2,\alpha}$ which we handle as before, saving at least $\alpha/2 + 1/2$ edges

in expectation. It follows that the expected total number of edges preserved in this subcase is $\alpha/2 + \beta + 3/2$.

The final subcase in which x and z are the same color y is the opposite color is totally symmetric. In this case the expected number of preserved edges is thus $\beta/2 + \alpha + 3/2$.

Let N_1 be the random variable equal to the number of edges conserved in H , then

$$\begin{aligned}\mathbb{E}(N_1) &= \frac{1}{4}\left(\frac{e(H)}{4} + \frac{3}{4}\right) + \frac{1}{4}\left(\frac{e(H)}{2} + \frac{1}{2}\right) + \frac{1}{4}\left(\frac{\alpha}{2} + \beta + \frac{3}{2}\right) + \frac{1}{4}\left(\frac{\beta}{2} + \alpha + \frac{3}{2}\right) \\ &= \frac{3}{16}e(H) + \frac{3}{8}(\alpha + \beta) + \frac{15}{16} \\ &= \frac{3}{16}e(H) + \frac{3}{16}(e(H) - 3) + \frac{15}{16} \\ &> \frac{3}{8}e(H).\end{aligned}$$

Case(H is of type 2): We will condition first on whether a and b are the same color or opposite and then on whether x and y are the same color or opposite.

Suppose first that a and b are opposite colors. Then all C_4 's lie in the subgraph induced by a, b, x and y . If x and y are the same color, no further edges need to be deleted. If x and y are opposite colors we must delete one additional edge. In either situation exactly $e(H)/2$ edges are preserved.

Now, assume that a and b are the same color, say red. Consider the subcase when x and y are also red, then all properly colored C_4 's must lie in a $K_{2,\gamma}$. By the reasoning we have used before, this implies that we can keep,

$$\frac{e(H) - 6}{4} + \frac{1}{2} = \frac{e(H)}{4} - 1,$$

edges in expectation.

If x and y are opposite colors, then 3 of the 6 edges in the K_4 defined by a, b, x and y remain. For each of the γ vertices which are blue we must delete an edge. Thus, we retain,

$$3 + \frac{e(H) - 6}{4} = \frac{e(H)}{4} + \frac{3}{2},$$

edges in expectation.

Finally, if x and y are both blue, then delete the edge $\{a, x\}$. By the same reasoning as the preceding subcase we retain,

$$3 + \frac{e(H) - 6}{4} = \frac{e(H)}{4} + \frac{3}{2},$$

edges in expectation. Letting N_2 be the random variable counting the number of preserved edges we have,

$$\begin{aligned}\mathbb{E}(N_2) &= \frac{1}{2}\frac{e(H)}{2} + \frac{1}{8}\left(\frac{e(H)}{4} - 1\right) + \frac{1}{4}\left(\frac{e(H)}{4} + \frac{3}{2}\right) + \frac{1}{8}\left(\frac{e(H)}{4} + \frac{3}{2}\right) \\ &= \frac{3}{8}e(H) + \frac{7}{16} \\ &\geq \frac{3}{8}e(H).\end{aligned}$$

Case(H is of type 3): H is either a K_5 , a K_5 minus an edge or a K_5 minus two nonadjacent edges. First, suppose H is a K_5 . There are three possibilities: all 5 vertices are the same color, there is a unique vertex of one color or there are two vertices of one color. These possibilities have probabilities $2/32, 10/32$ and $20/32$ respectively. In the first case we have 0 remaining edges and in the second we have 4. In the third we must delete 2 additional edges, again leaving a total of 4. Thus, if N_3 counts the expected number of edges remaining, we have

$$\mathbb{E}(N_3) = \frac{2}{32}0 + \frac{10}{32}4 + \frac{20}{32}4 = \frac{3}{8}e(H)$$

The analysis of K_5 minus one or two edges is similar.

References

- [1] Zoltan Füredi, Assaf Naor, and Jacques Verstraëte. On the Turán number for the hexagon. *Adv. Math.*, 203(2):476–496, 2006.
- [2] Ervin Györi. C_6 -free bipartite graphs and product representation of squares. *Discrete Math.*, 165/166:371–375, 1997. Graphs and combinatorics (Marseille, 1995).
- [3] Daniela Kühn and Deryk Osthus. Four-cycles in graphs without a given even cycle. *J. Graph Theory*, 48(2):147–156, 2005.