

ON THE EXTREMAL THEORY OF CONTINUED FRACTIONS

ALINA BAZAROVA¹, ISTVÁN BERKES², AND LAJOS HORVÁTH³

ABSTRACT. Letting $x = [a_1(x), a_2(x), \dots]$ denote the continued fraction expansion of an irrational number $x \in (0, 1)$, Khinchin proved that $S_n(x) = \sum_{k=1}^n a_k(x) \sim \frac{1}{\log 2} n \log n$ in measure, but not for almost every x . Diamond and Vaaler showed that removing the largest term from $S_n(x)$, the previous asymptotics will hold almost everywhere, showing the crucial influence of the extreme terms of $S_n(x)$ on the sum. In this paper we determine, for $d_n \rightarrow \infty$, $d_n/n \rightarrow 0$, the precise asymptotics of the sum of the d_n largest terms of $S_n(x)$ and show that the sum of the remaining terms has an asymptotically Gaussian distribution.

1. Introduction

For an irrational number $x \in (0, 1)$ let

$$x = \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{\ddots}}}$$

be the continued fraction expansion of x . Clearly

$$a_1(x) = [1/x], \quad a_{n+1}(x) = a_1(T^n x), \quad n \geq 1,$$

where the transformation $T : (0, 1) \rightarrow [0, 1)$ is defined by $Tx = \{1/x\}$; here $[\cdot]$ and $\{\cdot\}$ denote integral resp. fractional part. Let

$$\mu(E) = \frac{1}{\log 2} \int_E \frac{1}{1+x} dx$$

be the Gauss measure on the class \mathcal{B} of Borel subsets of $(0, 1)$. It is known (see e.g. [3]) that T is an ergodic transformation preserving the Gauss measure and thus with respect to the probability

2010 *Mathematics Subject Classification*. Primary 11K50, 60F05, 60G70.

Key words and phrases. Continued fraction expansion, extreme elements, mixing random variables, central limit theorem.

¹Graz University of Technology, Institute of Statistics, Kopernikusgasse 24, 8010 Graz, Austria. e-mail: bazarova@tugraz.at. Research supported by Austrian Science Fund (FWF) Grant W1230.

²Graz University of Technology, Institute of Statistics, Kopernikusgasse 24, 8010 Graz, Austria. e-mail: berkes@tugraz.at. Research supported by Austrian Science Fund (FWF) Grant P2304-N18 and OTKA grant K 108615.

³University of Utah, Department of Mathematics, Salt Lake City, UT 84112-0090, USA. e-mail: horvath@math.utah.edu. Research supported by NSF grant DMS-13-05858.

space $((0, 1), \mathcal{B}, \mu)$, $\{a_n(x), n \geq 1\}$ is a stationary ergodic sequence. Clearly, the set $\{a_1 = k\}$ is the interval $(1/(k+1), 1/k]$ and thus

$$\mu\{a_1 = k\} = \frac{1}{\log 2} \int_{1/(k+1)}^{1/k} \frac{1}{1+x} dx = \frac{1}{\log 2} \log \left\{ 1 + \frac{1}{k(k+2)} \right\} \sim \frac{1}{\log 2} \frac{1}{k^2}.$$

(We say that $a_k \sim b_k$ if $\lim_{k \rightarrow \infty} a_k/b_k = 1$.) Thus by the ergodic theorem we have for any function $F: \mathbb{N} \rightarrow \mathbb{R}$

$$(1.1) \quad \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{k=1}^N F(a_k(x)) = \frac{1}{\log 2} \sum_{j=1}^{\infty} F(j) \log \left\{ 1 + \frac{1}{j(j+2)} \right\} \quad \text{a.e.}$$

provided that the series on the right hand side converges absolutely.

The sequence $\{a_k(x), k \geq 1\}$ has remarkable mixing properties. Gauss noted that the distribution of a_k with respect to the uniform measure in $(0, 1)$ converges to μ and asked for the speed of convergence. Kusmin [19] showed that the convergence speed is $O(e^{-\lambda\sqrt{k}})$ and Lévy [21] improved this to $O(e^{-\lambda k})$. Lévy's result implies that the sequence $\{a_k(x), k \geq 1\}$ is ψ -mixing with exponential rate, i.e. for all $A \in \mathcal{F}_1^k, B \in \mathcal{F}_{k+n}^\infty$ we have

$$|\mu(A \cap B) - \mu(A)\mu(B)| \leq C e^{-\lambda n} \mu(A)\mu(B)$$

with positive absolute constants C, λ , where \mathcal{F}_r^s denotes the σ -field generated by the variables $\{a_k(x), r \leq k \leq s\}$.

Letting \mathbf{E} denote expectation with respect to μ , we have $\mathbf{E}a_1 = \infty$ and correspondingly for $F(x) = x$ the right hand side of (1.1) is $+\infty$. Thus the partial sums $\sum_{k=1}^N a_k(x)$ grow faster than N . Lévy [22] proved that

$$(1.2) \quad \frac{1}{N} \sum_{k=1}^N a_k(x) - \frac{\log N}{\log 2} \xrightarrow{d} G,$$

where \xrightarrow{d} means convergence in distribution in the probability space $((0, 1), \mathcal{B}, \mu)$ and G is the stable distribution with characteristic function

$$(1.3) \quad \exp \left(-it \log |t| - \frac{\pi |t|}{2 \log 2} \right).$$

Remainder term estimates for the convergence in (1.2) were obtained by Heinrich [14]. This implies that

$$(1.4) \quad \lim_{N \rightarrow \infty} \frac{1}{N \log N} \sum_{k=1}^N a_k(x) = \frac{1}{\log 2} \quad \text{in measure,}$$

a result obtained earlier by Khinchin [18]. Khinchin also noted that (1.4) cannot hold almost everywhere. Diamond and Vaaler [9] showed that the obstacle to a.e. convergence in (1.4) is the occurrence of one single large term in the sum $\sum_{k=1}^N a_k(x)$ and established an a.e. analogue of (1.4) by excluding the largest summand. They proved namely

$$(1.5) \quad \lim_{N \rightarrow \infty} \frac{1}{N \log N} S_N^{(1)}(x) = \frac{1}{\log 2} \quad \text{for almost all } x$$

where $S_N^{(d)}(x)$ denotes the sum $\sum_{k=1}^N a_k(x)$ after discarding its d largest summands. The proof shows that (1.5) remains valid if $S_N^{(1)}$ is replaced by $S_N^{(d)}$ for any fixed $d \geq 2$ and discarding more terms improves the rate of a.e. convergence in (1.5). An analogous result for the St. Petersburg game was proved by Csörgő and Simons [7]. For further analogies between continued fraction digits and the St. Petersburg game we refer to Vardi [33]. In view of these facts it is natural to ask what happens if from the sum $S_N = \sum_{k=1}^N a_k(x)$ we remove $d = d_N$ terms, where

$$(1.6) \quad d_N \rightarrow \infty, \quad d_N/N \rightarrow 0$$

so that the number of discarded terms is 'large', but is still negligible compared with N . The purpose of this paper is to answer this question. Let

$$(1.7) \quad m(t) = \frac{1}{\log 2} \sum_{1 \leq k \leq t} k \log \left(1 + \frac{1}{k(k+2)} \right), \quad t \geq 1.$$

We will prove the following result.

Theorem 1.1. *Let $d = d_N$ satisfy (1.6). Then we have*

$$(1.8) \quad \frac{S_N^{(d)} - Nm(\eta_{d,N})}{N/\sqrt{d}} \xrightarrow{d} N(0, (\log 2)^{-1})$$

where $\eta_{d,N}$ denotes the d -th largest of a_1, \dots, a_N and $N(\mu, \sigma^2)$ denotes the normal distribution with mean μ and variance σ^2 .

Theorem 1.1 reduces the asymptotic study of $S_N^{(d)}$ to that of $\eta_{d,N}$, which is a much simpler problem. We will show in (3.15) that $\eta_{d,N} \sim N/d$ in probability and since $m(t) \sim (\log 2)^{-1} \log t$ as $t \rightarrow \infty$, Theorem 1.1 can be rewritten equivalently as

$$(1.9) \quad S_N^{(d)} = Nm(\eta_{d,N}) + (N/\sqrt{d})\zeta_N = (1 + o_P(1)) \frac{1}{\log 2} N \log(N/d) + (N/\sqrt{d})\zeta_N,$$

where $\zeta_N \xrightarrow{d} N(0, 1/\log 2)$. Here and in the sequel, \xrightarrow{P} will denote convergence in probability and $o_P(1)$ a quantity converging to 0 in probability. Relation (1.9) shows that $Nm(\eta_{d,N})$ is the main term in an asymptotic expansion of $S_N^{(d)}$. As a comparison, write Lévy's limit theorem (1.2) in the form

$$(1.10) \quad S_N = \frac{1}{\log 2} N \log N + N\zeta_N^*,$$

where ζ_N^* converges in distribution to the Cauchy variable with characteristic function (1.3). In addition to the change of the order of magnitude of S_N caused by removing the d largest terms, note that the Cauchy fluctuations of S_N around $\frac{1}{\log 2} N \log N$ described by (1.10) changed to Gaussian fluctuations around $Nm(\eta_{d,N})$ in (1.9). An immediate consequence of relation (1.9) is

$$\frac{S_N^{(d)}}{N \log(N/d)} \xrightarrow{P} \frac{1}{\log 2}$$

under (1.6). If d grows slower than any power of N , i.e. $\log d / \log N \rightarrow 0$, then the last relation implies

$$\frac{1}{N \log N} S_N^{(d)} \xrightarrow{P} \frac{1}{\log 2}.$$

Thus in this case the order of magnitude of $S_N^{(d)}$ is the same as that of the complete sum S_N , i.e. the contribution of the d largest terms of S_N is still negligible compared to the whole sum. If $d \sim N^\gamma$ for some $0 < \gamma < 1$, then

$$\frac{1}{N \log N} S_N^{(d)} \xrightarrow{P} \frac{1 - \gamma}{\log 2}.$$

We thus see that the removal of a small portion of extreme elements of S_N changes the asymptotic order of magnitude of the sum, hence the role of large elements in S_N is very substantial.

In case of i.i.d. variables in the domain of attraction of a stable law with parameter $0 < \alpha < 2$, the effect of the extremal terms on the partial sums is well known. For positive variables Darling [8] showed (see also Arov and Bobrov [1]) that under some additional regularity assumptions the ratio of the sum and its largest term has a non-degenerate limit distribution if $0 < \alpha < 1$ and this holds also for $1 < \alpha < 2$ provided we center the partial sum by its mean. The case $\alpha = 1$ is critical and is not covered in [1], [8]. The sequence $\{a_k(x), k \geq 1\}$ in the continued fraction expansion corresponds to this case, except that the variables a_k are weakly dependent. Theorem 1.1 and its corollaries above show that the contribution of the d largest terms of S_N is negligible (in probability) compared with the total sum S_N if and only if $\log d / \log N \rightarrow 0$. In particular this holds for $d = 1$, i.e. in the case of the largest term. In the i.i.d. case, Csörgő, Horváth and Mason [6] also showed that removing the d largest and d smallest elements from the partial sum, where (1.6) holds, the remaining sum $S_N^{(d)}$ becomes asymptotically normal. Our Theorem 1.1 is a dependent analogue of this result for continued fractions. There is a large literature on the metric properties of continued fractions and using the exponential ψ -mixing property of the transformation T above, many classical limit theorems for partial sums of independent random variables have been extended to continued fractions. We refer to Doeblin [10], Gordin and Reznik [13], Ibragimov [15], Iosifescu [16], [17], Philipp [24], [26], Philipp and Stackelberg [27], Samur [29], [30], Stackelberg [31], Szewczak [32] and the references therein. Using the extremal theory of dependent processes, (see e.g. Leadbetter and Rootzen [20]), asymptotic properties of the (individual) extremes of $(a_1(x), \dots, a_n(x))$ can be established; limit theorems for the largest digit were obtained by Galambos [11], [12], Philipp [25]. Note that an analogue of Theorem 1.1 for different, less natural trimming of the partial quotients a_j was obtained in Philipp [26].

In Section 2, we will prove Theorem 1.1 in a probabilistic form and we will change the notation accordingly.

Theorem 1.2. *Let $\{X_j, j \geq 1\}$ be a strictly stationary sequence of positive, integer valued random variables with*

$$(1.11) \quad P(X_1 = k) \sim c_0 k^{-2} \quad \text{as } k \rightarrow \infty$$

for some constant $c_0 > 0$. Assume that $\{X_j, j \geq 1\}$ is ψ -mixing with rate $\psi(n) = Ce^{-\lambda n}$ for some $C > 0, \lambda > 0$. Let $\eta_{d,n}$ denote the d -th largest of X_1, \dots, X_n and assume that $d = d_n$ satisfies (1.6). Let $m(t) = \mathbb{E}X_1 I\{X_1 \leq t\}$ and

$$(1.12) \quad A_n = \sqrt{c_0} n / \sqrt{d}.$$

Then

$$(1.13) \quad \frac{1}{A_n} \sum_{i=1}^{[nt]} (X_i I\{X_i \leq \eta_{d,n}\} - m(\eta_{d,n})) \xrightarrow{\mathcal{D}[0,1]} W(t),$$

where W is the Wiener process.

Remark 1.1. If (X_n) is a sequence of positive random variables such that with probability one X_1, X_2, \dots are different, then the sum $\sum_{i=1}^{[nt]} X_i I\{X_i \leq \eta_{d,n}\}$ in (1.13) is obtained from $\sum_{i=1}^{[nt]} X_i$ by removing the $d-1$ largest terms and thus the conclusion of Theorem 1.2 for $t=1$ reduces to that of Theorem 1.1. However, for integer valued variables X_n , $\eta_{d,n}$ can appear in the sequence (X_1, \dots, X_n) more than once and in this case the number of terms of the sum $\sum_{i=1}^{[nt]} X_i$ exceeding $\eta_{n,d}$ can be smaller than $d-1$ and can actually be random. Thus, in a formal sense, Theorem 1.1 is not a special case of Theorem 1.2. However, using a simple perturbation argument Theorem 1.1 will be deduced from Theorem 1.2.

Let

$$U_n(t, s) = \sum_{i=1}^{[nt]} (X_i I\{X_i \leq s(n/d)\} - EX_i I\{X_i \leq s(n/d)\}) \quad (t \geq 0, s \geq 0).$$

We will derive Theorem 1.2 from the following two-dimensional limit theorem.

Theorem 1.3. *Under the assumptions of Theorem 1.2 we have*

$$(1.14) \quad \frac{1}{A_n} U_n(t, s) \longrightarrow W(t, s) \quad \text{weakly in } \mathcal{D}[0, 1] \times \mathcal{D}[1/2, 3/2],$$

where $\{W(t, s), t \geq 0, s \geq 0\}$ is a two-parameter Wiener process.

As we already noted, under (1.6) we have

$$\frac{\eta_{d,n}}{n/d} \longrightarrow 1 \quad \text{in probability.}$$

Since the limit process $W(t, s)$ in (1.14) has continuous trajectories a.s., Theorem 1.3 and Billingsley [4], p. 144-145 imply that

$$\frac{1}{A_n} U_n(t, \eta_{d,n}/(n/d)) \xrightarrow{\mathcal{D}[0,1]} W(t, 1)$$

which is exactly the functional central limit theorem (1.13).

2. Some lemmas

In the rest of the paper (X_k) denotes a sequence of random variables satisfying the conditions of Theorem 1.2 and $d = d_n$ denotes a sequence of positive integers satisfying (1.6). Moreover, c_0 denotes the constant in (1.11). Given a process $Y(s, t)$ defined on a rectangle $H = [a, b] \times [a', b']$, let $Y(H)$ denote the increment of Y over H .

Lemma 2.1. *Let $\{Y_n(t, s), n \geq 1\}$ be processes defined on a rectangle $[a, b] \times [a', b'] \subset [0, \infty)^2$ and assume that for some $\gamma > 0$*

$$(2.1) \quad \mathbb{E}|Y_n(B)|^\gamma |Y_n(C)|^\gamma \leq \mu(B)\mu(C),$$

where μ denotes area and B and C are rectangles of the form $[t_1, t_2] \times [s_1, s_2]$ having one common edge, but otherwise disjoint. Then the sequence $\{Y_n(t, s), n \geq 1\}$ is tight. If every $X_n(t, s)$ is piecewise constant in t , i.e. there exists a finite set $H_n \subset [a, b]$ such that $X_n(t, s)$ is constant in the

left closed intervals determined by the elements of $H_n \cup \{a\} \cup \{b\}$, then it suffices to verify (2.1) for rectangles $[t_1, t_2] \times [s_1, s_2]$ where $t_1, t_2 \in H_n$.

This is a special case of a general tightness condition in Bickel and Wichura [2].

Lemma 2.2. *Let X, Y be integrable random variables such that X is measurable with respect to $\sigma(X_1, \dots, X_k)$ and Y is measurable with respect to $\sigma(X_{k+n}, X_{k+n+1}, \dots)$. Then XY is also integrable and*

$$|\mathbf{E}XY - \mathbf{E}X\mathbf{E}Y| \leq \psi(n)\mathbf{E}|X|\mathbf{E}|Y|.$$

This follows from Theorem 3.10 in Bradley [5], p. 75.

Lemma 2.3. *Let \mathcal{G}_k denote the σ -field generated by X_k , let $n_1 < \dots < n_r$ be positive integers and let Y_1, \dots, Y_r be bounded r.v.'s such that Y_j is \mathcal{G}_{n_j} measurable ($j = 1, 2, \dots, r$). Then*

$$E|Y_1 \cdots Y_r| \leq C_r E|Y_1| \cdots E|Y_r|,$$

where $C_r = (1 + \psi(1))^r$.

Proof. This is immediate by induction upon observing that by the previous lemma we have for any $1 \leq j \leq r - 1$

$$E|Y_1 \cdots Y_{j+1}| \leq E|Y_1 \cdots Y_j|E|Y_{j+1}| + \psi(1)E|Y_1 \cdots Y_j|E|Y_{j+1}| = (1 + \psi(1))E|Y_1 \cdots Y_j|E|Y_{j+1}|.$$

Lemma 2.4. *For any $T \geq 3$ we have*

$$(2.2) \quad \mathbf{E}X_1 I\{X_1 \leq T\} \leq C_1 \log T.$$

Moreover, for any fixed $0 \leq s_1 < s_2$ we have

$$(2.3) \quad \mathbf{E}X_1^2 I\{s_1(n/d) < X_1 \leq s_2(n/d)\} \sim c_0(s_2 - s_1)(n/d) \quad \text{as } n \rightarrow \infty$$

and for any fixed $0 < s_1 < s_2$ and sufficiently large n

$$(2.4) \quad \mathbf{E}X_1 I\{s_1(n/d) < X_1 \leq s_2(n/d)\} \leq C_2(s_2 - s_1)/s_1.$$

Here C_1, C_2 are positive constants depending only on the sequence (X_k) .

This is immediate from (1.11).

Lemma 2.5. *Let*

$$X_{k,n}^{(s_1, s_2)} = X_k I\{s_1(n/d) < X_k \leq s_2(n/d)\} - \mathbf{E}X_k I\{s_1(n/d) < X_k \leq s_2(n/d)\}.$$

Then for any fixed $0 \leq t_1 < t_2 \leq 1$, $0 \leq s_1 < s_2 < \infty$ we have

$$(2.5) \quad \mathbf{E} \left(\sum_{k=nt_1+1}^{nt_2} X_{k,n}^{(s_1, s_2)} \right)^2 \sim c_0(n^2/d)(t_2 - t_1)(s_2 - s_1) \quad \text{as } n \rightarrow \infty$$

provided nt_1, nt_2 are integers. Moreover,

$$(2.6) \quad \mathbf{E} \left(\sum_{i=nt_1+1}^{nt_2} X_{i,n}^{(s_1, s_2)} \right) \left(\sum_{j=nt'_1+1}^{nt'_2} X_{j,n}^{(s'_1, s'_2)} \right) = o(n^2/d) \quad \text{as } n \rightarrow \infty$$

provided $0 \leq t_1 < t_2 \leq 1$, $0 \leq t'_1 < t'_2 \leq 1$, $0 \leq s_1 < s_2 < \infty$, $0 \leq s'_1 < s'_2 < \infty$, nt_1, nt_2, nt'_1, nt'_2 are integers and the intervals (nt_1, nt_2) and (nt'_1, nt'_2) are identical or disjoint and the same holds for the intervals (s_1, s_2) and (s'_1, s'_2) , but identity cannot hold at both places.

Proof. We have

$$\mathbf{E} \left(\sum_{k=nt_1+1}^{nt_2} X_{k,n}^{(s_1, s_2)} \right)^2 = n(t_2 - t_1) \mathbf{E} \left(X_{1,n}^{(s_1, s_2)} \right)^2 + R$$

where

$$R = 2 \sum_{j=2}^{nt_2-nt_1} (nt_2 - nt_1 - j + 1) \mathbf{E} \left(X_{1,n}^{(s_1, s_2)} X_{j,n}^{(s_1, s_2)} \right).$$

Using Lemmas 2.2 and 2.4 we get, using $n/d \rightarrow \infty$,

$$\begin{aligned} \mathbf{E} \left(X_{1,n}^{(s_1, s_2)} \right)^2 &= \mathbf{E} (X_1 I\{s_1(n/d) < X_1 \leq s_2(n/d)\})^2 - \mathbf{E}^2 (X_1 I\{s_1(n/d) < X_1 \leq s_2(n/d)\}) \\ &= c_0(1 + o(1))(n/d)(s_2 - s_1) + O(\log^2(n/d)) \sim c_0(n/d)(s_2 - s_1) \end{aligned}$$

and

$$|R| \leq 2n \sum_{j=2}^{nt_2-nt_1} \psi(j-1) \left(\mathbf{E} |X_{1,n}^{(s_1, s_2)}| \right)^2 \leq C_3 n \log^2(n/d) \sum_{j=2}^{\infty} e^{-\lambda j} = o(n^2/d),$$

proving (2.5).

To prove (2.6), consider a generic term

$$\begin{aligned} (2.7) \quad & \mathbf{E} X_{i,n}^{(s_1, s_2)} X_{j,n}^{(s'_1, s'_2)} \\ &= \mathbf{E} X_i X_j I\{s_1(n/d) < X_i \leq s_2(n/d)\} I\{s'_1(n/d) < X_j \leq s'_2(n/d)\} \\ & \quad - \mathbf{E} X_i I\{s_1(n/d) < X_i \leq s_2(n/d)\} \mathbf{E} X_j I\{s'_1(n/d) < X_j \leq s'_2(n/d)\} \end{aligned}$$

of the left hand side of (2.6). Fix $r \geq 0$ and sum those covariances in (2.7) where $j - i = r$ and $nt_1 + 1 \leq i \leq nt_2$, $nt'_1 + 1 \leq j \leq nt'_2$. Clearly, the case $r = 0$ can occur only if $(nt_1, nt_2) = (nt'_1, nt'_2)$, but in this case by the assumptions of the lemma (s_1, s_2) and (s'_1, s'_2) must be disjoint and thus the product of the two indicators in the second line of (2.7) is 0. Thus by the first statement of Lemma 2.4 the product expectation in the first line of (2.7) is $O(\log^2(n/d))$ and since the number of such terms in the expansion of (2.6) is at most n , the contribution of such terms in the sum in (2.6) is at most $O(n \log^2(n/d)) = o(n^2/d)$ by $n/d \rightarrow \infty$. For $r \geq 1$ the covariance in (2.7) is at most $\psi(r) O(\log^2(n/d))$ by Lemma 2.2 and the first statement of Lemma 2.4 and since for fixed r the number of pairs (i, j) is at most n , the contribution of all such terms for all $r \geq 1$ is at most $Cn \log^2(n/d) \sum_{r=1}^{\infty} \psi(r) = O(n \log^2(n/d)) = o(n^2/d)$, proving (2.6).

The following central limit theorem for triangular arrays of mixing sequences is Theorem 3 in Philipp [23].

Lemma 2.6. Let $\{x_{nk}, k = 1, 2, \dots, n, n = 1, 2, \dots\}$ be an array of mean zero random variables with finite fourth moments satisfying

$$(2.8) \quad \sup_{n,k} \mathbb{E}(x_{nk}^2) < \infty, \quad s_n^2 = \mathbb{E} \left(\sum_{k=1}^n x_{nk} \right)^2 \rightarrow \infty \quad (n \rightarrow \infty)$$

and assume that for any $n \geq 1$, $0 \leq m < m+r \leq n$ we have

$$(2.9) \quad \sum_{k=m+1}^{m+r} \|x_{nk}\|_4 \leq K \mathbb{E} \left(\sum_{k=m+1}^{m+r} x_{nk} \right)^2$$

for some constant $K > 0$ and

$$(2.10) \quad \sup_{B \in \mathcal{F}_{m+r,n}^{(n)}} |P(B|\mathcal{F}_{1,m}^{(n)}) - P(B)| \leq \begin{cases} 1 & \text{if } r \leq 4 \log s_n \\ s_n^2 e^{-\lambda r} & \text{if } 4 \log s_n < r \leq n \end{cases}$$

where $\mathcal{F}_{a,b}^{(n)}$ denotes the σ -algebra generated by the r.v.'s $\{x_{nk}, a \leq k \leq b \leq n\}$. Then

$$s_n^{-1} \sum_{k=1}^n x_{nk} \xrightarrow{d} N(0,1) \quad \text{as } n \rightarrow \infty.$$

As pointed out in the Introduction, the sequence $\{a_j, j \geq 1\}$ of partial quotients is ψ -mixing with exponential rate with respect to the Gauss measure. Thus it is also ϕ -mixing with exponential rate, i.e.

$$(2.11) \quad \phi(k) := \sup_{r \geq k} \sup_m \sup_{B \in \mathcal{F}_{m+r,\infty}} |P(B|\mathcal{F}_{1,m}) - P(B)| \leq C e^{-\lambda k}, \quad k \geq 1$$

for some positive constants C, λ . By an observation of Samur [29], p. 4 (see also Bradley [5]), we have $\phi(1) < 1$, and since $\phi(k)$ defined in (2.11) is nonincreasing, we have $\phi(k) < 1$ for all $k \geq 1$. This shows that condition (2.10) for the triangular array $\{x_{nk}, 1 \leq k \leq n, n \geq 1\}$ in (3.8) in Section 3 is satisfied. Actually, instead of Lemma 2.6 we could also use Corollary 4.6 in Samur [28] in the real valued case.

3. Proof of Theorem 1.3

Put

$$Q_n = \frac{1}{A_n} \sum_{m=1}^M \sum_{j=1}^J \mu_{m,j} U_n([t_{m-1}, t_m] \times [s_{j-1}, s_j])$$

and

$$Z = \sum_{m=1}^M \sum_{j=1}^J \mu_{m,j} W([t_{m-1}, t_m] \times [s_{j-1}, s_j])$$

for all $M \geq 1, J \geq 1$, real coefficients $\mu_{m,j}$, $0 = s_0 < s_1 < s_2 < \dots < s_J < \infty$, $0 = t_0 < t_1 < \dots < t_M = 1$. Clearly, Z is a normal random variable with mean zero and

$$(3.1) \quad \mathbb{E}Z^2 = \sum_{m=1}^M \sum_{j=1}^J \mu_{m,j}^2 (t_m - t_{m-1})(s_j - s_{j-1}).$$

We claim that

$$(3.2) \quad Q_n \xrightarrow{d} Z \quad \text{for all considered values of } M, J, \mu_{m,j}, t_m, s_j.$$

Since the processes U_n and W are equal to 0 on the boundary of the first quadrant, we have

$$U_n(t_m, s_j) = \sum_{p=1}^m \sum_{q=1}^j U_n([t_{p-1}, t_p] \times [s_{q-1}, s_q])$$

and the same relation holds for W . Thus (3.2) implies

$$\frac{1}{A_n} \sum_{m=1}^M \sum_{j=1}^J \mu_{m,j}^* U_n(t_m, s_j) \xrightarrow{d} \sum_{m=1}^M \sum_{j=1}^J \mu_{m,j}^* W(t_m, s_j)$$

for arbitrary real coefficients $\mu_{m,j}^*$ and this, by the Cramér-Wold device, implies the convergence of the finite-dimensional distributions in Theorem 1.3.

Clearly, $U_n([t_{m-1}, t_m] \times [s_{j-1}, s_j])$ equals

$$\sum_{k=[nt_{m-1}]+1}^{[nt_m]} X_k I\{s_{j-1}(n/d) < X_k \leq s_j(n/d)\} - \mathbb{E}X_k I\{s_{j-1}(n/d) < X_k \leq s_j(n/d)\}$$

and thus relation (3.2) is equivalent to

$$(3.3) \quad \frac{1}{A_n} \sum_{k=1}^n (z_{nk} - \mathbb{E}z_{nk}) \xrightarrow{d} N(0, \mathbb{E}Z^2),$$

where

$$(3.4) \quad z_{nk} = \sum_{j=1}^J \mu_{m,j} X_k I\{s_{j-1}(n/d) < X_k \leq s_j(n/d)\}, \quad [nt_{m-1}] + 1 \leq k \leq [nt_m].$$

Since the terms of the sum in (3.4) are random variables with disjoint support, by the second relation of Lemma 2.4 we have

$$\mathbb{E}z_{nk}^2 = (1 + o_n(1))c_0(n/d) \sum_{j=1}^J \mu_{m,j}^2 (s_j - s_{j-1}), \quad [nt_{m-1}] + 1 \leq k \leq [nt_m].$$

Also, by the first relation of Lemma 2.4 we have

$$\mathbb{E}|z_{nk}| = O(\log(n/d)),$$

where the constant in the O depends also on the $\mu_{m,j}, s_j$. Consequently, letting

$$(3.5) \quad B_m = c_0 \sum_{j=1}^J \mu_{m,j}^2 (s_j - s_{j-1}),$$

we get

$$(3.6) \quad \begin{aligned} & \text{Var } z_{nk} \\ &= (1 + o_n(1))c_0(n/d) \sum_{j=1}^J \mu_{m,j}^2 (s_j - s_{j-1}) = (1 + o_n(1))(n/d)B_m, \quad [nt_{m-1}] + 1 \leq k \leq [nt_m]. \end{aligned}$$

Further, Lemma 2.5 implies for $n \rightarrow \infty$

$$\mathbb{E}U_n([t_{m-1}, t_m] \times [s_{j-1}, s_j])^2 = (1 + o_n(1))c_0(t_m - t_{m-1})(s_j - s_{j-1})(n^2/d)$$

and

$$\mathbb{E}U_n([t_{m_1-1}, t_{m_1}] \times [s_{j_1-1}, s_{j_1}])U_n([t_{m_2-1}, t_{m_2}] \times [s_{j_2-1}, s_{j_2}]) = o_n(n^2/d)$$

provided the pairs (m_1, j_1) and (m_2, j_2) are different. Thus

$$(3.7) \quad \begin{aligned} & \mathbb{E} \left(\sum_{k=1}^n (z_{nk} - \mathbb{E}z_{nk}) \right)^2 = \mathbb{E} \left(\sum_{m=1}^M \sum_{j=1}^J \mu_{m,j} U_n([t_{m-1}, t_m] \times [s_{j-1}, s_j]) \right)^2 \\ &= \sum_{m_1, m_2=1}^M \sum_{j_1, j_2=1}^J \mu_{m_1, j_1} \mu_{m_2, j_2} \mathbb{E} [U_n([t_{m_1-1}, t_{m_1}] \times [s_{j_1-1}, s_{j_1}]) U_n([t_{m_2-1}, t_{m_2}] \times [s_{j_2-1}, s_{j_2}])] \\ &\sim c_0(n^2/d) \sum_{m=1}^M \sum_{j=1}^J \mu_{m,j}^2 (t_m - t_{m-1})(s_j - s_{j-1}) = c_0(n^2/d) \mathbb{E}Z^2 = A_n^2 \mathbb{E}Z^2. \end{aligned}$$

Let

$$B = \max_{1 \leq m \leq M} B_m, \quad B' = \min_{1 \leq m \leq M} B_m$$

where B_m is defined by (3.5). Clearly, without loss of generality we can assume that for each $1 \leq m \leq M$ at least one $\mu_{m,j}$ differs from 0, and thus B, B' are positive numbers depending on the numbers $c_0, s_j, \mu_{m,j}$. We now apply Lemma 2.6 with

$$(3.8) \quad x_{nk} = \frac{z_{nk} - \mathbb{E}z_{nk}}{\sqrt{Bn/d}}.$$

In view of (3.6) and (3.7), condition (2.8) is satisfied. Also, the sequence $\{X_j, j \geq 1\}$ in Theorem 1.3 is assumed to be ψ -mixing with exponential rate, and thus condition (2.10) of Lemma 2.6 is also satisfied. Finally, to verify (2.9), let us note that by the first relation of Lemma 2.4, we have

$$(3.9) \quad \sum_{k=u+1}^{u+r} \mathbb{E}|z_{nk} - \mathbb{E}z_{nk}| \leq C_4 r \log(n/d)$$

where C_4 is a positive constant depending on the $\mu_{m,j}, s_j$. On the other hand, using (3.6), the first relation of Lemma 2.4, exponential ψ -mixing and Lemma 2.2 we get

$$(3.10) \quad \mathbb{E} \left(\sum_{k=u+1}^{u+r} (z_{nk} - \mathbb{E}z_{nk}) \right)^2 \geq \sum_{k=u+1}^{u+r} \mathbb{E}(z_{nk} - \mathbb{E}z_{nk})^2 - 2 \sum_{k, \ell=u+1}^{u+r} |\text{Cov}(z_{nk}, z_{n\ell})|$$

$$\begin{aligned}
&\geq C_5(n/d)r - 2 \sum_{k,\ell=u+1}^{u+r} \psi(|k-\ell|) \mathbb{E}|z_{nk} - \mathbb{E}z_{nk}| \mathbb{E}|z_{n\ell} - \mathbb{E}z_{n\ell}| \\
&\geq C_5(n/d)r - C_6 \sum_{k,\ell=u+1}^{u+r} \psi(|k-\ell|) \log^2(n/d) \geq C_5(n/d)r - C_6 \log^2(n/d) \sum_{j=1}^{\infty} r\psi(j) \\
&\geq C_7(n/d)r
\end{aligned}$$

for $n \geq n_0$, where C_5, C_6, C_7 also depend on the $\mu_{m,j}, s_j$. Relation (2.9) now follows for $n \geq n_0$ from (3.8), (3.9) and (3.10). Applying Lemma 2.6 for the array $\{x_{nk}, 1 \leq k \leq n, n \geq n_0\}$ and using (3.7), the central limit theorem (3.3) follows.

Next we prove tightness in Theorem 1.3. Let

$$B_{11} = [t_1, t] \times [s_1, s], \quad B_{12} = [t_1, t] \times [s, s_2], \quad B_{21} = [t, t_2] \times [s_1, s].$$

where $0 \leq t_1 < t < t_2 \leq 1$, $1/2 \leq s_1 < s < s_2 \leq 3/2$. In view of Lemma 2.1, it suffices to show that

$$(3.11) \quad \mathbb{E} \left| \frac{1}{A_n} U_n(B_{11}) \right|^2 \left| \frac{1}{A_n} U_n(B_{ij}) \right|^2 \leq C^* \mu(B_{11}) \mu(B_{ij}),$$

holds for each $ij \in \{12, 21\}$ with some constant $C^* > 0$. Moreover, since $U_n(t, s)$ is constant on intervals $k/n \leq t < (k+1)/n$, by the last statement of Lemma 2.1 we may assume that nt, nt_1 and nt_2 are all integers. To prove (3.11), we introduce the notations

$$\begin{aligned}
X_i^{(1)} &= X_i I\{s_1(n/d) < X_i \leq s(n/d)\}, & m_i^{(1)} &= \mathbb{E}X_i^{(1)}, \\
X_i^{(2)} &= X_i I\{s(n/d) < X_i \leq s_2(n/d)\}, & m_i^{(2)} &= \mathbb{E}X_i^{(2)}.
\end{aligned}$$

Using Lemmas 2.3 and 2.5 and (1.12) we get

$$\begin{aligned}
&\mathbb{E} \left| \frac{1}{A_n} U_n(B_{11}) \right|^2 \left| \frac{1}{A_n} U_n(B_{21}) \right|^2 \\
&= \mathbb{E} \left(\frac{1}{A_n} \sum_{i=nt_1+1}^{nt} (X_i^{(1)} - m_i^{(1)}) \right)^2 \left(\frac{1}{A_n} \sum_{i=nt+1}^{nt_2} (X_i^{(1)} - m_i^{(1)}) \right)^2 \\
&\leq (1 + \psi(1))^2 \frac{1}{A_n^4} \mathbb{E} \left(\sum_{i=nt_1+1}^{nt} (X_i^{(1)} - m_i^{(1)}) \right)^2 \mathbb{E} \left(\sum_{i=nt+1}^{nt_2} (X_i^{(1)} - m_i^{(1)}) \right)^2 \\
&\leq C_8(t - t_1)(t_2 - t)(s - s_1)^2 = C_8 \mu(B_{11}) \mu(B_{21}).
\end{aligned}$$

for $n \geq n_0$. On the other hand,

$$\begin{aligned}
&\mathbb{E} \left| \frac{1}{A_n} U_n(B_{11}) \right|^2 \left| \frac{1}{A_n} U_n(B_{12}) \right|^2 \\
&= \frac{1}{A_n^4} \mathbb{E} \left(\sum_{i=nt_1+1}^{nt} (X_i^{(1)} - m_i^{(1)}) \right)^2 \left(\sum_{i=nt_1+1}^{nt} (X_i^{(2)} - m_i^{(2)}) \right)^2
\end{aligned}$$

$$(3.12) \quad = \frac{1}{A_n^4} \mathbb{E} \left(\sum_{i=nt_1+1}^{nt} Y_i^{(1)} \right)^2 \left(\sum_{i=nt_1+1}^{nt} Y_i^{(2)} \right)^2,$$

where we put

$$Y_i^{(1)} = X_i^{(1)} - m_i^{(1)}, \quad Y_i^{(2)} = X_i^{(2)} - m_i^{(2)}.$$

The expression in the third line of (3.12) equals the sum of all expressions

$$(3.13) \quad A_n^{-4} \mathbb{E}(Y_i^{(1)} Y_j^{(1)} Y_k^{(2)} Y_\ell^{(2)}),$$

where $nt_1 + 1 \leq i, j, k, \ell \leq nt$. The following facts can be verified by elementary calculations using Lemmas 2.2–2.4:

- (a) $\mathbb{E}|Y_i^{(1)}| \ll s - s_1, \quad \mathbb{E}|Y_i^{(2)}| \ll s_2 - s, \quad \mathbb{E}|Y_i^{(1)} Y_i^{(2)}| \ll (s - s_1)(s_2 - s)$
- (b) $\mathbb{E}(Y_i^{(1)})^2 \ll (n/d)(s - s_1), \quad \mathbb{E}(Y_i^{(2)})^2 \ll (n/d)(s_2 - s),$
- (c) $\mathbb{E}(Y_i^{(1)})^2 |Y_i^{(2)}| \ll (n/d)(s - s_1)(s_2 - s), \quad \mathbb{E}|Y_i^{(1)}| (Y_i^{(2)})^2 \ll (n/d)(s - s_1)(s_2 - s),$
- (d) $\mathbb{E}(Y_i^{(1)})^2 (Y_i^{(2)})^2 \ll (n/d)(s - s_1)(s_2 - s),$

where \ll means the same as the O notation, with an implied constant depending on the sequence (X_n) . We prove relation (d), the proof of (a), (b), (c) is similar (and simpler). We have

$$\begin{aligned} \mathbb{E}(Y_i^{(1)})^2 (Y_i^{(2)})^2 &= \mathbb{E} \left[(X_i^{(1)} - m_i^{(1)})^2 (X_i^{(2)} - m_i^{(2)})^2 \right] \\ &= \mathbb{E}(X_i^{(1)})^2 (X_i^{(2)})^2 - 2m_i^{(2)} \mathbb{E}(X_i^{(1)})^2 X_i^{(2)} + (m_i^{(2)})^2 \mathbb{E}(X_i^{(1)})^2 - 2m_i^{(1)} \mathbb{E}X_i^{(1)} (X_i^{(2)})^2 \\ &\quad + 4m_i^{(1)} m_i^{(2)} \mathbb{E}X_i^{(1)} X_i^{(2)} - 2m_i^{(1)} (m_i^{(2)})^2 \mathbb{E}X_i^{(1)} + (m_i^{(1)})^2 \mathbb{E}(X_i^{(2)})^2 - 2(m_i^{(1)})^2 m_i^{(2)} \mathbb{E}X_i^{(2)} \\ &\quad + (m_i^{(1)})^2 (m_i^{(2)})^2. \end{aligned}$$

Clearly $X_i^{(1)}$ and $X_i^{(2)}$ are supported on different sets and thus $X_i^{(1)} X_i^{(2)} = 0$. Thus among the 9 terms above, the first, second, fourth and fifth are equal to 0. Also, the second and third statement of Lemma 2.4 imply, in view of $1/2 \leq s_1 < s < s_2 \leq 3/2$,

$$\begin{aligned} m_i^{(1)} &= \mathbb{E}X_1^{(1)} \ll s - s_1, \quad m_i^{(2)} = \mathbb{E}X_i^{(2)} \ll s_2 - s \\ \mathbb{E}(X_i^{(1)})^2 &\ll (s - s_1)(n/d), \quad \mathbb{E}(X_i^{(2)})^2 \ll (s_2 - s)(n/d) \end{aligned}$$

for $n \geq n_0$. This shows that the remaining five terms of the sum above are $\ll (n/d)(s - s_1)(s_2 - s)$, proving statement (d) above. Statements (a), (b) and (c) can be proved similarly.

We can now estimate the expressions in (3.13). We will distinguish four cases according as i, j, k, ℓ are all different, or the number of different ones among them is 1, 2 or 3. Consider first the case when i, j, k, ℓ are all different, say $i < j < k < \ell$; let $r = j - i$. Applying Lemma 2.2 with $X = Y_i^{(1)}$, $Y = Y_j^{(1)} Y_k^{(2)} Y_\ell^{(2)}$ and using that $\mathbb{E}X = 0$, we get that the absolute value of the expression (3.13) is bounded by

$$A_n^{-4} \psi(r) \mathbb{E}|X| \mathbb{E}|Y| \leq C A_n^{-4} \psi(r) \mathbb{E}|Y_i^{(1)}| \mathbb{E}|Y_j^{(1)}| \mathbb{E}|Y_k^{(2)}| \mathbb{E}|Y_\ell^{(2)}| \leq C A_n^{-4} \psi(r) (s - s_1)^2 (s_2 - s)^2,$$

where we used Lemma 2.3 to estimate $\mathbb{E}|Y|$ and relation (a) above. Here, and in the rest of the tightness proof, C denotes (possibly different) constants depending only on the sequence (X_n) . Arguing similarly, but splitting the four-term product in (3.13) after the third term, we get the

same bound, except that $\psi(r)$ gets replaced by $\psi(r')$, where $r' = \ell - k$. Thus the absolute value of the expression in (3.13) is at most

$$CA_n^{-4}\psi(r)^{1/2}\psi(r')^{1/2}(s-s_1)^2(s_2-s)^2.$$

Fixing the pair (i, ℓ) and summing for (j, k) means summing for (r, r') and since $\sum_{n=1}^{\infty} \psi(n)^{1/2} < \infty$ and the pair (i, ℓ) can be chosen by at most $(nt-nt_1)^2$ different ways, it follows that the contribution of all terms (3.13) with $i < j < k < \ell$ is at most

$$\begin{aligned} CA_n^{-4}(nt-nt_1)^2(s-s_1)^2(s_2-s)^2 &\leq C(d^2/n^2)(t-t_1)^2(s-s_1)^2(s_2-s)^2 \\ &\leq C(t-t_1)^2(s-s_1)(s_2-s) = C\mu(B_{11})\mu(B_{12}), \end{aligned}$$

using (1.12) and $d/n \rightarrow 0$. The contribution of terms (3.13) where i, j, k, ℓ are different, but their order is different can be estimated similarly.

Next we consider the case when $i = j = k = \ell$. In this case the expression (3.13) becomes $A_n^{-4}E(Y_i^{(1)})^2E(Y_i^{(2)})^2$, which by the estimate in (d) above is at most $CA_n^{-4}(n/d)(s-s_1)(s_2-s)$. Since the number of choices for i is $nt-nt_1 \leq (nt-nt_1)^2$, the contribution of all such expressions is bounded by

$$CA_n^{-4}(n/d)(s-s_1)(s_2-s)(nt-nt_1)^2 \leq C(d/n)(s-s_1)(s_2-s)(t-t_1)^2 \leq C\mu(B_{11})\mu(B_{12}),$$

using again (1.12) and $d/n \rightarrow 0$.

Assume now that among i, j, k, ℓ there are two different ones, i.e. these numbers are pairwise equal or three are equal and the fourth is different. Starting with the case of two pairs, assume e.g. that $i = j$ and $k = \ell$, but $i \neq k$. In this case the expression (3.13) becomes $A_n^{-4}E(Y_i^{(1)})^2(Y_k^{(2)})^2$ which, in view of Lemma 2.3 and the estimate in (b) above is at most

$$CA_n^{-4}(n/d)^2(s-s_1)(s_2-s).$$

Since the number of choices for the pair (i, k) is at most $(nt-nt_1)^2$, using (1.12) it follows that the total contribution of all such terms (3.13) is at most

$$CA_n^{-4}(n/d)^2(s-s_1)(s_2-s)(nt-nt_1)^2 \leq C(s-s_1)(s_2-s)(t-t_1)^2 = C\mu(B_{11})\mu(B_{12}).$$

If $i = k, j = \ell$ and $i \neq j$, then the expression (3.13) becomes $A_n^{-4}EY_i^{(1)}Y_i^{(2)}Y_j^{(1)}Y_j^{(2)}$ which by Lemma 2.3 and the estimate in (a) above is bounded by

$$CA_n^{-4}E|Y_i^{(1)}Y_i^{(2)}|E|Y_j^{(1)}Y_j^{(2)}| \leq CA_n^{-4}(s-s_1)^2(s_2-s)^2.$$

Since the number of pairs (i, j) is $\leq (nt-nt_1)^2$, the contribution of such terms is at most

$$CA_n^{-4}(s-s_1)^2(s_2-s)^2(nt-nt_1)^2 \leq C(s-s_1)(s_2-s)(t-t_1)^2 = C\mu(B_{11})\mu(B_{12}).$$

Assume now that from the indices i, j, k, ℓ three are equal and the fourth one is different. Letting e.g. $i = j = k$ and $i \neq \ell$, the expression (3.13) becomes $A_n^{-4}E(Y_i^{(1)})^2Y_i^{(2)}Y_\ell^{(2)}$ which is, by Lemma 2.3 and the estimates (a) and (c) above is bounded by

$$CA_n^{-4}E(Y_i^{(1)})^2|Y_i^{(2)}|E|Y_\ell^{(2)}| \leq CA_n^{-4}(n/d)(s-s_1)(s_2-s)^2.$$

Since the number of pairs (i, ℓ) is $\leq (nt-nt_1)^2$, the total contribution of such terms is at most $C\mu(B_{11})\mu(B_{12})$.

Finally, if the number of different indices among i, j, k, l is 3, e.g. if $i = j < k < l$, then the expression (3.13) becomes $A_n^{-4} \mathbf{E}(Y_i^{(1)})^2 Y_k^{(2)} Y_l^{(2)}$ which by using $\mathbf{E}Y_\ell^{(2)} = 0$, Lemma 2.2, Lemma 2.3 and estimates (a) and (b) above, can be estimated by

$$(3.14) \quad CA_n^{-4} \psi(r) \mathbf{E}(Y_i^{(1)})^2 \mathbf{E}|Y_k^{(2)}| \mathbf{E}|Y_\ell^{(2)}| \leq CA_n^{-4} \psi(r) (n/d)(s-s_1)(s_2-s)^2,$$

where $r = \ell - k$. Since for fixed r the number of triples (i, k, ℓ) with $\ell - k = r$ is at most $(nt - nt_1)^2$, the contribution of such terms (3.13) is at most

$$CA_n^{-4} \psi(r) (n/d)(s-s_1)(s_2-s)^2 (nt - nt_1)^2 \leq C\psi(r)(s-s_1)(s_2-s)(t-t_1)^2$$

and summing for r we get again $\leq C\mu(B_{11})\mu(B_{12})$. The other cases (e.g. $i < j = k < \ell$, etc.) can be treated similarly and the proof of tightness in Theorem 1.3 is completed. This also completes the proof of the theorem.

We prove now, as claimed after Theorem 1.3, that

$$(3.15) \quad \frac{\eta_{d,n}}{n/d} \xrightarrow{P} 1$$

for $d = d_n \rightarrow \infty$, $d_n/n \rightarrow 0$. Fix $n \geq 1$, $1/2 < t < 2$ and let $T_k = I\{X_k \geq tn/d\}$, $1 \leq k \leq n$. Then by Lemma 2.2 and (1.11) we get

$$|\mathbf{E}T_1 T_k - \mathbf{E}T_1 \mathbf{E}T_k| \leq \psi(k-1) \mathbf{E}T_1 \mathbf{E}T_k \leq C_9 \exp(-\lambda k) (d/n)^2$$

and thus setting $\bar{T}_k = T_k - \mathbf{E}T_k$ we conclude that

$$\begin{aligned} \mathbf{E} \left(\sum_{k=1}^n \bar{T}_k \right)^2 &= n \mathbf{E}\bar{T}_1^2 + 2 \sum_{k=2}^n (n-k+1) \mathbf{E}\bar{T}_1 \bar{T}_k \\ &\leq n \left(\mathbf{E}\bar{T}_1^2 + 2 \sum_{k=2}^n |\mathbf{E}\bar{T}_1 \bar{T}_k| \right) \\ &\leq n \left(\mathbf{E}T_1^2 + C_{10} (d/n)^2 \sum_{k=2}^n \exp(-\lambda k) \right) \\ &\leq n (\mathbf{E}T_1 + C_{11} (d/n)^2) \\ &\leq C_{12} d. \end{aligned}$$

Hence Markov's inequality and $d = d_n \rightarrow \infty$ imply for any $\varepsilon > 0$

$$P \left\{ \sum_{k=1}^n \bar{T}_k \geq \varepsilon d \right\} \rightarrow 0,$$

and since $\mathbf{E}T_k = \mathbf{E}T_1 \sim d/(nt)$ by (1.11), it follows that

$$\#\{k \leq n : X_k \geq tn/d\} = \sum_{k=1}^n I\{X_k \geq tn/d\} \sim d/t \quad \text{in probability as } n \rightarrow \infty.$$

Thus for fixed $t > 1$ and n large, with probability tending to 1 the number of X_k 's, $1 \leq k \leq n$ exceeding tn/d is smaller than d , and thus $\eta_{d,n} \leq tn/d$. Similarly, for $t < 1$ and n large, with probability tending to 1 we have $\eta_{d,n} \geq tn/d$, and thus (3.15) is proved.

Proof of Remark 1.1. Let (X_n) be a sequence satisfying the assumptions of Theorem 1.2, and put $X'_n = X_n + 4^{-n}$. Letting $\eta'_{d,n}$ denote the d -th largest of X'_1, \dots, X'_n and $S_n^{(r)}$ and $S_n'^{(r)}$ denote the sums $\sum_{k=1}^n X_k$, $\sum_{k=1}^n X'_k$ after removing their r largest terms, it is easily seen that

$$(3.16) \quad |S_n^{(r)} - S_n'^{(r)}| \leq 2 \quad \text{for any } r \geq 1$$

and

$$(3.17) \quad n|m(\eta_{d,n}) - m(\eta'_{d,n})| = O_P(1).$$

Clearly, relation (1.11) will fail for the perturbed sequence (X'_n) , but as inspection shows, all the lemmas in the proof of Theorem 1.2 and the subsequent arguments remain valid, so conclusion (1.13) of the theorem remains valid if we replace X_i by X'_i and $\eta_{d,n}$ by $\eta'_{d,n}$. Since the X_n are integer valued, with probability one all the X'_j , $j = 1, 2, \dots$ are different, and thus the sum of the X'_j 's, $1 \leq j \leq n$ not exceeding $\eta'_{d,n}$ equals $S_n'^{(d-1)}$. Thus we have

$$(3.18) \quad \frac{S_n'^{(d-1)} - nm(\eta'_{d,n})}{n/\sqrt{d}} \xrightarrow{d} N(0, c_0).$$

In view of (3.16) and (3.17), we can drop the primes in (3.18) and since $S_n^{(d-1)} - S_n^{(d)} = \eta_{d,n} = O_P(n/d)$ by (3.15), the conclusion of Theorem 1.1 follows.

REFERENCES

- [1] Arov, D.Z. and Bobrov, A.A.: The extreme members of a sample and their role in the sum of independent variables. *Theor. Probability Appl.* 5 (1960), 377–396.
- [2] Bickel, P. J. and Wichura, M. J.: Convergence criteria for multiparameter stochastic processes and some applications. *Ann. Math. Statist.* 42 (1971), 1656–1670.
- [3] Billingsley, P.: *Ergodic theory and information*. Wiley, 1965.
- [4] Billingsley, P.: *Convergence of Probability Measures*. Wiley, New York 1968.
- [5] Bradley, R.: *Introduction to strong mixing conditions*, Vol I. Kendrick Press, 2007.
- [6] Csörgő, S., Horváth, L. and Mason, D.: What portion of the sample makes a partial sum asymptotically stable or normal? *Z. Wahrschein. verw. Gebiete* 72 (1986), 1–16.
- [7] Csörgő, S. and Simons, G.: A strong law of large numbers for trimmed sums, with applications to generalized St. Petersburg games. *Statist. Prob. Letters* 26 (1996), 65–73.
- [8] Darling, D.: The influence of the maximum term in the addition of independent random variables. *Trans. Amer. Math. Soc.* 73 (1952), 95–107.
- [9] Diamond, H. and Vaaler, J.: Estimates for partial sums of continued fraction partial quotients. *Pacific J. Math.* 122 (1986), 73–82.
- [10] Doeblin, W.: Remarques sur la théorie métrique des fractions continues. *Comp. Math.* 7 (1940), 353–371.
- [11] Galambos, J.: The distribution of the largest coefficient in continued fraction expansions. *Quart. J. Math. Oxford Ser.* 23 (1972), 147–151.
- [12] Galambos, J.: An iterated logarithm type theorem for the largest coefficient in continued fractions. *Acta Arith.* 25 (1973/74), 359–364.
- [13] Gordin, M.I. and Reznik, M.H.: The law of the iterated logarithm for the denominators of continued fractions. *Vestnik Leningrad Univ.* 25 (1970), 28–33. (In Russian)
- [14] Heinrich, L.: Rates of convergence in stable limit theorems for sums of exponentially ψ -mixing random variables with an application to metric theory of continued fractions. *Math. Nachr.* 131 (1987), 149–165.
- [15] Ibragimov, I.A.: A theorem from the metric theory of continued fractions. *Vestnik Leningrad Univ.* 1 (1960), 13–24. (In Russian)
- [16] Iosifescu, M.: A Poisson law for ψ -mixing sequences establishing the truth of a Doeblin's statement. *Rev. Roumaine Math. Pures Appl.* 22 (1977), 1441–1447.

- [17] Iosifescu, M.: A survey of the metric theory of continued fractions, fifty years after Doeblins 1940 paper. In: Probability Theory and Mathematical Statistics, vol. I, pp. 550–572. Vilnius, Mokslas (1990)
- [18] Khinchin, A. J.: Metrische Kettenbruchprobleme. *Compositio Math.* 1 (1935), 361–382.
- [19] Kusmin, R.: Sur un problème de Gauss. *Atti Congr. Intern. Bologna 1928.* 6, 83–89.
- [20] Leadbetter, M. and Rootzén, H.: Extremal theory for stochastic processes. *Ann. Probab.* 16 (1988), 431–478.
- [21] Lévy, P.: Sur les lois dont dépendent les quotients complets et incomplets d’une fraction continue. *Bull. Sci. Math. France* 57 (1929), 178–194.
- [22] Lévy, P.: Fractions continues aléatoires. *Rend. Circ. Mat. Palermo* 1 (1952), 170–208.
- [23] Philipp, W.: The remainder in the central limit theorem for mixing stochastic processes. *Ann. Math. Statist.* 40 (1969), 601–609.
- [24] Philipp, W.: Some metrical theorems in number theory II. *Duke Math. J.* 37 (1970), 447–458.
- [25] Philipp, W.: A conjecture of Erdős on continued fractions. *Acta Arith.* 28 (1975/76), 379–386.
- [26] Philipp, W.: Limit theorems for partial quotients of continued fractions. *Monatsh. Math.* 105 (1988), 195–206.
- [27] Philipp, W. and Stackelberg, O.: Zwei Grenzwertsätze für Kettenbrüche. *Math. Ann.* 181 (1969), 152–156.
- [28] Samur, J.: Convergence of sums of mixing triangular arrays of random vectors with stationary rows. *Ann. Probab.* 12 (1984), 390–426.
- [29] Samur, J.: On some limit theorems for continued fractions. *Trans. Amer. Math. Soc.* 316 (1989), 53–79.
- [30] Samur, J.: Some remarks on a probability limit theorem for continued fractions. *Trans. Amer. Math. Soc.* 348 (1996), 1411–1428.
- [31] Stackelberg, O.P.: On the law of the iterated logarithm for continued fractions. *Duke Math. J.* 33 (1966), 801–819.
- [32] Szewczak, Z.S.: On limit theorems for continued fractions. *J. Theor. Probab.* 22 (2009), 239–255.
- [33] Vardi, I.: The St. Petersburg game and continued fractions. *C.R. Acad. Sci. Paris Ser. I. Math.* 324 (1997), 913–918.