

Dear Author

Here are the proofs of your article.

- You can submit your corrections **online**, via **e-mail** or by **fax**.
- For **online** submission please insert your corrections in the online correction form. Always indicate the line number to which the correction refers.
- You can also insert your corrections in the proof PDF and **email** the annotated PDF.
- For **fax** submission, please ensure that your corrections are clearly legible. Use a fine black pen and write the correction in the margin, not too close to the edge of the page.
- Remember to note the **journal title**, **article number**, and **your name** when sending your response via e-mail or fax.
- **Check** the metadata sheet to make sure that the header information, especially author names and the corresponding affiliations are correctly shown.
- **Check** the questions that may have arisen during copy editing and insert your answers/corrections.
- **Check** that the text is complete and that all figures, tables and their legends are included. Also check the accuracy of special characters, equations, and electronic supplementary material if applicable. If necessary refer to the *Edited manuscript*.
- The publication of inaccurate data such as dosages and units can have serious consequences. Please take particular care that all such details are correct.
- Please **do not** make changes that involve only matters of style. We have generally introduced forms that follow the journal's style.
- Substantial changes in content, e.g., new results, corrected values, title and authorship are not allowed without the approval of the responsible editor. In such a case, please contact the Editorial Office and return his/her consent together with the proof.
- If we do not receive your corrections **within 48 hours**, we will send you a reminder.
- Your article will be published **Online First** approximately one week after receipt of your corrected proofs. This is the **official first publication** citable with the DOI. **Further changes are, therefore, not possible.**
- The **printed version** will follow in a forthcoming issue.

Please note

After online publication, subscribers (personal/institutional) to this journal will have access to the complete article via the DOI using the URL:

<http://dx.doi.org/10.1007/s00436-016-5215-y>

If you would like to know when your article has been published online, take advantage of our free alert service. For registration and further information, go to:

<http://www.link.springer.com>.

Due to the electronic nature of the procedure, the manuscript and the original figures will only be returned to you on special request. When you return your corrections, please inform us, if you would like to have these documents returned.

Metadata of the article that will be visualized in OnlineFirst

Please note: Images will appear in color online but will be printed in black and white.

1	Article Title	The life cycle of <i>Thelohanellos kitauei</i> (Myxozoa: Myxosporea) infecting common carp (<i>Cyprinus carpio</i>) involves aurantiactinomyxon in <i>Branchiura sowerbyi</i>	
2	Article Sub- Title		
3	Article Copyright - Year	Springer-Verlag Berlin Heidelberg 2016 (This will be the copyright line in the final PDF)	
4	Journal Name	Parasitology Research	
5		Family Name	Zhai
6		Particle	
7		Given Name	Yanhua
8		Suffix	
9		Organization	Huazhong Agricultural University
10		Division	Department of Aquatic Animal Medicine, College of Fisheries
11		Address	Wuhan
12		Organization	Freshwater Aquaculture Collaborative Innovation Center of Hubei Province
13	Corresponding Author	Division	
14		Address	Wuhan
15		Organization	Key Lab of Freshwater Animal Breeding, Ministry of Agriculture
16		Division	
17		Address	Wuhan
18		e-mail	zhaiyh@mail.hzau.edu.cn
19		Family Name	Zhao
20		Particle	
21	Author	Given Name	Dandan
22		Suffix	
23		Organization	Huazhong Agricultural University
24		Division	Department of Aquatic Animal Medicine, College of Fisheries
25		Address	Wuhan
26		Organization	Freshwater Aquaculture Collaborative Innovation Center of Hubei Province

27		Division	
28		Address	Wuhan
29		e-mail	
30		Family Name	Borkhanuddin
31		Particle	
32		Given Name	Muhammad Hafiz
33		Suffix	
34		Organization	Institute for Veterinary Medical Research, Centre for Agricultural Research
35	Author	Division	
36		Address	HAS, POB 18, Budapest H-1581
37		Organization	University Malaysia Terengganu
38		Division	School of Marine and Environmental Science
39		Address	Kuala Terengganu
40		e-mail	
41		Family Name	Wang
42		Particle	
43		Given Name	Weimin
44		Suffix	
45		Organization	Freshwater Aquaculture Collaborative Innovation Center of Hubei Province
46	Author	Division	
47		Address	Wuhan
48		Organization	Key Lab of Freshwater Animal Breeding, Ministry of Agriculture
49		Division	
50		Address	Wuhan
51		e-mail	
52		Family Name	Liu
53		Particle	
54		Given Name	Yang
55		Suffix	
56		Organization	Huazhong Agricultural University
57	Author	Division	Department of Aquatic Animal Medicine, College of Fisheries
58		Address	Wuhan
59		Organization	Freshwater Aquaculture Collaborative Innovation Center of Hubei Province
60		Division	

61		Address	Wuhan
62		Organization	Key Lab of Freshwater Animal Breeding, Ministry of Agriculture
63		Division	
64		Address	Wuhan
65		e-mail	
66		Family Name	Cech
67		Particle	
68		Given Name	Gábor
69		Suffix	
70	Author	Organization	Key Lab of Freshwater Animal Breeding, Ministry of Agriculture
71		Division	
72		Address	Wuhan
73		e-mail	
74		Family Name	Székely
75		Particle	
76		Given Name	Csaba
77		Suffix	
78	Author	Organization	Institute for Veterinary Medical Research, Centre for Agricultural Research
79		Division	
80		Address	HAS, POB 18, Budapest H-1581
81		e-mail	
82		Received	23 February 2016
83	Schedule	Revised	
84		Accepted	27 July 2016
85	Abstract	<i>Thelohanellus kitauei</i> is a freshwater myxosporean parasite causing intestinal giant cystic disease of common carp. To clarify the life cycle of <i>T. kitauei</i> , we investigated the oligochaete populations in China and Hungary. This study confirms two distinct aurantiactinomyxon morphotypes (Aurantiactinomyxon type 1 and Aurantiactinomyxon type 2) from <i>Branchiura sowerbyi</i> as developmental stages of the life cycle of <i>T. kitauei</i> . The morphological characteristics and DNA sequences of these two types are described here. Based on 18S rDNA sequence analysis, Aurantiactinomyxon type 1 (2048 bp) and Aurantiactinomyxon type 2 (2031 bp) share 99.2–99.4 %, 99.8–100 % similarity to the published sequences of <i>T. kitauei</i> , respectively. The 18S rDNA sequences of these two aurantiactinomyxon morphotypes share 99.4 % similarity, suggesting intraspecific variation within the	

taxon, possibly due to geographic origin. Phylogenetic analyses demonstrate the two aurantiactinomyxon types clustered with *T. kitauei*. Regardless, based on 18S rDNA synonymy, it is likely that Aurantiactinomyxon type 1 and 2 are conspecific with *T. kitauei*. This is the fourth elucidated two-host life cycle of *Thelohanellus* species and the first record of *T. kitauei* in Europe.

86	Keywords separated by ' - '	<i>Thelohanellus kitauei</i> - Life cycle - Aurantiactinomyxon - 18S rDNA - <i>Branchiura sowerbyi</i> - China - Hungary
87	Foot note information	

The life cycle of *Thelohanellus kitauei* (Myxozoa: Myxosporea) infecting common carp (*Cyprinus carpio*) involves aurantiactinomyxon in *Branchiura sowerbyi*

Dandan Zhao^{1,2} · Muhammad Hafiz Borkhanuddin^{3,4} · Weimin Wang^{2,5} · Yang Liu^{1,2,5} · Gábor Cech⁵ · Yanhua Zhai^{1,2,5} · Csaba Székely³

Received: 23 February 2016 / Accepted: 27 July 2016
© Springer-Verlag Berlin Heidelberg 2016

Abstract *Thelohanellus kitauei* is a freshwater myxosporean parasite causing intestinal giant cystic disease of common carp. To clarify the life cycle of *T. kitauei*, we investigated the oligochaete populations in China and Hungary. This study confirms two distinct aurantiactinomyxon morphotypes (Aurantiactinomyxon type 1 and Aurantiactinomyxon type 2) from *Branchiura sowerbyi* as developmental stages of the life cycle of *T. kitauei*. The morphological characteristics and DNA sequences of these two types are described here. Based on 18S rDNA sequence analysis, Aurantiactinomyxon type 1 (2048 bp) and Aurantiactinomyxon type 2 (2031 bp) share 99.2–99.4 %, 99.8–100 % similarity to the published sequences of *T. kitauei*, respectively. The 18S rDNA sequences of these two aurantiactinomyxon morphotypes share 99.4 % similarity, suggesting intraspecific variation within the taxon, possibly due to geographic origin. Phylogenetic analyses demonstrate the two aurantiactinomyxon types clustered with *T. kitauei*. Regardless, based on 18S rDNA synonymy, it is likely that Aurantiactinomyxon type 1 and 2 are conspecific with *T. kitauei*. This is the fourth elucidated

two-host life cycle of *Thelohanellus* species and the first record of *T. kitauei* in Europe.

Keywords *Thelohanellus kitauei* · Life cycle · Aurantiactinomyxon · 18S rDNA · *Branchiura sowerbyi* · China · Hungary

Introduction

Myxozoans, a group of obligate parasitic metazoans, play a significant pathogenic role in aquatic vertebrates (mostly fish) worldwide. The myxozoan life cycles are complex and mostly involve both vertebrate and invertebrate hosts, which was first demonstrated in 1984 for *Myxobolus cerebralis* (Wolf & Markiw 1984). Since the pioneering discovery, approximately 50 myxozoan species are confirmed to follow this life cycle pattern that involves the alternation of a myxosporean stage developing in fish with an actinosporean stage in annelid worms (Székely et al. 2014; Eszterbauer et al. 2015).

Initially, life cycle studies of myxozoans were solely performed based on experimental infections. However, most of these studies only replicate partial life cycles (typically fish to worm transmission). It is difficult to conduct holistic transmission experiments for the following reasons: uncertainty of susceptible hosts, deficiency of knowledge of the intrapiscine developmental pathways, and appropriate conditions for infection (Eszterbauer et al. 2015). Along with increasing importance of molecular methods in parasitological studies, the small subunit ribosomal RNA gene (18S rDNA) has been applied as a molecular marker for identifying actinosporean-myxosporean pairs (Bartholomew et al. 1997; Holzer et al. 2004; Eszterbauer et al. 2006; Zhai et al. 2012; Borkhanuddin et al. 2014; Rosser et al. 2015). Compared with the complex, time-consuming and labor intensive nature of

✉ Yanhua Zhai
zhaiyh@mail.hzau.edu.cn

¹ Department of Aquatic Animal Medicine, College of Fisheries, Huazhong Agricultural University, Wuhan, China
² Freshwater Aquaculture Collaborative Innovation Center of Hubei Province, Wuhan, China
³ Institute for Veterinary Medical Research, Centre for Agricultural Research, HAS, POB 18, H-1581 Budapest, Hungary
⁴ School of Marine and Environmental Science, University Malaysia Terengganu, Kuala Terengganu, Malaysia
⁵ Key Lab of Freshwater Animal Breeding, Ministry of Agriculture, Wuhan, China

63 experimental infections, researchers are more inclined to
 64 identify actinosporean-myxosporean pairs using molecular
 65 analysis.

66 *Thelohanellus* Kudo, 1933 (Myxosporea, Bivalvulida) is
 67 the sixth most speciose genus within the phylum Myxozoa.
 68 At present, at least 109 *Thelohanellus* spp. have been
 69 described (Yuan et al. 2015). The majority of *Thelohanellus*
 70 spp. are seemingly innocuous to their hosts, while others can
 71 be severely pathogenic. One of the more well-researched
 72 members of the genus is *Thelohanellus kitauei*, which has
 73 been implicated in fish kills in common carp (*Cyprinus*
 74 *carpio*) in Asia and can cause considerable economic losses
 75 (Liu et al. 2011; Seo et al. 2012; Zhai et al. 2016). While
 76 investigations into the host specificity, infection site tropism
 77 and genome of *T. kitauei* have taken place, information
 78 regarding the extrapiscine developmental stage has remained
 79 elusive (Shin et al. 2014; Yang et al. 2014). As such, the
 80 source of infection is unknown, although it is presumed the
 81 life cycle of *T. kitauei* is similar to other myxozoans, involving
 82 an actinosporean stage in an alternate invertebrate host
 83 (Seo et al. 2012).

84 Historically, *T. kitauei* has been reported from common
 85 carp in East Asia, mainly referring to Japan, Korea, and
 86 China (Egusa & Nakajima 1981; Seo et al. 2012; Zhai et al.
 87 2016). To date, *T. kitauei* has not been reported from Europe
 88 or the Americas, even though the common carp is globally
 89 distributed.

90 In the present study, we performed investigations of the
 91 actinosporean fauna at Lake Balaton and Kis-Balaton
 92 Reservoir of Hungary and freshwater fish ponds in China,
 93 and examined the morphology and 18S rDNA sequences of
 94 actinosporean stages. The aim of this study was to confirm the
 95 identity of two aurantiactinomaxon morphotypes from
 96 *Branchiura sowerbyi* found in different biotopes as the
 97 counterpart actinosporean stages of *T. kitauei* based on
 98 morphological and molecular analyses.

99 **Materials and methods**

100 **Sample collection**

101 Oligochaetes were harvested from different sites of Lake
 102 Balaton (Keszthely, Tihany, Balatonvilágos, Balatonszemes,
 103 Zala channel, Siófok) and Kis-Balaton Reservoir, western part
 104 of Hungary (hereinafter referred to as 'natural waters') and
 105 cultured fish ponds located at Dongxi Lake and Datong
 106 Lake of Hubei Province, China (hereinafter referred to as 'fish
 107 ponds'). Sampling from natural waters was carried out from
 108 April to October, 2011 and April to September, 2012, while
 109 samples were collected in the fish ponds from August to
 110 December, 2014 and March to July 2015. Oligochaete

samples were taken one to three times each month during
 the sampling period.

For the natural waters, sediment was collected near water
 vegetation at about 0.5 to 1 m depth. At each sampling
 occasion, as much as 40–60 l of mud volume was sieved in
 situ through 1000 µm mesh size net. Oligochaetes trapped
 together with debris, vegetation roots and decayed particles
 were then transferred to the laboratory with minimal lake
 water. For the fish ponds, sediment was sampled by spade
 from the accessible shallow water along the ponds. Then,
 the mud samples were sieved and washed gently with pond
 water through a 450-µm mesh sieve. The material remaining
 in the sieve was transferred into plastic boxes containing a
 small quantity of pond water and transported to the laboratory.

On arrival, the sediments were aerated and supplied with
 additional fresh dechlorinated tap water. Oligochaetes were
 hand-sorted from the retained material in trays filled with
 dechlorinated tap water. Oligochaetes of natural waters were
 identified according to the key of Timm (1999), while
 oligochaetes in the fish ponds were identified morphologically
 following the guidelines of Wang (2002).

Collection and morphological identification of actinospores

Oligochaetes were separated and placed into wells of 24 cell-
 well plates with 2 ml dechlorinated tap water or distilled water
 according to the methods of Yokoyama et al. (1991). Plates
 with worms were held at ambient temperatures and the water
 in the plates was examined daily for the presence of
 actinospores by inverted microscopy. Observed actinospores
 were harvested upon release. Photomicrographs of actinospores
 from natural waters were taken from fresh material using both
 bright-field and phase contrast illumination and a DP-20 digital
 camera mounted on an Olympus BH-2 microscope. Similarly,
 photomicrographs of actinospores from fish ponds were taken
 with a Zeiss Axio Imager A2 fluorescence microscope
 equipped with Andor Clara CCD camera. Line drawings were
 made based on these photomicrographs.

Morphological measurements were taken from a variable
 number of spores, depending on availability from one infected
 oligochaete. Spore measurements of actinosporean types were
 performed according to the guidelines of Lom et al. (1997).
 The number of germ cells was determined by placing
 actinospores on glass slides and pressing gently on the spores
 with a coverslip to mechanically release and disrupt the
 sporoplasm. All measurements were in micrometers (µm).

Molecular methods

Actinospores harvested from oligochaetes collected from
 natural waters were initially preserved in 80 % ethanol.
 Genomic DNA was extracted using the DNeasy™ tissue kit

(animal tissue protocol; QIAGEN, Germany) according to the manufacturer's instructions. A semi-nested PCR system was used for amplification. The first round of PCR reaction was carried out with universal eukaryotic primers ERIB1 and ERIB10 (Barta et al. 1997). In the second round amplification, two semi-nested PCR reactions were performed using two myxozoan primer pairs, MyxospecF (Fiala 2006)-ERIB10 and ERIB1-TKR1 (Seo et al. 2012), respectively. The first round PCR was carried out in a 25- μ l reaction mixture comprising 2 μ l of extracted genomic DNA, 5 μ l of 1 mM deoxy-ribonucleotide triphosphates (dNTPs; MBI Fermentas), 0.325 μ M of each primer, 2.5 μ l of 10 \times DreamTaq buffer (MBI Fermentas), 0.1 μ l of DreamTaq polymerase (2 U; MBI Fermentas) and 15 μ l of water. The following profile was used to amplify the 18S rDNA region: an initial denaturation step at 95 $^{\circ}$ C for 3 min, followed by 35 cycles at 95 $^{\circ}$ C for 1 min, 55 $^{\circ}$ C for 1 min, 72 $^{\circ}$ C for 2 min, and was completed with terminal extension at 72 $^{\circ}$ C for 7 min. For the second round of semi-nested PCR reactions, 1 μ l of the initial amplified product was used as the template. Compared with the first round PCR, the second round of PCR was performed in a 50- μ l reaction mixture and the quantities of each ingredient were doubled. Amplification conditions in the second round followed this profile: 95 $^{\circ}$ C for 3 min, then 35 cycles at 95 $^{\circ}$ C for 50 s, 50 $^{\circ}$ C for 50 s, 72 $^{\circ}$ C for 1 min 40 s, and terminated with an extension period at 72 $^{\circ}$ C for 7 min. The PCR products were electrophoresed in 1 % agarose gels in 1 \times TAE buffer gel stained with ethidium bromide. Amplified DNA was purified with the EZ-10 Spin Column PCR Purification Kit (Bio Basic Inc). Purified PCR products were sequenced with myxozoan specific primers listed by Székely et al. (2014) and CR1F (Székely et al. 2015a, 2015b), using the BigDye Terminator v3.1 Cycle Sequencing Kit (Life Technologies) with an ABI PRISM $^{\circ}$ 3100 Genetic Analyser (Life Technologies).

Genomic DNA of actinospores harvested from oligochaetes in fish culture ponds was extracted using a TransDirect $^{\text{TM}}$ Animal Tissue PCR Kit (TransGen Biotech, Beijing) according to the manufacturer's instructions. The PrimeSTAR $^{\circ}$ Max DNA Polymerase (TaKaRa) was used to ensure high fidelity sequence. The 18S rDNA was amplified with universal eukaryotic primer pairs 18e (Hillis & Dixon 1991) and 18r (Whipps et al. 2003). PCR was performed in a 50- μ l reaction mixture comprising 25 μ l 2 \times PrimeSTAR Max Premix, 0.4 μ M of each primer, 16 μ l distilled water and approximate 100–150 ng of genomic DNA. Amplification was carried out using a ProFlex $^{\text{TM}}$ PCR System (ABI, America) and the amplification profile was as follows: 98 $^{\circ}$ C for 10 s, 55 $^{\circ}$ C for 15 s, and terminal elongation at 72 $^{\circ}$ C for 1 min for 35 cycles. The PCR products were electrophoresed through a 1 % agarose gel in 1 \times Tris-Acetate-EDTA (TAE) buffer and purified using the Gel Extraction Kit (CWBIO, Beijing). Purified PCR fragments

were cloned into pMD-19T vector system (TaKaRa). Positive clones were selected and sequenced with an ABI PRISM $^{\circ}$ 3730XL DNA sequencer (Applied Biosystems Inc., Foster, USA). The contiguous sequences were assembled according to the corresponding chromatograms with the SeqMan $^{\text{TM}}$ utility of the Lasergene software package (DNA Star, Madison, Wisconsin) and submitted to the National Center for Biotechnology Information (NCBI) nucleotide database.

Phylogenetic analyses

Myxozoan nucleotide sequences used in phylogenetic analyses were chosen based on BLAST searches. The highly similar sequences (>80 % match) that were at least 1500 bp in length ($n=34$) were downloaded from the NCBI nucleotide database. The dataset was aligned with the software MAFFT v. 7.271 (Katoh & Standley 2013), and manually corrected using the BioEdit sequence alignment editor program (Hall 1999). Hypervariable or ambiguous regions were deleted to ensure comparison of homologous positions. Phylogenetic analyses were conducted by Bayesian (BI) and maximum likelihood (ML) methods. The best-fit evolutionary model for BI and ML analyses was determined by MrModeltest v. 2.3 (Nylander 2004), which identified the optimal evolutionary model as the general time reversible model (GTR+I+G) using the Akaike information criteria. Bayesian analysis was conducted in MrBayes v. 3.1 (Ronquist & Huelsenbeck 2003). The length of chains was of 2,000,000 generations with sampling each 100 generations, and the prior 5000 were discarded as burn-in. ML analysis was performed using PhyML v. 3.0 (Guindon et al. 2010). Nucleotide frequencies were estimated from the data (A=0.2548, C=0.2029, G=0.2703, T=0.2720); six rates of nucleotide substitution were [AC]=1.1156, [AG]=3.4511, [AT]=1.3843, [CG]=0.7126, [CT]=5.4490, [GT]=1.0000; proportion of invariable sites=0.2339; gamma distribution shape parameter=0.3271. Bootstrap confidence values were calculated with 100 replicates. The resulting topologies were annotated with MEGA v. 6.06 software package (Tamura et al. 2013) and compared with each other. *Ceratonova shasta* was selected as outgroup.

Results

Oligochaetes identified as possible hosts in the present study mainly consisted of *B. sowerbyi*, *Isochaetides michaelsoni*, *Limnodrilus hoffmeisteri*, *Nais* sp., *Dero* sp., *Aelosoma* sp. and *Bothrioneurum vej dovskyanum*. From *B. sowerbyi* (434 and 7321 specimens collected from Hungary and China, respectively), two distinct aurantiactinomyxon morphotypes (Aurantiactinomyxon type 1 and Aurantiactinomyxon type 2) corresponded to known *T. kitauei* deposited in the GenBank

261 database. Morphometrics of relevant aurantiactinomyxon
262 types are listed in Table 1. The genetic similarities of
263 aurantiactinomyxon types with *T. kitauei* deposited in the
264 GenBank are summarized in Table 2.

265 **Description of aurantiactinomyxon morphotypes**

266 *Aurantiactinomyxon type 1 nov. (Fig. 1a–d)*

267
268 Description: Spore body is spherical in apical view but with
269 three smooth indentations at the edge of the structure, 19.7
270 (17.3–23.3) µm in diameter. The caudal processes are of equal
271 length, triangular shaped, tapering to pencil point tips, and
272 curve downwards, measuring 20.4 (18.7–23.3) µm long and
273 8.9 (7.4–10.0) µm wide at the base. Three polar capsules
274 are located in the middle of the spore body and are pyriform
275 in side view, measuring 3.4 µm long, 2.8 µm wide.
276 Flattened sporoplasmic body reveals at least 28 germ cells
277 (most likely 32).

278 Host: *Branchiura sowerbyi* Beddard, 1892

279 Locality: Kis-Balaton Reservoir, Hungary

280 Date of collection: July, 2011

281 Prevalence: 0.01 % (1 out of 7755)

282 GenBank accession no.: KU664643

283 Remarks: Aurantiactinomyxon type 1 described here closely
284 resembled the Aurantiactinomyxon of *Thelohanellus*
285 *nikolskii* (Székely et al. 1998) and Aurantiactinomyxon type
286 ‘B2’ of Eszterbauer et al. (2006), but the caudal processes of the
287 latter two were shorter. Furthermore, our aurantiactinomyxon
288 also showed similar morphological characteristics with
289 Aurantiactinomyxon type 2 of El-Mansy et al. (1998).

290 However, the latter possessed slightly bigger caudal pro-
291 cesses (20.4 vs 22.6 µm in mean length and 8.9 vs 11.7 µm
292 in mean width) and was collected from *Limnodrilus* sp.
293 (Table 1). No pansporocysts could be detected from the
294 infected oligochaete. By amplification and sequencing, a
295 total of 2048 bases of 18S rDNA was generated from
296 Aurantiactinomyxon type 1. On the basis of the DNA se-
297 quences, Aurantiactinomyxon type 1 showed maximum
298 identity with *T. kitauei*. The contiguous sequence fragment
299 presented a similar percentage of 99.4, 99.4, 99.3, and 99.2 %
300 to the sequences of *T. kitauei* available in GenBank: JQ690367,
301 KR872638, HM624024 and GQ396677, respectively
302 (Table 2).

303 *Aurantiactinomyxon type 2 nov. (Fig. 2a–d)*

304
305 Description: Spore body is typically trefoil-shaped with
306 obvious interlobular retractions in apical view, 20.9 (19.3–
307 22.1) µm in diameter, and in side view it is ellipsoidal, 18.4
308 (17.6–23.0) µm in length, 22.3 (21.6–23.0) µm in width (mea-
309 sured from 8 spores). Three equal-sized triangular-sepal-like
310 caudal processes entirely embrace the spore body, and curve
311 downward with pointed ends in side view, measuring 19.7
312 (17.9–22.3) µm long, 11.6 (9.8–13.0) µm wide at the base.
313 In addition, the caudal processes seem to be composed of two
314 parts, that is, one is round and the other pointed at the end. The
315 nuclei of valve cell locate either distally or proximally in cau-
316 dal processes. Three polar capsules are spherical in apical
317 view, 2.7 (2.6–2.9) µm in diameter and pyriform in side view,
318 measuring 3.0 (2.8–3.3) µm long, 2.4 (2.2–2.6) µm wide.
319 Number of secondary cells is 32.

Q5 t1.1 **Table 1** Comparison of morphological measurements of the newly identified and the previously reported aurantiactinomyxon types

t1.2	Actinospore (Reference)	Caudal processes		Spore body	Polar capsules	No. of germ cells	Host	
		L	W					
t1.4	Aurantiactinomyxon type 1 (present study)	20.4 (18.7–23.3)	8.9 (7.4–10.0)	D: 19.7 (17.3–23.3)	L: 3.4 W: 2.8	>26 (c. 32)	<i>Branchiura sowerbyi</i>	t1.5
t1.6	Aurantiactinomyxon type 2 (present study)	19.7 (17.9–22.3)	11.6 (9.8–13.0)	D: 20.9 (19.3–22.1)	L: 3.0 (2.8–3.3) W: 2.4 (2.2–2.6) D: 2.7 (2.6–2.9)	32	<i>B. sowerbyi</i>	t1.7 t1.8
t1.9	Aurantiactinomyxon of <i>Thelohanellus nikolskii</i> (Székely et al. 1998)	13.4 (11.3–15.5)	9.0 (8.5–9.6)	D: 21.1 (21–21.2)	D: 2.1 (2.0–2.2)	16	<i>Tubifex tubifex</i>	
t1.10	Aurantiactinomyxon type 2 (El-Mansy et al. 1998)	22.6	11.7	D: 21.1	L: 2.8 W: 2.0	n.d.	<i>Limnodrilus</i> sp.	t1.11
t1.12	Aurantiactinomyxon ‘B2’ (Eszterbauer et al. 2006)	16 (14–20)	8.4 (7–10.6)	D: 19 (18–21)	L: 2.6 (2–4.1)	n.d.	<i>B. sowerbyi</i>	
t1.13	Aurantiactinomyxon type JD (Xi et al. 2015)	21.7 (20.0–24.4)	14.0 (11.2–16.4)	L: 15.6 W: 21.2 (17.1–24.0)	D: 2.3 (2.0–2.8)	>30	<i>B. sowerbyi</i>	t1.14

All measurements are in µm and ranges are given in parentheses
L length, W width, D diameter, n.d. no data

Table 2 Genetic similarities of aurantiactinomyxon types from Kis-Balaton Reservoir (Hungary) and fish pond of Honghu (China) with *Thelohanellos kitauei* deposited in the GenBank. Length of 18S rDNA sequences and GenBank accession numbers are shown in parentheses

Myxozoan	1	2	3	4	5	6
1, Aurantiactinomyxon type 1 (2048 bp, KU664643)	100 %					
2, Aurantiactinomyxon type 2 (2031 bp, KU664644)	99.4 %	100 %				
3, <i>T. kitauei</i> (2051 bp, JQ690367)	99.4 %	99.9 %	100 %			
4, <i>T. kitauei</i> (2048 bp, HM624024)	99.3 %	99.8 %	99.9 %	100 %		
5, <i>T. kitauei</i> (2031 bp, KR872638)	99.4 %	99.8 %	99.7 %	99.6 %	100 %	
6, <i>T. kitauei</i> (1561 bp, GQ396677)	99.2 %	100 %	99.9 %	99.7 %	99.6 %	100 %

Host: *Branchiura sowerbyi* Beddard, 1892
Locality: Datong Lake, Honghu City, Hubei Province, China
Date of collection: May, 2015
Prevalence: 0.04 % (3 out of 7755)
GenBank accession no.: KU664644
Remarks: The Aurantiactinomyxon type 2 presented here was nearly similar to the Aurantiactinomyxon ‘B2’ of Eszterbauer et al. (2006), but caudal processes of the latter were relatively smaller, 19.7 (17.9–22.3) vs 16 (14–20) µm in length and 11.6 (9.8–13.0) vs 8.4 (7–10.6) µm in width, and spore body of the latter was incompletely encircled within the base of caudal processes. The shape and measurements of this type also closely resembled the Aurantiactinomyxon type JD of Xi et al. (2015) except the different shape ends of caudal processes (triangular vs rounded). The partial 18S rDNA sequence of Aurantiactinomyxon type 2 was obtained from 3 clones, and the sequences had no variation among 3 clones

chosen. The contiguous sequence (2031 bp) was processed to remove vector sequence and subsequently deposited in GenBank database. Based on DNA sequence analysis, Aurantiactinomyxon type 2 showed the highest genetic similarity with *T. kitauei* (JQ690367, HM624024, KR872638 and GQ396677), reaching a similarity percentage of 99.8–100 % (Table 2).

Phylogenetic analyses

Phylogenetic inferences were established based on the two newly obtained 18S rDNA sequences and 34 myxozoan sequences, which consisted of actinospores and their closest related myxospores retrieved from GenBank. Phylogenetic trees constructed by BI and ML analyses had similar topological structure though with different support values at some branch nodes (Fig. 3). Phylogenetic analyses displayed that the two newly identified aurantiactinomyxon types clustered

Fig. 1 Aurantiactinomyxon type 1 nov. **a** Line drawing of apical view of a mature spore. **b** Line drawing of side view of a mature spore. **c** Apical view of a waterborne spore. **d** Side view of a waterborne spore. Scale bars = 20 µm





Fig. 2 Aurantiactinomyxon type 2 nov. **a** Line drawing of apical view of a mature spore. **b** Line drawing of side view of a mature spore. **c** Apical view of a waterborne spore. **d** Side view of a waterborne spore. Scale bars = 20 μ m



354 with *T. kitauei* and formed a sister relationship with
 355 *Thelohanellus hovorkai*, which integrated *Thelohanellus*
 356 *wuhanensis* and *T. nikolskii* to jointly constitute an indepen-
 357 dent *Thelohanellus* clade. The phylogenetic trees also showed
 358 *Thelohanellus* species had a close relationship with some
 359 *Myxobolus* species, which were consistent with several previ-
 360 ous reports (Shin et al. 2014; Yuan et al. 2015).


361 Discussion

362 The current study revealed the oligochaete *B. sowerbyi* is
 363 an extrapiscine host in the life cycle of *T. kitauei* and
 364 demonstrated two phenotypic aurantiactinomyxon, namely
 365 Aurantiactinomyxon type 1 and Aurantiactinomyxon type
 366 2, as the developmental stages of the life cycle of *T. kiatauei*
 367 by morphological and molecular analyses (Fig. 4).

368 In terms of morphology, Aurantiactinomyxon type 1 and
 369 Aurantiactinomyxon type 2 had obvious distinctions. The spore
 370 body of Aurantiactinomyxon type 2 was typically trefoil-
 371 shaped with conspicuous interlobular retractions in apical view
 372 while that of Aurantiactinomyxon type 1 was spherical despite
 373 three inconspicuous indentations at the edge of the structure. In
 374 addition, the caudal processes of Aurantiactinomyxon type 2

were unique and seemed to be composed of two parts: one is
 375 round and the other pointed at the end. It is noticeable that the
 376 special structure of caudal processes is only reported in the
 377 Aurantiactinomyxon type 2 and type 4 of Özer et al. (2002).
 378 At DNA level, Aurantiactinomyxon type 2 presented here
 379 showed maximum identity with *T. kitauei* and shared a similar
 380 percentage of 99.8–100 % to the sequences of *T. kitauei* avail-
 381 able in GenBank (Table 2), which strongly suggests that
 382 Aurantiactinomyxon type 2 and *T. kitauei* are conspecific.
 383 The 18S rDNA sequence of Aurantiactinomyxon type 1 was
 384 99.2–99.4 % similar to *T. kitauei* deposited in the GenBank and
 385 that of two aurantiactinomyxon types described here
 386 exhibited 99.4 % similarities to each other (Table 2),
 387 suggesting that this taxon have intraspecific variability,
 388 possibly due to different sampling locations. Hallett et al.
 389 (2004) reported that different morphometrical triactinomyxon
 390 types exhibited by morphological similarity were actually the
 391 same genotype. This was later supported by Eszterbauer
 392 et al. (2006), who similarly reported the same genotype
 393 can represent multiple morphotypes, suggesting that
 394 actinospore classification based solely on traditional mor-
 395 phological features can lead to erroneous associations.
 396 Therefore, DNA sequence analysis is recommended in
 397 species descriptions.
 398

Fig. 3 Phylogenetic positions of Aurantiactinomyxon type 1 and Aurantiactinomyxon type 2 based on 18S rDNA data analyzed using Bayesian analysis (BI). Numbers at nodes represent bootstrap support values of Bayesian posterior probabilities and maximum likelihood (ML). GenBank accession numbers are in parentheses. *Dotted box* represents two *Myxobolus* species are separated from the major *Myxobolus* clade. *Ceratonova shasta* is taken as outgroup. Aurantiactinomyxon type 1 and Aurantiactinomyxon type 2 examined in this study are in **bold**



In North China, common carp are the most prevalent cultured fish species with the highest production. *T. kitauei*, the pathogenic agent of intestinal giant-cystic disease of common carp, has the potential to cause catastrophic fish kills




Fig. 4 Schematic illustration of *T. kitauei* life cycle: Aurantiactinomyxon type 1 (A1) and Aurantiactinomyxon type 2 (A2) actinospores infect the vertebrate host common carp (V) in which then myxospores (M) infecting the invertebrate host *B. sowerbyi* (I) develop

and severe economic losses in the aquaculture industry of North China (Chen & Ma 1998). Interestingly, the myxospore stage of *T. kitauei* has not been reported from Europe. Molnár (2009) hypothesized that European common carp originated from the Far-Eastern Amur-Chinese geographical region. During the long migration from China to Europe, the common carp lost its original parasite fauna. Then, after introduction of the Amur wild common carp and the color carp from Asia to Europe, several ‘old’ parasites of the common carp have been introduced and are gradually expanding. This would suggest that more recently described parasites of carp, at present known only in China, Japan and the Amur Basin might also make their way to Europe (Székely et al. 2015a, 2015b). This is the first report on the actinosporean stage of *T. kitauei* from Europe based on morphological, molecular and phylogenetic analyses. The Aurantiactinomyxon type 1 identified in the current study indicates that further investigations are urgently needed to find the myxosporean stage of *T. kitauei* in fish host, most probably, common carp in Europe.

So far, only three life cycles of *Thelohanellus* species have been elucidated, all with *B. sowerbyi* as invertebrate hosts. *T. hovorkai* and *T. nikolskii* involve aurantiactinomyxon as intraoligochaete developmental stages (Yokoyama 1997; Székely et al. 1998). *Thelohanellus wangi* involves

neoactinomyxon as extrapiscine alternating stage (Xi et al. 2015). This study confirms two distinct aurantiactinomyxon morphotypes as developmental stages of the life cycle of *T. kitauei* by morphological and molecular analyses and constitutes the fourth description of life cycle of *Thelohanellus* species.

Acknowledgments This project has received funding from the European Union's Horizon 2020 research and innovation program under grant agreement no. 634429 (ParaFishControl), OTKA K 100132, Nature Science Foundation of China (31502209), New Century Excellent Talents in University (NCET-12-0866), and China Postdoctoral Science Foundation (2014M550399). The authors thank Hao Wu, Junfa Yuan, Qinxian Guo, Jianchun Shao, Zhe Wang, Dan Li, and Yongliang Shen (Huazhong Agricultural University, China) for collecting sediment samples.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Ethical approval All applicable international, national, and/or institutional guidelines for the care and use of animals were followed. All procedures performed in studies involving animals were in accordance with the ethical standards of the institution or practice at which the studies were conducted.

Informed consent Informed consent was obtained from all individual participants included in the study. The publication only reflects the views of authors, and the European Commission cannot be held responsible for any use which may be made of the information contained therein.

References

Barta JR, Martin DS, Liberator PA, Dshkevich M, Anderson JW, Feigner SD, Elbrecht A, Perkins-Barrow A, Jenkins MC, Danforth HD, Ruff MD, Profous-Juchelka H (1997) Phylogenetic relationships among eight *Eimeria* species infecting domestic fowl inferred using complete small subunit ribosomal DNA sequences. *J Parasitol* 83: 262–271. doi:10.2307/3284453

Bartholomew JL, Whipple MJ, Stevens DG, Fryer JL (1997) The life cycle of *Ceratomyxa shasta*, a myxosporean parasite of salmonids, requires a freshwater polychaete as an alternate host. *J Parasitol* 83: 859–868. doi:10.2307/3284281

Borkhanuddin MH, Cech G, Molnár K, Németh S, Székely C (2014) Description of raabeia, synactinomyxon and neoactinomyxon developing stages of myxosporeans (Myxozoa) infecting *Isochaetides michaelsoni* Lastockin (Tubificidae) in Lake Balaton and Kis-Balaton Water Reservoir, Hungary. *Syst Parasitol* 88: 245–259. doi:10.1007/s11230-014-9496-1

Chen QL, Ma CL (1998) Fauna Sinica: Myxozoa, Myxosporea. Science Press (In Chinese), Beijing

Egusa S, Nakajima K (1981) A new myxozoa *Thelohanellus kitauei*, the cause of intestinal giant cystic disease of carp. *Fish Pathol* 15: 213–218. doi:10.3147/jsfp.15.213

El-Mansy A, Székely C, Molnár K (1998) Studies on the occurrence of actinosporian stages of myxosporeans in Lake Balaton, Hungary, with

the description of triactinomyxon, raabeia and aurantiactinomyxon types. *Acta Vet Hung* 46:437–450

Eszterbauer E, Marton S, Rácz OZ, Letenyey M, Molnár K (2006) Morphological and genetic differences among actinosporian stages of fish-parasitic myxosporeans (Myxozoa): difficulties of species identification. *Syst Parasitol* 65:97–114. doi:10.1007/s11230-006-9041-y

Eszterbauer E, Atkinson S, Diamant A, Morris D, El-Matbouli M, Hartikainen H (2015) Myxozoan life cycles: practical approaches and insights. In: Okamura B, Gruhl A, Bartholomew JL (eds) Myxozoan evolution, ecology and development. Springer International Publishing, Switzerland. doi:10.1007/978-3-319-14753-6_10

Fiala I (2006) The phylogeny of Myxosporea (Myxozoa) based on small subunit ribosomal RNA gene analysis. *Int J Parasitol* 36:1521–1534. doi:10.1016/j.ijpara.2006.06.016

Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, Gascuel O (2010) New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Syst Biol* 59:307–321. doi:10.1093/sysbio/syq010

Hall TA (1999) BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symp Ser* 41:95–98

Hallett SL, Atkinson SD, Erséus C, El-Matbouli M (2004) Molecular methods clarify morphometric variation in triactinomyxon spores (Myxozoa) released from different oligochaete hosts. *Syst Parasitol* 57:1–14. doi:10.1023/B:SYPA.000010682.90311.91

Hillis DM, Dixon MT (1991) Ribosomal DNA: molecular evolution and phylogenetic inference. *Q Rev Biol* 66:411–453. doi:10.1086/417338

Holzer AS, Sommerville C, Wootten R (2004) Molecular relationships and phylogeny in a community of myxosporeans and actinosporians based on their 18S rDNA sequences. *Int J Parasitol* 34:1099–1111. doi:10.1016/j.ijpara.2004.06.002

Katoh K, Standley DM (2013) MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol Biol Evol* 30:772–780. doi:10.1093/molbev/mst010

Liu Y, Whipps CM, Liu WS, Zeng LB, Gu ZM (2011) Supplemental diagnosis of a myxozoan parasite from common carp *Cyprinus carpio*: synonymy of *Thelohanellus xinyangensis* with *Thelohanellus kitauei*. *Vet Parasitol* 178:355–359. doi:10.1016/j.vetpar.2011.01.008

Lom J, McGeorge J, Feist S, Morris D, Adams A (1997) Guidelines for the uniform characterisation of the actinosporian stages of parasites of the phylum Myxozoa. *Dis Aquat Org* 30:1–9. doi:10.3354/dao030001

Molnár K (2009) Data on the parasite fauna of the European common carp *Cyprinus carpio carpio* and Asian common carp *Cyprinus carpio haematopterus* support an Asian ancestry of the species. *AACL Bioflux* 2:391–400

Nylander JAA (2004) MrModeltest v2. Program distributed by the author. Evolutionary Biology Centre, Uppsala University, Uppsala

Özer A, Wootten R, Shinn AP (2002) Survey of actinosporian types (Myxozoa) belonging to seven collective groups found in a freshwater salmon farm in northern Scotland. *Folia Parasitol* 49:189–210. doi:10.14411/fp.2002.036

Ronquist F, Huelsenbeck JP (2003) MrBayes 3: bayesian phylogenetic inference under mixed models. *Bioinformatics* 19:1572–1574. doi:10.1093/bioinformatics/btg180

Rosser TG, Griffin MJ, Quiniou SMA, Khoo LH, Greenway TE, Wise DJ, Pote LM (2015) Small subunit ribosomal RNA sequence links the myxospore stage of *Henneguya mississippiensis* n. sp. from channel catfish *Ictalurus punctatus* to an actinosporian released by the benthic oligochaete *Dero digitata*. *Parasitol Res* 114: 1595–1602. doi:10.1007/s00436-015-4345-y

- 546 Seo JS, Jeon EJ, Kim MS, Woo SH, Kim JD, Jung SH, Park M, Jee BY, 585
- 547 Kim JW, Kim YC, Lee EH (2012) Molecular identification and real- 586
- 548 time quantitative PCR (qPCR) for rapid detection of *Thelohanellus* 587
- 549 *kitauei*, a myxozoan parasite causing intestinal giant cystic disease 588
- 550 in the Israel carp. Korean J Parasitol 50:103–111. doi:10.3347/ 589
- 551 kjp.2012.50.2.103 590
- 552 Shin SP, Nguyen VG, Jeong JM, Jun JW, Kim JH, Han JE, Baek GW, 591
- 553 Park SC (2014) The phylogenetic study on *Thelohanellus* species 592
- 554 (Myxosporea) in relation to host specificity and infection site 593
- 555 tropism. Mol Phylogenet Evol 72:31–34. doi:10.1016/j. 594
- 556 ympev.2014.01.002 595
- 557 Székely C, El-Mansy A, Molnár K, Baska F (1998) Development of 596
- 558 *Thelohanellus hovorkai* and *Thelohanellus nikolskii* (Myxosporea: 597
- 559 Myxozoa) in oligochaete alternate hosts. Fish Pathol 33:107–114. 598
- 560 doi:10.3147/jsfp.33.107 599
- 561 Székely C, Borkhanuddin MH, Cech G, Kelemen O, Molnár K (2014) 600
- 562 Life cycles of three *Myxobolus* spp. from cyprinid fishes of Lake 601
- 563 Balaton, Hungary involve triactinomyxon-type actinospores. 602
- 564 Parasitol Res 113:2817–2825. doi:10.1007/s00436-014-3942-5 603
- 565 Székely C, Cech G, Chaudhary A, Borzák R, Singh HS, Molnár K 604
- 566 (2015a) Myxozoan infections of the three Indian major carps in fish 605
- 567 ponds around Meerut, UP, India, with descriptions of three new 606
- 568 species, *Myxobolus basuhalidari* sp. n., *M. kalavatieae* sp. n. and 607
- 569 *M. meerutensis* sp. n., and the redescription of *M. catlae* and 608
- 570 *M. bhadrensis*. Parasitol Res 114:1301–1311. doi:10.1007/s00436- 609
- 571 014-4307-9 610
- 572 Székely C, Molnár K, Cech G (2015b) Description of *Myxobolus* 611
- 573 *balatonicus* n. sp. (Myxozoa: Myxobolidae) from the common carp 612
- 574 *Cyprinus carpio* L. in Lake Balaton. Syst Parasitol 91:71–79. 613
- 575 doi:10.1007/s11230-015-9560-5 614
- 576 Tamura K, Stecher G, Peterson D, Filipski A, Kumar S (2013) MEGA6: 615
- 577 molecular evolutionary genetics analysis version 6.0. Mol Biol Evol 616
- 578 30:2725–2729. doi:10.1093/molbev/mst197 617
- 579 Timm T (1999) A guide to the Estonian Annelida. Estonian Academy 618
- 580 Publishers, Tartu-Tallinn 619
- 581 Wang HZ (2002) Studies on taxonomy, distribution and ecology of 620
- 582 microdrile oligochaetes of china, with descriptions of two new spe- 621
- 583 cies from the vicinity of the great wall station of China, Antarctica. 622
- 584 Higher Education (HEP), Beijing 623
- 625 624
- Whipps CM, Adlard RD, Bryant MS, Lester RJ, Findlav V, Kent ML 585
- (2003) First report of three *Kudoa* species from eastern Australia: 586
- Kudoa thyrsites* from mahi mahi (*Coryphaena hippurus*), *Kudoa* 587
- amamiensis* and *Kudoa minithyrsites* n. sp. from sweeper 588
- (*Pemppheris ypsilychnus*). J Eukaryot Microbiol 50:215–219. 589
- doi:10.1111/j.1550-7408.2003.tb00120.x 590
- Wolf K, Markiw ME (1984) Biology contravenes taxonomy in the 591
- Myxozoa: new discoveries show alternation of invertebrate and ver- 592
- tebrate hosts. Science 225:1449–1452. doi:10.1126/science. 593
- 225.4669.1449 594
- Xi BW, Zhou ZG, Xie J, Pan LK, Yang YL, Ge XP (2015) Morphological 595
- and molecular characterization of actinosporeans infecting oligo- 596
- chaete *Branchiura sowerbyi* from Chinese carp ponds. Dis Aquat 597
- Org 114:217–228. doi:10.3354/dao02859 598
- Yang Y, Xiong J, Zhou Z, Huo F, Miao W, Ran C, Liu Y, Zhang J, Feng J, 599
- Wang M, Wang M, Wang L, Yao B (2014) The genome of the 600
- myxosporean *Thelohanellus kitauei* shows adaptations to nutrient 601
- acquisition within its fish host. Genome Biol Evol 6:3182–3198. 602
- doi:10.1093/gbe/evu247 603
- Yokoyama H (1997) Transmission of *Thelohanellus hovorkai* Achmerov, 604
- 1960 (Myxosporea: Myxozoa) to common carp *Cyprinus carpio* 605
- through the alternate oligochaete host. Syst Parasitol 36:79–84. 606
- doi:10.1023/A:1005752913780 607
- Yokoyama H, Ogawa K, Wakabayashi H (1991) A new collection meth- 608
- od of actinosporeans. a probable infective stage of myxosporeans to 609
- fishes from tubificids and experimental infection of goldfish with the 610
- actinosporean, *Raabeia* sp. Fish Pathol 26:133–138. doi:10.3147/ 611
- jsfp.26.133 612
- Yuan S, Xi BW, Wang JG, Xie J, Zhang JY (2015) *Thelohanellus wangi* n. 613
- sp. (Myxozoa, Myxosporea), a new gill parasite of allogynogenetic 614
- gibel carp (*Carassius auratus gibelio* Bloch) in China, causing severe 615
- gill myxosporidiosis. Parasitol Res 114:37–45. doi:10.1007/s00436- 616
- 014-4157-5 617
- Zhai YH, Zhou L, Gui JF (2012) Identification and characterization of 618
- one novel type of triactinospomyxon with short spore axis. Parasitol 619
- Res 110:2385–2393. doi:10.1007/s00436-011-2775-8 620
- Zhai Y, Gu Z, Guo Q, Wu Z, Wang H, Liu Y (2016) New type of 621
- pathogenicity of *Thelohanellus kitauei* Egusa & Nakajima, 1981 622
- infecting the skin of common carp *Cyprinus carpio* L. Parasitol Int 623
- 65:78–82. doi:10.1016/j.parint.2015.10.010 624

AUTHOR QUERIES

AUTHOR PLEASE ANSWER ALL QUERIES.

- Q1. Please check if the affiliations are presented correctly.
- Q2. City has been provided. Please check if it is correct.
- Q3. Please check if the section headings are assigned to appropriate levels.
- Q4. Székely et al. 2015 has been changed to Székely et al. 2015a, 2015b as per the reference list. Please check if okay.
- Q5. Please check if the tables are presented correctly.
- Q6. Please check if the figure captions are presented correctly.

UNCORRECTED PROOF