

Croat Med J. 2016;57:100-9
doi: 10.3325/cmj.2016.57.100

The effects of estrogen on the α_2 -adrenergic receptor subtypes in rat uterine function in late pregnancy *in vitro*

Judit Hajagos-Tóth¹, Judit Bóta¹, Eszter Ducza¹, Adrienn Csányi¹, Zita Tiszai¹, Anna Borsodi², Reza Samavati², Sándor Benyhe², Róbert Gáspár¹

¹Department of Pharmacodynamics and Biopharmacy, Faculty of Pharmacy, University of Szeged, Szeged, Hungary

²Institute of Biochemistry, Biological Research Centre, Hungarian Academy of Sciences, Szeged, Hungary

Aim To assess the effect of 17 β -estradiol pretreatment on the function and expression of α_2 -adrenergic receptors (ARs) subtypes in late pregnancy in rats.

Methods Sprague-Dawley rats (n=37) were treated with 17 β -estradiol for 4 days starting from the 18th day of pregnancy. The myometrial expression of the α_2 -AR subtypes was determined by real time polymerase chain reaction and Western blot analysis. *In vitro* contractions were stimulated with (-)-noradrenaline, and its effect was modified with the selective antagonists BRL 44408 (α_{2A}), ARC 239 ($\alpha_{2B/C}$), and spiroxatrine (α_{2A}). The cyclic adenosine monophosphate (cAMP) accumulation was also measured. The activated G-protein level was investigated by guanosine 5'-O-[gamma-thio]triphosphate (GTP γ S) binding assay.

Results 17 β -estradiol pretreatment decreased the contractile effect of (-)-noradrenaline via the α_2 -ARs, and abolished the contractile effect via the α_{2B} -ARs. All the α_2 -AR subtypes' mRNA was significantly decreased. 17 β -estradiol pretreatment significantly increased the myometrial cAMP level in the presence of BRL 44408 ($P=0.001$), ARC 239 ($P=0.007$), and spiroxatrine ($P=0.045$), but did not modify it in the presence of spiroxatrine + BRL 44408 combination ($P=0.073$). It also inhibited the G-protein-activating effect of (-)-noradrenaline by 25% in the presence of BRL 44408 + spiroxatrine combination.

Conclusions The expression of the α_2 -AR subtypes is sensitive to 17 β -estradiol, which decreases the contractile response of (-)-noradrenaline via the α_{2B} -AR subtype, and might cause changes in G-protein signaling pathway. Estrogen dysregulation may be responsible for preterm labor or uterine inertia via the α_2 -ARs.

Received: January 8, 2016

Accepted: March 22, 2016

Correspondence to:

Róbert Gáspár
Szeged, H-6701
P.O. Box 121, Hungary
gaspar@pharm.u-szeged.hu

In spite of the numerous attempts to explore it, the exact action mechanism and risk of preterm birth still remains one of the biggest challenges in obstetrics and gynecology and a major contributor to perinatal mortality and morbidity, affecting around 9% of births in developed countries (1-4). On the other hand, weak contractions and poor labor outcomes also represent a problem mainly among obese women, increasing the number of cesarean deliveries (5).

Uterine contractility is regulated by several factors, such as the adrenergic system (6) and female sexual hormones (7,8). Progesterone was demonstrated to increase the synthesis of β_2 -ARs during pregnancy (9-11) and the number of activated G-proteins (12,13), which is why it can be combined with β_2 -AR agonists in threatening preterm labor. Myometrial α_1 -AR expression is influenced by female sexual steroid hormones, mainly estrogens. 17β -estradiol decreases the expression of the α_{1A} -ARs, but does not influence the expression of α_{1D} -ARs (14). However, the effect of estrogens on the myometrial α_2 -AR subtypes is unknown. Considering the fact that estrogens play a major role in myometrial contractions during human parturition (15,16), it is important to know if they have a direct influence on the α_2 -AR subtypes, which are also involved in the mechanism of uterine contractions (17).

The α_2 -ARs have been divided into (18,19) α_{2A} , α_{2B} , and α_{2C} subtypes. All three receptor subtypes are coupled to the pertussis toxin-sensitive G_i -protein α -subunit (20) and decrease the activity of adenylyl cyclase (AC) and voltage-gated Ca^{2+} currents, at the same time activating the receptor-operated K^+ currents (21). The stimulation of these receptors leads to presynaptic feedback inhibition of (-)-noradrenaline release on the adrenergic neurons (18), and mediates a variety of cell functions, such as vasoconstriction, increased blood pressure, and nociception. Furthermore, all three α_2 -AR subtypes were identified in both pregnant and non-pregnant myometrium and were shown to take part in both increased and decreased myometrial contractions (22,23). Under certain circumstances, α_2 -ARs can couple not only to G_i -proteins but to G_s -proteins, resulting in the activation of AC (24). On the other hand, pregnancy has been proved to induce a change in the G_i/G_s -activating property of the α_2 -ARs in rats, resulting in a differential regulation of myometrial AC activity in mid-pregnancy vs term (25). The α_{2B} -ARs were shown to predominate and mediate contraction in last-day-pregnant animals by decreasing the intracellular cAMP level, while α_{2A} - and α_{2C} -ARs mediate only weak contractions by

increasing the cAMP level, which can be regarded as relaxation as they are compared with the effect of (-)-noradrenaline (23).

Since female sexual steroid hormones play an important role in the regulation of the adrenergic receptor system (26), the effect of estrogen on different α_2 -AR subtypes has been investigated. The mRNA expression of the α_{2A} -ARs in the spinal cord was increased after estrogen pretreatment (27), which could contribute to the higher prevalence of pain syndromes in women. On the other hand, estrogen was shown to increase the smooth muscle expression of α_{2C} -ARs and therefore the cold-induced constriction of cutaneous arteries (28). In addition, it was shown to stimulate the (-)-noradrenaline release in the hypothalamus due to the decreased coupling of the α_2 -adrenoceptors to G protein (29).

Since there are no available data on the effects of 17β -estradiol on the myometrial functions of different α_2 -AR subtypes, the aim of this study was to clarify the changes in expression and function of the α_{2A} , α_{2B} , and α_{2C} -AR subtypes after 17β -estradiol pretreatment on the last day of pregnancy in rats by using RT-PCR and Western blot analysis. Since the changes in the intracellular cAMP are crucial in the control of smooth muscle contractions and relaxations, our further aim was to measure the cAMP release after 17β -estradiol pretreatment in the presence of the subtype-specific α_2 -AR antagonists. We also investigated the changes in the G-protein activation of α_2 -ARs using GTP γ S binding assay.

MATERIALS AND METHODS

The animal experimentation was carried out with the approval of the Hungarian Ethics Committee for Animal Research (permission number: IV/198/2013). The animals were treated in accordance with the European Communities Council Directives (86/609/ECC) and the Hungarian Act for the Protection of Animals in Research (XXVIII. tv. 32.5).

Housing and handling of the animals

Sprague-Dawley rats were obtained from the INNOVO Ltd (Gödöllő, Hungary) and were housed under controlled temperature (20-23°C), in humidity (40%-60%) and light (12 h light/dark regime) regulated rooms. The animals were fed standard rodent pellet diet (INNOVO Ltd, Isaszeg, Hungary), with tap water available *ad libitum*.

Mating of the animals

Mature female (180-200 g, n=58) and male (240-260 g, n=12) Sprague-Dawley rats were mated in a special mating cage with a time-controlled electrically movable metal door separating the rooms for male and female animals. Since rats are usually active at night, the door was opened before dawn. Within 4-5 hours after the possibility of mating, female rats with the presence of copulation plug or a sperm-positive vaginal smear (search was performed under a microscope at a magnification of 1200 times) were separated. The day of copulation was considered as the first day of pregnancy.

In vivo sexual hormone treatments of the rats

The 17 β -estradiol (Sigma Aldrich, Budapest, Hungary) pre-treatment of the pregnant animals was started on the day 18 of pregnancy. The compound was dissolved in olive oil. The animals were injected subcutaneously with 5 μ g/kg of 17 β -estradiol once a day for 4 days (30). On the day 22, the uterine samples were collected and the contractility and molecular pharmacological studies were carried out.

RT-PCR studies

Tissue isolation: Rats (250-300 g) were sacrificed by CO₂ asphyxiation. Newborn rats were sacrificed by immediate cervical dislocation. The uterine tissues from pregnant animals (n=5 in each experiment) (tissue between two implantation sites) were rapidly removed and placed in RNAlater Solution (Sigma-Aldrich). The tissues were frozen in liquid nitrogen and stored at -70°C until total RNA extraction.

Total RNA preparation from tissue: Total cellular RNA was isolated by extraction with guanidinium thiocyanate-acid-phenol-chloroform according to Chomczynski and Sacchi (31). After precipitation with isopropanol, the RNA was washed with 75% ethanol and then re-suspended in diethyl pyrocarbonate-treated water. RNA purity was controlled at an optical density of 260/280 nm with BioSpec Nano (Shimadzu, Japan); all samples exhibited an absorbance ratio in the range 1.6-2.0. RNA quality and integrity were assessed by agarose gel electrophoresis.

Reverse transcription and amplification of the PCR products was performed by using the TaqMan RNA-to-CTTM 1-Step Kit (Life Technologies, Budapest, Hungary) and the ABI StepOne Real-Time cycler. RT-PCR amplifications were

performed as follows: 48°C for 15 min and 95°C for 10 min, followed by 40 cycles at 95°C for 15 sec, and 60°C for 1 min. The generation of specific PCR products was confirmed by melting curve analysis. Table 1 shows the assay IDs for the used primers. The amplification of β -actin served as an internal control. All samples were run in triplicates. The fluorescence intensities of the probes were plotted against PCR cycle numbers. The amplification cycle displaying the first significant increase in the fluorescence signal was defined as the threshold cycle (Ct).

TABLE 1. Assay IDs of the applied primers

TaqMan assays	Assay ID (Life Technologies, Budapest, Hungary)
α_{2A} -AR	Rn00562488_s1
α_{2B} -AR	Rn00593312_s1
α_{2C} -AR	Rn00593341_s1
β -actin	Rn00667869_m1

Western blot analysis

20 μ g of protein per well was subjected to electrophoresis on 4%-12% NuPAGE Bis-Tris Gel in XCell SureLock Mini-Cell Units (Life Technologies) (n=5 for each α_2 -AR subtype antagonists). Proteins were transferred from gels to nitrocellulose membranes, using the iBlot Gel Transfer System (Life Technologies). The antibody binding was detected with the WesternBreeze Chromogenic Western blot immunodetection kit (Life Technologies). The blots were incubated on a shaker with α_{2A} -AR, α_{2B} -AR, α_{2C} -AR, and β -actin polyclonal antibody (Santa Cruz Biotechnology, Santa Cruz, CA, USA, 1:200) in the blocking buffer. Images were captured using the EDAS290 imaging system (Csertex Ltd, Budapest, Hungary), and the optical density of each immunoreactive band was determined with Kodak 1D Images analysis software. Optical densities were calculated as arbitrary units after local area background subtraction.

Isolated organ studies

The uteri were removed from the 22-day pregnant rats (250-350 g) (n=8 in each experiment). 5 mm-long muscle rings were sliced from both horns of the uterus and mounted vertically in an organ bath containing 10 mL de Jongh solution (composition: 137 mM NaCl, 3 mM KCl, 1 mM CaCl₂, 1 mM MgCl₂, 12 mM NaHCO₃, 4 mM NaH₂PO₄, 6 mM glucose, pH=7.4). The temperature of the organ bath was maintained at 37°C, and carbogen (95% O₂ + 5% CO₂) was perfused through the bath. After mounting, the rings were allowed to equilibrate for approximately 60 min be-

fore experiments were started, with a buffer change every 15 min. The initial tension of the preparation was set to about 1.5 g and the tension dropped to about 0.5 g by the end of the equilibration period. The tension of the myometrial rings was measured with a gauge transducer (SG-02; Experimetria Ltd, Budapest, Hungary) and recorded with a SPEL Advanced ISOSYS Data Acquisition System (Experimetria Ltd). In the following step contractions were elicited with (-)-noradrenaline (10^{-8} to $10^{-4.5}$ M) and cumulative concentration-response curves were constructed in each experiment in the presence of doxazosin (10^{-7} M) and propranolol (10^{-5} M) in order to avoid α_1 -adrenergic and β -adrenergic actions. Selective α_2 -AR subtype antagonists (each 10^{-7} M), propranolol, and doxazosin were left to incubate for 20 minutes before the administration of contracting agents. Following the addition of each concentration of (-)-noradrenaline, recording was performed for 300 s.

Statistical analysis

Concentration-response curves were fitted and areas under curves (AUC) were evaluated and analyzed statistically with the Prism 4.0 (Graphpad Software Inc. San Diego, CA, USA) computer program. From the AUC values, maximum possible effect (E_{max}) and half maximum effective concentration (EC_{50}) values were calculated. ANOVA Dunnett test or two-tailed unpaired *t* test were used. $P < 0.05$ was considered as a level of significance.

Measurement of uterine cAMP accumulation

Uterine cAMP accumulation was measured with a commercial cAMP Enzyme Immunoassay Kit (Cayman Chemical, Ann Arbor, MI, USA). Uterine tissue samples (control and 17β -estradiol treated) from 22-day-pregnant rats ($n = 6$ in each experiment) were incubated in an organ bath (10 mL) containing de Jongh solution (37°C, perfused with carbogen). Isobuthylmethylxanthine (10^{-3} M), doxazosin (10^{-7} M), propranolol (10^{-5} M) and the investigated subtype-selective α_2 -AR antagonists (each 10^{-7} M) were incubated with the tissues for 20 minutes, and (-)-noradrenaline (3×10^{-6} M) were added to the bath for 10 minutes. At the end of (-)-noradrenaline incubation period, forskolin (10^{-5} M) was added for another 10 min. After stimulation, the samples were immediately frozen in liquid nitrogen and stored until the cAMP extraction (32). Frozen tissue samples were then ground, weighed, homogenized in 10 volumes of ice-cold 5% trichloroacetic acid and centrifuged at 1000g for 10 min. The supernatants were extracted with

3 volumes of water-saturated diethyl ether. After drying, the extracts were stored at -70°C until cAMP assay. Tissue cAMP levels were expressed in pmol/mg tissue.

GTP γ S binding assay

The uteri were removed ($n = 5$ in each experiment) and homogenized in 20 volumes (w/v) of ice-cold buffer (10 mM Tris-HCl, 1 mM EDTA, 0.6 mM MgCl_2 , and 0.25 M sucrose, pH 7.4) with an Ultra Turret T25 (Janke & Kunkel, Staufen, Germany) homogenizer, and the suspension was then filtered on four layers of gauze and centrifuged (40,000g, 4°C, 20 min). After centrifugation, the pellet was resuspended in a 5-fold volume of buffer. The protein contents of the samples were diluted to 10 mg protein/sample. Membrane fractions were incubated in a final volume of 1 mL at 30°C for 60 min in Tris-EGTA buffer (pH 7.4) composed of 50 mM Tris-HCl, 1 mM EGTA, 3 mM MgCl_2 , 100 mM NaCl, containing 20 MBq/0.05 cm^3 [^{35}S]GTP γ S (0.05 nM) (Sigma Aldrich) together with increasing concentrations (10^{-9} - 10^{-5} M) of (-)-noradrenaline. BRL 44408, ARC 239, and spiroxatrine were used in a fixed concentration of 0.1 μM . For the blocking of α_1 - and β -ARs, doxazosin and propranolol were used in a fixed concentration of 10 μM . The determination of total and non-specific binding, filtration, washing procedure, and radioactivity detection were performed (33). The [^{35}S]GTP γ S binding experiments were performed in triplicate and repeated at least three times. G_i protein was inhibited with pertussis toxin (Sigma Aldrich) at a concentration of 500 ng/mL after the addition of protein and GDP to the Tris-EGTA buffer 30 min before [^{35}S]GTP γ S.

RESULTS

RT-PCR and Western blot studies

The mRNA expression of all α_2 -AR subtypes (Figure 1A, 1C, 1E) was significantly decreased ($P < 0.048$) after 17β -estradiol pretreatment compared to non-treated uteri ($P < 0.001$). Western blot analysis at the level of protein expression revealed significant decrease ($P < 0.027$) in each α_2 -AR subtype, corresponding to the PCR results (Figure 2A-F).

Isolated organ studies

In the 22-day-pregnant myometrium, (-)-noradrenaline in the concentration range of 10^{-8} to $10^{-4.5}$ M increased ($P = 0.001$) myometrial contractions (Figure 3A). After 17β -estradiol pretreatment, the myometrial contracting effect

of (-)-noradrenaline was decreased ($P=0.005$). The EC_{50} and E_{max} values of the curves are shown in Table 2.

In the presence of the α_{2A} -AR antagonist BRL 44408, 17 β -estradiol pretreatment increased the (-)-noradrenaline evoked contractions compared to the 17 β -estradiol-treated control ($P=0.004$) (Figure 3B). However, it decreased ($P=0.029$) the myometrial contracting effect of (-)-noradrenaline compared to the BRL 44408-treated control (Table 2).

In the presence of the $\alpha_{2B/C}$ -AR antagonist ARC 239, 17 β -estradiol pretreatment decreased the myometrial contractions compared to the 17 β -estradiol-treated control ($P=0.007$) (Figure 3B) and decreased it ($P=0.045$) compared to the ARC 239-treated control (Table 2).

In the presence of spiroxatrine, 17 β -estradiol increased the maximum contracting effect of (-)-noradrenaline compared to the 17 β -estradiol-treated control ($P<0.001$) (Figure 3B), but decreased it ($P=0.003$) compared to the spiroxatrine-treated control (Table 2).

TABLE 2. Changes in the uterus-contracting effect of (-)-noradrenaline (EC_{50} and E_{max} values) in the absence of α_2 -antagonists, or in the presence of an α_{2A} -antagonist, an $\alpha_{2B/C}$ -antagonist, an α_{2C} -antagonist, or α_{2A} -antagonist and α_{2C} -antagonist in the 22-day-pregnant rat after 17 β -estradiol pretreatment ($n=8$ in each experiment)

	EC_{50} (M \pm SD)	E_{max} (% \pm SD)
Control		
non-treated	$2.6 \times 10^{-6} \pm 6.6 \times 10^{-6}$	274.1 ± 57.8
17 β -estradiol pretreated	$1.5 \times 10^{-6} \pm 1.8 \times 10^{-5}$ ns	88.7 ± 35.5 **
BRL 44408		
non-treated	$1.8 \times 10^{-6} \pm 1.6 \times 10^{-5}$	364.3 ± 83.4
17 β -estradiol pretreated	$2.9 \times 10^{-6} \pm 7.1 \times 10^{-6}$ ns	202.0 ± 59.9 *
ARC 239		
non-treated	$1.2 \times 10^{-6} \pm 2.9 \times 10^{-6}$	147.1 ± 82.0
17 β -estradiol pretreated	$3.5 \times 10^{-6} \pm 7.8 \times 10^{-5}$ ns	55.9 ± 36.7 *
Spiroxatrine		
non-treated	$1.6 \times 10^{-6} \pm 1.2 \times 10^{-5}$	382.4 ± 103.5
17 β -estradiol pretreated	$1.4 \times 10^{-6} \pm 1.5 \times 10^{-6}$ ns	183.7 ± 53.6 *
Spiroxatrine + BRL 44408		
non-treated	$2.9 \times 10^{-6} \pm 1.9 \times 10^{-6}$	444.6 ± 79.7
17 β -estradiol pretreated	$1.1 \times 10^{-6} \pm 4.6 \times 10^{-6}$ ns	111.4 ± 59.0 ***

EC_{50} – the concentration of (-)-noradrenaline alone or in the presence of an α_2 -AR antagonist which elicits half of the maximum contracting effect of (-)-noradrenaline. E_{max} – the maximum contracting effect of (-)-noradrenaline alone or in the presence of an α_2 -AR antagonist. ns – not significant. * $P<0.05$; ** $P<0.01$; *** $P<0.001$. Significance levels were calculated in comparison with non-treated values.

In the presence of the combination of BRL 44408 and spiroxatrine, 17 β -estradiol did not change the maximum myometrial contracting effect of (-)-noradrenaline compared to the 17 β -estradiol-treated control (Fig-




FIGURE 1. Changes in the myometrial mRNA expression of the α_{2A} - (A), α_{2B} - (B), and α_{2C} - adrenergic receptors (ARs) (C) after 17 β -estradiol pretreatment ($n=5$). The statistical analyses were carried out with a two-tailed unpaired t test. ** $P=0.005$; *** $P<0.001$.




FIGURE 2. Changes in the α_2 -adrenergic receptor (AR) levels in the 22-day pregnant rat myometrium after 17 β -estradiol pretreatment (n=5). The α_2 -AR and β -actin Western blot products for α_{2A} - (B), α_{2B} - (D), and α_{2C} -ARs (F). The 70, 62, and 60 kDa proteins relate to α_{2A} -, α_{2B} -, and α_{2C} -ARs and β -actin, respectively. The antibody binding was expressed as optical density (OD) data (A) for α_{2A} -, (C) for α_{2B} -, and (E) for α_{2C} -ARs. The y-axis shows the ratio of α_2 -AR/ β -actin protein optical density. The statistical analyses were carried out with a two-tailed unpaired t test. * $P < 0.027$; *** $P < 0.001$

ure 3B), but decreased it ($P < 0.001$) compared to the BRL 44408+spiroxatrine treated control (Table 2).

cAMP studies

17 β -estradiol pretreatment increased the myometrial cAMP level ($P = 0.007$) (Figure 4) produced in the presence of (-)-noradrenaline. 17 β -estradiol pretreatment also increased the myometrial cAMP level in the presence of (-)-noradrenaline and BRL 44408 ($P = 0.001$), ARC 239 ($P = 0.007$), and spiroxatrine ($P = 0.045$). However, it did not change the cAMP level in the presence of the spiroxatrine + BRL 44408 combination.

[35 S]-GTP γ S binding assay studies

In the presence of BRL 44408, (-)-noradrenaline increased the [35 S]GTP γ S binding, which was significantly decreased after 17 β -estradiol pretreatment ($P = 0.038$). In the presence of pertussis toxin, the [35 S]GTP γ S binding-stimulating effect of (-)-noradrenaline ceased, and 17 β -estradiol pretreatment did not modify this effect (Figure 5A).

In the presence of ARC 239, (-)-noradrenaline increased ($P < 0.001$) the [35 S]GTP γ S binding similarly to 17 β -estradiol pretreatment. In the presence of pertussis toxin, (-)-noradrenaline slightly decreased the [35 S]GTP γ S bind-

ing, which was not changed after 17 β -estradiol pretreatment (Figure 5B).

In the presence of spiroxatrine, (-)-noradrenaline increased the [35 S]GTP γ S binding ($P < 0.001$), which was slightly decreased ($P = 0.037$) after 17 β -estradiol pretreatment. In the presence of pertussis toxin, however, (-)-noradrenaline decreased the [35 S]GTP γ S binding below the basal level from a concentration of 1×10^{-9} M ($P < 0.001$). In the presence of pertussis toxin, 17 β -estradiol pretreatment abolished the [35 S]GTP γ S binding-inhibitory effect of (-)-noradrenaline (Figure 5C).

In the presence of spiroxatrine+BRL 44408 combination, (-)-noradrenaline inhibited the [35 S]GTP γ S binding, and 17 β -estradiol further inhibited the [35 S]GTP γ S binding of (-)-noradrenaline and abolished the dose-dependency of noradrenalin action. In the presence of pertussis toxin, the spiroxatrine+BRL 44408 combination dose-dependently inhibited the [35 S]GTP γ S binding of (-)-noradrenaline, similarly to 17 β -estradiol pretreatment (Figure 5D).

DISCUSSION

Since estrogens and the adrenergic system play a major role in myometrial contractions during human gestation, the main focus of our study was to clarify the ef-

fects of estrogen on the α_2 -AR subtypes in late pregnant uterine function *in vitro*. The estrogen- α_2 -AR connection was investigated via the effects of subtype-selective antagonists after 17 β -estradiol pretreatment on the (-)-noradrenaline-stimulated contractions. The experiments were carried out in the presence of the α_1 -AR blocker doxazosin and the β -AR blocker propranolol in order to avoid α_1 - or β -adrenergic actions.




FIGURE 3. Effects of the subtype-selective α_{2A} -adrenoceptor antagonist BRL 44408, the $\alpha_{2B/C}$ -adrenoceptor antagonist ARC 239, and the α_C -adrenoceptor antagonist, spiroxatrine on the (-)-noradrenaline-evoked contractions in the 22-day-pregnant rat myometrium (A) and after 17 β -estradiol pretreatment (B) (n=8). The studies were carried out in the presence of the β -adrenoceptor antagonist, propranolol (10^{-5} M) and the α_1 -adrenoceptor antagonist, doxazosin (10^{-7} M) in each case. The change in contraction was calculated using the area under the curve and expressed in % \pm SEM. The statistical analyses were carried out with the ANOVA Dunnett test. * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$.

17 β -estradiol pretreatment decreased the mRNA and protein expression of the myometrial α_2 -AR subtypes and (-)-noradrenaline-evoked myometrial contraction via the α_2 -ARs, which is similar to our earlier findings with α_{1A} -ARs (14). According to these findings, we can claim that estrogen differently affects the expression of the α_2 -ARs in various tissues, as it increases the expression of the receptors in the spinal cord and cutaneous arteries (27,28).

In isolated organ bath studies, 17 β -estradiol pretreatment decreased (-)-noradrenaline-evoked myometrial contractions via the α_2 -ARs, although it did not modify the myometrial relaxing effect via the α_{2A} -ARs. However, it abolished the myometrial contraction-increasing effect via the α_{2B} -ARs. Since there are no available antagonists to produce only α_{2C} -AR stimulation (ie, $\alpha_{2A/B}$ -AR blockers), we can only presume that 17 β -estradiol did not modify the myometrial relaxing effect via the α_{2C} -ARs.

To explain why weaker myometrial contractions via the α_{2B} -AR subtype occurred after 17 β -estradiol pretreatment, we measured the myometrial cAMP level, as the changes in the cAMP level are involved in the myometrial effect of the α_2 -ARs. 17 β -estradiol pretreatment increased the myometrial cAMP level, which also proves the decreased myometrial contracting effect of (-)-noradrenaline via the α_2 -ARs. It did not modify the cAMP level via the α_{2A} -ARs, which is in accordance with our previous study (23). However, it increased the myometrial cAMP level via the α_{2B} -ARs, which can explain the weaker myometrium contracting effect of (-)-noradrenaline.




FIGURE 4. Effects of the subtype-selective α_{2A} -adrenoceptor antagonist, BRL 44408, the $\alpha_{2B/C}$ -adrenoceptor antagonist, ARC 239, and the α_{2C} -adrenoceptor antagonist, spiroxatrine on the myometrial cAMP level (pmol/mg tissue \pm standard deviation) in the presence of 3-isobutyl-1-methylxanthine (IBMX) (10^{-3} M) and forskolin (10^{-5} M) (control) in the 22-day-pregnant rat (n=6) after 17 β -estradiol pretreatment. The statistical analyses were carried out with ANOVA followed by Dunnett's Multiple Comparison Test. * $P = 0.046$, ** $P < 0.007$, *** $P = 0.001$.




FIGURE 5. Changes induced by various concentrations of noradrenaline in $[^{35}\text{S}]\text{GTP}\gamma\text{S}$ binding in the presence of subtype-selective α_{2A} -antagonist BRL 44408 (A), the $\alpha_{2B/C}$ -antagonist ARC 239 (B), the α_{2C} -antagonist spiroxatrine (C), and the BRL 44408-spiroxatrine combination (D) following pretreatment with 17 β -estradiol ($n=5$). In all cases, the β -adrenoceptors and α_1 -adrenoceptors were inhibited by propranolol and doxazosin. Basal refers to the level of $[^{35}\text{S}]\text{GTP}\gamma\text{S}$ binding without substance. The statistical analyses were carried out with a two-tailed unpaired t test. * $P < 0.038$; ** $P < 0.004$, *** $P < 0.001$.

The α_2 -ARs can couple not only to the G_i protein α -subunit, but under certain circumstances, also to G_s proteins (24). Estrogen was also shown to decrease the coupling of the α_2 -adrenoceptors to G protein (29). To find an explanation for the cAMP changes, we measured the myometrial $[^{35}\text{S}]\text{GTP}\gamma\text{S}$ binding of the α_2 -AR subtypes after 17 β -estradiol pretreatment and in the presence of pertussis toxin, whose inhibitory action is specific for the G_i protein. In the presence of pertussis toxin, 17 β -estradiol did not modify the $[^{35}\text{S}]\text{GTP}\gamma\text{S}$ binding of the α_{2A} -ARs, but it reversed the effect of (-)-noradrenaline on $[^{35}\text{S}]\text{GTP}\gamma\text{S}$ binding via the α_{2A} - and α_{2B} -ARs (with spiroxatrine). These findings show that 17 β -estradiol modifies the coupling of the α_{2B} -ARs, but does not change the G protein binding of the α_{2A} -ARs. To prove this hypothesis, we measured the myometrial $[^{35}\text{S}]\text{GTP}\gamma\text{S}$ binding of the α_{2B} -AR subtype in the presence of

spiroxatrine+BRL 44408. 17 β -estradiol decreased the amount of activated G-protein, which is probably a consequence of 17 β -estradiol-induced uncoupling of α_{2B} -ARs from the G proteins (29). This process did not change myometrial contraction as compared with the hormone-treated control.

In the light of our results, we conclude that the functions of the α_2 -AR subtypes are influenced by the female sexual steroid, 17 β -estradiol. It decreases the expressions of the α_2 -AR subtypes and increases uterine cAMP level. It does not modify the myometrial relaxing effect via the α_{2A} - and α_{2C} -ARs. In case of these receptors we suppose that the 17 β -estradiol treatment mainly induces the activation of $\beta\gamma$ subunit of G_i protein, increasing the smooth muscle cAMP level (17). In case of α_{2B} -ARs, 17 β -estradiol alters the

myometrial contracting effect of (-)-noradrenaline by reduced coupling of the receptor to G_i protein.

A limitation of our study is that we did not carry out any studies on human myometrium, and there might be differences in the function of the rat and human myometrial α₂-AR subtypes. However, our present findings give a better understanding on the complex physiology of changes during pregnancy, as estrogen is the predominantly expressed hormone during human parturition at term (15,34), which, together with the α₂-ARs, plays an essential role in myometrial contractility. It was also demonstrated that estrogen level in the amniotic fluid was elevated in uterine inertia (35), which might be caused by the decreased myometrial contractility via the α₂-AR subtypes. Therefore, estrogen level dysregulation during pregnancy might change the function of the α₂-AR subtypes and result in either preterm labor or labor delay. We would like to extend these preclinical studies for premature birth models in rats. We suppose that either subtype-specific agonists or antagonists can be used as a target for drugs against abnormal myometrial contractility.

Acknowledgment The study was supported by Cedars Sinai Medical Center's International Research and Innovation in Medicine Program, the Association for Regional Cooperation in the Fields of Health, Science and Technology (RECOOP HST Association) and the participating Cedars-Sinai Medical Center - RECOOP Research Centers (CRRC).

Funding The study was supported by a grant from the National Research, Development and Innovation Office (NKFI), Budapest, Hungary; OTKA-108518.

Ethical approval received from the Hungarian Ethics Committee for Animal Research (registration number: IV/198/2013).

Declaration of authorship JHT wrote the manuscript and performed the experiments. JB performed the contractility studies and cAMP determination. ED performed the RT-PCR and Western blot studies. AC performed the RT-PCR studies. ZT performed the cAMP determination. RS performed the [³⁵S]-GTPγS binding assay studies. AB designed [³⁵S]-GTPγS binding assay studies. SB performed and supervised the [³⁵S]-GTPγS binding assay studies. RG supervised and organized all experiments, wrote the manuscript, and is the corresponding author.

Competing interests All authors have completed the Unified Competing Interest form at www.icmje.org/coi_disclosure.pdf (available on request from the corresponding author) and declare: no support from any organization for the submitted work; no financial relationships with any organizations that might have an interest in the submitted work in the previous 3 years; no other relationships or activities that could appear to have influenced the submitted work.

References

- Flenady V, Wojcieszek AM, Papatsonis DN, Stock OM, Murray L, Jardine LA, et al. Calcium channel blockers for inhibiting preterm labour and birth. *Cochrane Database Syst Rev*. 2014;6:CD002255. [Medline:24901312](#)
- Tan MY, To M. Recent advances in the prevention and management of preterm birth. *F1000Prime Rep*. 2015;7:40. [Medline:26097713](#)
- doi:10.12703/P7-40
- Illanes SE, Pérez-Sepúlveda A, Rice GE, Mitchell MD. Preterm labour: association between labour physiology, tocolysis and prevention. *Expert Opin Investig Drugs*. 2014;23:759-71. [Medline:24717074](#) [doi:10.1517/13543784.2014.905541](#)
- Arora CP, Kacerovsky M, Zinner B, Ertl T, Ceausu I, Rusnak I, et al. Disparities and relative risk ratio of preterm birth in six Central and Eastern European centers. *Croat Med J*. 2015;56:119-27. [Medline:25891871](#) [doi:10.3325/cmj.2015.56.119](#)
- Fyfe EM, Anderson NH, North RA, Chan EH, Taylor RS, Dekker GA, et al. Risk of first-stage and second-stage cesarean delivery by maternal body mass index among nulliparous women in labor at term. *Obstet Gynecol*. 2011;117:1315-22. [Medline:21606741](#) [doi:10.1097/AOG.0b013e318217922a](#)
- Borda E, Sauvage J, Sterin-Borda L, Gimeno MF, Gimeno AL. Adrenoceptors involved in the contractile activity of isolated pregnant rat uterus. *Eur J Pharmacol*. 1979;56:61-7. [Medline:223858](#) [doi:10.1016/0014-2999\(79\)90433-3](#)
- Chaud M, Gonzalez ET, Franchi AM, Gimeno MF, Gimeno AL. Sex hormones and the motility of and prostaglandin output from, uterine horns of immature rats. *Prostaglandins Leukot Med*. 1984;15:35-44. [Medline:6591213](#) [doi:10.1016/0262-1746\(84\)90054-4](#)
- Clabaut M, Duclos R, Chartrel N, Adjroud O, Acritopoulou-Fourcroy S, Collado H, et al. Variation of myometrial activities and steroid sexual hormones following bilateral ovariectomy in the rat at midpregnancy. *Acta Physiol Hung*. 1988;71:477-83. [Medline:2849861](#)
- Roberts JM, Riemer RK, Bottari SP, Wu YY, Goldfien A. Hormonal regulation myometrial adrenergic responses: the receptor and beyond. *J Dev Physiol*. 1989;11:125-34. [Medline:2553793](#)
- Dowell RT, Forsberg AL, Kauer CD. Decreased ovarian blood flow may confound the tocolytic effect of ritodrine. *Gynecol Obstet Invest*. 1994;37:168-71. [Medline:8005545](#) [doi:10.1159/000292551](#)
- Engstrom T, Vilhardt H, Bratholm P, Christensen NJ. Desensitization of beta2-adrenoceptor function in non-pregnant rat myometrium is modulated by sex steroids. *J Endocrinol*. 2001;170:147-55. [Medline:11431147](#) [doi:10.1677/joe.0.1700147](#)
- Roberts JM, Riemer RK, Bottari SP, Wu YY, Goldfien A. Hormonal regulation myometrial adrenergic responses: the receptor and beyond. *J Dev Physiol*. 1989;11:125-34. [Medline:2553793](#)
- Gáspár R, Ducza E, Mihályi A, Márki A, Kolarovszki-Sipiczki Z, Páldy E, et al. Pregnancy-induced decrease in the relaxant effect of terbutaline in the late-pregnant rat myometrium: role of G-protein activation and progesterone. *Reprod*. 2005;130:113-22. [Medline:15985637](#) [doi:10.1530/rep.1.00490](#)
- Bóta J, Hajagos-Tóth J, Ducza E, Samavati R, Borsodi A, Benyhe S, et al. The effects of female sexual hormones on the expression and function of α1A- and α1D-adrenoceptor subtypes in the late-pregnant rat myometrium. *Eur J Pharmacol*. 2015;769:117-84.

- Medline:26593425 doi:10.1016/j.ejphar.2015.11.015
- 15 Kamel RM. The onset of human parturition. *Arch Gynecol Obstet.* 2010;281:975-82. Medline:20127346 doi:10.1007/s00404-010-1365-9
 - 16 Renthall NE, Williams KC, Montalbano AP, Chen CC, Gao L, Mendelson CR. Molecular regulation of parturition: a myometrial perspective. *Cold Spring Harb Perspect Med.* 2015;5(11). Medline:26337112 doi:10.1101/cshperspect.a023069
 - 17 Zhou XB, Wang GX, Huneke B, Wieland T, Korth M. Pregnancy switches adrenergic signal transduction in rat and human uterine myocytes as probed by BKCa channel activity. *J Physiol.* 2000;524:339-52. Medline:10766916 doi:10.1111/j.1469-7793.2000.t01-1-00339.x
 - 18 Knaus AE, Muthig V, Schickinginger S, Moura E, Beetz N, Gilsbach R, et al. Alpha2-adrenoceptor subtypes—unexpected functions for receptors and ligands derived from gene-targeted mouse models. *Neurochem Int.* 2007;51:277-81. Medline:17664025 doi:10.1016/j.neuint.2007.06.036
 - 19 Civantos Calzada B, Alexandre de Artiñano A. Alpha-adrenoceptor subtypes. *Pharmacol Res.* 2001;44:195-208. Medline:11529686 doi:10.1006/phrs.2001.0857
 - 20 Karim F, Roerig SC. Differential effects of antisense oligodeoxynucleotides directed against Gza and Goa on antinociception produced by spinal opioid and α 2 adrenergic receptor agonists. *Pain.* 2000;87:181-91. Medline:10924811 doi:10.1016/S0304-3959(00)00279-7
 - 21 Wang Q. α 2-adrenergic receptors. *Primer on the Autonomic Nervous System.* 2012;(10)55-58.
 - 22 Bouet-Alard R, Mhaouty-Kodja S, Limon-Boulez I, Coudouel N, Maltier JP, Legrand C. Heterogeneity of alpha 2-adrenoceptors in human and rat myometrium and differential expression during pregnancy. *Br J Pharmacol.* 1997;122:1732-8. Medline:9422821 doi:10.1038/sj.bjpp.0701555
 - 23 Gáspár R, Gál A, Gálík M, Ducza E, Minorics R, Kolarovszki-Sipiczki Z, et al. Different roles of alpha2-adrenoceptor subtypes in non-pregnant and late-pregnant uterine contractility in vitro in the rat. *Neurochem Int.* 2007;51:311-8. Medline:17664026 doi:10.1016/j.neuint.2007.06.029
 - 24 Offermanns S. G-proteins as transducers in transmembrane signalling. *Prog Biophys Mol Biol.* 2003;83:101-30. Medline:12865075 doi:10.1016/S0079-6107(03)00052-X
 - 25 Mhaouty S, Cohen-Tannoudji J, Bouet-Alard R, Limon-Boulez I, Maltier JP, Legrand C. Characteristics of the alpha 2/beta 2-adrenergic receptor-coupled adenylyl cyclase system in rat myometrium during pregnancy. *J Biol Chem.* 1995;270:11012-6. Medline:7738044 doi:10.1074/jbc.270.18.11012
 - 26 Marshall JM. Effects of ovarian steroids and pregnancy on adrenergic nerves of uterus and oviduct. *Am J Physiol.* 1981;240:C165-74. Medline:7015875 doi:10.1113/jphysiol.1981.sp013899
 - 27 Thompson AD, Angelotti T, Nag S, Mokha SS. Sex-specific modulation of spinal nociception by alpha2-adrenoceptors: differential regulation by estrogen and testosterone. *Neurosci.* 2008;153:1268-77. Medline:18434028 doi:10.1016/j.neuroscience.2008.03.008
 - 28 Eid AH, Maiti K, Mitra S, Chotani MA, Flavahan S, Bailey SR, et al. Estrogen increases smooth muscle expression of alpha2C-adrenoceptors and cold-induced constriction of cutaneous arteries. *Am J Physiol Heart Circ Physiol.* 2007;293:H1955-61. Medline:17644575 doi:10.1152/ajpheart.00306.2007
 - 29 Ansonoff MA, Etgen AM. Receptor phosphorylation mediates estradiol reduction of alpha2-adrenoceptor coupling to G protein in the hypothalamus of female rats. *Endocr.* 2001;14:165-74. Medline:11394633 doi:10.1385/ENDO:14:2:165
 - 30 Hódi A, Földesi I, Ducza E, Hajagos-Tóth J, Seres AB, Klukovits A, et al. Tocopherol inhibits the relaxing effect of terbutaline in the respiratory and reproductive tracts of the rat: the role of the oxidative stress index. *Life Sci.* 2014;105:48-55. Medline:24780318 doi:10.1016/j.lfs.2014.04.023
 - 31 Chomczynski P, Sacchi N. Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal Biochem.* 1987;162:156-9. Medline:2440339 doi:10.1016/0003-2697(87)90021-2
 - 32 Hajagos-Tóth J, Hódi Á, Seres AB, Gáspár R. Effects of d- and l-limonene on the pregnant rat myometrium in vitro. *Croat Med J.* 2015;56:431-8. Medline:26526880 doi:10.3325/cmj.2015.56.431
 - 33 Zádor F, Kocsis D, Borsodi A, Benyhe S. Micromolar concentrations of rimonabant directly inhibits delta opioid receptor specific ligand binding and agonist-induced G-protein activity. *Neurochem Int.* 2014;67:14-22. Medline:24508403 doi:10.1016/j.neuint.2013.12.005
 - 34 Romero R, Scoccia B, Mazor M, Wu YK, Benveniste R. Evidence for a local change in the progesterone/estrogen ratio in human parturition at term. *Am J Obstet Gynecol.* 1988;159:657-60. Medline:2971319 doi:10.1016/S0002-9378(88)80029-2
 - 35 Yan J, Chen W. Observations on serum and amniotic fluid hormone levels and ultrastructure of uterine myometrium before and during term labor. *Zhonghua Fu Chan Ke Za Zhi.* 1995;30:391-4. Medline:7587572