

EFFECTS OF STARCHY AND β -GLUCAN ADDITIVES ON FLOUR, DOUGH, AND BREAD PARAMETERS

S. GAVURNÍKOVÁ^{a,b}, M. HAVRENTOVÁ^a and J. KRAIC^{a,c*}

^aPlant Production Research Center Piešťany, Bratislavská cesta 122, 92168 Piešťany. Slovakia

^bConstantine the Philosopher University, Trieda Andreja Hlinku 1, 94974 Nitra. Slovakia

^cUniversity of SS. Cyril and Method, Námestie J. Herdu 2, 91701 Trnava. Slovakia

(Received: 25 July 2012; accepted: 4 February 2013)

Composite flours were formulated from wheat flour and additives containing high amylose starch, resistant starches of RS₂ and RS₃ types, and barley β -glucan. Different parameters of flours, doughs, and final breads were evaluated. Almost all composite flours had significantly worse parameters as flour and dough in comparison to control. Sensory parameters of breads were also lower, though loaves supplemented with up to 15% (w/w) of high amylose starch (Hylon® VII), RS₂ (Hi-maize™ 260), and RS₃ (Novelose® 330) were considered as acceptable, with higher content of RS observed. Loaves with β -glucan (Barliv™ barley betafiber) were not acceptable either in sensory or technological parameters.

Keywords: wheat, flour, dough, bread, resistant starch, β -D-glucan, functional food

Both the sufficiency and quality of food influence the physiological state of the consumer. A relevant trend is the supplementation of foods with compounds uncovered by the daily nutrition and the production of functional foods. Foods are functional when it is possible to evidently demonstrate their beneficial effect in amounts normally consumable in the daily diet. From practical point of view, the functional food could be food with compounds added or replaced to achieve the benefit. Food additives usually change technological parameters, content of nutrients, and final product traits; in wheat it is gluten quality, rheological parameters, and bread acceptability. Attractive additives from nutritional point of view are resistant starches (RS) and β -glucans (BG). RSs are resistant to absorption in the small intestine and provide substrate for microbial fermentation in the large intestine resulting in beneficial short-chain fatty acids. ENGLYST and co-workers (1992) described RS₁ (unavailable for enzymatic degradation in gastrointestinal tract), RS₂ (naturally resistant to degradation), RS₃ (created by hydrothermal processing), and RS₄ (chemically modified). β -glucan affects human body as an activator of immune processes (ESTRADA et al., 1997) or an agent reducing the risk of cardio-vascular diseases and gastrointestinal problems (MÄLKKI & VIRTANEN, 2001; KEOGH et al., 2003).

Different additives supplied to basic wheat flour essentially influence parameters of flour, dough, and the final product. Therefore, the aim of this study was to evaluate the effect of different additives (high amylose starch and resistant starches from maize, barley β -glucan) on parameters of composite flours, dough, and bread.

* To whom correspondence should be addressed.

Phone: +421 33 7722330; fax: +421 33 7726306; e-mail: kraic@vurv.sk

1. Materials and methods

The basic component of experimental flours was soft bread wheat flour (type T 650 from the mill Penam a. s., Trnava, Slovakia) containing proteins 11.3%, carbohydrates 73.3%, lipids 1.5%, ash $\leq 0.78\%$ with wet gluten $\geq 26\%$ and falling number ≥ 170 s. Commercial additives were resistant starches Hi-maize™ 260 (RS₂) and Novelse® 330 (RS₃), high amylose (min. 68%) maize starch Hylon® VII (all from National Starch & Chemical Co., Bridgewater, USA), and Barliv™ barley betafiber (concentrated barley β -glucan) (Cargill Inc., Minneapolis, USA). The weight ratios (w/w) of additives in composite flours were 5–20%.

Wet gluten content and gluten swelling (STN, 1988), total ash (STN ISO, 2006a), Zeleny's sedimentation index (STN ISO, 2000), falling number (STN ISO, 2006b), farinographic parameters (ICC, 1992), and final bread, made from flour 250 g, yeasts 12.5 g, sugar 2.5 g, salt 3.75 g, fat 2.5 g, and water, were evaluated. Flour and dough were blended (mixer RM 800 A-B, RM Gastro s.r.o., Veselí n. L., Czech Republic), fermented for 20 min at 32 ± 1 °C, rolled, leavened for 10 min, left rising for 25 min at 32 ± 1 °C, and baked for 20 min at 230 °C (Domino modular oven, Marton, Slovakia). Loaves were evaluated for specific loaf volume and weight. Sensory parameters of breads, crust (colour, thickness, and firmness), crumb (colour, hardness, size, and uniformity of crumb porosity), aroma, and taste were evaluated 2 h after baking by the 5-point hedonic scale (POKORNÝ, 1997) by seven educated panellists. The moisture content on the 1st and 2nd d after baking was evaluated according to STN (1990) after pre-drying up to 45 °C, melting, and drying at 130 °C for 60 min. The total, soluble, and resistant starch, and β -glucan content were determined by kits K-TSTA, K-RSTAR, K-BGLU (Megazyme Int. Ireland Ltd., Wicklow, Ireland) based on relevant official methods A.A.C.C. (2000; 2002; 2003), A.O.A.C. (1995; 1996; 2002), and ICC Standard (1998), respectively.

Data were evaluated by analysis of variance (ANOVA), multiple comparisons by LSD test by software Statgraphic Plus 7.0. Superscripts used in tables and figures mean statistically significant differences ($P < 0.05$).

2. Results and discussion

2.1. Qualitative parameters of composite flours

Additions of Hylon® VII (HAS), Hi-maize™ 260 (RS₂), Novelse® 330 (RS₃), and Barliv™ barley betafiber (BG) changed the basic parameters of flours. The total ash content significantly decreased with the increased ratio of HAS and both RSs and increased with the increased ratio of BG in the flour reflecting content of incombustible compounds supplied by the additives. All agents significantly decreased wet gluten content as a consequence of the dilution of glutinous proteins, though not below 25 ml in case of either starchy agent. BG affected the wet gluten content more negatively and in flours with $\geq 15\%$ of BG gluten was not possible to be determined. The gluten swelling significantly decreased at the addition of 20% of RS₃ and at all ratios of RS₂. Only $\geq 15\%$ of HAS increased gluten swelling significantly. BG decreased gluten swelling and at $\geq 15\%$ this parameter was not possible to be determined. The reason could be the change of the viscoelastic structure of gluten to a gel-like state due to the presence of β -glucan (IRAKLI et al., 2004). RS₃ significantly increased the sedimentation index. RS₂ significantly and HAS linearly decreased the sedimentation index, though neither decreased its value below 22 ml (minimal value required by norm STN, 2003). Low

concentration (5%) of BG increased the sedimentation index, but higher (15–20%) concentration decreased its value below 22 ml. Composite flours with all added agents had significantly lower falling number than basic wheat flour. Addition of 20% RS₂ decreased falling number under the critical value (220 s, STN 46 1100-2., 2003). All ratios of BG affected the falling number very negatively ($\geq 10\%$ below 220 s).

2.2. Rheological parameters

Increased ratios of all additives increased water absorption related to the high water binding capacity of non-starch polysaccharides (SKENDI et al., 2010), similarly to fibres due to the high number of hydroxyl-groups in their structure (WANG et al., 2002).

The dough development time and dough stability were reduced by all added starchy agents. The addition of 20% RS₂ and $\geq 15\%$ HAS had the highest negative effect on dough stability as a consequence of the dilution effect of gluten-creating proteins and the weakening of gluten structure. On the contrary, addition of BG significantly increased dough development time and decreased dough stability in comparison to control. SKENDI and co-workers (2010) published that addition of β -glucan to weak wheat flour increased development time and improved dough stability, but in strong flour both parameters were increased. MOHAMED and co-workers (2005) mentioned that increasing of β -glucan content did not significantly reduce dough stability. The difference could be explained by different origin, molecular weight, solubility, and concentration of β -glucan, and also by the different origin (cultivar) and type of basic wheat flour. The dough softening parameters generally increased with the addition of HAS, RS, and BG. The higher was their amount the higher were the dough softening parameters. The effect of RS₂ was higher than that of RS₃. The farinograph quality numbers were lower at all ratios of added HAS and RS and also at $\geq 10\%$ of BG than those of the control. Rheological evaluation confirmed the behaviour of non-interacting polysaccharides and the transition of dough properties from the desired viscoelastic to gel-like structure.

2.3. Bread quality parameters

The weight of loaves with added HAS, RS, and BG were statistically significantly higher than the control due to higher dough water absorption. The specific volumes of loaves in all samples were lower than in control, the lowest with the addition of HAS, the highest with the addition of $\geq 10\%$ of RS₃. Loaf volume reduction was expected as amylolytic enzymes cannot digest RS to sugars available for fermentation and CO₂ production. Addition of BG had the strongest negative effect on weight and volume and loaves with $\geq 10\%$ of BG were non-acceptable. JACOBS and co-workers (2008) reported that 1% of β -glucan in wheat flour reduced loaf volume by 15%. Each percentage of added BG in our study reduced loaf volume by $\approx 3.5\%$.

All breads containing HAS, RS, and BG had higher water content in crumb on the 1st and 2nd day after baking in comparison to control. Water loss from the 1st to 2nd d was 4.0% in the control bread, 1.0–2.6% in breads with HAS, 0.0–4.0% with RS₂, and 1.4–4.1% with RS₃. Lower water loss resulted in finer bread crust. Breads with HAS and low amounts of RS and BG maintained water better than control wheat bread and were fresh for a longer time.

The sensory evaluation revealed that additions of HAS and RS influenced crust and crumb colour. The higher the ratios of HAS or RS the brighter both crust and crumb were. At $\geq 15\%$ of HAS or RS crumbs were less elastic, tougher, more sticky, the taste and aroma also changed. Significant differences were in the sensory evaluation of breads supplemented with $\geq 15\%$ of RS₃, 20% of RS₂, and with all ratios of BG. All were less accepted by panellists,

but they expressed that bread containing RS₃ up to 15% were very tasty. Addition of 5% of HAS decreased sensory parameters only slightly, higher ratios did not change sensory quality, nevertheless breads with $\geq 15\%$ of HAS were less tasty. SANZ and co-workers (2009) performed similar experiment with RS₂ and RS₃ in muffins, where RS₃ had higher impact on the texture, and muffins were less accepted, but RS₂ supplementation did not cause significant differences in the acceptability of the muffins. ÖZTURK and co-workers (2009) tested the addition of three agents containing RS₃ and found that loaf volume was reduced less significantly by Novelose® 330, and the RS₃ agents had no negative influence on crumb colour and bread appearance.

None of the loaves with BG were accepted by panellists due to sticky crumb, taste, and appearance. Limited potential of β -glucan extract supplemented into wheat bread has been reported by HAGER and co-workers (2011). However, β -glucan added to composite flour in its natural form could have different effect than as extract or commercial agent. KNUCKLES and co-workers (1997) evaluated sensory characteristics of wheat pasta and bread supplemented with fractions of barley flour fortified with β -glucan in different ratios. The supplementation increased water absorption, dough development and blending time, however, breads with $\leq 20\%$ of barley flour with β -glucan were acceptable for consuming.




Fig. 1. Sensory parameters of breads baked from composite flours supplemented with agents containing HAS, RS, and BG. ▲: Hylon VII; ■: Hi-maize 260; ◆: Novelose 330; ×: Barliv barley betafiber

2.4. Total, soluble, and resistant starch and β -glucan content in breads

Contents of resistant, soluble, and total starch in flours and breads are shown in Tables 1 and 2. The RS content was significantly increased in flours and breads by the addition of starchy agents. The highest RS contents were in flours supplemented with HAS but the baking process reduced it in the breads. The addition of RS₂ and RS₃ increased the RS content in flour, which was enhanced by the baking process. The supplementation with RS₃ increased

the content of RS in bread more than 5 times compared to flour. Similar trend was detected by HUNG and co-workers (2005) at the addition of high amylose wheat flour. The content of RS in foods was generally in range 0–4%. Breads non-supplemented with RS contain 1–2.5% of RS (TAS & EL, 2000) but daily intake of RS should be ≥ 17 g (LONDON, 2007). This relates to 200 g of bread with added 15% of Novelose® 330 in our study.

Soluble starch content decreased with the addition of HAS, RS₂, and RS₃ in flours and more or less also in breads. The total starch content in flours and breads increased ambiguously with the supplementation.

The content of β -glucan in flours after addition of BG was in the range of 4.6–13.5% (Table 3) and in baked breads decreased only slightly due to hydrolysis or reduction of some dietary fibre by the high temperature during baking. Consumption of β -glucan reducing total blood cholesterol should be ≥ 3 g daily (FDA, 1997). The β -glucan content in regular wheat bread is about 1.4% in dry bread weight or 0.63% in fresh bread (HAGER et al., 2011). Approximately 100 g of bread made in our study with the addition of 5% of BG to basic wheat flour contains three grams of β -glucan.

Table 1. Content of resistant, soluble, and total starch in flours

Flours	Resistant starch (%)		Soluble starch (%)		Total starch (%)	
	x	SD	x	SD	x	SD
Control wheat flour	0.39	± 0.00 ^a	73.44	± 0.22 ^g	73.83	± 0.22 ^{ab}
5% Novelose® 330	2.66	± 0.01 ^b	70.57	± 0.13 ^{ef}	73.23	± 0.13 ^a
10% Novelose® 330	4.84	± 0.04 ^d	70.41	± 0.36 ^{de}	75.25	± 0.34 ^{bcd}
15% Novelose® 330	7.19	± 0.14 ^e	68.06	± 0.25 ^b	75.25	± 0.16 ^{bcd}
20% Novelose® 330	9.06	± 0.12 ^f	68.24	± 0.36 ^b	77.30	± 0.44 ^f
5% Hi-maize™ 260	2.68	± 0.09 ^b	72.55	± 0.80 ^g	75.23	± 0.87 ^{bc}
10% Hi-maize™ 260	5.19	± 0.15 ^d	71.50	± 0.65 ^f	76.68	± 0.77 ^{def}
15% Hi-maize™ 260	7.53	± 0.02 ^e	69.45	± 0.04 ^{cd}	76.97	± 0.06 ^{ef}
20% Hi-maize™ 260	8.58	± 0.13 ^f	67.80	± 0.75 ^b	76.38	± 0.88 ^{cdef}
5% Hylon® VII	3.67	± 0.14 ^c	69.88	± 0.88 ^{de}	73.55	± 0.82 ^a
10% Hylon® VII	7.18	± 0.19 ^e	68.44	± 0.22 ^{bc}	75.62	± 0.38 ^{cde}
15% Hylon® VII	11.05	± 0.09 ^g	68.16	± 0.00 ^b	79.21	± 0.09 ^g
20% Hylon® VII	13.54	± 0.42 ^h	62.54	± 0.80 ^a	76.07	± 0.72 ^{cdef}

Superscripts represent statistically significant differences at $P < 0.05$.

3. Conclusions

Analysed qualitative parameters of flour, rheological parameters of dough, quality parameters of bread, contents of total, soluble, and resistant starch and β -glucan were influenced by the addition of commercial agents containing high amylose starch, resistant starches, and barley

Table 2. Content of resistant, soluble, and total starch in baked breads

Samples	Resistant starch (%)		Soluble starch (%)		Total starch (%)	
	x	SD	x	SD	x	SD
Control wheat flour	2.07 \pm 0.03 ^a		69.67 \pm 0.19 ⁱ		71.74 \pm 0.17 ^{bc}	
5% Novelose [®] 330	4.53 \pm 0.02 ^d		67.09 \pm 0.19 ^{fg}		71.62 \pm 0.17 ^b	
10% Novelose [®] 330	6.60 \pm 0.07 ^e		65.48 \pm 0.25 ^c		72.08 \pm 0.28 ^{bc}	
15% Novelose [®] 330	8.75 \pm 0.29 ⁱ		66.36 \pm 0.20 ^{def}		75.11 \pm 0.46 ^e	
20% Novelose [®] 330	10.81 \pm 0.16 ^l		61.75 \pm 0.45 ^a		72.56 \pm 0.28 ^c	
5% Hi-maize [™] 260	3.95 \pm 0.03 ^c		66.26 \pm 0.09 ^{cde}		70.21 \pm 0.06 ^a	
10% Hi-maize [™] 260	5.95 \pm 0.04 ^f		65.85 \pm 0.42 ^{cd}		71.80 \pm 0.41 ^{bc}	
15% Hi-maize [™] 260	8.00 \pm 0.04 ⁱ		68.01 \pm 0.19 ^h		76.01 \pm 0.23 ^{ef}	
20% Hi-maize [™] 260	9.53 \pm 0.11 ^k		64.01 \pm 0.36 ^b		73.53 \pm 0.46 ^d	
5% Hylon [®] VII	3.58 \pm 0.05 ^b		67.04 \pm 0.45 ^{efg}		70.62 \pm 0.50 ^a	
10% Hylon [®] VII	5.02 \pm 0.05 ^e		67.34 \pm 0.42 ^{gh}		72.37 \pm 0.47 ^{bc}	
15% Hylon [®] VII	7.33 \pm 0.08 ^h		69.13 \pm 0.16 ⁱ		76.46 \pm 0.14 ^f	
20% Hylon [®] VII	7.83 \pm 0.02 ⁱ		67.44 \pm 0.82 ^{gh}		75.27 \pm 0.84 ^e	

Superscripts represent statistically significant differences at $P < 0.05$

Table 3. Content of β -glucan in flours and baked breads

Samples	β -glucan in flours (%)		β -glucan in loaves (%)	
	x	SD	x	SD
Control wheat flour	0.61 \pm 0.02 ^a		0.26 \pm 0.03 ^a	
5% Barliv [™] barley betafiber	4.60 \pm 0.02 ^b		3.99 \pm 0.07 ^b	
10% Barliv [™] barley betafiber	8.40 \pm 0.04 ^c		7.86 \pm 0.35 ^c	
15% Barliv [™] barley betafiber	11.26 \pm 0.03 ^d		10.20 \pm 0.00 ^d	
20% Barliv [™] barley betafiber	13.52 \pm 0.04 ^e		-	

Superscripts represent statistically significant differences at $P < 0.05$

β -glucan. Composite flours with added starchy agents up to 15% (w/w) satisfied the quality required for bakery utilization and were accepted by sensory evaluation. Higher ratios reduced gluten quality, sedimentation index, or falling number. Addition of barley β -glucan decreased all parameters even at a low ratio (5%). Higher ratio was not acceptable from technology or sensory points of view. All starchy and β -glucan agents increased water absorption of dough, reduced dough stability, consequently significantly increased weight and reduced volume of loaves, but water retention in bread crumb was higher. The highest content of RS in final bread was in loaves with added RS₃ (Novelose[®] 330). According to our study, the daily intake of RS could be provided by the consumption of 200 g of bread with

15% of Novelose® 330. Suggested daily intake of β -glucan needed for serious clinical reduction of total blood cholesterol could be received by the consumption of 100 g of bread with 5% of Barliv™ barley betafiber, if the technological and sensory parameters are overlooked.

*

This work was supported by Operational Programme Research and Development “Development of new types of crops with genetically improved traits of economic importance” (ITMS 26220220027) from European Regional Development Fund.

References

- A.A.C.C. (2000): *Total starch assay procedure*. AACC Inc., St. Paul, Minnesota, Method 76-13.01.
- A.A.C.C. (2002): *Resistant starch in starch samples and plant materials*. AACC Inc., St. Paul, Minnesota, Method 32-40.01.
- A.A.C.C. (2003): *Beta-glucan content of barley and oats - Rapid enzymatic procedure*. AACC Inc., St. Paul, Minnesota, Method 32-23.
- A.O.A.C. (1995): Official Method 995.16. *Beta-D-glucan in barley and oats*. AOAC Inc., Gaithersburg, Maryland.
- A.O.A.C. (1996): Official Method 996.11. *Starch (Total) in cereal products*. AOAC Inc., Gaithersburg, Maryland.
- A.O.A.C. (2002): Official Method 2002.02. *Resistant starch in starch and plant materials. Enzymatic digestion. First action 2002*. AOAC Inc., Gaithersburg, Maryland.
- ENGLYST, H.N., KINGSMAN, S.M. & CUMMINGS, J.H. (1992): Classification and measurement of nutritionally important starch fractions. *Eur. J. Clin. Nutr.*, 46, 30–50.
- ESTRADA, A., YUN, C.H., VAN KESSEL, A., LI, B., HAUTA, S. & LAARVELD, B. (1997): Immunomodulatory activities of oat beta-glucan in vitro and in vivo. *Microbiol. Immunol.*, 41, 991–998.
- FDA (1997): Food labeling: health claims; oats and coronary heart disease. *Federal Register The Daily Journal of the United States Government*, 23, 3584–3601.
- HAGER, A.S., RYAN, L.A.M., SCHWAB, C., GÄNZLE, M.G., O'DOHERTY, J.V. & ARENDT, E.K. (2011): Influence of the soluble fibres inulin and oat β -glucan on quality of dough and bread. *Eur. Food Res. Technol.*, 232, 405–413.
- HUNG, P.V., YAMAMORI, M. & MORITA, N. (2005): Formation of enzyme-resistant starch in bread as affected by high-amylose wheat flour substitutions. *Cereal Chem.*, 82, 690–694.
- ICC (1998): *Determination of β -glucan in barley, oat and rye*. ICC, Vienna, Standard No. 166.
- IRAKLI, M., BILLADERIS, C.G., IZYDORCZYK, M.S. & PAPADOYANNIS, I.N. (2004): Isolation, structural features and rheological properties of water-extractable β -glucans from different Greek barley cultivars. *J. Sci. Food Agric.*, 84, 1170–1178.
- JACOBS, M.S., IZYDORCZYK, M.S., PRESTON, K.R. & DEXTER, J.E. (2008): Evaluation of baking procedures for incorporation of barley roller milling fractions containing high levels of dietary fibre into bread. *J. Sci. Food Agric.*, 88, 558–568.
- KEOGH, G.F., GARTH, J.S., MULVEY, T.B., MCARDLE, B.H., COLES, G.D., MONRO, J.A. & POPPITT, S.D. (2003): Randomized controlled crossover study of the effect of a highly β -glucan-enriched barley on cardiovascular disease risk factors in mildly hypercholesterolemic men. *Am. J. Clin. Nutr.*, 78, 711–718.
- KNUCKLES, B.E., HUDSON, C.A., CHIU, M.M. & SAYRE, R.N. (1997): Effect of β -glucan barley fractions in high-fiber bread and pasta. *Cereal Foods World*, 42, 94–99.
- LANDON, S. (2007): Resistant starch review. http://www.resistantstarch.com/NR/rdonlyres/46E5AB79-DABE-4ACA-96A8-1117C99B2ED6/0/Resistant_Starch_Review_Australia.pdf
- MÄLKKI, Y. & VIRTANEN, E. (2001): Gastrointestinal effects of oat bran and oat gum. *Fd Sci. Technol.*, 34, 337–347.
- MOHAMED, A.A., RAYAS-DUARTE, P., XU, J.Y., PALMQUIST, E. & INGLETT, G.E. (2005): Hard red winter wheat/Nutrim-OB alkaline fresh noodles: Processing and texture analysis. *J. Food Sci.*, 71, S1–S7.
- OZTURK, S., KOKSEL, H. & PERRY, K.W. (2009): Farinograph properties and bread quality of flours supplemented with resistant starch. *Int. J. Food Sci. Nutr.*, 60, 449–457.
- POKORNÝ, J. (1997): *Metody senzorické analýzy potravin a stanovení senzorické jakosti*. (Methods of sensory food analysis and evaluation of sensory quality.) Ústav zemědělských a potravinářských informací, Praha, 196 pages.
- SANZ, T., SALVADOR, A., BAIXAULI, R. & FISZMAN, S.M. (2009): Evaluation of four types of resistant starch in muffins. II. Effects in texture, colour and consumer response. *Eur. Food Res. Technol.*, 229, 197–204.

- SKENDI, A., BILLADERIS, C.G., PAPAGEORGIOU, M. & IZYDORCZYK, M.S. (2010): Effects of two barley β -glucan isolates on wheat flour dough and bread properties. *Fd Chem.*, 119, 1159–1167.
- STN (1988): *Skúšanie obilnín, strukovín a olejnin. Skúšanie obilnín. Stanovenie mokrého lepku. Stanovenie ťažnosti lepku. Stanovenie napúchavosti lepku* (Testing of cereals, legumes and oil-bearing crops. Cereal testing. Determination of gluten ductility and swelling capacity.) No. 46 1011-9. SÚTN, Bratislava.
- STN (1990): *Metódy skúšania pekárskeho výrobkov. Stanovenie vlhkosti*. (Methods of testing for baked products. Determination of moisture content.) No. 56 0116-3. SÚTN, Bratislava.
- STN (2003): *Potravinárske obilniny. Časť 2: Zrno potravinárskej pšenice letnej* (Food cereals. Part 2: Food wheat.), No. 46 1100-2, SÚTN, 8 pages.
- STN ISO (2000): *Pšenica. Stanovenie sedimentačného indexu. Skúška podľa Zelenyho*. (Wheat. Determination of the sedimentation index. Zeleny test.), No. 5529. SÚTN, Bratislava, 8 pages.
- STN ISO (2006a): *Obilniny a mleté výrobky z obilnín. Stanovenie celkového popola*. (Cereals, pulses and by-products. Determination of ash yield by incineration.), No. 2171. SÚTN, Bratislava, 8 pages.
- STN ISO (2006b): *Obilniny. Pšenica, raž, respektíve múky, tvrdá pšenica a tvrdá pšenica semolina. Stanovenie čísla poklesu podľa Hagberga-Perten*. (Wheat, rye and their flours, durum wheat and durum wheat semolina. Determination of the falling number according to Hagberg-Perten.) No. 3093. SÚTN, Bratislava, 20 pages.
- TAS, A.A. & EL, S.N. (2000): Determination of nutritionally important starch fractions of some Turkish breads. *Fd Chem.*, 70, 493–497.
- WANG, J., ROSELL, C.M. & DE BARBER, C.B. (2002): Effect of the addition of different fibres on wheat dough performance and bread quality. *Fd Chem.*, 79, 221–226.