

## BIOLOGICAL CONTROL OF AGRICULTURAL PESTS BY FILAMENTOUS FUNGI

L. MANCZINGER

*Department of Microbiology, Attila József University, Szeged, Hungary*

Agricultural use of chemical pesticides has polluted the environment and resulted in resistance among the target organisms. The chemical strategies of pest control are dangerous to both the nontarget organisms in natural habitats and human health. Biological control is an attractive less dangerous possibility for controlling plant pathogens.

Some methods of biological control are becoming now commercially available against plant parasitic fungi, nematods and insects. Among filamentous fungi many candidates with biocontrol potential can be found. Fungal biocontrol agents are less effective and reliable than the synthetic pesticides therefore their use in the agricultural practice requires genetic improvement.

### Mycofungicides

There are many fungal genera, which contain species, which effectively antagonize and/or parasitize plant parasitic fungi both in the rhizosphere and phyllosphere [1].

The most effective species can be found in *Trichoderma*, *Gliocladium*, *Fusarium*, *Talaromyces*, *Ampelomyces* and *Coniothyrium* genera.

Mycoparasitic *Trichoderma* and *Gliocladium* strains could be used to control a wide range of plant-pathogenic fungi e.g. *Pythium*, *Rhizoctonia*, *Fusarium* and *Botrytis*.

The ecological, physiological and molecular aspects of biocontrol by these fungi were recently reviewed [2–5, 77–78, 80].

Alternative mechanisms may be responsible for the control effects of these species. Some strains of them are able to produce antifungal antibiotics [6–7], which may have synergistic effect with extracellular fungal-cell-wall degrading extracellular enzymes [8].

The production of cell-wall hydrolyzing enzymes seems to be the most important factor of antagonism. Both *Trichoderma* and *Gliocladium* mycoparasitic strains in some cases constitutively secrete high amounts of  $\beta$ -1,3-glucanases [9], chitinases [10–11] and proteases [12]. The other important phenomenon of mycoparasitic strains is the capability for the production of appressoria which penetrate the cell wall of the target fungus and

LÁSZLÓ MANCZINGER  
Department of Microbiology, Attila József University  
P.O. Box 533, H-6701 Szeged, Hungary

grow inside the target cells. The production of appressoria is an inductive process which require the presence of specific lectins in the cell wall of the target fungus [13]. There are reports that the cell-wall-degrading enzymes act synergistically in the cell-wall hydrolyzing process. Several different enzymes, chitinases,  $\beta$ -1,3-glucanases,  $\beta$ -1,6-glucanases and proteases have been purified from the inductive ferment broths of *Trichoderma* [14–17] and *Gliocladium* [8] mycoparasitic strains. From the purified enzymes chitinases and  $\beta$ -1,3-glucanases showed the most intensive spore-germination inhibitory effect.

The purified degradative enzymes acted synergistically not only with each other, but also with antibiotics and chemical pesticides [18–19]. Many *Trichoderma* and *Gliocladium* biocontrol strains could be originally resistant against a wide variety of chemical fungicides frequently used for controlling phytopathogenic fungi.

Combining fungicide resistant biocontrol strains with chemical fungicides could results in decreased fungicide level in soils. This is the base of integrated pest management which cold be very effective, but less dangerous to the ecosystem and human health than the chemical control alone [3].

In some cases the biocontrol strains are also sensitive to the fungicide to be applied (e.g. to the frequently used fungicide benomyl); in this case resistant strains could be isolated by induced mutagenesis [20].

An effective pest control may require  $10^5$ – $10^6$  propagula to be present in one gram soil in the case of *Trichoderma* and *Gliocladium* [2]. These high amounts of conidia to be present are not desirable either of ecological and economical or public health of aspects.

To solve this problem the biocontrol strains of *Trichoderma* and *Gliocladium* should breed for more effective.

Strains, which do not require a long inductive process for production of degradative enzymes, could be more effective biocontrol agents. Generally the biocontrol strains produce  $\beta$ -1,3-glucanases constitutively, some elements of chitinase systems require induction and the most effective proteases are also produced inductively [12, 21].

Only a limited number of publications deal with mycoparasitic strains obtained by mutagenesis. Some of these investigations report on random isolate tests following only mutagenetic treatments, others publish on obtaining fungicide resistant mutants which could be used in integrated pest management and only some of them deal with true targeted mutagenetic improvement of the enzyme secretion abilities of the strains [22–23].

The mutagenetic methods of breeding are worth to bigger attention as these strains could get registration more easily for on field use from environmental protection agencies than the strains, which are produced by protoplast fusion, transformation or via gene cloning.

Protoplast fusion is a quick and easy method for combining the advantageous properties of distinct promising strains. It was successfully applied for the breeding of *T. harzianum* biocontrol strains [24–25]. The parasexual cycle of *Trichoderma* has some interesting phenomenons, which were first published in *T. reesei* [26]. The parameiotic behaviour was later also published for *T. harzianum* and *T. viride* [27]. Protoplast could be easily produced from the biocontrol *Trichoderma* strains and their induced fusion resulted in genetic recombinants with elevated biocontrol abilities in many instances [28]. Many publications deal with the transformation methodology of *Trichoderma* and

*Gliocladium* strains. Some of the most frequently applied methods use fungicide resistance, others auxotrophic-complementing or acetamid-catabolic genes for transformation [29, 67–68, 75–76]. Transformation for benomyl resistance interestingly never resulted in high transformation frequencies even if homologous system was used. The use of hygromycin B selection systems resulted in the highest transformation frequencies. These frequencies were even more elevated if spermin was present in the transformation mixture [29]. Many *Gliocladium* and *Trichoderma* strains are originally resistant to hygromycin B. In these cases the acetamidase or auxotrophic complementation methods could be used.

Many publications are dealing with the cloning of structural genes of enzymes important in the mycoparasitic process. Most frequently chitinase genes have been cloned [30–35].

In some cases protease [36] and a  $\beta$ -1,3-glucanase [37] genes were also successfully cloned from mycoparasitic strains.

The cloned genes in some cases with new constitutive promoters were transformed into *Trichoderma* strains. More efficient biocontrol strains could be found among the transformants [38].

There are many ecological factors which could disturb the effectiveness of biocontrol strains in soils: e.g. other *Trichoderma* strains [39], bacteria [40], low temperature [41] and ionic composition. A more intensive investigation of these fields is very important for the effective practical use of the wild type and the breded strains.

In on field investigation the monitoring of the biocontrol strains in soils is also very important.

For these purposes resistance markers, RFLP analysis, DNA fingerprinting, PCR and RAPD techniques could be used efficiently [42–44].

**Table I**

*Commercial biocontrol products for use against plant-pathogenic fungi*

Biocontrol fungus	Product Name
<i>Ampelomyces quisqualis</i>	AQ10
<i>Coniothyrium minitans</i>	Contans
<i>Fusarium oxysporum</i>	Biofox C, Fusaclean
<i>Gliocladium virens</i>	SoilGard (Formerly GlioGard)
<i>Phlebia gigantea</i>	Rotstop, P.g. Suspension
<i>Pythium oligandrum</i>	Polygandron
<i>Trichoderma harzianum</i> and other <i>Trichoderma</i> species	Bio-Fungus, Binab T, RootShield, T-22G, T-22 Planter Box, Bio-Trek, Promote, Supresivit, Trichodex, Trichopel, Trichoject, Trichodowels, Trichoseal, Trichoderma 2000

Besides *Trichoderma* and *Gliocladium* species apathogenic *Fusarium* strains against phytopathogenic *Fusaria* [45], *Coniothyrium* strains against *Sclerotinia* [46] and *Ampelomyces* strains against powdery mildews [47] are used successfully for biocontrol, but few publications are dealing with their biochemical and genetic properties.

Many biocontrol filamentous fungi were patented and their registered on field use were permitted by the environmental agencies. Some important commercially available preparations are listed in Table I.

### Myconematocides and mycoinsecticides

There are three types of nematophagous fungi. The first is able to produce trapping hyphae e.g. *Arthrobotrys* species [48], the conidia of second type are germinating on the surface of the nematode e.g. *Verticillium* [49] species, and in the third case the conidia are germinating in the intestinal system of the nematod e.g. *Harposporium* [50] species.

The trapping method of biocontrol is not reliable, as many environmental factors may disturb the differentiation of the trapping apparatus [51].

An effective and widespread nematode biocontrol could be accomplished by the use of distinct strains of *Verticillium lecanii*, *V. chlamydosporum* and *Paecilomyces lilacinus* [52–53].

The most important factor of pathogenesis is the extracellular protease system both in *Arthrobotrys*, *Verticillium* and *Paecilomyces* species [54–56]. *Verticillium*, *Arthrobotrys* and *Paecilomyces* species besides their nematocidal effect, might also be effective against fungi and insects.

Many filamentous fungal genera contain insect parasitizing species. From them the species belonging to the *Beauveria*, *Metarhizium*, *Verticillium* and *Paecilomyces* are the most promising [79]. The effectiveness of the strains is related to the secretion of extracellular chitinases and proteinases; these enzyme systems are well characterized in the most strains [57].

The nematocide and insecticide strains were bred by mutagenesis, protoplast fusion [58] and transformation [59–60]. In some cases structural genes of the degradative enzymes have been cloned [61–62].

There are possibilities for integrated control with chemical insecticides and fungicides by applying fungicide resistant strains.

The modern molecular methods, RFLP, PCR and RAPD for typing, identification and on field monitoring of strains are widely and successfully used [63–64].

Commercially *Beauveria bassiana*, *Verticillium lecanii*, *Metarhizium anisopliae* and *Paecilomyces lilacinus* are available. There are several products that contain *B. bassiana*, including Naturalis® and Mycotrol®.

### Hungarian research activities

In Hungary research activities related to fungal biological control are mainly limited to the *Trichoderma* genus. In his pioneering works Vajna has determined the distribution of *Trichoderma* species in Hungary and described some antagonistic isolates [39, 65].

Manczinger in Department of Microbiology, Attila József University, Szeged, investigated the catabolic abilities of many Hungarian isolates and detected species within the species aggregates with the use of assimilation spectra [66], also he explored the transient nature of the somatic diploid state in *Trichoderma* following protoplast fusion [26]. Recently in the microbial ecological working group of this department new biopesticid strains were produced by mutagenic treatment with excellent antagonistic properties [23] and for some *T. harzianum* and *T. viride* strains genetic transformation systems were established [29, 67–68]. Naár in the Department of Plant Science, Esterházy Teachers Training College, Eger investigates the ecological behavior of *Trichoderma* strains in distinct soil types, the effect of heavy metals to biocontrol *Trichoderma* strains and the possibilities for integrated pest management with *Trichoderma* strains [69–72]. Turóczi and others in the Department of Plant Pathology, Plant Protection Institute, Hungarian Academy of Sciences, Budapest and in the Agricultural Biotechnology Center, Gödöllő, respectively, in collaboration investigate the molecular characterization possibilities of *Trichoderma* strains [73–74] and recently they successfully cloned and transformed back a chitinase gene in *T. hamatum* [75].

### REFERENCES

1. Lumsden,R.D.: Mycoparasitism of soilborne plant pathogens. In Carroll,G.C., Wicklow,D.T. (eds): The fungal community. Second edition. Marcel Dekker, New York. 1992. p. 275.
2. Adams,P.B.: The potential of mycoparasites for biological control of plant diseases. Annu Rev Phytopathol **28**, 59 (1990).
3. Sivan,A., Chet,I.: Microbial control of plant diseases. In Mitchell,R. (ed.): Environmental Microbiology. Wiley-Liss Inc., New York. 1992. p. 335.
4. Goldman,G.H., Hayes,C., Harman,G.E.: Molecular and cellular biology of biocontrol by *Trichoderma* spp. Trends Biotechnol **12**, 478 (1994).
5. Samuels,G.J.: *Trichoderma*: a review of biology and systematics of the genus. Mycol Res **100**, 923 (1996).
6. Claydon,N., Allan,M., Hanson,J.R., Avent,A.G.: Antifungal alkyl pyrones of *Trichoderma harzianum*. Trans Brit Mycol Soc **88**, 503 (1987).
7. Wilhite,S.E., Straney,D.C.: Timing of gliotoxin biosynthesis in the fungal biological control agent *Gliocladium virens* (*Trichoderma virens*). Appl Microbiol Biotechnol **45**, 513 (1996).
8. Di Pietro,A., Lorito,M., Hayes,C.K., Broadway,R.M., Harman,G.E.: Endochitinase from *Gliocladium virens*: isolation, characterization, and synergistic antifungal activity in combination with gliotoxin. Phytopathology **83**, 308 (1993).
9. Thrane,C., Tronsmo,A., Jensen,D.F.: Endo-1,3-beta-glucanase and cellulase from *Trichoderma harzianum*: purification and partial characterization, induction of and biological activity against plant pathogenic *Pythium* spp. Eur J Plant Pathol **103**, 331 (1997).

10. de la Cruz,J., Rey,M., Lora,J.M., Hidalgo-Gallego,A., Dominguez,F., Pintor-Toro,J.A., Llobell,A., Benitez,T.: Carbon source control on beta-glucanases, chitobiase and chitinase from *Trichoderma harzianum*. *Arch Microbiol* **159**, 316 (1993).
11. Ulhoa,C.J., Peberdy,J.F.: Regulation of chitinase synthesis in *Trichoderma harzianum*. *J General Microbiol* **137**, 2163 (1991).
12. Haran,S., Schickler,H., Chet,I.: Molecular mechanisms of lytic enzymes involved in the biocontrol activity of *Trichoderma harzianum*. *Microbiology* **142**, 2321 (1996).
13. Inbar,J., Chet,I.: Biomimics of fungal cell wall recognition by use of lectin coated nylon fibers. *J Bacteriol* **174**, 1055 (1992).
14. Lorito,M., Harman,G.E., Hayes,C.K., Broadway,R.M., Tronsmo,A., Woo,S.L., Di Pietro,A.: Chitinolytic enzymes produced by *Trichoderma harzianum*: antifungal activity of purified endochitinase and chitobiosidase. *Phytopathology* **83**, 302 (1993).
15. Harman,G.E., Hayes,C.K., Lorito,M., Broadway,R.M., Di Pietro,A., Peterbauer,C., Tronsmo,A.: Chitinolytic enzymes of *Trichoderma harzianum*: purification of chitobiosidase and endochitinase. *Phytopathology* **86**, 313 (1993).
16. Lima,L.H.C., Ulhoa,C.J., Fernandes,A.P., Felix,C.R.: Purification of a chitinase from *Trichoderma* sp. and its action on *Sclerotium rolfsii* and *Rhizoctonia solani* cell walls. *J Gen Appl Microbiol* **43**, 31 (1997).
17. Pintor-Toro,J.A., Benitez,T., Llobell,A.: Purification and characterization of an endo-beta-1,6-glucanase from *Trichoderma harzianum* that is related to its mycoparasitism. *J Bacteriol* **177**, 1864 (1995).
18. Schirmböck,M., Lorito,M., Wang,Y.L., Hayes,C.K., Arisan-Atac,I., Scala,F., Harman,G.E., Kubicek,C.P.: Parallel formation and synergism of hydrolytic enzymes and peptaibol antibiotics, molecular mechanisms involved in the antagonistic action of *Trichoderma harzianum* against phytopathogenic fungi. *Appl Environ Microbiol* **60**, 4364 (1994).
19. Lorito,M., Peterbauer,C., Hayes,C.K., Harman,G.E.: Synergistic interaction between fungal cell wall degrading enzymes and different antifungal compounds enhances inhibition of spore germination. *Microbiology* **140**, 623 (1994).
20. Ahmad,J.S., Baker,R.: Rhizosphere competence of benomyl-tolerant mutants of *Trichoderma* spp. *Can J Microbiol* **34**, 694 (1988).
21. van Tilburg,A.U.B., Thomas,M.D.: Production of extracellular proteins by the biocontrol fungus *Trichoderma virens*. *App Environ Microbiol* **59**, 236 (1993).
22. Sinha,I., Upadhyay,R.S.: Modification of biocontrol potential of *Trichoderma viride*. *Acta Phytopathol Entomol Hung* **28**, 209 (1993).
23. Manczinger,L., Antal,Zs., Ferenczy,L.: Breeding of mycoparasitic *Trichoderma* strains by UV-mutagenesis. Poster Abstracts, IUMS Congresses, Praha. 1994.
24. Sivan,A., Harman,G.E.: Improved rhizosphere competence in a protoplast fusion progeny of *Trichoderma harzianum*. *J Gen Microbiol* **137**, 23 (1991).
25. Stasz,T.E.: Genetic improvement of fungi by protoplast fusion for biological control of plant pathogens. *Can J Plant Pathol* **12**, 322 (1990).
26. Manczinger,L., Ferenczy,L.: Somatic cell fusion of *Trichoderma reesei* resulting in new genetic combinations. *Appl Microbiol Biotechnol* **22**, 72 (1985).
27. Stasz,T.E., Harman,G.E.: Nonparental progeny resulting from protoplast fusion in *Trichoderma* in the absence of parosexuality. *Exp Mycol* **14**, 145 (1990).
28. Pe'er,S., Chet,I.: *Trichoderma* protoplast fusion: a tool for improving biocontrol agents. *Can J Microbiol* **36**, 6 (1990).
29. Manczinger,L., Komonyi,O., Antal,Zs., Ferenczy,L.: A method for high-frequency transformation of *Trichoderma viride*. *J Microbiol Methods* **29**, 207 (1997).
30. Garcia,I., Lora,J.M., de la Cruz,J., Benitez,T., Llobel,A., Pintor-Toro,J.A.: Cloning and characterization of a chitinase (CHIT42) cDNA from the mycoparasitic fungus *Trichoderma harzianum*. *Curr Genet* **27**, 83 (1994).

31. Carsolio,C., Gutierrez,A., Jimenez,B., van Montagu,M., Herrera-Estrella,A.: Characterization of *ech-42*, a *Trichoderma harzianum* endochitinase gene expressed during mycoparasitism. Proc Natl Acad Sci USA **91**, 10903 (1994).
32. Hayes,C.K., Klemsdal,S., Lorito,M., Di Pietro,A., Peterbauer,C., Nakas,J.P., Tronsmo,A., Harman,G.E.: Isolation and sequence of an endochitinase-encoding gene from a cDNA library of *Trichoderma harzianum*. Gene **138**, 143 (1994).
33. Peterbauer,C.K., Lorito,M., Hayes,C.K., Harman,G.E., Kubicek,C.P.: Molecular cloning and expression of the *nag1* gene (N-acetyl-beta-D-glucosaminidase-encoding gene) from *Trichoderma harzianum* P1. Curr Genet **30**, 325 (1996).
34. Draborg,H., Christgau,S., Halkier,T., Rasmussen,G., Dalbøge,H., Kauppinen,S.: Secretion of an enzymatically active *Trichoderma harzianum* endochitinase by *Saccharomyces cerevisiae*. Curr Genet **29**, 404 (1966).
35. Draborg,H., Kauppinen,S., Dalbøge,H., Christgau,S.: Molecular cloning and expression in *Saccharomyces cerevisiae* of two exochitinases from *Trichoderma harzianum*. Biochem Mol Biol Int **36**, 781 (1995).
36. Geremia,R.A., Goldman,G.H., Jacobs,D., Ardiles,W., Vila,S.B., Van-Montagu,M., Herrera-Estrella,A.: Molecular characterization of the proteinase-encoding gene, *prb1*, related to mycoparasitism by *Trichoderma harzianum*. Mol Microbiol **8**, 603 (1993).
37. de la Cruz,J., Pintor-Toro,J.A., Benitez,T., Llobell,A., Romero,L.C.: A novel endo-beta-1,3-glucanase, BGN13.1, involved in the mycoparasitism of *Trichoderma harzianum*. J Bacteriol **177**, 6937 (1995).
38. Flores,A., Chet,H., Herrera-Estrella,A.: Improved biocontrol activity of *Trichoderma harzianum* by over-expression of the proteinase-encoding gene *prb-1*. Curr Genet **31**, 30 (1997).
39. Vajna,L.: Mutual parasitism between *Trichoderma hamatum* and *Trichoderma pseudokoningii*. Phytopathol Z **113**, 300 (1985).
40. Upadhyay,R.S., Visintin,L., Jayawal,R.K.: Environmental factors affecting the antagonism of *Pseudomonas cepacia* against *Trichoderma viride*. Can J Microbiol **37**, 880 (1991).
41. Tronsmo,A., Dennis,C.: Effect of temperature on antagonistic properties of *Trichoderma* species. Tr Br Mycol Soc **71**, 469 (1978).
42. Schlick,A., Kuhls,K., Meyer,W., Lieckfeldt,E., Borner,T., Messner,K.: Fingerprinting reveals gamma-ray induced mutations in fungal DNA: implications for identification of patent strains of *Trichoderma harzianum*. Curr Genet **26**, 74 (1994).
43. Arisan-Atac,I., Heidenreich,E., Kubicek,C.P.: Randomly amplified polymorphic DNA fingerprinting identifies subgroups of *Trichoderma viride* and other *Trichoderma* sp. capable of chestnut blight biocontrol. FEMS Microbiol Lett **126**, 249 (1995).
44. Bulat,S.A., Lübeck,M., Mironenko,N., Jensen,D.F., Lübeck,P.S.: UP-PCR analysis and ITS1 ribotyping of strains of *Trichoderma* and *Gliocladium*. Mycol Res **102**, 933 (1998).
45. Mandeel,Q., Baker,R.: Mechanisms involved in biological control of *Fusarium* wilt of cucumber with strains of nonpathogenic *Fusarium oxysporum*. Phytopathology **81**, 462 (1991).
46. McQuilken,M.P., Whipps,J.M.: Production, survival and evaluation of solid-substrate inocula of *Coniothyrium minitans* against *Sclerotinia sclerotiorum*. Eur J Plant Pathol **101**, 101 (1995).
47. Falk,S.P., Gadoury,D.M., Cortesi,P., Pearson,R.C., Seem,R.C.: Parasitism of *Uncinula necator* cleistothecia by the mycoparasite *Ampelomyces quisqualis*. Phytopathology **85**, 794 (1995).
48. den Belder,E., Jansen,E.: Saprophytic and predacious abilities in *Arthrobotrys oligospora* in relation to dead and living root-knot nematodes. Fundamental Appl Nematol **17**, 432 (1994).
49. Kerry,B.R.: Ecological considerations for the use of the nematophagous fungus, *Verticillium chlamydosporium*, to control plant parasitic nematodes. Can J Botany **73**, S65 (1995).
50. Glockling,S.L., Dick,M.W.: A new species of *Harposporium* infecting *Rhabditis nematodes* with microspores. Mycol Res **98**, 854 (1994).
51. Quinn,M.A.: The influence of saprophytic competition on nematode predation by nematode-trapping fungi. J Invertebr Pathol **49**, 170 (1987).

52. Meyer,S.L.F., Huettel,R.N.: Application of a sex pheromone, pheromone analogs, and *Verticillium lecanii* for management of *Heterodera glycines*. *J Nematology* **28**, 36 (1996).
53. Gaspard,J.T., Jaffee,B.A., Ferris,H.: *Meloidogyne incognita* survival in soil infested with *Paecilomyces lilacinus* and *Verticillium chlamydosporium*. *J Nematology* **22**, 176 (1990).
54. Tunlid,A., Jansson,S.: Proteases and their involvement in the infection and immobilization of nematodes by the nematophagous fungus *Arthrobotrys oligospora*. *Appl Environ Microbiol* **57**, 2868 (1991).
55. Butt,T.M., Keen,J.N., Kerry,B.R., Peberdy,J.F.: The subtilisins of the invertebrate mycopathogens *Verticillium chlamydosporium* and *Metarhizium anisopliae* are serologically and functionally related. *FEMS Microbiol Lett* **126**, 227 (1995).
56. Bonants,P.J.M., Fitters,P.F.L., Thijss,H., den Belder,E., Waalwijk,C., Henfling,J.W.D.M.: A basic serine protease from *Paecilomyces lilacinus* with biological activity against *Meloidogyne hapla* eggs. *Microbiology* **141**, 775 (1995).
57. St Leger,R.J., Joshi,L., Roberts,D.W.: Adaptation of proteases and carbohydrases of saprophytic, phytopathogenic and entomopathogenic fungi to the requirements of their ecological niches. *Microbiology* **143**, 1983 (1997).
58. Couteaudier,Y., Viaud,M., Riba,G.: Genetic nature, stability, and improved virulence of hybrids from protoplast fusion in *Beauveria*. *Microb Ecol* **32**, 1 (1996).
59. Bogo,M.R., Vainstein,M.H., Aragao,F.J.L., Rech,E., Schrank,A.: High frequency gene conversion among benomyl resistant transformants in the entomopathogenic fungus *Metarhizium anisopliae*. *FEMS Microbiol Lett* **142**, 123 (1996).
60. Goettel,M.S., Leger,R.J.S., Bhairi,S., Jung,M.K., Oakley,B.R., Roberts,D.W., Staples,R.C.: Pathogenicity and growth of *Metarhizium anisopliae* stably transformed to benomyl resistance. *Curr Genet* **17**, 129 (1990).
61. St Leger,R.J., Bidochka,M.J.: Cloning of a cuticle-degrading protease from the entomopathogenic fungus, *Beauveria bassiana*. *FEMS Microbiol Lett* **125**, 211 (1995).
62. Smithson,S.L., Paterson,I.C., Bailey,A.M., Screen,S.E., Hunt,B.A., Cobb,B.D., Cooper,R.M., Charnley,A.K., Clarkson,J.M.: Cloning and characterisation of a gene encoding a cuticle-degrading protease from the insect pathogenic fungus *Metarhizium anisopliae*. *Gene* **166**, 161 (1995).
63. Leal,S.C.M., Bertioli,D.J., Butt,T.M., Peberdy,J.F.: Characterization of isolates of the entomopathogenic fungus *Metarhizium anisopliae* by RAPD-PCR. *Mycol Res* **98**, 1077 (1994).
64. Hegedus,D.D., Khachatourians,G.G.: Identification and differentiation of the entomopathogenic fungus *Beauveria bassiana* using polymerase chain reaction and single-strand conformation polymorphism analysis. *J Invertebr Pathol* **67**, 289 (1996).
65. Vajna,L.: *Trichoderma* species in Hungary. *Acta Phytopathol* **18**, 291 (1983).
66. Manczinger,L., Polner,G.: Cluster analyses of carbon source utilization patterns of *Trichoderma* isolates. *System Appl Microbiol* **9**, 214 (1987).
67. Manczinger,L., Antal,Zs., Ferenczy,L.: Isolation of uracil auxotrophic mutants of *Trichoderma harzianum* and their transformation with heterologous vectors. *FEMS Microbiol Lett* **130**, 59 (1995).
68. Antal,Zs., Manczinger,L., Ferenczy,L.: Transformation of a mycoparasitic *Trichoderma harzianum* strain with the *arg B* gene of *Aspergillus nidulans*. *Biotechn Techn* **11**, 205 (1997).
69. Naár,Z., Kecskés,M.: Factors influencing the colonization of different *Trichoderma* species in the soil. (in Hungarian) *Agrokémia és Talajtan* **44**, 221 (1995).
70. Naár,Z., Váradyi,Gy., Kucsma,N., Kecskés,M.: Cadmium susceptibility of *Trichoderma* strains. (in Hungarian) *Agrokémia és Talajtan* **44**, 457 (1995).
71. Naár,Z., Kecskés,M.: A method for selecting trichoderma strains antagonistic against *Sclerotinia minor*. *Microbiol Res* **150**, 239 (1995).
72. Naár,Z., Kecskés,M.: Factors influencing the competitive saprophytic ability of *Trichoderma* species. *Microbiol Res* **153**, 119 (1998).

73. Fekete,Cs., Weszely,T., Hornok,L.: Assignment of a PCR-amplified chitinase sequence cloned from *Trichoderma hamatum* to resolved chromosomes of potential biocontrol species of *Trichoderma*. FEMS Microbiol Lett **145**, 385 (1996).
74. Turóczki,G., Fekete,Cs., Kerényi,Z., Nagy,R., Pomázi,A., Hornok,L.: Biological and molecular characterisation of potential biocontrol strains of *Trichoderma*. J Basic Microbiol **36**, 63 (1996).
75. Giczey,G., Kerényi,Z., Dallmann,G., Hornok,L.: Homologous transformation of *Trichoderma hamatum* with an endochitinase encoding gene, resulting in increased levels of chitinase activity. FEMS Microbiol Lett **165**, 247 (1998).
76. Ulhoa,C.J., Vainstein,M.H., Peberdy,J.F.: Transformation of *Trichoderma* species with dominant selectable markers. Curr Genet **21**, 23 (1992).
77. Esposito,E., Silva,M.: Systematics and environmental application of the genus *Trichoderma*. Crit Rev Microbiol **24**, 89 (1998).
78. Kubicek,C.P.: *Trichoderma* and *Gliocladium*: Enzymes, biological control, and commercial applications. Taylor and Francis, 1998.
79. Leathers,T.D., Gupta,S.C., Alexander,N.J.: Mycopesticides : status, challenges and potential. J Ind Microbiol **12**, 69 (1993).
80. Handelsman,J., Stabb,E.V.: Biocontrol of soilborne plant pathogens. Plant Cell **8**, 1855 (1996).