

New Data to the Knowledge on the Eriophyoid Fauna on Grasses in Hungary (Acari: Eriophyoidea)

G. GÓLYA, E. KOZMA and M. SZABÓ

Department of Plant Protection, Faculty of Agriculture and Environmental Sciences,
Szent István University, H-2100 Gödöllő, P.O. Box 303, Hungary

Aculodes dubius (Nalepa), *Aculodes mckenziei* (Keifer) and *Aceria tosichella* Keifer were recorded for the first time in Hungary. *Festuca pratensis*, *Phleum phleoides* and *Bromus inermis* were found as new host-plants of *Aculodes dubius*, although other plant species in these genera are known to be hosts of this mite. *Aceria tulipae* (Keifer) was identified on *Phleum pratense* and *Aceria tenuis* (Nalepa) on *Phleum phleoides*, which were also new host-plants for them.

Keywords: Graminea, *Aculodes*, *Aceria*.

The eriophyoid mites are frequent on monocotyledons and dicotyledons, and the number of known species increases continuously. However, the researchers mainly give report on the eriophyoid fauna of woody plants in Hungary (Balás, 1939; Dellei and Szendrey, 1988, 1989; Györffy, 1987, 1992; Ripka, 1997; Ripka and de Lillo, 1997). In addition to the description of several mites, Farkas (1965, 1966) gave short characterization on the species occurred on monocotyledons in Central Europe. The imperfect knowledge of the Hungarian eriophyoid fauna on grasses needs further examinations.

Materials and Methods

In 2000 and 2001 investigations were carried out to study the eriophyoid fauna on grasses in Gödöllő and Környe. Whole plants were collected, and the occurrence of mites were examined by Leica Wild mounting microscope. The cleaning of mites was performed by lactic acid, the staining of cuticula by Keifer's intermediate mixture (1954) and the preserving by Keifer's permanent medium (1954). The solution of Heinze (Boczek, 1999) was also used for mounting. The identification of mites was carried out by using Olympus BX-41 microscope with phase contrast and immersion lens. The genera were identified by the keys of Boczek et al. (1989). The species were recognized by using the catalogue of Davis et al. (1982) and Amrine and Stasny (1994), and on the basis of the original and cited descriptions (Keifer, 1938, 1944, 1966, 1969, 1970; Farkas, 1965, 1966; Proeseler, 1972; Jeppson et al., 1975).

Results

From grasses five eriophyoid species were identified (see *Table 1*). Three of them – *Aculodes dubius*¹ (Nalepa), *Aculodes mckenziei*² (Keifer) and *Aceria tosichella* Keifer – were recorded for the first time in Hungary. *Festuca pratensis*, *Phleum phleoides* and *Bromus inermis* were found as new host-plants of *Aculodes dubius*, although other plant species in these genera are known to be hosts of this mite. *Aceria tulipae*³ (Keifer) was found on *Phleum pratense* and *Aceria tenuis*⁴ (Nalepa) on *Phleum phleoides*, which were also new host-plants for them.

Table 1

Eriophyoid mites collected from grasses

Plant species	Mite species
<i>Festuca pratensis</i>	<i>Aculodes dubius</i> (Nalepa)
<i>Festuca rubra</i>	<i>Aculodes dubius</i> (Nalepa) <i>Aculodes mckenziei</i> (Keifer)
<i>Lolium perenne</i>	<i>Aceria tulipae</i> (Keifer) <i>Aculodes dubius</i> (Nalepa)
<i>Phleum pratense</i>	<i>Aceria tulipae</i> (Keifer) <i>Aculodes dubius</i> (Nalepa)
<i>Phleum phleoides</i>	<i>Aceria tenuis</i> (Nalepa) <i>Aculodes dubius</i> (Nalepa)
<i>Arrhenatherum elatius</i>	<i>Aceria tosichella</i> Keifer <i>Aculodes dubius</i> (Nalepa) <i>Aculodes mckenziei</i> (Keifer)
<i>Bromus inermis</i>	<i>Aculodes dubius</i> (Nalepa) <i>Aculodes mckenziei</i> (Keifer)
<i>Triticum aestivum</i>	<i>Aculodes mckenziei</i> (Keifer)

Discussion

Aculodes dubius was a new species on *Festuca pratensis*, *Phleum phleoides* and *Bromus inermis* and in the Hungarian fauna. This mite also occurred on *Festuca rubra* and *Lolium perenne*. According to the catalogue of Amrine and Stasny (1994), this species was originally described on *Helichotrichon pratensis* = *Avena pratensis*, then it was also found on *Bromus mollis* and *Bromus sterilis*. Davis et al. (1982) mentioned *Aculodes dubius* on

¹ Syn.: *Phytocoptes*, *Phyllocoptes*, *Aculops*

² Syn.: *Vasates*

³ Syn.: *Eriophyes*

⁴ Syn.: *Phytoptus*, *Eriophyes*

Agropyron repens and *Bromus mollis*. However, Skoracka (1999) observed this species on even more plants, on *Agrostis vulgaris*, *Alopecurus aequalis*, *Alopecurus pratensis*, *Arrhenatherum elatius*, *Bromus mollis*, *Bromus sterilis*, *Bromus tectorum*, *Corynephorus canescens*, *Dactylis glomerata*, *Festuca arundinacea*, *Festuca rubra*, *Holcus lanatus*, *Lolium perenne*, *Phleum pratense*, *Tymotka kolankowata*, *Poa annua* and *Poa pratensis*.

Aculodes mckenziei was identified for the first time in Hungary on *Festuca rubra*, *Arrhenatherum elatius*, *Bromus inermis* and *Triticum aestivum*. In 1944 Keifer described the species on the leaves of *Elymus triticoides* and named *Vasates mckenziei*. In the catalogue of Davis et al. (1982) and Amrine and Stasny (1994) *Agropyron repens*, *Agropyron smithi*, *Arundo donax*, *Distichlis stipata* and *Triticum* spp. are among the host-plants of this mite. According to Skoracka's (1999) results, the *Aculodes mckenziei* occurs on *Agropyron repens*, *Agrostis alba*, *Arrhenatherum elatius*, *Bromus mollis*, *Bromus sterilis*, *Bromus inermis*, *Calamagrostis epigeios*, *Dactylis glomerata*, *Festuca arundinacea*, *Festuca rubra*, *Phalaris arundinacea*, *Poa pratensis* and *Puccinellia capillaris* in Poland.

Aceria tosichella was also recorded for the first time in Hungary. It was described by Keifer (1969) on *Triticum sativa*, then Skoracka (1999) also found the species on *Agropyron repens*, *Festuca arundinacea*, *Festuca rubra*, *Agrostis vulgaris*, *Arrhenatherum elatius* and *Bromus inermis*.

Aceria tulipae is common in the Hungarian fauna. It was collected on *Lolium perenne* and *Phleum pratense*. This mite has many host-plants, but the *Phleum pratense* was new among them. Originally, Keifer (1938) described the species as *Eriophyes tulipae* on tulip bulbs, but later it was ranked among the members of *Aceria* genus (Keifer 1944). Farkas (1965, 1966) mentioned the occurrence of *Aceria tulipae* on onion, *Tulipa gesneriana* and Liliaceae. On the basis of Davis et al. (1982) and Amrine and Stasny (1994), the host-plants of this species are *Allium cepa*, *Allium sativum*, *Ornithogalum* spp., *Agropyron* spp., *Avena pratensis*, *Bromus* spp., *Digitaria sanguinalis*, *Elymus canadensis*, *Festuca* spp., *Hordeum* spp., *Lolium* spp., *Muhlenbergia racemosa*, *Setaria* spp., *Triticum* spp., *Xerophyllum tenax* and *Zea mays*.

Aceria tenuis also common in Hungary, but the *Phleum phleoides* – on which it were found – was a new host for this mite. According to Davis et al. (1982) and Amrine and Stasny (1994), *Aceria tenuis* can be found on *Agropyron repens* var. *maritimum*, *Agrostis* spp., *Aira* spp., *Alopecurus* spp., *Anthoxanthum odoratum*, *Avena pratensis*, *Avena pubescens*, *Briztia media*, *Bromus* spp., *Calamagrostis* spp., *Cynosurus cristatus*, *Dactylis glomerata*, *Deschampsia caespitosa*, *Deschampsia flexuosa*, *Elymus arenarius*, *Elymus repens*, *Festuca* spp., *Lolium perenne*, *Melica nutans*, *Melica uniflora*, *Milium effusum*, *Molinia coerulea*, *Phalaris arundinacea*, *Phleum pratense*, *Poa* spp., *Sesleria coerulea*, *Sieglindia decumbens* and *Triticum* spp.

Literature

- Amrine, J. W., Jr. and Stasny, T. A. (1994): Catalog of the Eriophyoidea (Acarina: Prostigmata). Indira Publishing House, West Bloomfield, Michigan, 531 pp.
- Balás, G. (1939): Levélatka okozta növénybetegségek Magyarországon (Plant diseases caused by gall mites in Hungary). Magyar Királyi Kertészeti Tanintézet Közleménye 5, 52–70.

- Boczek, J. (1999): Zarys Akarologie Rolniczej. Wyd. PWN. 358 pp.
- Boczek, H. J., Shevtchenko, V. G. and Davis, R. (1989): Generic Key to World Fauna of Eriophyid Mites (Acarina: Eriophyoidea). Warsaw Agricultural University Press, Warsaw, 192 pp.
- Davis, R., Flechtmann, C. H. W., Boczek, J. H. and Barke, H. E. (1982): Catalogue of eriophyid mites (Acari: Eriophyoidea). Warsaw Agricultural University Press, Warsaw, 254 pp.
- Dellei, A. and Szendrey, L. (1988): Fitofág és ragadozó atkafajok előfordulása Heves megye szőlőültetvényeiben (Occurrence of herbivorous and predatory mite species in the vineyards of Heves county). *Növényvédelem* 24, 112–115.
- Dellei, A. and Szendrey, L. (1989): Fitofág és ragadozó atkafajok előfordulása Heves megye gyümölcsöseiben (Occurrence of herbivorous and predatory mite species in the orchards of Heves county). *Növényvédelem* 25, 437–442.
- Farkas, H. (1965): Spinnentiere, Eriophyidae (Gallmilben). In: *Die Tierwelt Mitteleuropas, Band 3, Lief. 3*, Verlag von Quelle and Meyer, Leipzig, 155 pp.
- Farkas, H. (1966): Gubacsatkák (Gall mites) – Eriophyidae. In: *Magyarország állatvilága (Fauna Hungariae) 15*, Akadémiai Kiadó, Budapest, 164 pp.
- Győrfy, M. J. (1987): Veszprém megyei szőlőültetvényekben élő atkafajok dominanciaviszonyai (Dominance between the mite species in the vineyards of Heves county). *Növényvédelem* 23, 202–204.
- Győrfy, M. J. (1992): Új levélatkafaj a hazai faunában (New eriophyoid species in the Hungarian fauna). *Növényvédelem* 28, 320–322.
- Jeppson, L. R., Keifer, H. H. and Baker, E. W. (1975): *Mites Injurious to Economic Plants*. University California Press, Los Angeles, 614 pp.
- Keifer, H. H. (1938): Eriophyid Studies I. *The Bulletin Department of Agriculture State of California* 27, 181–206.
- Keifer, H. H. (1944): Eriophyid Studies XIV. *The Bulletin Department of Agriculture State of California* 43, 121–131.
- Keifer, H. H. (1954): Eriophyid Studies XXII. *The Bulletin Department of Agriculture State of California* 33, 18–38.
- Keifer, H. H. (1966): Eriophyid Studies B-21. *California Department of Agriculture*, 1–24.
- Keifer, H. H. (1969): *ARS-USDA Eriophyid Studies C-3.*, 1–24.
- Keifer, H. H. (1970): *ARS-USDA Eriophyid Studies C-4.*, 1–24.
- Proeseler, G. (1972): Ein Beitrag zur Morphologie und Ökologie der Gallmilben (Eriophyoidea) an Gramineen. *Biol. Zbl.* 91, 477–492.
- Ripka G. (1997): A díszfák és díszcserjék levéltetű- és atkafaunája (Aphid and mite fauna of ornamental trees and shrubs). PhD. thesis. *Kertészeti és Élelmiszeripari Egyetem, Rovartani Tanszék, Budapest*, iv + 209 pp.
- Ripka, G. and de Lillo, E. (1997): New data to the knowledge on the eriophyoid fauna in Hungary (Acari: Eriophyoidea). *Folia Entomologica Hungarica* 28, 147–157.
- Skoracka, A (1999): Występowanie i preferencje pokarmowe szpecieli (Acari: Eriophyoidea) na trawach w Polsce. *Materia y XXVI Sympozjum Akarologicznego "Akarologia polska u progu nowego tysiąclecia"*, Kazimierz Dolny, 333–339.

Acta Phytopathologica et Entomologica Hungarica


Editor-in-Chief

Z. KIRÁLY

Editorial Board

P. Benedek, T. Érsek, L. Hornok, J. Horváth, Z. Klement, T. Kőmíves,
Z. Mészáros, Gy. Sáringer, M. Tóth, F. Virányi

Volume 37



Akadémiai Kiadó, Budapest
2002