

Photoactivation of the BLUF protein PixD Probed by the Site-Specific Incorporation of Fluorotyrosine Residues

Agnieszka A. Gil, Sergey P. Laptanok, James N. Iuliano, Andras Lukacs, Anil Verma, Christopher R. Hall, EunBin Yoon, Richard Brust, Gregory M. Greetham, Michael Towrie, Jarrod B. French, Stephen R. Meech, and Peter J Tonge

J. Am. Chem. Soc., **Just Accepted Manuscript** • DOI: 10.1021/jacs.7b07849 • Publication Date (Web): 06 Sep 2017

Downloaded from <http://pubs.acs.org> on September 29, 2017

Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.



1
2
3
4
5
6
7
8

Photoactivation of the BLUF Protein PixD Probed by the Site-Specific Incorporation of Fluorotyrosine Residues

9 Agnieszka A. Gil[‡], Sergey P. Laptanok^{§,⊥}, James N. Iuliano[‡], Andras Lukacs[#], Anil Verma[†],
10
11 Christopher R. Hall[§], EunBin Yoon[‡], Richard Brust[‡], Gregory M. Greetham[†], Michael Towrie[†],
12
13 Jarrod B. French^{‡,¥}, Stephen R. Meech^{§,*}, Peter J. Tonge^{‡,*}
14
15
16
17

18 [‡]Department of Chemistry, and [¥]Biochemistry & Cell Biology, Stony Brook University, Stony
19 Brook, New York 11794-3400, USA; [§]School of Chemistry, University of East Anglia, Norwich
20 Research Park, Norwich NR4 7TJ, UK; [#]Department of Biophysics, Medical School, University
21 of Pecs, Pecs, Hungary; [†]Central Laser Facility, Harwell Science and Innovation Campus,
22 Didcot, Oxon OX11 0QX, UK.
23
24
25
26
27
28
29

30 [⊥]Current address (SPL): Biological and Environmental Science and Engineering Division, King
31 Abdullah University of Science and Technology, P.O. Box 4700, Thuwal 23955-6900, Kingdom
32 of Saudi Arabia.
33
34
35
36
37
38
39

40 *Authors to whom correspondence should be addressed: Email: s.meech@uea.ac.uk (SRM);
41 peter.tonge@stonybrook.edu (PJT)
42
43

44 KEYWORDS PixD, BLUF, Ultrafast infrared, Fluorotyrosine, FAD, Photoactivation, kinetics
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Abstract

The flavin chromophore in blue light using FAD (BLUF) photoreceptors is surrounded by a hydrogen bond network that senses and responds to changes in the electronic structure of the flavin on the ultrafast time scale. The hydrogen bond network includes a strictly conserved Tyr residue, and previously we explored the role of this residue, Y21, in the photoactivation mechanism of the BLUF protein AppA_{BLUF} by the introduction of fluorotyrosine (F-Tyr) analogs that modulated the pK_a and reduction potential of Y21 by 3.5 pH units and 200 mV, respectively. Although little impact on the forward (dark to light adapted form) photoreaction was observed, the change in Y21 pK_a led to a 4,000-fold increase in the rate of dark state recovery. In the present work we have extended these studies to the BLUF protein PixD, where, in contrast to AppA_{BLUF}, modulation in the Tyr (Y8) pK_a has a profound impact on the forward photoreaction. In particular, a decrease in Y8 pK_a by 2 or more pH units prevents formation of a stable light state, consistent with a photoactivation mechanism that involves proton transfer or proton coupled electron transfer from Y8 to the electronically excited FAD. Conversely, the effect of pK_a on the rate of dark recovery is markedly reduced in PixD. These observations highlight very significant differences between the photocycles of PixD and AppA_{BLUF}, despite their sharing highly conserved FAD binding architectures.

Introduction

Flavins are ubiquitous biological cofactors and are found in ~1% of all proteins, usually either as flavin mononucleotide (FMN) or flavin adenine dinucleotide (FAD).¹ The chemistry associated with FMN and FAD is performed by the 7,8-dimethylisoalloxazine ring, a planar tricyclic aromatic system which can undergo 1 or 2 electron reduction and also participate in the formation of covalent adducts with reactants and amino acid side chains.¹⁻³ The broad distribution of flavoproteins across biological space includes flavin-dependent photoreceptors which utilize the visible electronic transition of the oxidized isoalloxazine ring, centered at around 450 nm, to sense and respond to blue-light.⁴ Structural changes initiated by light absorption on the picosecond time scale lead to macromolecular reorganization and biological output on the millisecond to second time scale.⁵ Of central importance in photoreceptor biology is the understanding of the underlying structural dynamics that link events separated by more than ten decades of time. Indeed, the mechanism of photoreceptor signaling is an area of intense experimental and theoretical activity,⁶⁻¹² stimulated by the recent interest in deploying flavin-dependent photoreceptors as optogenetic devices that enable the *in vivo* control of biology through non-invasive light signaling.¹³ However, despite the unique qualities and potential applications of flavoprotein photoreceptors, in many cases the detailed mechanism of photocycling still remains to be elucidated.

Blue-light sensing flavoproteins are found in animals, plants, fungi, and bacteria,¹⁴⁻¹⁷ and include the photolyase/cryptochromes, the light-oxygen-voltage (LOV) domain proteins, and the blue light using FAD (BLUF) domain proteins.^{15, 18} For both the photolyase/cryptochrome and LOV domain photosensors, photoexcitation leads to changes to the flavin that recapitulate common themes in flavin-dependent enzyme reactions: an alteration in oxidation state of the

flavin or the formation of a covalent adduct. In contrast, a unifying chemical mechanism linking flavin excitation to photoreceptor activation is yet to emerge for the BLUF protein family. While it is clear that the formation of a flavin adduct is not involved in BLUF domain activation, the role of electron transfer and accompanying changes in flavin redox state remains controversial, with both dark adapted and light adapted forms containing a fully oxidized flavin.

In the BLUF photoreceptors,¹⁹ the flavin is surrounded by a hydrogen bonding network that includes conserved Tyr and Gln residues, both of which are essential for photoactivity (**Figure 1**). In some BLUF protein structures a conserved Trp is within hydrogen bonding distance of the network, and indeed the variable location of the Trp side chain, between this Trp_{in} conformation and a Trp_{out} conformation, supports proposed mechanisms in which a light-induced change in Trp conformation is on the reaction coordinate for photoactivation.^{7, 20-21}




Figure 1. The Hydrogen Bonding Network and Primary Photoactivation Mechanism

The isoalloxazine ring is surrounded by a conserved hydrogen bonded network that includes a Tyr Y21(8), Gln Q63(50) and Asn N45(32). Also shown is a Trp W104(91) which is shown hydrogen bonded to the Gln residue in the dark state but not in the light state. Movement of the Trp from a Trp_{in} to a Trp_{out} conformation is thought to be a central component of the BLUF photoactivation mechanism although we note that in the X-ray crystal structures of PixD in the

1
2
3 dark (PDB: 2HFN)²² and a stable lit states (PDB: 3MZI),⁷ the W91 sidechain is rotated out of the
4
5 binding pocket. Photoactivation involves rotation of the Gln side-chain, and in the above
6
7 mechanism keto-enol tautomerism of the Gln side-chain is shown preceding side-chain
8
9 rotation.^{10, 23-25} Residues are numbered based on the sequence of AppA with the residue number
10
11 in PixD given in parentheses.
12
13
14
15
16

17 All active BLUF domain proteins exhibit a characteristic ~10 nm red-shift in the
18
19 electronic spectrum of the oxidized flavin upon blue light excitation,²⁶ which has been linked to
20
21 the formation of an additional hydrogen bond to the flavin C2=O carbonyl group from the side
22
23 chain of the conserved Gln.^{5, 23, 27-29} This, in turn, is thought to result from light-induced rotation
24
25 of the Gln side chain, a proposal that remains a central tenet of BLUF protein photoactivation.<sup>5,
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60</sup>

30-32 Based on ultrafast infrared spectroscopy, we have proposed that keto-enol tautomerism of
the Gln side chain precedes rotation,²³ a hypothesis that is supported by theoretical and
experimental studies.^{9-10, 24} Gln rotation may trigger a change in Trp conformation, and we and
others have demonstrated that mutagenesis of the Trp uncouples the light-induced changes in the
flavin absorption spectrum and hydrogen-bonding from the global changes in BLUF protein
structure that accompany photoactivation.^{8, 33}

The light-driven changes in the hydrogen bond network presumably result from
alterations in the electronic structure of the flavin caused by photoexcitation. Based on
experiments with PixD (Slr1694), which controls phototaxis in the cyanobacterium
Synechocystis sp. PCC 6903 through a light-dependent interaction with PixE, it has been shown
that electron transfer from the conserved Tyr (Y8 in PixD) to the photoexcited isoalloxazine ring
of the flavin results in formation of an anionic semiquinone radical (FAD^{•-}).^{25, 34-36} This electron

1
2
3 transfer is proposed to be followed by the sequential transfer of a proton leading to the neutral
4
5 flavin semiquinone, FADH^\bullet , in which Y8 operates as both the electron as well as a proton donor
6
7 to the flavin (i.e. $\text{FAD}^* \rightarrow \text{FAD}^{\bullet-} \rightarrow \text{FADH}^\bullet$).³⁴⁻³⁷ However, we have previously shown that a
8
9 metastable radical intermediate state is not formed during photoactivation of the BLUF domain
10
11 proteins AppA_{BLUF} and BlsA, whereas radical formation was clearly observed in PixD.³⁴ We
12
13 further confirmed that electron transfer did not occur during the photoactivation of AppA_{BLUF}
14
15 using the site-specific incorporation of fluorotyrosine (F-Tyr) analogs into position Y21.²¹ The
16
17 observation of radical formation in PixD,^{34,38} and in the BLUF protein PapB,³⁹ thus suggests that
18
19 significant differences exist in the mechanism of photoactivation within the BLUF protein
20
21 family.
22
23
24
25
26

27 To gain insight into the structural basis for the significant variation in photoactivation
28
29 mechanism, we superimpose the structures of AppA_{BLUF} and PixD in **Figure 2**. Analysis of the
30
31 flavin binding pocket reveals that the conserved Tyr and Gln residues, as well as N45 (N32)
32
33 occupy essentially identical locations in both photoreceptors. Thus, at least on this basis, Y21 in
34
35 AppA_{BLUF} should be able to fulfill the same function in electron and proton transfer as Y8 does
36
37 in PixD. However, two differences can be observed. In AppA_{BLUF}, the FAD C2=O group makes
38
39 a single hydrogen bond to H44, whereas in PixD two hydrogen bonds are present to N31 (the
40
41 homolog of H44) and to R65, which is a His (H78) in AppA_{BLUF}. We note that the homolog of
42
43 N31 (H44) in BlsA is a Phe (F32),⁴⁰ and given that radical formation is not observed in either
44
45 AppA_{BLUF} or BlsA, it is possible that alterations in the environment of C2=O may modulate the
46
47 BLUF photocycle. The other major difference is the position of the conserved Trp, which is in
48
49 the Trp_{in} conformation in AppA_{BLUF} (W104) and in the Trp_{out} conformation in PixD (W91).
50
51 Since the X-ray structure of BlsA is not available, a comparison with this protein cannot be
52
53
54
55
56
57
58
59
60

drawn. However, it is possible that the position of the Trp influences the ability of the Tyr to participate in proton coupled electron transfer (PCET), as proposed recently based on computational studies.¹¹




Figure 2: The Flavin Binding Pocket of AppA_{BLUF} and PixD

The structures of AppA_{BLUF} (PDB: 1YRX; yellow) and PixD (PDB: 2HFO; slate) have been overlaid to compare and contrast the hydrogen bonding network that surrounds the isoalloxazine ring of the FAD chromophore. The numbering for AppA_{BLUF} is used for residues that are conserved between the two proteins with the residue number in PixD given in parentheses. Superposition was performed using pymol,⁴¹ using the main chain C_α atoms of 1YRX (117 residues) and Chain A from 2HFO (150 residues) and gave an overall rmsd of 0.7 over 76 atoms. Hydrogen bond distances are in Å.

1
2
3
4
5
6 In the current work, we have probed the specific role of the Y8 hydroxyl group in PixD
7
8 photoactivation by replacing this residue with F-Tyr analogs that modulate the pK_a and reduction
9
10 potential by 3.5 pH units and 200 mV, respectively. Although little impact was observed on the
11
12 forward light driven reaction in AppA_{BLUF},²¹ the F-Tyr analogues are shown here to have a
13
14 profound impact on the photoactivation mechanism of PixD. In wild-type PixD there is evidence
15
16 from ultrafast infrared spectroscopy for the sequential formation of FAD^{•-} followed by FADH[•]
17
18 on the reaction coordinate for light state formation. In 3-F-Tyr (3-FY8) PixD, sequential
19
20 electron and proton transfer is no longer observed, with FAD^{•-} and FADH[•] appearing
21
22 simultaneously, while the proteins substituted with 2,3-diF-Tyr (2,3-F₂Y8), 3,5-diF-Tyr (3,5-
23
24 F₂Y8), or 2,3,5,-triF-Tyr (2,3,5-F₃Y8) do not progress to a stable light state. In contrast, for the
25
26 light to dark recovery, the F-Tyr analogs resulted in a dramatically enhanced recovery rate in
27
28 AppA_{BLUF}, an effect which is here shown to be much less marked in PixD. These observations
29
30 highlight fundamental differences between the photocycles of PixD and AppA_{BLUF}.
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Experimental Methods

Materials

3-Fluorophenol and 2,6-difluorophenol were purchased from Sigma-Aldrich. 2,3-Difluorophenol was purchased from Acros Organics. 2,3,6-Trifluorophenol was purchased from Oakwood Chemical. Pyridoxal-5'-phosphate, and flavin adenine dinucleotide were purchased from Sigma-Aldrich.

F-Tyr synthesis using Tyrosine Phenol Lyase (TPL)

Tyrosine phenol lyase (TPL) was prepared and used for the synthesis of the F-Tyr analogs as previously described.²¹

Expression and purification of PixD

Expression of PixD was performed using a plasmid containing the gene for PixD cloned into the pOPINE vector so that the expressed protein included a C-terminal His-tag. BL21(DE3) *Escherichia coli* cells were transformed by the PixD C-6xHis construct (pOPINE vector) and plated on LB-Agar containing 50 mg/L ampicillin (Amp). Overnight cultures were prepared by inoculating 10 mL Luria Broth (LB) media containing 50 mg/L Amp with a single colony and incubating the culture at 37 °C at 250 rpm. Subsequently 5 mL of the overnight culture was used to inoculate 1L of LB media containing 50 mg/L Amp in a 4L flask. The culture was shaken at 37 °C and 250 rpm until the cells reached an OD₆₀₀ of about 0.8, after which the temperature was lowered to ~ 18 °C. After ~30 min, 0.8 mM isopropyl β-D-1-thiogalactopyranoside (IPTG) was added to induce protein expression, and the culture was then incubated for ~ 16 hr in the dark. Cells were harvested by centrifugation at 4,225 x g and 4 °C and stored in the dark at -20 °C

1
2
3 until needed. Purification of PixD was performed under minimal illumination. The cell pellets
4
5 were thawed and resuspended in 40 mL of 50 mM NaH₂PO₄, pH 8.0 buffer containing 10 mM
6
7 NaCl (buffer A). Then, 200 μL of 50 mM phenylmethanesulphonylfluoride (PMSF) was added
8
9 together with 14 μL of β-mercaptoethanol, and the cells were lysed using a high-pressure cell
10
11 disruptor (Constant Systems, Model E1061) at 30 kpsi whilst maintaining the cell disruptor
12
13 vessel at 4 °C. Cell debris was then removed by ultracentrifugation at 126,264 x g and 4 °C for
14
15 90 min, after which the supernatant was incubated with 10 mg of FAD for 45 min on ice. PixD
16
17 was then purified using Ni-NTA affinity chromatography. After applying the supernatant to a 25
18
19 mL column containing 5 mL of resin, the column was washed with 100 mL of buffer A. A
20
21 second wash was performed with 100 mL of buffer A containing 10 mM imidazole, followed by
22
23 100 mL of buffer A containing 20 mM imidazole. Protein was eluted using buffer A containing
24
25 250 mM imidazole. The eluted protein was dialyzed against buffer A overnight. The purity of
26
27 the protein was determined by SDS-PAGE electrophoresis. To exchange the protein into D₂O,
28
29 the samples were first frozen in liquid N₂ and lyophilized overnight, followed by 2-3 cycles of
30
31 resuspension in D₂O and lyophilization.
32
33
34
35
36
37
38
39
40

41 **Incorporation of F-Tyr residues using orthogonal aminoacyl-tRNA synthetases**

42
43 F-Tyr analogs 3-FY, 2,3-F₂Y, 3,5-F₂Y and 2,3,5-F₃Y were incorporated into position 8 of
44
45 PixD using two orthogonal polyspecific aminoacyl-tRNA synthetases, E3 and E11, generously
46
47 provided by Professor Joanne Stubbe.⁴² Site-directed mutagenesis was first used to convert the
48
49 codon for Y8 to the amber codon (TAG) in PixD using the primers in **Table S1**. A C-terminal
50
51 6xHis-tag PixD construct was used (pOPINE C-6His), so that only protein with a F-Tyr
52
53 incorporated at position 8 was able to bind to the Ni-NTA affinity purification resin. The PixD
54
55
56
57
58
59
60

1
2
3 plasmid with the Y8-TAG mutation was co-transformed into BL21(DE3) *E. coli* cells together
4
5 with either the E3 or E11 pEVOL plasmid, and plated on LB agar containing both Amp (50
6
7 mg/L) and Cam (50 mg/L) to select cells harboring both plasmids. A colony from the
8
9 LB/Amp/Cam plate was used to start a 10 mL overnight culture, which was subsequently used to
10
11 inoculate 500 mL 2x YT media containing Amp and Cam. The culture was grown until an
12
13 OD₆₀₀ of ~ 0.4, and a F-Tyr analog dissolved in NaOH was added to the media to give a final
14
15 concentration of ~1 mM. After 30 min, 0.05% w/v arabinose was added to induce expression of
16
17 the E3 or E11 synthetase. The culture was incubated at 37 °C (250 rpm) until the OD₆₀₀ reached
18
19 ~1.0, and then 0.8 mM IPTG was added to the media. After incubating overnight at 30 °C (250
20
21 rpm), the cells were harvested and protein was purified using the same protocol as that described
22
23 for the purification of PixD above. Incorporation of F-Tyr amino acids was analyzed by ESI
24
25 MS/MS mass spectrometry of tryptic digests using a Thermo Scientific Orbitrap Fusion Lumos
26
27 instrument.
28
29
30
31
32
33
34
35

36 **UV-Vis Spectroscopy**

37
38 Absorption spectra of all protein samples were obtained at 20 °C using a Cary 100 Bio
39
40 UV-Vis spectrometer. The concentration of the protein samples was kept at around 80 μM in
41
42 either H₂O or D₂O buffer (50 mM NaH₂PO₄, 10 mM NaCl, pH/D 8.0). Dark-adapted spectra
43
44 were obtained first and then the samples were illuminated with ~500 mW of 455 (± 10) nm light
45
46 for ~ 1 min in order to generate the light state.
47
48
49
50

51 **Time-resolved UV-Vis spectroscopy**

1
2
3 Absorption spectra of protein samples were obtained using an Ocean Optics USB2000+
4 spectrometer described previously.²¹ Spectra of dark adapted PixD and the F-Tyr-substituted
5 variants were first obtained, and then the sample window (5 cm² area) was irradiated with ~500
6 mW of 455 (± 10) nm light until the photostationary state was generated. Conversion to the light
7 state was monitored by taking absorption spectra of the samples in the IR cell. Recovery of the
8 dark state was then monitored after illumination was discontinued.
9
10
11
12
13
14
15
16
17
18
19

20 **Ultrafast Time-Resolved Infrared Spectroscopy**

21
22 Ultrafast time-resolved IR (TRIR) spectra were measured at the STFC Central Laser
23 Facility with ~ 100 fs time resolution.⁴³ TRIR spectra were acquired at 20 °C from 1300 – 1800
24 cm⁻¹ at a resolution of 3 cm⁻¹ per pixel. Data were obtained using a 50 μm path length flow cell
25 of volume < 1 ml, which was also rastered in the excitation beam in order to minimize unwanted
26 secondary photochemistry processes such as photobleaching, photodegradation and
27 photoconversion. The excitation beam of the 450 nm 100 fs 5 kHz pulses was focused to a spot
28 size of ~ 100 μm and the pulse energy was kept below 400 nJ to avoid buildup of the light state.
29 Transient difference spectra (pump on – pump off) were recorded using the IR probe at time
30 delays between 1 ps and 2 ns. After the measurements were recorded, the extent of
31 photoconversion was shown to be negligible using absorbance spectroscopy. Spectra were
32 calibrated relative to the IR transmission of a pure *cis* stilbene standard sample placed at the
33 sample position. TRIR spectra of light-adapted PixD were recorded while continuously
34 illuminating the sample cell with ~200 mW of 450 nm light from an LED. The cell was rastered
35 during data collection.
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Time Resolved Multiple Probe Spectroscopy (TRMPS)

TRMPS spectra were also obtained at the STFC Central Laser Facility.⁴⁴ The TRMPS method has been described,⁴⁴ and previously used by us to analyze the photoactivation of BLUF and LOV domain proteins.⁸ Spectra were measured at 20 °C between 100 fs to 200 μ s after excitation of FAD at 450 nm. A rastered flow cell was used, and data were acquired using a 450 nm pump operated at 0.6-0.8 μ J per pulse and a repetition rate of 1 kHz. After the measurements were recorded, the extent of photoconversion was shown to be negligible using absorbance spectroscopy. The spectral resolution was 3 cm^{-1} and the temporal resolution was 200 fs. A typical measurement was acquired during 45 min of data collection. All samples were prepared at 1-2 mM concentration in D₂O buffer (50 mM NaH₂PO₄, 10 mM NaCl, pD 8.0). Spectra were calibrated relative to the IR transmission of a pure *cis* stilbene standard sample placed at the sample position. Data were analyzed globally using the sequential model with Glotaran software package.⁴⁵

Results and Discussion

TRIR of the Wild-type PixD Dark State

Complex multiexponential kinetics have been previously reported during the decay of the transient electronic spectrum of PixD, consistent with the sequential formation of radical intermediates during photoactivation.^{34-35, 37, 46-47} Kennis and co-workers used transient electronic spectroscopy to observe the formation of radical states during photoactivation of PixD,^{38, 47} while our recent publication confirmed the observation of a flavin radical as well as complex kinetics leading to light state formation in PixD, and contrasted this with the absence of observable radical intermediates in AppA_{BLUF}, which had overall slower photoactivation kinetics.³⁴ The initial step in the PixD photocycle was characterized using ultrafast time-resolved infrared (TRIR) spectroscopy which probes changes in the infrared spectrum of both the chromophore and protein following photoexcitation. In the present work, we first reexamine the TRIR spectra of wild-type PixD, acquired at higher S/N than in our previous study, as a foundation for exploring the impact of F-Tyr substitution on the PixD photocycle.

The newly measured TRIR spectra (**Figure 3A**) were subjected to global analysis using Glotaran,⁴⁵ in which the kinetic traces at all measured wavenumbers were simultaneously fit to a sum of exponential decays. The resulting decay associated spectra, generated by assuming parallel decay of all the components, are presented in the supplementary information (**Figure S1**). The corresponding evolution associated spectra (EAS) of wild-type PixD are shown in **Figure 3B**, where the EAS assume a sequential reaction scheme: for the wild-type photoreceptor, four time constants are needed to adequately describe the data (**Table 1**).




Figure 3. TRIR Spectra of the Wild-type PixD Dark State. (A) TRIR spectra of wild-type PixD (B) EAS of wild-type PixD from a global fit of the TRIR data in (A). Transients assigned

to FADH[•] (1535 cm⁻¹) and FAD^{•-} (1515 cm⁻¹) are shown in the inset. (C) Kinetic traces of the excited state decay (1380 cm⁻¹) and the ground state recovery (1547 cm⁻¹, shown inverted for comparison). The overlay demonstrates slower recovery of the ground state compared to the decay of the excited state. (D) Kinetic traces show the rise and decay of bands assigned to FAD radicals in PixD. Raw data are shown as solid lines while global fitting results are shown as dashed lines.

Table 1. Time constants from TRIR Data for PixD and n-FY8 substituted PixD^a

	τ_1 /ps	τ_2 /ps	τ_3 /ps	τ_4 /ps	$\langle \tau \rangle$ /ps ^d
PixD^b	2.5±0.1	20±0.5	110±2	525±15	96
3-FY8 PixD^b	4.5±0.3	33±1	217±5	1300±100	347
2,3-F₂Y8 PixD^c	3±1	60±1	1395±45	infinite	917
3,5-F₂Y8 PixD^c	2.5±0.5	54±1	965±20	infinite	640
2,3,5-F₃Y8 PixD^c	4.5±1	48±1	1030±30	infinite	730
PixD Light State^b	3±0.2	24±0.5	563±7	infinite	23

^aUnless noted, the TRIR data are from the dark state of the wild-type and F-Tyr PixD variants. Data were globally analyzed using Glotaran.⁴⁵ ^bFour time constants were needed to adequately describe the data for the dark and light states of PixD and 3-FY8 PixD. ^cVery long kinetics were observed for the 2,3-F₂Y8, 3,5-F₂Y8 and 2,3,5-F₃Y8 substituted proteins, and only three time constants were needed to describe the data for these proteins, followed by an ‘infinite’ lifetime extending beyond the range of the TRIR measurement.

^dMean lifetime of the excited state decay calculated from the global analysis at 1380 cm⁻¹.

Major negative bands (bleaches) are observed at 1700, 1638, 1580, and 1547 cm⁻¹ in **Figure 3A, B** which are assigned to the C4=O and C2=O carbonyl groups of the flavin (1700 and 1638 cm⁻¹) and C-N vibrations of the isoalloxazine ring.^{23, 48-49} In addition to these bleaches, which appear instantaneously after photoexcitation (<100 fs), positive bands (transients) are also

1
2
3 observed at 1420 and 1380 cm^{-1} which correspond to the excited state (ES) of the flavin (FAD^*).
4
5 The change in intensity of the 1380 and 1547 cm^{-1} bands as a function of time can thus be used
6
7
8 to quantify excited state decay and ground state recovery, respectively. In **Figure 3C** it can be
9
10 seen that the excited state decays more rapidly than recovery of the ground state, consistent with
11
12 a mechanism that involves excited state relaxation to an intermediate species before ground state
13
14 recovery.
15

16
17 Previously we employed model studies to assign a transient at 1528 cm^{-1} in the TRIR
18
19 spectrum of PixD to the formation of an FAD radical.^{34, 50} The improvement in signal quality of
20
21 the current data enables us to clearly observe two features in this region at around 1515 and 1535
22
23 cm^{-1} (**Figure 3A, B**), which have different kinetics (**Figure 3D**) and are assigned to $\text{FAD}^{\bullet-}$ and
24
25 FADH^{\bullet} , respectively, based on TRIR spectra of the anionic and neutral semiquinones formed by
26
27 model flavins and glucose oxidase.⁵⁰ The 1515 cm^{-1} transient appears on a timescale of 2.5 ps
28
29 and is assigned to formation of the $\text{FAD}^{\bullet-}$. As the transient at $\sim 1515 \text{ cm}^{-1}$ is decaying on the 20
30
31 ps timescale, a new transient is formed around 1535 cm^{-1} which we associate with the formation
32
33 of FADH^{\bullet} by proton transfer (**Figure 3A, B**). The transient at 1535 cm^{-1} decays in a bi-
34
35 exponential manner with time constants of 110 ps and 525 ps, leading ultimately to the formation
36
37 of the PixD light state. The final TRIR spectrum (2 ns) is characteristic of the light state and
38
39 shows a transient at 1690 cm^{-1} which can be assigned to the C4=O flavin carbonyl group, red
40
41 shifted from the position observed in the dark state (1700 cm^{-1}) as a consequence on an increase
42
43 in hydrogen bonding concomitant with photoactivation.^{48, 51-53} In **Figure 3** it can be seen that
44
45 FADH^{\bullet} forms before complete decay of the excited state, FAD^* . This, together with the
46
47 observation of multiexponential kinetics, suggests the presence of an inhomogeneous population
48
49 of states imperfectly resolved in the sequential analysis, possibly resulting from a distribution of
50
51
52
53
54
55
56
57
58
59
60

1
2
3 ground state conformations around the flavin chromophore. Both electron and proton transfer
4 reactions are very distance dependent and sensitive to their environment, so photokinetics will
5 reveal inhomogeneity which is not apparent in crystallography. However, there is clear evidence
6 for FAD radical formation in wild-type PixD as reported previously,³⁴⁻³⁵ and significantly, we
7 are now able to resolve the radical formation into distinct electron and proton transfer phases.
8
9

10
11 To support the assignments discussed above, we also obtained TRIR spectra of the Y8F
12 PixD mutant (**Figure S2**). Although Y8F PixD is photoinactive, excited state quenching also
13 occurs, and is similar in magnitude to that observed for wild-type PixD, and again at a faster rate
14 than the rate of ground state recovery (1380 cm^{-1} v 1547 cm^{-1}). However, in the region where we
15 observe features associated with FAD radicals in wild-type PixD, only one transient is observed,
16 at 1520 cm^{-1} . Based on the possibility that the 1520 cm^{-1} band might be comprised of unresolved
17 contributions from $\text{FAD}^{\bullet-}$ and FADH^{\bullet} , we measured the kinetics at 1515 and 1530 cm^{-1} .
18 However, no difference was observed in the evolution of the signal at these two frequencies,
19 suggesting that only one transfer process occurs in the Y8F mutant. Since the source of the
20 electron is most likely W91, and since this residue cannot donate a proton, we conclude that in
21 Y8F the 1520 cm^{-1} band is associated with the flavin semiquinone anion, supporting the
22 assignment of the band at 1515 cm^{-1} in wild-type PixD to $\text{FAD}^{\bullet-}$.
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45

46 **TRIR of the Wild-type PixD Light State**

47
48 We also obtained TRIR spectra of PixD in the light state (**Figure 4**). A bleach is formed
49 instantaneously at 1685 cm^{-1} , which is assigned to the C4=O of the flavin. The position of this
50 band is consistent with the transient observed at 1690 cm^{-1} in the final EAS of the dark-adapted
51 protein. The assignment of the 1685 cm^{-1} band is also consistent with the proposed mechanism
52
53
54
55
56
57
58
59
60

of BLUF protein photoactivation, in which an additional hydrogen bond is formed to the C4=O in the light state, and the red shift in the C4=O frequency from 1700 cm^{-1} has been observed previously in PixD,⁴⁸ as well as in other BLUF proteins, such as AppA_{BLUF}.^{23, 51-53}




Figure 4. TRIR of the Wild-type PixD Light State.

(A) EAS of light adapted PixD generated by constant illumination with 450 nm light focused on the sample in the TRIR experiment. (B) Comparison of the kinetics of the bands at 1512, 1525, cm^{-1} . Raw data are shown as solid lines while global fitting results are shown as dashed lines.

The excited state of the light state decays significantly faster than the corresponding excited state of the dark state and indeed has completely decayed during the time scale of the TRIR experiment, where three time constants are now needed to adequately describe the data

1
2
3 (Table 1). It has been proposed that in the light state of PixD a concerted electron proton
4 transfer (CEPT) takes place directly leading to the formation of FADH^\bullet .^{35,37} In Figure 4, bands
5 at 1512, and 1525 cm^{-1} are observed within 3 ps, which are assigned to the FAD radical.
6
7
8
9
10 However, unlike the dark state, the bands associated with the radical do not evolve at different
11 rates but have the same kinetics, consistent with a concerted process such as CEPT. Thus,
12
13 contrary to the dark state, where we observe $\text{FAD}^* \rightarrow \text{FAD}^{\bullet-} \rightarrow \text{FADH}^\bullet$, in the light state FAD^*
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
decays directly to FADH^\bullet . These data agree well with the mechanism proposed by Mathes et
al.³⁷

Effect of Y8 fluorination on PixD photoactivation

Y8 in PixD is essential for photoactivity and has been proposed to function as both the
electron and proton donor in the concerted formation of the FAD radical state (PCET). To probe
the role of Y8 in PCET, we site-specifically replaced this residue with 3-FY8, 2,3-F₂Y8, 3,5-
F₂Y8, and 2,3,5-F₃Y8 in order to modulate both the redox potential and $\text{p}K_a$. The F-Tyr analogs
were synthesized and incorporated into the protein,²¹ and ESI MS/MS analysis did not detect any
native Tyr in the F-Tyr substituted proteins indicating that the F-Tyr content of each variant was
 $\geq 99\%$ (Figure S3).

Electronic Spectrum: We first examined the electronic absorption spectrum of each PixD variant.
In Figure 5A the dark state spectrum of each F-Tyr substituted PixD is compared with wild-type
PixD and FAD in solution. The electronic spectra of the proteins are essentially identical and are
characterized by a long wavelength shoulder that is not resolved in the solution spectrum of

FAD. Thus, we can conclude that F-Tyr substitution has not perturbed the FAD binding pocket in a way that alters the electronic spectrum.




Figure 5. Electronic Spectra of Wild-type PixD and the F-Tyr PixD Variants (A)

Comparison of the dark state absorbance spectra of wild-type PixD and the F-Tyr PixD variants.

(B) Comparison of the light state absorbance spectra of wild-type PixD and the F-Tyr PixD

variants acquired immediately after ~ 10 s of ~ 500 mW illumination of the entire 5 cm^2 of the sample window with 450 nm light.

1
2
3 We next analyzed the change in the spectrum caused by photoexcitation. In BLUF
4 proteins, irradiation with blue light results in a ~10 nm red shift in the 450 nm FAD peak, as is
5
6 observed for wild-type PixD in **Figure 5B**. Light induced changes are also observed for all the
7
8 F-Tyr PixD variants with the exception of 2,3,5-F₃Y8-PixD, although the magnitude of the red
9
10 shift resolved in these steady-state measurements is apparently lower for 3-FY8, 2,3-F₂Y8 and
11
12 3,5-F₂Y8 PixD.
13
14
15
16
17
18
19

20 *TRIR and TRMPS of the Forward Photoreaction:* The TRIR spectra of the F-Tyr-substituted
21
22 dark adapted PixD proteins are shown in **Figure 6** where it can be seen that the major bleaches
23
24 and transients observed in the initial (2 ps) spectrum of the wild-type PixD are conserved
25
26 amongst all the substituted proteins. The similarity of the 2 ps TRIR spectra (**Figure 6**) across
27
28 all proteins indicates that the ground state environment of the flavin has not been perturbed by
29
30 modification of Y8, consistent with the similarity in the dark state absorption spectra (**Figure 5**).
31
32 However, a small (4 cm⁻¹) increase in the C4=O flavin carbonyl model, from 1700 to 1704 cm⁻¹,
33
34 is observed in the spectra of the F-Tyr PixD proteins, which may indicate that the hydrogen bond
35
36 to the C4=O carbonyl is slightly weakened due to the incorporation of F-Tyr analogs, although
37
38 the effect is not a function of pK_a.
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 6: TRIR Spectra Wild-type PixD and the F-Tyr PixD Variants. TRIR Spectra of wild-type PixD and the F-Tyr PixD variants at 2, 40, 200, 1000, and 3400 ps for (A) wild-type, (B) 3-FY8 PixD, (C) 2,3-F₂Y8 PixD, (D) 3,5-F₂Y8 PixD and (E) 2,3,5-F₃Y8 PixD.

1
2
3 The TRIR spectra at later times demonstrate that alteration in the reduction potential
4 and/or pK_a of Y8 has a dramatic impact on the forward photoreaction (**Figure 6**), in contrast to
5 the BLUF protein AppA_{BLUF} where replacement of the conserved Y21 with F-Tyr analogs results
6 in only a small ~3-fold change in the kinetics for photoactivation over the entire pK_a range.²¹ 3-
7 FY8 PixD exhibits the highest similarity to the wild-type protein although the rate of spectral
8 evolution is described by significantly longer time constants compared to wild-type PixD (**Table**
9 **1**). The light state, characterized by the transient at 1690 cm^{-1} forms within 230 ps, although a
10 considerable amount of the excited state is still present at this stage, again consistent with the
11 presence of multiple conformations and an associated distribution of rates. The transients at
12 1515 and 1535 cm^{-1} are also still observed for 3-FY8 PixD suggesting that both the electron and
13 proton transfer steps occur, albeit more slowly than for wild-type PixD. This observation is
14 consistent with the studies of Kennis and coworkers, who also observed an increase in the
15 average excited state lifetime in 3-FY8 PixD,³⁵ a result which broadly recapitulates similar
16 studies on ribonucleotide reductase, where an increase in reduction potential leads to a decrease
17 in the rate of ribonucleotide reduction.⁵⁴

18
19
20 In our studies on AppA_{BLUF}, we considered the impact of F-Tyr substitution on the TRIR
21 spectra using Marcus theory to analyze a sequential scheme involving charge separation (k_{cs})
22 followed by charge recombination (k_{cr}) and concluded that either $k_{cs} \ll k_{cr}$ since no radical
23 intermediate could be detected, or that the reaction proceeded without population of a distinct
24 charge transfer intermediate. Consistent with the absence of electron transfer in the AppA_{BLUF}
25 mechanism, F-Tyr substitution only had a small (3-fold) effect on the overall rate of light state
26 formation.²¹ Although the situation is more complex in PixD, since both electron and proton
27 transfer are clearly involved, in contrast to AppA_{BLUF}, the data do indicate that the rate of charge
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 separation must be faster than recombination, since a flavin radical clearly forms on
4 photoactivation (**Figure 3**). In addition, the observation that the rate of radical formation and
5 decay are both decreased by fluorination suggests that both k_{cs} and k_{cr} are in the Marcus inverted
6 region.
7
8
9
10

11
12 The TRIR spectra of 2,3-F₂Y8, 3,5-F₂Y8, and 2,3,5-F₃Y8 PixD differ more dramatically
13 from the wild-type data than those of 3-FY8 PixD (**Figure 6**). The most significant differences
14 include the lack of a distinct 1535 cm⁻¹ transient, assigned to FADH[•], and the lack of the 1690
15 cm⁻¹ transient assigned to formation of the light state. TRMPS spectra (**Figure S4**) show that no
16 further evolution occurs following the final TRIR spectrum, and the absence of the 1690 cm⁻¹
17 transient therefore shows that F-Tyr substitution prevents the formation of a detectable
18 population of light state, which contrasts sharply with AppA_{BLUF}, where light state formation
19 persists for all F-Tyr analogs. In addition, the kinetics for the spectral evolution that can be
20 observed for 2,3-F₂Y8, 3,5-F₂Y8, and 2,3,5-F₃Y8 PixD are significantly slower compared to 3-
21 FY8 and wild-type PixD, and only three distinct lifetimes are now required to fit the nanosecond
22 domain TRIR data (**Table 1**).
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37

38 The transient assigned to FAD^{•-} is still observed at 1515 cm⁻¹ for 2,3-F₂Y8, 3,5-F₂Y8 and
39 2,3,5-F₃Y8 PixD (**Figure 6**). Analysis of the kinetics for this band show both rise and decay
40 components (**Figure S5**), consistent with the assignment to an intermediate, although differences
41 between the rates of excited state decay (1380 cm⁻¹) and ground state recovery (1550 cm⁻¹)
42 (**Figure S6 and S7**) are less obvious for the di- and tri-substituted F-Tyr PixD variants than for
43 the wild-type or 3-FY8 PixD. Finally, in the TRMPS data for 3,5-F₂Y8, and 2,3,5-F₃Y8 PixD
44 (**Figure S4**), a feature around 1490 cm⁻¹ can be observed, indicative of triplet state formation that
45 is absent in wild-type PixD; this result is consistent with an increase in lifetime of the singlet
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 excited state, allowing intersystem crossing to compete. Due to the slower excited state decay in
4
5 3-FY8, 2,3-F₂Y8, 3,5-F₂Y8, and 2,3,5-F₃Y8 PixD (**Figure S6**) we also observe a significant
6
7 increase in flavin fluorescence with an ~3-fold increase in quantum yield between the wild-type
8
9 protein and the F-Tyr substituted PixD variants (**Table S2**).
10
11

12
13 Taken together, the data suggest that the change in reduction potential and/or pK_a of Y8
14
15 upon di- and tri-fluoro substitution apparently halts the photocycle after $FAD^{\bullet-}$ is formed. A
16
17 simple explanation for the lack of further progression of the photocycle is that the pK_a of 2,3-
18
19 F₂Y8, 3,5-F₂Y8, and 2,3,5-F₃Y8 has dropped sufficiently that this residue is no longer protonated
20
21 and thus cannot transfer a proton to $FAD^{\bullet-}$, supporting the importance of PCET in the
22
23 photoactivation of PixD. The proton transfer is evidently critical for trapping the light state in
24
25 which Q50 has rotated to H-bond to the C4=O carbonyl group.
26
27
28
29
30
31

32 **Recovery of the Dark State**

33
34 The electronic spectra of wild-type PixD and the F-Tyr PixD variants indicate that
35
36 photoexcitation leads to the formation of a red-shifted absorption spectrum for all proteins except
37
38 2,3,5-F₃Y8 PixD (**Figure 5**), which is evidence for the formation of the PixD light state in those
39
40 cases. However, under the same illumination conditions the red shift is clearly less marked for
41
42 2,3-F₂Y8 and 3,5-F₂Y8 PixD than for wild-type and 3-FY8 PixD. This is consistent with the
43
44 slower excited state decay, indicating a reduced quenching rate constant for light state formation,
45
46 thus allowing other processes (such as intersystem crossing) to compete. One apparent paradox
47
48 is that for all PixD variants, except 2,3,5-F₃Y8 PixD, the red shift in the absorption spectra
49
50 indicates light-state formation, whilst, based on the TRIR spectra, we concluded that the
51
52 photocycle for 2,3-F₂Y8 and 3,5-F₂Y8, as well as 2,3,5-F₃Y8 PixD stalled after the initial
53
54
55
56
57
58
59
60

1
2
3 electron transfer event. This suggests that the light state populated during the continuous
4 illumination, generating the red-shifted electronic spectra, is not sufficiently stable to result in an
5 observable signal in the single-shot TRIR spectra of 2,3-F₂Y8 and 3,5-F₂Y8 PixD. To test this
6 hypothesis, we obtained TRIR spectra under continuous illumination (**Figure 7**). In the TRIR
7 spectra of both wild-type PixD and 3-FY8 PixD, continuous illumination at 450 nm leads to the
8 observation of a bleach at 1686-1689 cm⁻¹ which is assigned to the C4=O carbonyl group of the
9 light state, consistent with the transient observed at ~1690 cm⁻¹ in the pump-probe TRIR spectra
10 (**Figures 3, 4 and 6**). However, in 2,3-F₂Y8 PixD two bands are observed, at 1686 and 1700 cm⁻¹
11 while in the 3,5-F₂Y8 and 2,3,5-F₃Y8 PixD proteins the major band in this region is observed at
12 1704 cm⁻¹, with a barely resolvable shoulder at around 1686 cm⁻¹ in the spectrum of 3,5-F₂Y8
13 PixD. These data indicate that the light state can be partially populated in 2,3-F₂Y8 PixD, and
14 possibly in 3,5-F₂Y8 PixD, under continuous illumination, consistent with the steady state
15 spectra; however, such a light state may be unstable with respect to reversion to the dark state.
16 One plausible explanation is the generation of an increasing fraction of a population of PixD
17 which is incapable of light state formation with decreasing Y8 pK_a.




Figure 7: TRIR Spectra of Wild-type PixD and the F-Tyr PixD Variants Under Continuous Illumination. TRIR spectra were obtained whilst continuously illuminating the protein samples with ~200mW of 450 nm light (L). The top spectrum is the TRIR spectrum of wild-type PixD obtained using the standard pump-probe approach (dark state, D). A key observation is the change in position of the C4=O bleach from 1701 cm^{-1} in the PixD dark state to 1686 cm^{-1} upon formation of the light state. Continuous illumination of the F-Tyr substituted PixD variants results in full conversion of 3-FY8 PixD, but only partial population of the light state in 2,3-F₂Y8 PixD, and possibly in 3,5-F₂Y8 PixD. In addition, there is no light state signature in 2,3,5-F₃Y8 PixD.

Based on the conclusion that the alteration in the electronic spectra upon photoexcitation represented light state formation, we monitored the time-dependent recovery of the absorption spectra to quantify the rate of dark state recovery. It has already been noted that the rate of

1
2
3 recovery in wild-type PixD is substantially greater than for AppA_{BLUF} (Table 2). For 3-FY8 and
4
5 2,3-F₂Y8 PixD this rate is further accelerated 6 and 7 fold, consistent with formation of a light
6
7 state which is increasingly unstable with respect to dark state recovery in F-Tyr PixD. Further,
8
9 as was observed for AppA_{BLUF}, reduction in the Y8 pK_a monotonically accelerates the rate of
10
11 dark state recovery (**Table 2**). However, the relative increase in the rate of recovery is much less
12
13 marked in PixD: in H₂O, the rate increases 970-fold between wild-type AppA_{BLUF} and 3,5-F₂Y21
14
15 AppA_{BLUF},²¹ compared to only 15-fold for PixD, while in D₂O the change in rate is 850-fold for
16
17 AppA_{BLUF} and 6-fold for PixD. The corresponding Brønsted plots for PixD have shallower
18
19 slopes of 0.43 in H₂O and 0.28 in D₂O (**Figure 8**), compared to slopes of ~1 for AppA_{BLUF}.²¹
20
21 Thus, alteration of the Y8 pK_a has a smaller effect on the rate of dark state recovery in PixD
22
23 compared to AppA_{BLUF}. However, the observation of large normal solvent isotope effects on the
24
25 rate of recovery for both PixD (2-11) and AppA_{BLUF} (2-9), together with Brønsted α values of
26
27 0.43 and 1, respectively, indicates that two proteins have similar recovery mechanisms which
28
29 differ primarily in the degree of proton transfer from the conserved Tyr (Y21 or Y8) in the rate
30
31 limiting transition state.
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 2. Dark state recovery rate constants^a

	pK_a^c	PixD			AppA _{BLUF} ^b		
		k_{H_2O} (s ⁻¹)	k_{D_2O} (s ⁻¹)	sie ^d	k_{H_2O} (s ⁻¹)	k_{D_2O} (s ⁻¹)	sie ^d
Wild-type	9.9	0.038 ± 0.004	0.010 ± 0.0003	3.7 ± 0.1	0.00065 ± 0.00006	0.00013 ± 0.00001	5.0 ± 1
3-FY	8.4	0.22 ± 0.012	0.023 ± 0.0008	9.5 ± 0.1	0.034 ± 0.002	0.0039 ± 0.0001	8.8 ± 0.7
2,3-F₂Y	7.8	0.27 ± 0.013	0.047 ± 0.0065	5.8 ± 0.1	0.22 ± 0.01	0.040 ± 0.001	5.5 ± 0.2
3,5-F₂Y	7.2	0.58 ± 0.06	0.058 ± 0.0036	9.9 ± 0.2	0.63 ± 0.02	0.11 ± 0.01	5.8 ± 0.3
2,3,5-F₃Y^e	6.4	ND	ND	ND	2.6 ± 0.1	1.15 ± 0.03	2.2 ± 0.1

^aRecovery rate constants were obtained from the change in absorption spectra once irradiation at 455 nm had been terminated by fitting the data to a single exponential function. Errors are based on measurements made in triplicate or quadruplicate.

^bAppA_{BLUF}. Recovery data are for the BLUF domain of AppA and are taken from Gil et al.²¹

^cThe pK_a is that for the amino acid in solution.⁵⁴

^dsie, solvent isotope effect.

^eND, not detected. For 2,3,5-F₃Y8 PixD no light state was detected.




Figure 8. Rate of Dark State Recovery for Wild-type PixD and the F-Tyr PixD Variants.

(A) Brønsted plots in H_2O and D_2O . The log of recovery rate constant $-\log(k)$ plotted against corresponding $-pK_a$. The rate increases 15.4-fold from wild-type (pK_a 9.9) to 3,5-F₂Y8 PixD (pK_a 7.2) with a slope of 0.43 ± 0.04 (R^2 0.99), in D_2O the rate increases 5.6-fold from wild-type to 3,5-F₂Y8 PixD with a slope of 0.28 ± 0.03 (R^2 0.972). (B) Comparisons of the dark state recovery rate for PixD and AppA_{BLUF}, data measured in H_2O , the values for AppA_{BLUF} taken from Gil et al.²¹

Comparison of the PixD and AppA_{BLUF} Photocycles

In the BLUF proteins only the forward photoreaction is light driven, while the recovery of the dark state occurs in the absence of illumination. Thus, it may be assumed that photoactivation and dark state recovery follow different reaction pathways. In both PixD and AppA_{BLUF}, the conserved Tyr (Y8 or Y21) is essential for reactivity, and in both cases replacement of this Tyr with, for example Phe, leads to a loss of photoactivity. However, despite these overall similarities, profound differences are found in the photocycles of PixD and AppA_{BLUF}. In PixD there is clear evidence for coupled electron and proton transfer resulting in the intermediate formation of a flavin radical during photoactivation. In contrast, radical formation is not apparent in the photoactivation of AppA_{BLUF}. Modulation of the pK_a and reduction potential of the conserved Tyr adds fine detail that allows a deeper comparison of AppA_{BLUF} and PixD. Thus, in addition to the lack of radical formation in AppA_{BLUF}, replacement of Y21 with F-Tyr analogs has little effect on AppA_{BLUF} activation. In contrast, and in agreement with a radical mechanism, changes to the pK_a and reduction potential of Y8 have a major impact on PixD activation, where 2,3-F₂Y8, 3,5-F₂Y8 and 2,3,5-F₃Y8 PixD are seemingly unable to proceed past the initial electron transfer event to create a significant long-lived population of light state, and 2,3,5-F₃Y8 PixD is unable to populate the light state at all, even under continuous illumination.

In contrast to major differences in the mechanism of photoactivation, the recovery of the dark state in AppA_{BLUF} and PixD appear to follow essentially the same mechanism, although there are large absolute differences in the recovery rates. Large normal isotope effects are observed on the rate of recovery for both proteins, and in each case the increase in acidity of the conserved Tyr results in an increase in the observed rate constant, albeit that the change in rate

1
2
3 between wild-type and 3,5-F₂Tyr proteins (pK_a 9.9 to 7.2) is 15-fold for PixD compared to 970-
4 fold for AppA_{BLUF}. In AppA_{BLUF}, a Brønsted analysis indicated that complete transfer of a
5 proton occurred in the rate limiting transition state on the reaction coordinate for dark state
6 recovery. Based on this result, we proposed a mechanism of recovery in which proton transfer
7 from Y21 to Q63 played a key role in keto-enol tautomerism and subsequent rotation of Q63.²¹
8 In PixD, the Brønsted coefficient (α) is 0.43 compared to 1 in AppA_{BLUF} indicating an earlier
9 transition state in the recovery of dark adapted PixD and thus that proton transfer involving Y8
10 has a less significant role in the mechanism of recovery. In addition, the differences in α indicate
11 that the transition state in AppA_{BLUF} is similar in structure to that of the dark state whereas the
12 transition state in PixD more closely resembles the photoactivated state of the protein.⁵⁵
13
14
15
16
17
18
19
20
21
22
23
24
25
26

27 It is difficult to rationalize the different roles for the conserved Tyr in AppA_{BLUF} and
28 PixD, given the similarity in the position of Y21 and Y8 from the X-ray structures. However,
29 the flavin binding pockets in the two proteins are not strictly conserved. For instance, in PixD
30 the flavin C2=O group is hydrogen-bonded to two amino acids, N31 and R65, whereas in AppA
31 the C2=O is only hydrogen bonded to H44, the homolog of N31. Analysis of existing data for
32 other BLUF proteins suggests that the presence of two hydrogen bonds to the C2=O may
33 correlate with radical formation. The N31, R65 hydrogen bonding motif is repeated in TePixD
34 (Tll0078),⁵⁶ which also forms a radical during the photocycle.⁵⁷ In contrast, while N31 is
35 replaced by an Arg in BlrB, a second hydrogen bond donor to C2=O is not present,⁶ and there is
36 no evidence for radical formation during photoactivation of this BLUF protein.⁵⁸ X-ray
37 structures are not available for BlsA and PapB, however sequence alignment suggests that two
38 hydrogen bonds may be present in PapB, where N31 is conserved and R65 is replaced by a Lys,
39 while only one hydrogen bond is possible in BlsA where the residue homologous to N31 is a
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Phe. Again, while radical formation is detected during the photocycle of PapB,³⁹ there is no
4
5 evidence for a radical intermediate in BlsA.³⁴ Secondly, as noted in the introduction, W91 is
6
7 predominantly in the Trp_{out} conformation in the PixD X-ray structures, in contrast to AppA
8
9 where both Trp_{in} and Trp_{out} conformations are populated.^{5, 31} Although the X-ray structures do
10
11 not rule out the possibility that the Trp side chain can adopt multiple conformations, including
12
13 Trp_{in} and Trp_{out}, Mehlhorn et al,⁵⁹ have suggested that light-induced changes in the BLUF β -
14
15 sheet may be communicated from the flavin via another residue (N32) rather than W91, based on
16
17 the supposition that W91 remains in the Trp_{out} conformation throughout the PixD photocycle.
18
19 Thus, if W91 is not hydrogen bonded to Q50 in the dark state, then keto-enol tautomerism and
20
21 rotation of Q50 may play a less critical role in stabilizing the PixD light state, hence reducing the
22
23 impact of the Y8 pK_a on dark state recovery. Finally, it should be noted that the studies on
24
25 AppA_{BLUF}, PixD, PapB and BlsA have been conducted in the absence of their output partners,
26
27 PpsR, PixE and PapA,⁶⁰ respectively (the output partner for BlsA has not yet been identified),
28
29 and indeed in AppA_{BLUF}, without the C-terminal domain of the full length AppA protein. Thus it
30
31 remains to be seen how protein-protein interactions in physiologically relevant complexes of the
32
33 BLUF proteins might affect the photocycles that have been so thoroughly characterized in the
34
35 isolated BLUF domains. Our observation that relatively small changes in structure modify the
36
37 mechanism of photoactivation suggest that this might be a fruitful line of research.
38
39
40
41
42
43
44
45
46
47

48 **Conclusion**

49
50 The role of Y8 in the photocycle of the BLUF protein PixD has been probed via a
51
52 combination of unnatural amino acid mutagenesis and ultrafast infrared spectroscopy. In the
53
54 wild-type protein, the TRIR spectra are consistent with a photoactivation mechanism in which
55
56
57
58
59
60

1
2
3 electron-proton transfer from Y8 occurs sequentially, with electron transfer (2.5 ps) to the
4
5 excited FAD* first generating FAD^{•-} followed by proton transfer (20 ps) to generate the neutral
6
7 semiquinone FADH[•]. This mechanism is quite different to that operating in AppA_{BLUF}, where
8
9 no radical intermediates are observed and the excited state decay is longer. Replacement of Y8
10
11 with F-Tyr analogs that increase the reduction potential of the phenolic group reduce the rate of
12
13 electron transfer. In addition, for the F-Tyr analogs with pK_a values in solution below 8 (2,3-
14
15 F₂Y8, 3,5-F₂Y8 and 2,3,5-F₃Y8 PixD), no progression of the photocycle is observed after
16
17 formation of the FAD anionic semiquinone FAD^{•-}. Assuming that the F-Tyr analogs have
18
19 similar pK_a values in the protein, this observation confirms that Y8 is the source of both electron
20
21 and proton in the PCET mechanism, since at pH 8 the Tyr analogs will be deprotonated.
22
23 Although no stable light state can be detected in the TRIR spectra of 2,3-F₂Y8, 3,5-F₂Y8 and
24
25 2,3,5-F₃Y8 PixD, a photostationary population of light state can be generated in 2,3-F₂Y8 and
26
27 3,5-F₂Y8 PixD indicating that the reaction coordinate leading to the light state is still accessible,
28
29 but that the light state reversion to the ground state is accelerated. In contrast to the profound
30
31 impact on the forward photoreaction, F-Tyr substitution has a more modest effect on the rate of
32
33 dark state recovery. These observations differ fundamentally from the impact of F-Tyr
34
35 substitution on the photocycle of AppA_{BLUF} where the change in pK_a and reduction potential have
36
37 only a minor effect on excited state decay, but increase the rate of dark state recovery 4,000-fold.
38
39 Thus, despite high levels of structural conservation in the FAD binding pocket, profound
40
41 differences can exist between the photophysics of BLUF proteins. For AppA_{BLUF} and PixD, the
42
43 identity of the residues that bind to the C2=O carbonyl group and/or the position of the
44
45 conserved Trp (W91 and W104) in the two proteins may play a fundamental role in determining
46
47 the function of the conserved Tyr in the photocycle.
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 ASSOCIATED CONTENT
4
5

6 **Supporting Information Available:** ESI MS/MS analysis of fluorotyrosine incorporation into
7
8 PixD. Additional TRMPS and TRIR datasets. This material is available free of charge via the
9
10 Internet at <http://pubs.acs.org>.
11
12
13

14
15
16
17
18 AUTHOR INFORMATION
1920
21 **Corresponding Authors**
22

23
24 *PJT: Telephone: (631) 632-7907; Fax: (631) 632-7960; Email: peter.tonge@stonybrook.edu;
25

26 SRM:44(0)1603 593141; Fax: 44(0)1603 592004; Email: s.meech@uea.ac.uk
27
28
29
30
31
32

33 ACKNOWLEDGMENT
34
35

36 Funded by the EPSRC (EP/G002916 to SRM) and NSF (CHE-1223819 to PJT). We are grateful
37
38 to STFC for access to the ULTRA laser facility. We are grateful to Professor Ray Owens for
39
40 assistance in protein preparation and access to the Oxford Protein Production Facility – UK, and
41
42 to Arthur Makarenko at CSHL for assistance with the mass spectrometry analysis. JI was
43
44 supported by an NIH Chemistry-Biology Interface training grant (T32GM092714). AL is a
45
46 Bolyai Janos Research Fellow and was supported by OTKA NN113090.
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

1. Piano, V.; Palfey, B. A.; Mattevi, A., *Trends Biochem Sci* **2017**, *42*, 457.
2. Massey, V., *FASEB J* **1995**, *9*, 473.
3. Ghisla, S.; Massey, V., *Eur J Biochem* **1989**, *181*, 1.
4. Heelis, P. F., *Chem Soc Rev* **1982**, *11*, 15.
5. Anderson, S.; Dragnea, V.; Masuda, S.; Ybe, J.; Moffat, K.; Bauer, C., *Biochemistry* **2005**, *44*, 7998.
6. Jung, A.; Domratcheva, T.; Tarutina, M.; Wu, Q.; Ko, W. H.; Shoeman, R. L.; Gomelsky, M.; Gardner, K. H.; Schlichting, I., *Proc Natl Acad Sci U S A* **2005**, *102*, 12350.
7. Yuan, H.; Dragnea, V.; Wu, Q.; Gardner, K. H.; Bauer, C. E., *Biochemistry* **2011**, *50*, 6365.
8. Brust, R.; Lukacs, A.; Haigney, A.; Addison, K.; Gil, A.; Towrie, M.; Clark, I. P.; Greetham, G. M.; Tonge, P. J.; Meech, S. R., *J Am Chem Soc* **2013**, *135*, 16168.
9. Udvarhelyi, A.; Domratcheva, T., *J Phys Chem B* **2013**, *117*, 2888.
10. Domratcheva, T.; Hartmann, E.; Schlichting, I.; Kottke, T., *Sci Rep* **2016**, *6*, 22669.
11. Goyal, P.; Hammes-Schiffer, S., *Proc Natl Acad Sci U S A* **2017**, *114*, 1480.
12. Hall, C. R.; Heisler, I. A.; Jones, G. A.; Frost, J. E.; Gil, A. A.; Tonge, P. J.; Meech, S. R., *Chem Phys Lett* **2017**, *683*, 365.
13. Stierl, M.; Stumpf, P.; Udvari, D.; Gueta, R.; Hagedorn, R.; Losi, A.; Gartner, W.; Petereit, L.; Efetova, M.; Schwarzel, M.; Oertner, T. G.; Nagel, G.; Hegemann, P., *J Biol Chem* **2011**, *286*, 1181.
14. Gomelsky, M.; Klug, G., *Trends Biochem Sci* **2002**, *27*, 497.
15. Masuda, S., *Plant Cell Physiol* **2013**, *54*, 171.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
16. Cao, Z.; Livoti, E.; Losi, A.; Gartner, W., *Photochem Photobiol* **2010**, *86*, 606.
17. Tang, Y.; Cao, Z.; Livoti, E.; Krauss, U.; Jaeger, K. E.; Gartner, W.; Losi, A., *Photochem Photobiol Sci* **2010**, *9*, 47.
18. Kennis, J. T.; Groot, M. L., *Curr Opin Struct Biol* **2007**, *17*, 623.
19. van der Horst, M. A.; Hellingwerf, K. J., *Acc Chem Res* **2004**, *37*, 13.
20. Okajima, K.; Yoshihara, S.; Fukushima, Y.; Geng, X.; Katayama, M.; Higashi, S.; Watanabe, M.; Sato, S.; Tabata, S.; Shibata, Y.; Itoh, S.; Ikeuchi, M., *J Biochem* **2005**, *137*, 741.
21. Gil, A. A.; Haigney, A.; Laptinok, S. P.; Brust, R.; Lukacs, A.; Iuliano, J. N.; Jeng, J.; Melief, E. H.; Zhao, R. K.; Yoon, E.; Clark, I. P.; Towrie, M.; Greetham, G. M.; Ng, A.; Truglio, J. J.; French, J. B.; Meech, S. R.; Tonge, P. J., *J Am Chem Soc* **2016**, *138*, 926.
22. Yuan, H.; Anderson, S.; Masuda, S.; Dragnea, V.; Moffat, K.; Bauer, C., *Biochemistry* **2006**, *45*, 12687.
23. Stelling, A. L.; Ronayne, K. L.; Nappa, J.; Tonge, P. J.; Meech, S. R., *J Am Chem Soc* **2007**, *129*, 15556.
24. Domratcheva, T.; Grigorenko, B. L.; Schlichting, I.; Nemukhin, A. V., *Biophys. J.* **2008**, *94*, 3872.
25. Lukacs, A.; Haigney, A.; Brust, R.; Zhao, R. K.; Stelling, A. L.; Clark, I. P.; Towrie, M.; Greetham, G. M.; Meech, S. R.; Tonge, P. J., *J Am Chem Soc* **2011**, *133*, 16893.
26. Masuda, S.; Bauer, C. E., *Cell* **2002**, *110*, 613.
27. Kraft, B. J.; Masuda, S.; Kikuchi, J.; Dragnea, V.; Tollin, G.; Zaleski, J. M.; Bauer, C. E., *Biochemistry* **2003**, *42*, 6726.
28. Masuda, S.; Hasegawa, K.; Ishii, A.; Ono, T. A., *Biochemistry* **2004**, *43*, 5304.
29. Hasegawa, K.; Masuda, S.; Ono, T. A., *Biochemistry* **2004**, *43*, 14979.

- 1
2
3 30. Unno, M.; Masuda, S.; Ono, T. A.; Yamauchi, S., *J Am Chem Soc* **2006**, *128*, 5638.
4
5
6 31. Jung, A.; Reinstein, J.; Domratcheva, T.; Shoeman, R. L.; Schlichting, I., *J. Mol. Biol.* **2006**,
7
8 *362*, 717.
9
10 32. Grinstead, J. S.; Hsu, S. T.; Laan, W.; Bonvin, A. M.; Hellingwerf, K. J.; Boelens, R.;
11
12 Kaptein, R., *Chembiochem* **2006**, *7*, 187.
13
14
15 33. Masuda, S.; Hasegawa, K.; Ono, T. A., *Plant Cell Physiol* **2005**, *46*, 1894.
16
17
18 34. Lukacs, A.; Brust, R.; Haigney, A.; Laptanok, S. P.; Addison, K.; Gil, A.; Towrie, M.;
19
20 Greetham, G. M.; Tonge, P. J.; Meech, S. R., *J Am Chem Soc* **2014**, *136*, 4605.
21
22
23 35. Mathes, T.; van Stokkum, I. H.; Stierl, M.; Kennis, J. T., *J Biol Chem* **2012**, *287*, 31725.
24
25
26 36. Laptanok, S. P.; Lukacs, A.; Brust, R.; Haigney, A.; Gil, A.; Towrie, M.; Greetham, G. M.;
27
28 Tonge, P. J.; Meech, S. R., *Faraday Discuss* **2015**, *177*, 293.
29
30
31 37. Mathes, T.; Zhu, J. Y.; van Stokkum, I. H. M.; Groot, M. L.; Hegemann, P.; Kennis, J. T. M.,
32
33 *J Phys Chem Lett* **2012**, *3*, 203.
34
35
36 38. Gauden, M.; van Stokkum, I. H. M.; Key, J. M.; Luhrs, D. C.; Van Grondelle, R.; Hegemann,
37
38 P.; Kennis, J. T. M., *Proc Natl Acad Sci U S A* **2006**, *103*, 10895.
39
40
41 39. Fujisawa, T.; Takeuchi, S.; Masuda, S.; Tahara, T., *J Phys Chem B* **2014**, *118*, 14761.
42
43
44 40. Brust, R.; Haigney, A.; Lukacs, A.; Gil, A.; Hossain, S.; Addison, K.; Lai, C. T.; Towrie, M.;
45
46 Greetham, G. M.; Clark, I. P.; Illarionov, B.; Bacher, A.; Kim, R. R.; Fischer, M.;
47
48 Simmerling, C.; Meech, S. R.; Tonge, P. J., *J Phys Chem Lett* **2014**, *5*, 220.
49
50
51 41. The PyMOL Molecular Graphics System, Version 1.8 Schrodinger, LLC.
52
53
54
55
56
57
58
59 42. Minnihan, E. C.; Young, D. D.; Schultz, P. G.; Stubbe, J., *J Am Chem Soc* **2011**, *133*, 15942.
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
43. Greetham, G. M.; Burgos, P.; Cao, Q.; Clark, I. P.; Codd, P. S.; Farrow, R. C.; George, M. W.; Kogimtzis, M.; Matousek, P.; Parker, A. W.; Pollard, M. R.; Robinson, D. A.; Xin, Z. J.; Towrie, M., *Appl Spectrosc* **2010**, *64*, 1311.
44. Greetham, G. M.; Sole, D.; Clark, I. P.; Parker, A. W.; Pollard, M. R.; Towrie, M., *Rev Sci Instrum* **2012**, *83*, 103107.
45. Snellenburg, J. J.; Liptonok, S. P.; Seger, R.; Mullen, K. M.; van Stokkum, I. H. M., *J Stat Softw* **2012**, *49*, 1.
46. Bonetti, C.; Mathes, T.; van Stokkum, I. H. M.; Mullen, K. M.; Groot, M. L.; van Grondelle, R.; Hegemann, P.; Kennis, J. T. M., *Biophys. J.* **2008**, *95*, 4790.
47. Bonetti, C.; Stierl, M.; Mathes, T.; van Stokkum, I. H.; Mullen, K. M.; Cohen-Stuart, T. A.; van Grondelle, R.; Hegemann, P.; Kennis, J. T., *Biochemistry* **2009**, *48*, 11458.
48. Hasegawa, K.; Masuda, S.; Ono, T. A., *Plant Cell Physiol* **2005**, *46*, 136.
49. Haigney, A.; Lukacs, A.; Zhao, R. K.; Stelling, A. L.; Brust, R.; Kim, R. R.; Kondo, M.; Clark, I.; Towrie, M.; Greetham, G. M.; Illarionov, B.; Bacher, A.; Romisch-Margl, W.; Fischer, M.; Meech, S. R.; Tonge, P. J., *Biochemistry* **2011**, *50*, 1321.
50. Lukacs, A.; Zhao, R. K.; Haigney, A.; Brust, R.; Greetham, G. M.; Towrie, M.; Tonge, P. J.; Meech, S. R., *J Phys Chem B* **2012**, *116*, 5810.
51. Laan, W.; van der Horst, M. A.; van Stokkum, I. H.; Hellingwerf, K. J., *Photochem Photobiol* **2003**, *78*, 290.
52. Unno, M.; Sano, R.; Masuda, S.; Ono, T. A.; Yamauchi, S., *J Phys Chem B* **2005**, *109*, 12620.
53. Masuda, S.; Hasegawa, K.; Ono, T. A., *Biochemistry* **2005**, *44*, 1215.

- 1
2
3 54. Seyedsayamdost, M. R.; Yee, C. S.; Reece, S. Y.; Nocera, D. G.; Stubbe, J., *J Am Chem Soc*
4
5 **2006**, *128*, 1562.
6
7
8 55. Szabo, A., *Proc Natl Acad Sci U S A* **1978**, *75*, 2108.
9
10 56. Kita, A.; Okajima, K.; Morimoto, Y.; Ikeuchi, M.; Miki, K., *J Mol Biol* **2005**, *349*, 1.
11
12 57. Nagai, H.; Fukushima, Y.; Okajima, K.; Ikeuchi, M.; Mino, H., *Biochemistry* **2008**, *47*,
13 12574.
14
15
16 58. Mathes, T.; van Stokkum, I. H.; Bonetti, C.; Hegemann, P.; Kennis, J. T., *J Phys Chem B*
17 **2011**, *115*, 7963.
18
19
20 59. Mehlhorn, J.; Lindtner, T.; Richter, F.; Glass, K.; Steinocher, H.; Beck, S.; Hegemann, P.;
21 Kennis, J. T.; Mathes, T., *J Phys Chem Lett* **2015**, *6*, 4749.
22
23
24
25 60. Kanazawa, T.; Ren, S.; Maekawa, M.; Hasegawa, K.; Arisaka, F.; Hyodo, M.; Hayakawa, Y.;
26 Ohta, H.; Masuda, S., *Biochemistry* **2010**, *49*, 10647.
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table of Contents Graphic

