

Article

Mitragynine/corynantheidine pseudoindoxyls as opioid analgesics with mu agonism and delta antagonism which do not recruit β -arrestin-2

András Váradi, Gina F Marrone, Travis C Palmer, Ankita Narayan, Márton R Szabó, Valerie Le Rouzic, Steven G Grinnell, Joan J Subrath, Evelyn Warner, Sanjay Kalra, Amanda Hunkele, Jeremy Pagirsky, Shainnel O Eans, Jessica M Medina, Jin Xu, Ying Xian Pan, Attila Borics, Gavril W. Pasternak, Jay P. McLaughlin, and Susruta Majumdar

J. Med. Chem., **Just Accepted Manuscript** • DOI: 10.1021/acs.jmedchem.6b00748 • Publication Date (Web): 24 Aug 2016

Downloaded from <http://pubs.acs.org> on August 24, 2016

Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.



ACS Publications

Mitragynine/corynantheidine pseudoindoxyls as opioid analgesics with mu agonism and delta antagonism, which do not recruit β -arrestin-2

András Váradi^a, Gina F. Marrone^a, Travis C. Palmer^a, Ankita Narayan^a, Márton R. Szabó^c, Valerie Le Rouzic^a, Steven G. Grinnell^a, Joan J. Subrath^a, Evelyn Warner^a, Sanjay Kalra^a, Amanda Hunkele^a, Jeremy Pagirsky^a, Shainnel O. Eans^b, Jessica M. Medina^b, Jin Xu^a, Ying-Xian Pan^a, Attila Borics^c, Gavril W Pasternak^a, Jay P. McLaughlin^b and Susruta Majumdar^{a,*}

^aMolecular Pharmacology & Chemistry Program and Department of Neurology, Memorial Sloan Kettering Cancer Center, New York, NY 10065

^bDepartment of Pharmacodynamics, University of Florida, Gainesville, FL 032610

^cInstitute of Biochemistry, Biological Research Centre, Hungarian Academy of Sciences, Szeged, Hungary H-6726

ABSTRACT

Natural products found in *Mitragyna speciosa*, commonly known as *kratom*, represent diverse scaffolds (indole, indolenine and spiro pseudoindoxyl) with opioid activity, providing opportunities to better understand opioid pharmacology. Herein, we report the pharmacology and SAR studies both *in vitro* and *in vivo* of mitragynine pseudoindoxyl (**3**), an oxidative rearrangement product of the corynanthe alkaloid mitragynine. **3** and its corresponding corynantheidine analogs show promise as potent analgesics with a mechanism of action that includes mu opioid receptor agonism-delta opioid receptor antagonism. *In vitro*, **3** and its analogs were potent agonists in [³⁵S]GTPγS assays at the mu opioid receptor but failed to recruit β -arrestin-2, which is associated with opioid side effects. Additionally, **3** developed analgesic tolerance more slowly than morphine, showed limited physical dependence, respiratory depression, constipation, and displayed no reward or aversion in CPP/CPA assays, suggesting that analogs might represent a promising new generation of novel pain relievers.

INTRODUCTION

Opioids, including morphine, are clinically used for the treatment of moderate to severe chronic pain. However, despite their proven efficacy, mu opioid receptor agonists have problematic side effects such as tolerance, physical dependence, and substance abuse.¹ Agonists selective for other opioid receptors produce analgesia, but with their own liabilities.²⁻⁴ The ultimate goal of opioid-related drug development has been to design and synthesize potent antinociceptive agents that are devoid of adverse side effects. Many approaches have been taken over the years, starting with the development of partial agonists or mixed agonist/antagonists.⁵⁻⁹ A more recent approach takes advantage of biased agonism, in which distinct downstream signaling pathways are activated by different agonists working through the same receptor.^{10, 11} It has been proposed that ligands biased against recruiting β -arrestin-2, or showing preference for activating specific G-protein mediated signal transduction pathways, will demonstrate diminished side effects.^{12, 13} Oliceridine (TRV130)^{14, 15} is an example of a mu-opioid receptor biased agonist which has recently entered phase-III clinical trials, showing separation between antinociception and some opioid-related side effects. 6'-Guanidinonaltrindole (6'-GNTI),¹⁶ 22-thiocyanatosalvinorin A (RB-64),¹⁷ and two new classes of kappa opioid ligands from the Aube group have also recently been reported in the opioid literature as biased kappa opioid receptor agonists.^{18, 19}

Natural products have provided many lead compounds leading to the design of new pharmaceuticals. Natural products and their derivatives account for approximately 50% of approved drugs.²⁰ Morphine, the most commonly employed opioid, and thebaine, the structure on which the vast majority of semi-synthetic opiates is based, are natural alkaloids found in the poppy plant, *Papaver somniferum*. While opioid chemistry has traditionally been dominated by

thebaine-derived alkaloids isolated from poppy, there are a growing number of opioid natural products derived from structures other than the traditional morphinan scaffold and thus structurally not closely related to morphine. These include analogs of salvinorin A²¹⁻²³ such as herkinorin²⁴ and thiocyanatosalvinorin A¹⁷, which have been developed as mu- and kappa-opioid receptor biased agonists, respectively, while some peptide analogs such as cyclo[Phe-d-Pro-Phe-Trp] (CJ-15,208)²⁵ are being developed as analgesics and medications against cocaine abuse (Figure 1). Mitragynine (indole core) (**1**) and its congeners isolated from the Southeast Asian plant *Mitragyna speciosa*, commonly known as *kratom*, are monoterpene indole alkaloids structurally not closely related to morphine.²⁶ In addition to its traditional use, *kratom* has become a quickly emerging substance of abuse. It is currently legal in many parts of the world, and kratom leaves are available for purchase over the internet. Case studies of fatalities resulting from overdose have been published, although the risk posed by mitragynine remains uncertain, given that simultaneous use or contamination with other substances (including opioids) that may have been involved in the reported deaths.²⁷⁻³⁰ Both mitragynine (**1**) and its naturally occurring oxidation product, 7-OH mitragynine (indolenine core) (**2**), are opioid antinociceptive agents that have been examined both *in vitro* and *in vivo*.³¹⁻³⁹ Mitragynine pseudoindoxyl (**3**), a rearrangement product of **2** with a spiro-pseudoindoxyl core, was first isolated in 1974 by Zarembo et al. as a microbial metabolite of **1** by the fungus *Helminthosporium sp.*⁴⁰ Yamamoto et al. reported that it acted non-selectively on mu- and delta-opioid receptors while its kappa-opioid receptor affinity was negligible.⁴¹ In later publications by Takayama et al., the *in vivo* supraspinal analgesic properties of **3** were briefly discussed.³⁹

In this work, we report the *in vitro* and *in vivo* pharmacology and structure-activity relationships (SAR) of mitragynine pseudoindoxyl (**3**). We demonstrate for the first time that **3**

and its C-9 substituted derivatives, corynantheidine pseudoindoxyls, are systemically active mixed mu-opioid receptor agonist/delta-opioid receptor antagonist compounds *in vitro*, and produce potent antinociception *in vivo*. Characterization of **3** demonstrated opioid-mediated analgesia devoid of any place-conditioning effects, and a side effect profile far superior to clinically used mu opioid based antinociceptive agents.

RESULTS

Chemistry:

1 was extracted from dry kratom powder using a modified protocol reported by Ponglux et al.⁴² Compounds **2** and **3** were synthesized from **1** as shown in Scheme 1.^{39, 43} To better understand the pharmacology of this template, SAR studies were carried out by modifying the C-9 and N-1 (indole nitrogen) positions. Six analogs with various substituents in the C-9 positions and 2 analogs at N-1 were synthesized. C-9-substituted corynantheidine pseudoindoxyl derivatives were synthesized starting from **2**. To gain access to the C-9 position on the pseudoindoxyl scaffold, **2** was converted to 9-hydroxycorynantheidine pseudoindoxyl (**4**) using AlCl₃ and ethanethiol in DCM. This intermediate was converted to its triflate (**5**) using triflic anhydride and pyridine, which was subsequently used as the precursor for further reactions.

The pseudoindoxyl of corynantheidine (**6**) was synthesized using palladium-catalyzed removal of the triflate ester by formic acid. The synthesis of the nitrile **7** was accomplished in a palladium-catalyzed reaction of **5** with Zn(CN)₂. Compounds **8** and **9** were obtained via Suzuki coupling reactions of **5** and the appropriate boronic acids. **3** was alkylated in the N-1 position with benzyl bromide and iodomethane to synthesize **11** and **12**, respectively.

In vitro pharmacology:

Initial investigations used *in vitro* radioligand binding assays with cell lines stably expressing murine opioid receptors (Figure 2, Table 1). Mitragynine (**1**) showed poor affinity at all opioid receptors, whereas 7-hydroxymitragynine (**2**) showed moderate affinity at the mu opioid receptor clone MOR-1, 5-fold higher than **1**, and was considerably more potent at the expressed delta opioid receptor clone DOR-1 than **1**.⁴⁴ Mitragynine pseudoindoxyl (**3**) displayed the highest overall binding affinity for MOR-1 and DOR-1 (K_i 0.8 nM and 3 nM, respectively) while also showing a moderate affinity for KOR-1. These data suggest that conversion of the indole to indolenine ring to the spiro-pseudoindoxyl core dramatically increases affinity for opioid receptors. The binding affinities of **3** at MOR-1 and DOR-1 were comparable to the prototypic mu ligands morphine and DAMGO, and delta ligands DPDPE and NTI. Alkyl substitution at the N-1 position of the template (compounds **11**, **12**) eliminated opioid affinity, suggesting that the unsubstituted indole NH is important for receptor binding. Substitutions at C-9, however, yielded potent derivatives. All six C-9-modified compounds (**4**, **6-9**, **10**) maintained the high affinity at both MOR-1 and DOR-1 sites as observed with **3** previously. **2** and **3** were also screened across a panel of other non-opioid drug targets using the PDSP screening facility at NIMH.⁴⁵ **2** exhibited no affinity appreciable affinity at these receptors ($K_i > 10 \mu\text{M}$). **3** had poor affinity at α_{2A} adrenergic receptor, α_{2C} adrenergic receptor and 5HT₇ (Table S1).

In [³⁵S]GTP γ S functional assays using opioid transfected cell lines, **1** was a partial agonist with moderate potency at MOR-1, and a weak antagonist at both DOR-1 and KOR-1 (Table 2). **2** was a partial agonist at MOR-1, 4-fold more potent than **1**, and a weak KOR-1 and DOR-1 antagonist.⁴⁴ Analog **3** was a potent full agonist at MOR-1 and an antagonist at both DOR-1 and KOR-1. Most C-9-modified derivatives (**4**, **6**, **7** and **9**) were MOR-1 agonists and

DOR-1 antagonists in functional assays with the exception of compound **8**, which was a dual MOR-1/DOR-1 agonist (Table 2).

Both DAMGO and endomorphin-2 effectively recruited β -arrestin-2 in CHO cells expressing MOR-1, as measured with the DiscoverX PathHunter assay. In contrast, compounds **1-4** and **6-9** failed to recruit β -arrestin-2 at concentrations as high as 10 μ M (Figure 3A). Both **2** and **3** reduced DAMGO-induced β -arrestin-2 stimulation in these cells in a concentration-dependent manner (Figure 3B)⁴⁴ consistent with their respective binding affinities at the receptor. Thus, **3** and its analogs potently stimulated [³⁵S]GTP γ S binding without stimulating β -arrestin-2 recruitment. However, their antagonism of DAMGO stimulation of β -arrestin-2 recruitment revealed that they could still bind to both G-protein and arrestin functional receptor configurations.

Antinociception:

The antinociceptive effect of mitragynine and its derivatives was evaluated *in vivo* in mice using the radiant heat tail flick assay (Table 1). After subcutaneous administration, **1** produced antinociception with an ED₅₀ (and 95%CI) value of 166 mg/kg (101, 283), 66-fold less active than morphine. On the other hand, **2** was about 5-fold more potent than morphine and 350-fold more potent than **1** (Figure S1), similar to literature values.³²⁻³⁴ Compound **3** was 1.5-fold more potent than morphine after intracerebroventricular administration (icv, Figure 4A and S2), and 3-fold more potent following subcutaneous (sc, Figure 4B and S4A and S4B) administration. Compound **3** has a shorter duration of antinociceptive effect than morphine with a peak effect at 15 min (Figure S3). Compound **3** proved equally active in CD1, C57BL/6 and 129Sv6 strains of mice subcutaneously (Figure S5). Furthermore, **3** also was active orally, with an ED₅₀ (95% CI) value of 7.5 (4.3-13) mg/kg (Figure 4C). The C-9 derivatives (**4** and **6-10**) also produced

antinociception following systemic administration, with ED₅₀ values comparable to **3** (Table 1). Compound **3** was also active in the hot plate assay of antinociception, with an ED₅₀ (95% CI) value of 0.99 (0.75–1.3) mg/kg (Figure 4D and S4C), comparable to morphine (ED₅₀ = 1.7 (1.3, 2.4) mg/kg) (Figure S4D).

Opioid receptor antagonism:

Based on the high affinity and favorable mixed mu agonist/delta antagonist profile *in vitro*, we examined the activity of compound **3** in greater detail. Naloxone and the mu-selective antagonist β -FNA effectively reversed **3**-induced antinociception, whereas the delta-selective antagonist NTI and the kappa antagonist norBNI did not. Yohimbine, an α_2 antagonist, had no effect on the antinociception of **3** (Figure 5A).

To examine the selectivity of antinociception further, we used an antisense oligodeoxynucleotide mapping paradigm. The activity of the oligodeoxynucleotide antisense probes for MOR-1, KOR-1 and DOR-1 has been established previously.^{46–48} Targeting exon 1 of MOR-1, the antisense oligodeoxynucleotide lowered the analgesic actions of morphine, reproducing earlier studies (Figure S6).⁴⁶ Similarly, the responses of **3** were lowered (Figure 5B). The specificity of the response was established by the inactivity of the control mismatch. Similarly, downregulation of exon 3 of DOR-1⁴⁸ and exon 2 of KOR-1⁴⁹ with antisense oligodeoxynucleotides attenuated antinociception produced by the prototypic delta agonist DPDPE and the prototypic kappa agonist U50,488H, in accordance with previous studies (Figure S6). However, these oligodeoxynucleotides targeting kappa and delta receptors did not alter the antinociception of **3** (Figure 5B).

The mu opioid receptor *Oprm1* creates an array of splice variants through alternative splicing with patterns conserved from rodents to humans.^{50, 51} The major sets of variants are full-

length 7 transmembrane domain (7TM) variants associated with exon 1 (E1). A second set of variants of truncated six transmembrane domain splice variants (6TM) is generated by an alternative promoter associated with exon 11 (E11) of the *Oprm1* gene.^{52, 53} MOR-1 KO mice were used to establish the contributions of 7TM E1-MOR-1 and 6TM E11-MOR-1 variants to **3** antinociception.

Two different types of MOR-1 KO mice were utilized: exon 11 (E11) MOR-1 KO mice, which lack the 6TM E11splice variants of MOR-1 but retain expression of 7TM E1 splice variants of MOR-1, and total mu opioid receptor knockout in which both exons 1 and 11 were disrupted (E1/E11) to eliminate all 7TM, and 6TM mu opioid receptor variants of the *Oprm1* gene. Morphine antinociception has previously been demonstrated to be independent of the E11-associated 6TM splice variants of MOR-1, maintaining full analgesic activity in the E11 KO⁵⁴, but its antinociception was completely eliminated in E1/E11 MOR-1 KO mice,⁵⁵ suggesting 7TM E1-MOR-1 variant as the primary mechanism of analgesic action. Compound **3** showed similar antinociceptive responses as morphine. Compound **3** antinociception was similar in wild-type ($ED_{50} = 0.83$ mg/kg (0.37-1.9)) and exon 11 KO C57/BL6 mice ($ED_{50} = 1.4$ mg/kg (0.34-5.8)) in a tail flick assay (Figure 5D). However, antinociception of compound **3** was eliminated in E1/E11 MOR-1 KO mice ($ED_{50} > 30$ mg/kg), indicating a mu opioid receptor mechanism.⁵⁵ Taken together with the antisense results, these *in vivo* findings indicate that **3** analgesia is mediated by 7TM E1-MOR-1 receptors.

Side effect profile:

We next evaluated **3** in mouse models of antinociceptive tolerance, dependence, respiratory depression, and inhibition of GI transit (Figure 6). Mice developed antinociceptive tolerance to morphine after twice daily administration for 5 days (5 mg/kg/injection, sc). In

contrast, antinociceptive tolerance to **3** developed far more slowly, with twice daily administration of **3** (at an equianalgesic dose as morphine) requiring 29 days instead of 5 days (Figure 6A). After 5 days, the morphine ED₅₀ value was shifted 6-fold to 12.1 (7.6-19.4) mg/kg from 2.0 (1.2-3.3) mg/kg. In contrast, after 5 days, the ED₅₀ value for **3** was shifted less than 2-fold to 1.1 (0.66-2.0) mg/kg. After 29 days, the ED₅₀ value for **3** was shifted 6-fold (4.5 mg/kg (2.7-7.7) mg/kg) (Figure S7). An independent sample t-test was used to test for the group difference between **3** and morphine on day 5. The mean difference of 53.97 %MPE between the two groups was found to be statistically significant (p-value<0.0001) at the 5% level of significance. This difference is likely to range between 42.27 and 65.67 %MPE as measured by a 95% confidence interval. To highlight **3**'s ability to sustain antinociception over repeated dosing, the composite areas under the curve (AUCs) were calculated using the trapezoidal rule on the mean response across 1 to 29 days for the **3** group, and 1 to 5 days for the morphine group, respectively. The AUC for the morphine group was 205.9, while for the **3** group it was 6-fold larger, 1239.

Time action studies revealed that **3** has a shorter duration of action than morphine (Figure S3). Since the shorter duration of action of **3** led to a decreased drug exposure, we also examined tolerance in a different dosing paradigm in which **3** was given four times per day to provide similar drug exposures for morphine and **3**. In this paradigm, **3** failed to show a significant decreased effect over 5 days, whether examined at a fixed dose over the 5 days (Figure S8A) or dose-response curves at Day 5 (Figure S8B). After 5 days, the ED₅₀ value for **3** was shifted from 0.69 (0.46, 1.0) to 1.6 (0.97-2.6) mg/kg (Figure S8C), a shift similar to that observed when **3** was given twice a day (ED₅₀ = 1.1 (0.66-2.0)) on day 5 (Figure S7).

Physical dependence was assessed in mice treated repeatedly with **3** at twice its ED₅₀ dose (1.5 mg/kg) twice daily for 5, 22 or 29 days by administration of the opioid antagonist naloxone. Mice showed only minimal signs of withdrawal following administration of naloxone with 12±4.1, 13±4.4 and 14±6.0 jumps on average, respectively (Figure 6B). These values were not significantly different from saline but differed significantly from mice treated with morphine for 5 days, which showed 77±7.2 jumps (Figure 6B). In summary, compound **3** demonstrated limited antinociceptive tolerance and physical dependence in comparison with morphine following chronic administration.

Differences in effect between **3** and morphine extended to other opioid effects. Mu selective agonists inhibit gastrointestinal transit, a major component of constipation. At a dose twice its analgesic ED₅₀ value (5 mg/kg, sc), morphine almost totally eliminated transit (Figure 6C). An equianalgesic dose of **3** (1.5 mg/kg, sc) also lowered gastrointestinal transit, but not nearly as much as morphine. The effect on GI transit plateaued, with a greater dose (4 mg/kg, sc) also showing no further inhibition (Figure 6C).

Morphine dose-dependently reduced the respiratory rate in mice, with a decrease of approximately 30% by a dose twice its analgesic ED₅₀ value (5 mg/kg, sc) by approximately 50% after a higher dose corresponding to 4-fold its analgesic ED₅₀ value. In contrast, **3** showed no respiratory depression at ~twice its antinociceptive ED₅₀ dose in C57BL/6 mice (1.2 mg/kg, sc). Although a higher dose (3 mg/kg, sc), transiently lowered the respiratory rate by approximately 15%, this was still significantly less than morphine (Figure 6D).

Compound **3** also failed to show either rewarding or aversive behavior in a conditioned place preference paradigm. In this study, morphine produced significant conditioned place preference (CPP; $F_{(4,180)}=5.62$, $p=0.003$; two-way ANOVA with Tukey HSD *post-hoc* test) and

U50,488H produced conditioned place aversion (CPA), but **3** demonstrated neither preference or aversive behavior at doses 2-fold or 5-fold its analgesic ED₅₀ value (n.s.; Tukey HSD *post hoc* test; Figure 6E). Overall, these results demonstrate that **3** produces potent opioid receptor-mediated antinociception both centrally and systemically, yet shows a separation of antinociception from some classic opioid side effects such as antinociceptive tolerance, dependence, and conditioned place preference. Furthermore, **3** shows a lower propensity to cause respiratory depression and constipation compared with the canonical opioid, morphine.

Modeling:

In silico docking studies were carried out to unravel potential differences in receptor interactions of **1-3**. The results of *in silico* prediction of inhibitory constants are listed in Table 3 for all three mitragynine compounds and the opioid receptors. In general, ligands of the lowest energy complexes were located in the binding pocket observed in the crystal structures. Inhibitory constants (K_i) calculated for the lowest energy complexes follow the trend observed in the experiments, but the range of values is much more narrow compared to experimental data, suggesting much lower selectivity of both the ligands and the receptors. The lowest energy complexes which are considered to reflect specific binding between **1-3** and the receptors are shown in Figures 7 and S14. These *in silico* K_i values reproduce experimental data with much higher accuracy compared to the ones calculated for the lowest energy complexes in the first pass, where all non-specific hits were included (for explanation, see the Methods). Receptor side chains in contact with the bound **1-3** are also depicted in Figure 7 and listed in Table 3.

DISCUSSION

Although initially described in the scientific literature as early as 1974, very little was known about the pharmacology of **3** prior to this study. Its chemical structure suggested it as an

excellent starting point for semi-synthetic diversification. Thus, we subjected **3** to detailed pharmacological analysis. In opioid receptor-transfected CHO cell lines, **3** had a high affinity for MOR-1 and DOR-1 sites with a moderate affinity at KOR-1. After subcutaneous administration, **3** was a potent antinociceptive agent in two thermal pain models: radiant heat tail flick and hot plate. It was also active in the mouse tail flick assay when administered supraspinally and orally. We established the agonist selectivity of **3** through pharmacological and genetic approaches, demonstrating that its antinociception was mediated by mu opioid receptors and not kappa, delta, and/or α_2 adrenergic receptors. Mu antagonists attenuate the antinociception but it was insensitive to kappa and delta antagonists. Clonidine, an α_2 agonist, is a potent analgesic used for the treatment of various pain conditions.⁵⁶ Yohimbine, an α_2 antagonist did not affect the antinociception of **3**, therefore, its antinociception is likely not related to adrenergic pathways. The non-reversal of analgesia also rules out a role of α_{2A} and α_{2C} adrenergic receptors in mediating analgesia of **3** although the drug had some affinity for these receptors in our initial screening. Antisense downregulation of E1-MOR-1 variants attenuated antinociception while downregulation of DOR-1 and KOR-1 failed to modify **3** antinociception. In mice lacking E11-MOR-1 variants (E11 MOR-1 KO mice) compound **3**, like morphine, still exhibited antinociception comparable to wild-type mice while in mice lacking all mu opioid receptor splice variants (E1/E11 MOR-1 KO mice), **3** antinociception was completely eliminated. The results from antisense and KO mice experiments implicate traditional 7TM mu opioid receptor aka E1-MOR-1 variants in **3** antinociception.

Of interest, **3** signaling failed to recruit β -arrestin-2 alone and antagonized both DAMGO-induced β -arrestin-2 recruitment and stimulation of [³⁵S]GTP γ S binding. To the best of our knowledge, **3** is the first example of a mixed activity mu opioid agonist/delta antagonist

ligand which does not recruit β -arrestin-2. Prior evidence in the literature suggests that failure to recruit β -arrestin-2 and delta antagonism may both be successful in separating antinociception from unwanted side effects. Consistent with this, **3** displayed a robust antinociceptive effect in mice and without conditioned place preference or aversion. Two different dosing paradigms (two and four times per day) revealed the slow development of tolerance and a marked decrease in jumping following challenge with naloxone, which is characteristic of physical dependence. Furthermore, **3** showed no respiratory depression at twice its analgesic ED₅₀ dose, and far less constipation than morphine.

A number of mixed mu agonist-delta antagonist ligands have been reported in the literature. DIPP-NH₂[Ψ],⁵⁷ 5''-(4-chlorophenyl)-6,7-didehydro-4,5 α -epoxy-3-hydroxy-17-methylpyrido[2',3':6,7]morphinan (SoRi20411),⁵ and 14-alkoxy pyridomorphinans⁵⁸ are potent delta antagonists and mu agonists which produce analgesia with reduced tolerance when given supraspinally. It must be noted that a majority of these studies monitored the development of tolerance through the accepted practice of administering the ED₈₀ antinociceptive dose of the test drug twice daily for 5-7 days. The current study tested this approach even more rigorously, yet still found significant reductions in antinociceptive tolerance following administration of **3** for 29 days. DIPP-NH₂[Ψ] also shows no physical dependence in treated mice. 4a,9-Dihydroxy-7a-(hydroxymethyl)-3-methyl-2,3,4,4a,5,6-hexahydro-1H-4,12-methanobenzofuro[3,2-e]isoquinolin-7(7aH)-one (UMB425),⁵⁹ and the cyclic peptide analog of KSK103 (C-terminal Ser(β -Glc)NH₂)⁶ and ([Dmt¹]DALDA \rightarrow CH₂CH₂NH \leftarrow TICP[Ψ]), which connects Dmt¹-DALDA (mu agonist) with the delta antagonist TICP[Ψ] through a spacer, also show reduced acute antinociceptive tolerance compared to morphine when given systemically.⁶⁰ Administration of delta antagonists or induced downregulation of delta opioid receptors has been reported to

1
2
3 prevent morphine tolerance without sacrificing analgesic potency.^{61, 62} Genetic disruption of the
4
5 opioid system led to similar observations. In DOR-1 antisense knockdown mouse models,
6
7 tolerance and acute dependence to morphine were eliminated⁶³ while DOR-1 knockout animals
8
9 did not develop antinociceptive tolerance to morphine.⁶⁴ The role of delta-opioid receptors in
10
11 blocking mu opioid-mediated CPP is still unclear⁶⁵. CPP is retained unchanged or abolished
12
13 depending on assay conditions in DOR-1 KO mice, while morphine reportedly was more
14
15 rewarding in a study with β -arrestin-2 KO mice.⁶⁶ Additional study of this phenomenon should
16
17 be possible as additional novel ligands with suitable MOR agonist/DOR antagonist activity
18
19 profiles such as **3** become available.
20
21
22
23

24
25 Although GPCRs are mostly studied in relation to their ability to activate G-proteins,
26
27 these transmembrane receptors are capable of recruiting β -arrestins to initiate separate cellular
28
29 signal transduction pathways.⁶⁷ β -Arrestin-2 activation has been implicated in the mechanism of
30
31 receptor desensitization and the occurrence of deleterious side effects.⁶⁸ Similarly to G-protein
32
33 signaling, β -arrestin-2 activation is ligand- and receptor-dependent. Different ligands are able to
34
35 stabilize GPCRs in a variety of conformations, resulting in the differential activation (bias) of G-
36
37 protein and β -arrestin-2-mediated signaling pathways. Therefore, ligands interacting with the
38
39 receptor do not simply have a linear effect on efficacy but also affect the functional quality of the
40
41 downstream pathways.⁶⁹ Biased signaling of opioid receptors has been studied in detail by Bohn
42
43 and co-workers. Compared with wild-type mice, morphine displayed enhanced antinociception
44
45 and significantly attenuated respiratory depression, and inhibition of GI transit in β -arrestin-2
46
47 KO mice.⁷⁰⁻⁷² These results suggest that a fully G-protein biased opioid ligand that does not
48
49 activate β -arrestin-2 signaling may be able to separate antinociception from some opioid adverse
50
51 effects. An important example of G-protein biased opioids is oliceridine, a synthetic mu agonist
52
53
54
55
56
57
58
59
60

currently in clinical trials as an alternative to morphine and fentanyl for the treatment of chronic pain. Oliceridine is a potent analgesic that causes less respiratory depression and constipation than morphine at equianalgesic doses in humans.¹⁵

Taken together, these previous observations may explain why **3** is able to separate antinociception from typical opioid side effects resulting in low risk of developing tolerance while showing limited physical dependence, respiratory depression and inhibition of GI transit. However, the relative contributions of its mu agonist/delta antagonist activity and its inability to recruit β -arrestin-2 to this advantageous pharmacological profile are not clear.

In order to investigate the SAR of the mitragynine pseudoindoxyl scaffold, semi-synthetic analogs were made starting from **1** (C-9 analogs) and **3** (N-1 analogs). Receptor affinities were not significantly affected by modifications at the C-9 position, although 9-*O*-acetylation slightly lowered mu and delta affinities. Compounds **4** & **6-9** retained high, subnanomolar affinity at cells expressing MOR-1 with little change in affinity in DOR-1 cells. None of the derivatives stimulated β -arrestin-2 activation. According to previous literature reports, C-9 modifications of mitragynine altered the efficacy at mu receptors. Replacing the C-9 methoxy group with H yields corynantheidine, a mu antagonist, whereas C-9 *O*-demethylation yields 9-OH corynantheidine, a partial agonist in in vitro assays.^{39, 73} According to our studies, the SAR of the C-9-modified pseudoindoxyl scaffold is quite distinct. Various substituents can be tolerated at this position, maintaining full mu agonism. Neither C-9 *O*-demethylation (**4**), nor the removal of the methoxy group (**6**) affected the efficacies at mu as **4** and **6** were mu full agonists. However, the activity at delta receptors is differentially affected by varying substituents. Compounds **4**, **6**, **7**, **9**, **10** retained delta antagonism, while the 9-phenyl analog, **8**, was a delta agonist. **8** was a dual mu-delta agonist with similar intrinsic activity and potency at

both receptors. Substitution and the introduction of bulky groups at N-1 (the indoxyl nitrogen) were not tolerated. Both the *N*-benzyl (**11**) and *N*-methyl (**12**) derivatives showed diminished affinities at all three opioid receptors compared to **3**. *In vivo*, the C-9 analogs were active after systemic administration. The 9-OH derivative (**4**) was more potent than **3**. Removal of the methoxy group (**6**) increased analgesic potency. The analgesic potencies of the corresponding, C-9 mitragynine analogs (corynantheidine and 9-OH corynantheidine) is not reported in the literature. These results, in addition to the *in vitro* data, suggest that the SAR of compounds of the pseudoindoxyl scaffold differs from that of the natural *Mitragyna* alkaloids. Substitution of the C-9 methoxy group with –CN, phenyl and furan-3-yl groups afforded products (**7-9**, respectively) roughly equipotent to **3**. Acetylation of **4** had a slightly negative impact on the analgesic potency (**10**).

In comparison to the mu opioid receptor-bound morphinans⁷⁴⁻⁷⁶, *in silico* modeling suggests the mitragynine derivatives (**1-3**) are likely to have a different binding pose. The salt bridge between Asp¹⁴⁷ and the tertiary amine of the ligand and the participation of the phenolic OH in a water molecule assisted hydrogen bonded polar network of Tyr¹⁴⁸, Lis²³³ and His²⁹⁷ were described as the main, conserved interactions between morphinan ligands and the binding pocket.⁷⁷ The presence of a salt bridge between Asp¹⁴⁷ and the tertiary amine of **1** and its derivatives was a filtering criterion of docking results, therefore, it is present in the docked complexes of all three mitragynine compounds. The β-methoxy acrylate moiety in the mitragynine compounds docked in the mu receptor pocket occupied the same space as the phenol moiety of the morphinan scaffold in the crystallographic structures.^{75, 77}

The main difference between the mu receptor-bound **1**, **2** and **3** is that the oxidation products (**2** and **3**) seem to participate in the polar network formed between Tyr¹⁴⁸, Lys²³³ and

His²⁹⁷ by contributing their methoxy (C-9) and methyl ester groups and replacing H-bond assisting water molecules observed in the crystal structure of the agonist-bound mu receptor (Table 3).⁷⁶ On the other hand, the beta-methoxy methyl acrylate moiety of **1** forms polar interactions with Gln¹²⁴ and Tyr¹²⁸ and do not take part in the aforementioned polar network (Figure S9 and Table 3). In addition to the salt bridge between Asp¹⁴⁷ in the mu receptor crystal and the tertiary amine, it seems that there is a hydrogen bond present, formed with the 7-OH group of **2** (Figure S10). There is no significant difference in the relative orientation between the receptor-bound **2** and **3** (Figure S11). Differences in the calculated binding free energies and K_i values are most likely to emerge from the type and number of receptor contacts formed by these two compounds in those particularly similar docked orientations. Such differences in interactions are possibly due to the different heterocyclic scaffolds of **2** and **3**. Compared to its derivatives, **1** adopts a different orientation when bound to the delta receptor and forms fewer contacts with the residues constituting the binding pocket, resulting in a loss of affinity for this receptor. Admittedly, the relative orientation and the number and type of contacts formed between **2** and **3** with the delta receptor and the calculated in silico K_i values are highly similar, rendering it difficult to give an accurate explanation for the one order of magnitude difference in the experimentally determined binding affinities. A key difference between the two bound geometries is that the beta-methoxy methyl acrylate moiety of **2** is positioned in a hydrophobic environment of Ile²⁷⁷, Leu³⁰⁰, and Ile³⁰⁴, (Figure S12 and Table 3) while in the case of **3** this group is projected into a more hydrophilic part of the binding pocket lined with Asn¹³¹, Trp²⁷⁴, and His²⁷⁸ (Figure S13). Furthermore, the 7-OH group of **2** was found to interact with Asp¹²⁸ (Figure S12) similarly to that observed when docked to the mu (Figure S10). In addition, the 9-methoxy group of **2** formed contact with the phenolic OH of Tyr¹²⁹. In the delta-bound **3**

1
2
3 complex, both the 9-methoxy group and the carbonyl group of the pseudoindoxyl moiety were
4
5 calculated to be in favorable position for H-bonding with Tyr¹²⁹ (Figure 7 and S13).
6
7

8 9 CONCLUSIONS

10
11 In summary, we report for the first time the detailed *in vitro* and *in vivo* studies on
12
13 mitragynine pseudoindoxyl. Mitragynine pseudoindoxyl is a mu agonist/delta antagonist opioid
14
15 with a signaling bias for G-protein-mediated signaling pathways *in vitro*, and which produced
16
17 potent antinociception *in vivo*. Perhaps owing to its mixed mu agonism/delta antagonism activity,
18
19 mitragynine pseudoindoxyl may avoid some of the major problems of opioid therapy as we
20
21 observed no reward or aversion, and diminished antinociceptive tolerance, physical dependence,
22
23 respiratory depression, and GI transit inhibition in mouse models. Upon chemical modification
24
25 of this scaffold, key SAR features distinct from the mitragynine template were revealed. Among
26
27 analogs modified at the C-9 position, compounds with differential efficacies within *in vitro*
28
29 functional assays and improved *in vivo* potencies were identified. Docking studies to opioid
30
31 receptors revealed the characteristic binding modes of mitragynine-type derivatives. It is hoped
32
33 that these studies will contribute to improved understanding of the mechanism of action of
34
35 compounds related to mitragynine, while holding the promise to provide novel antinociceptive
36
37 drug candidates based on the mitragynine/corynantheidine pseudoindoxyl template that separate
38
39 antinociception from the potential for abuse and other side effects due to their unique
40
41 pharmacological properties. Observations reported in this paper and past studies on opioid
42
43 ligands suggest a dual mechanism (mu agonism/delta antagonism coupled with β -arrestin2- non-
44
45 recruitment) which may account for the separation of side effects from antinociception seen with
46
47 this template. The potential contributions from each of these two mechanisms will be explored in
48
49 future studies on this template.
50
51
52
53
54
55
56
57
58
59
60

EXPERIMENTAL SECTION

Drugs and Chemicals: Opiates were provided by the Research Technology Branch of the National Institute on Drug Abuse (Rockville, MD). IBNtxA and [125 I]BNtxA were synthesized in our laboratory as previously described.⁷⁸⁻⁸⁰ Na 125 I and [35 S]GTP γ S were purchased from Perkin-Elmer (Waltham, MA). Selective opioid antagonists were purchased from Tocris Bioscience. Miscellaneous chemicals and buffers were purchased from Sigma-Aldrich. Kratom “Red Indonesian Micro Powder” was purchased from Moon Kratom (Austin, TX).

Mice: Male CD1 mice (20-32 g) were obtained from Charles River Laboratories, C57BL/6J mice (20–32 g each) were obtained from Jackson Laboratories (Bar Harbor, ME). Exon-11 KO⁷⁹ and Exon-1/Exon-11 KO mice⁵⁵ were bred in our laboratory. All mice used throughout the manuscript were opioid naïve. All mice were maintained on a 12-hour light/dark cycle with Purina rodent chow and water available ad libitum, and housed in groups of five until testing. All animal studies were preapproved by the Institutional Animal Care and Use Committees of the Memorial Sloan Kettering Cancer Center or University of Florida, in accordance with the 2002 National Institutes of Health Guide for the Care and Use of Laboratory Animals.

Radioligand Competition Binding Assays: [125 I]IBNtxA binding was carried out in membranes prepared from Chinese Hamster Ovary (CHO) cells stably expressing murine clones MOR-1, DOR-1, and KOR-1, as previously described.^{7-9, 80} Binding was performed at 25°C for 90 min. Binding in MOR-1/CHO was carried out in 50 mM potassium phosphate buffer with 5 mM MgSO₄ and 20 μ g/ml protein while binding in KOR-1/CHO and DOR-1/CHO was carried out in 50 mM potassium phosphate pH=7.0 buffer and 40 μ g/ml protein. After the incubation,

the reaction was filtered through glass-fiber filters (Whatman Schleicher & Schuell, Keene, NH) and washed three times with 3 mL of ice-cold 50 mM Tris-HCl, pH 7.4, on a semiautomatic cell harvester. Nonspecific binding was defined by the addition of levallorphan (8 μ M) to matching samples and was subtracted from total binding to yield specific binding. K_i values were calculated by nonlinear regression analysis (GraphPad Prism, San Diego, CA). Protein concentrations were determined using the Lowry method with BSA as the standard.⁸¹

[³⁵S]GTP γ S Functional Assay: [³⁵S]GTP γ S binding was performed on membranes prepared from transfected cells stably expressing opioid receptors in the presence and absence of the indicated compound for 60 min at 30 °C in the assay buffer (50 mM Tris-HCl, pH 7.4, 3 mM MgCl₂, 0.2 mM EGTA, and 10 mM NaCl) containing 0.05nM [³⁵S]GTP γ S; 2 μ g/ml each leupeptin, pepstatin, aprotinin, and bestatin; and 30 μ M GDP, as previously described.⁸² After the incubation, the reaction was filtered through glass fiber filters (Whatman Schleicher & Schuell, Keene, NH) and washed three times with 3 mL of ice-cold buffer (50 mM Tris-HCl, pH 7.4) on a semi-automatic cell harvester. Filters were transferred into vials with 3 mL of Liquiscint (National Diagnostics, Atlanta, GA), and the radioactivity in vials was determined by scintillation spectroscopy in a Tri-Carb 2900TR counter (PerkinElmer Life and Analytical Sciences). Basal binding was determined in the presence of GDP and the absence of drug. Data was normalized to 1000 nM DAMGO, DPDPE, and U50,488 for MOR-1, DOR-1 and KOR-1 binding, respectively. EC₅₀, IC₅₀, and %E_{max} values were calculated by nonlinear regression analysis (GraphPad Prism, San Diego, CA).

β-arrestin-2 Recruitment Assay: β-arrestin-2 recruitment was determined using the PathHunter enzyme complementation assay (DiscoverX, Fremont, CA) using modified MOR-1 expressed in CHO cells (DiscoverX). Cells were plated at a density of 2500 cells/well in a 384-well plate as described in the manufacturer's protocol. The following day, cells were treated with the indicated compound for 90 minutes at 37 °C followed by incubation with PathHunter detection reagents for 60 minutes. Chemiluminescence was measured with an Infinite M1000 Pro plate reader (Tecan, Männedorf, Switzerland). For the antagonist dose response assay, the cells were incubated with the antagonist for 30 minutes at 37 °C prior to the addition of agonist. Following antagonist treatment, the cells were treated with 10μM DAMGO for 90 minutes at 37°C and chemiluminescence was detected using the PathHunter detection reagents.

Antinociception: Tail flick antinociception was determined using the radiant heat tail flick technique using an Ugo Basile model 37360 instrument as previously described.^{8,9} The intensity was set to achieve a baseline between 2 and 3 seconds. Baseline latencies were determined before experimental treatments for all mice. Tail flick antinociception was assessed quantally as a doubling or greater of the baseline latency, with a maximal 10 second latency to minimize damage to the tail. Data were analyzed as percent maximal effect, %MPE, and was calculated according to the formula: % MPE [(observed latency – baseline latency)/(maximal latency – baseline latency)] x 100. Compounds were injected subcutaneously (s.c.) or intracerebroventricularly (i.c.v.), and antinociception was assessed 15 min later at the peak effect. Intracerebroventricular dosing (i.c.v.) was carried out as previously described.⁸³ Briefly, the mice were anesthetized with isoflurane. A small incision was made, and synthetic opiate

analog (2 μ L/mouse) was injected using a 10 μ L Hamilton syringe fitted to a 27-gauge needle. Injections were made into the right lateral ventricle at the following coordinates: 2 mm caudal to bregma, 2 mm lateral to sagittal suture, and 2 mm in depth. Mice were tested for antinociception 15 minutes post injection. For oral (p.o.) studies, mice were fasted for 18 h with access to water before administering the drug by oral gavage. For the antagonism studies β -FNA (40 mg/kg, s.c.) and norbinaltorphimine (norBNI, 10 mg/kg, s.c.) were administered 24 hours before **3**. Naltrindole (NTI, 0.5 mg/kg, s.c.) was administered 15 min before **3**. Antinociception also was assessed using the hot plate test.⁵⁴ The hot plate (Ugo Basile 35100) consisted of a metal surface (55 °C) with a transparent plexiglass cylinder to contain the mouse. The latency to lick a hind paw or shake/flutter when the mouse was placed on the hot plate was measured, with a maximal latency of 30 seconds to avoid tissue damage. Baseline latencies were taken for each mouse prior to any drug administration. Mice were tested for analgesia with cumulative subcutaneous doses of the drug until the mouse can withstand the maximal latency. Once the mouse reached the maximal latency, the mouse was no longer given higher doses. *In vivo* experiments were evaluated using GraphPad Prism, San Diego, CA as described above.

Antisense Assays: Antisense (AS) and mismatch (MIS) oligodeoxynucleotides were designed based on the published sequences of the mouse mu opioid receptor gene (*Oprm1*), delta opioid receptor gene (*Oprd1*), and kappa opioid receptor gene (*Oprk1*) (Table 3). These probes have been previously described and validated.^{49, 84-86} Antisense oligodeoxynucleotide injection: Groups of mice received the stated antisense by icv administration (5-10 μ g) or mismatch (5-10 μ g) oligodeoxynucleotide i.c.v. under light isoflurane anesthesia on days 1, 3 and 5, as previously described.⁸⁵ Tail flick antinociception was tested on day 6. Control groups received

no injection prior to testing. On test day, mice received **3** (1.5 mg/kg, s.c.), morphine (0.75 µg, i.c.v.), DPDPE (10 µg, i.c.v.), or U50,488H (5 mg/kg, s.c.). All experiments were performed 3 times with similar results observed with each determination.

Table 4. Sequences of Antisense (AS) and mismatch (MIS) oligodeoxynucleotides.

Target	Antisense	Mismatch Control
MOR-1 exon 1	CGCCCCAGCCTCTTCCTCT	CGCCCCGACCTCTTCCCTT
DOR-1 exon 3	AGGGGAAGGTCGGGTAGG	GAGGAGAGGTGCGTGGAG
KOR-1 exon 2	CGCCCCAGCCTCTTCCTCT	CTCCGCGCTCTCACCTCT

Respiratory Depression Assessment: Respiratory rate was assessed in awake, freely moving, adult male C57BL/6 mice with the MouseOx pulse oximeter system (Starr Life Sciences) as described previously.⁷⁹ Each animal was habituated to the device for 30 min and then tested. A 5-s average breath rate was assessed at 5-min intervals. A baseline for each animal was obtained over a 25-min period before drug injection, and testing began at 15 min post-injection and continued for a period of 35 min. Groups of mice (n = 5) were treated s.c. with either morphine (5 or 10 mg/kg) or **3** (1.2 or 3 mg/kg). Groups were compared with repeated-measures ANOVA followed by Tukey’s multiple-comparison test.

GI transit: Gastrointestinal transit was determined as previously described.⁸⁷ Animals received the indicated drug followed by a charcoal meal (2.5% gum tragacanth in 10% activated charcoal in water) by gavage. Animals were sacrificed 30 min later, and the distance traveled by charcoal was measured. Significance was determined by ANOVA followed by Tukey’s multiple-comparison test.

Conditioned Place Preference/Aversion: Mice were conditioned with a counterbalanced place conditioning paradigm using similar timing as detailed previously.⁸ The amount of time subjects spent in each of three compartments was measured over a 30 min testing period. Prior to place conditioning, the animals (n=95) did not demonstrate significant differences in their time spent exploring the left (543 ± 13 s) vs. right (571 ± 12 s) compartments ($p = 0.15$; Student's *t*-test), resulting in a combined preconditioning response of -0.1 ± 19 s. During each of the next two days, mice were administered vehicle (0.9% saline) and consistently confined in a randomly assigned outer compartment for 40 min, half of each group in the right chamber, half in the left chamber. Four h later, mice were administered morphine (10 mg/kg, i.p.), U50,488H (30 mg/kg, i.p.), cocaine (10 mg/kg, i.p.), or **3** (1.3 or 3.2 mg/kg, i.p.) and confined to the opposite compartment for 40 min. Conditioned place preference data is presented as the difference in time spent in drug- and vehicle-associated chambers, and were analyzed via repeated measures two-way ANOVA with the difference in time spent on the treatment- vs. vehicle-associated side as the dependent measure and conditioning status as the between-groups factor. Where appropriate, Tukey's HSD or Sidak's multiple comparison *post-hoc* tests were used to assess group differences. Effects were considered significant when $p < 0.05$. All effects are expressed as mean \pm SEM.

In silico docking. Full sequence target structures of the human mu opioid and delta opioid receptors for docking studies were built and used as described elsewhere⁹, using crystal structures of the homologous murine opioid receptors^{74, 75} as templates (PDB codes: 4DKL and 4EJ4, respectively). The x-ray structure of the human kappa receptor (PDB code: 4DJH)⁸⁸ was used as docking target after missing side chains were added. Dockings were performed with the

Autodock 4.2 software. Side chains in contact with the bound ligands observed in the crystal complexes of the mu, delta and kappa opioid receptors were kept flexible as well as all ligand torsions. Mitragynine and its natural derivatives were docked using the Lamarckian genetic algorithm in an 80 Å x 80 Å x 80 Å grid volume with 0.375 Å spacing. This docking volume is large enough to cover the whole receptor region accessible from the extracellular side. (Figure S14) In this sense, blind docking studies were performed and 1000 dockings were done for all compounds and receptor models. The resultant ligand-receptor complexes were clustered and ranked according to the corresponding binding free energies, which were also used to calculate inhibitory constants according to the following equation: $\Delta G = RT \ln K_i$.

Chemistry

General Methods: All chemicals were purchased from Sigma-Aldrich Chemicals, and were used without further purification. Reactions were carried out in flame-dried reaction flasks under Ar. Reaction mixtures were purified by Silica Flash chromatography on E. Merck 230–400 mesh silica gel 60 using a Teledyne ISCO CombiFlash R_f instrument with UV detection at 280 and 254 nm. RediSep R_f silica gel normal phase columns were used. The yields reported are isolated yields. IR spectra were recorded on a Bruker Optics Tensor 27 FTIR spectrometer with peaks reported in cm⁻¹. NMR spectra were recorded on Bruker Avance III 500, Avance III 600 with DCH CryoProbe instruments. NMR spectra were processed with MestReNova software (ver. 10.0.2.). Chemical shifts are reported in parts per million (ppm) relative to residual solvent peaks rounded to the nearest 0.01 for proton and 0.1 for carbon (CDCl₃ ¹H: 7.26, ¹³C: 77.3; CD₃OD ¹H: 3.31, ¹³C: 49.0; DMSO-*d*₆ ¹³C: 39.5). Peak multiplicity is reported as follows: s – singlet, d – doublet, t – triplet, q – quartet, m – multiplet. Coupling constants (*J*) are expressed in Hz. Mass spectra were obtained at the MSKCC Analytical Core Facility on a Waters Acuity

SQD LC-MS by electrospray (ESI) ionization. High-resolution mass spectra were obtained on a Waters Acuity Premiere XE TOF LC-MS by electrospray ionization. Accurate masses are reported for the molecular ion $[M+H]^+$. Purity ($\geq 95\%$) was confirmed using HPLC: Waters 1525 Binary Pump, Waters 2489 UV/vis Detector, Waters XBridge C18 column ($5\mu\text{m} \times 150 \times 4.6$ mm), mobile phase: solvent A: water with 0.1% TFA; solvent B: acetonitrile with 0.1% TFA. Gradient: 5-95% acetonitrile/water. Flow rate: 1ml/min.

Isolation of mitragynine (1) from *Mitragyna speciosa* (kratom): Kratom “Red Indonesian Micro Powder” was purchased from Moon Kratom (Austin, TX). Mitragynine (1) was extracted from the powdered leaves by a modified method (added new step: petroleum ether extraction of the acidic aqueous phase) from that reported by Ponglux et al.⁴² Kratom powder (450 g) was extracted by refluxing with MeOH (5 x 500 mL) for 40 min. The suspension was filtered after each extraction and the solvent evaporated. The residue was resuspended in 20% acetic acid solution (2 L) and rinsed with petroleum ether (3 x 500 mL). The aqueous layer was then cooled on ice bath and basified (pH \sim 9) with 50% aqueous NaOH solution. The basified suspension was extracted with DCM (4 x 1 L). The combined organic layers were dried over Na_2SO_4 and filtered. The solvent was evaporated and the residue purified using flash column chromatography (gradient: 0-50% EtOAc in hexanes). The major constituent was **1** (yield 5.59 ± 0.59 g (1.24%); smaller quantities of speciogynine and paynantheine were also isolated.

(E)-Methyl-2-((2S,3S,12bS)-3-ethyl-8-methoxy-1,2,3,4,6,7,12,12b-octahydroindolo[2,3-a]quinolizin-2-yl)-3-methoxyacrylate (Mitragynine, 1): IR (NaCl): 3363, 2950, 2796, 1698, 1643, 1570, 1508, 1435, 1310, 1275, 1255, 1148, 1106, 769, 734. ^1H NMR (600 MHz, Chloroform-*d*) δ 7.74 (s, 1H), 7.43 (s, 1H), 6.99 (t, $J = 7.9$ Hz, 1H), 6.90 (d, $J = 8.0$ Hz, 1H), 6.45 (d, $J = 7.7$ Hz, 1H), 3.87 (s, 3H), 3.72 (s, 3H), 3.71 (s, 3H), 3.18 – 3.08 (m, 2H), 3.06 –

2.99 (m, 2H), 3.00 – 2.93 (m, 1H), 2.94 – 2.90 (m, 1H), 2.57 – 2.42 (m, 3H), 1.83 – 1.75 (m, 2H), 1.62 (dt, $J = 11.5, 3.2$ Hz, 1H), 1.24 – 1.16 (m, 1H), 0.87 (t, $J = 7.4$ Hz, 3H). ^{13}C NMR (151 MHz, CDCl_3) δ 169.45, 160.75, 154.69, 137.41, 133.90, 121.98, 117.82, 111.67, 108.03, 104.37, 99.91, 61.74, 61.46, 57.94, 55.52, 53.98, 51.57, 40.87, 40.12, 30.14, 24.14, 19.28, 13.07. HRMS calcd for $\text{C}_{23}\text{H}_{30}\text{N}_2\text{O}_4$ (MH^+), 399.2284; found 399.2285.

(E)-Methyl-2-((2S,3S,7aS)-3-ethyl-7a-hydroxy-8-methoxy-1,2,3,4,6,7,7a,12b-octahydroindolo[2,3-a]quinolizin-2-yl)-3-methoxyacrylate (7-OH mitragynine, 2): Mitragynine (**1**, 2.00 g, 5.02 mmol) was dissolved in acetonitrile (150 mL), then water (50 mL) was added. The resulting suspension was cooled to 0 °C, and the following solution was added slowly over the course of several minutes: PIFA (2.16 g, 1.1 equiv) in 22 mL acetonitrile. The reaction mixture was stirred at 0 °C for 1 hour, then saturated aqueous NaHCO_3 solution was added and the mixture extracted with EtOAc. The organic phase was rinsed with brine (60 mL), dried over anhydrous Na_2SO_4 , and then it was evaporated under reduced pressure. The residue was dissolved in DCM and purified using flash column chromatography (gradient: 0-75% EtOAc in hexanes). The fractions containing the product were evaporated to yield 1075 mg (57%) of **2** as a light brown amorphous powder. IR (NaCl): 3436, 2952, 1702, 1645, 1599, 1487, 1461, 1436, 1270, 1246, 1145, 1078, 795, 738. ^1H NMR (600 MHz, Chloroform- d) δ 7.44 (s, 1H), 7.34 (t, $J = 8.0$ Hz, 1H), 7.24 (d, $J = 7.6$ Hz, 1H), 6.78 (d, $J = 8.3$ Hz, 1H), 3.91 (s, 3H), 3.80 (s, 3H), 3.70 (s, 3H), 3.31 (dd, $J = 11.1, 2.6$ Hz, 1H), 3.03 (ddt, $J = 11.5, 5.5, 2.8$ Hz, 2H), 2.84 – 2.75 (m, 3H), 2.67 (ddd, $J = 12.3, 4.3, 2.6$ Hz, 1H), 2.53 – 2.46 (m, 1H), 1.98 – 1.93 (m, 1H), 1.87 (ddd, $J = 14.6, 12.2, 4.3$ Hz, 1H), 1.70 – 1.54 (m, 3H), 1.26 – 1.23 (m, 1H), 0.81 (t, $J = 7.3$ Hz, 3H). ^{13}C NMR (151 MHz, CDCl_3) δ 181.25, 169.44, 160.94, 156.02, 154.67, 131.50, 126.24, 114.46,

111.38, 109.65, 69.62, 62.00, 60.73, 58.32, 55.92, 51.54, 50.39, 40.67, 39.32, 36.22, 26.38, 19.10, 13.02. HRMS calcd for $C_{23}H_{30}N_2O_5$ (MH⁺), 415.2233; found 415.2248.

(E)-Methyl-2-((1'S,6'S,7'S)-6'-ethyl-4-methoxy-3-oxo-3',5',6',7',8',8a'-hexahydro-2'H-spiro[indoline-2,1'-indolizine]-7'-yl)-3-methoxyacrylate (Mitragynine pseudoindoxyl, 3): 7-OH-mitragynine (**2**, 200 mg, 0.48 mmol) was dissolved in dry toluene (6 mL) and Zn(OTf)₂ (350 mg, 2 equiv) was added. The reaction was stirred in a sealed tube for 2 hrs at 110 °C. To the cooled mixture was added 10 mL sat. aqueous NaHCO₃ solution and water (20 mL). Extracted with EtOAc (30 mL). The organic layer was rinsed with brine (20 mL) and dried over anhydrous Na₂SO₄. After evaporation of the solvent under reduced pressure, the residue was redissolved in DCM and purified using flash column chromatography (gradient: 1-5% MeOH in DCM) to yield: 78 mg (39%) of **3** as a yellow amorphous powder. NMR was identical to that reported in the literature.³⁹ IR (NaCl): 3350, 2947, 2794, 1687, 1615, 1502, 1343, 1269, 1246, 1148, 1079, 757. ¹H NMR (600 MHz, Chloroform-*d*) δ 7.32 (t, *J* = 8.1 Hz, 1H), 7.27 (s, 1H), 6.40 (d, *J* = 8.1 Hz, 1H), 6.13 (d, *J* = 8.1 Hz, 1H), 5.13 (s, 1H), 3.89 (s, 3H), 3.66 (s, 3H), 3.62 (s, 3H), 3.15 – 3.07 (m, 2H), 2.76 (dt, *J* = 11.9, 3.5 Hz, 1H), 2.38 – 2.29 (m, 2H), 2.29 – 2.18 (m, 1H), 2.14 (dt, *J* = 10.2, 6.3 Hz, 1H), 1.93 – 1.84 (m, 1H), 1.63 (dt, *J* = 11.3, 6.8 Hz, 1H), 1.49 (d, *J* = 11.3 Hz, 1H), 1.18 (ddd, *J* = 13.2, 7.8, 2.9 Hz, 1H), 1.11 (dd, *J* = 11.3, 3.6 Hz, 1H), 0.84 (t, *J* = 7.4 Hz, 3H). ¹³C NMR (151 MHz, CDCl₃) δ 199.73, 169.05, 162.27, 160.40, 158.74, 138.85, 111.85, 109.96, 103.95, 99.21, 75.37, 73.38, 61.61, 55.86, 54.96, 53.35, 51.36, 40.28, 38.57, 35.25, 23.95, 19.47, 13.11. HRMS calcd for $C_{23}H_{30}N_2O_5$ (MH⁺), 415.2233; found 415.2216.

(E)-Methyl-2-((1'S,6'S,7'S)-6'-ethyl-4-hydroxy-3-oxo-3',5',6',7',8',8a'-hexahydro-2'H-spiro[indoline-2,1'-indolizine]-7'-yl)-3-methoxyacrylate (9-OH-corynantheidine

pseudoindoxyl, 4): 7-OH mitragynine (**2**, 400 mg, 0.97 mmol) was dissolved in dry DCM (20 mL), then AlCl_3 (1.29 g, 10 equiv) was added. The mixture was cooled to 0°C and ethanethiol (1.39 mL, 20 equiv) was added. The mixture was stirred at rt for 5 hr. Water (30 mL) was slowly added, then it was separated from the organic layer. The organic layer was rinsed with brine (30 mL), then separated and dried over Na_2SO_4 . Evaporated under reduced pressure. The residue was redissolved in DCM and purified using flash column chromatography (gradient: 1-3% MeOH in DCM) to yield 342 mg (89%) of **4** as a bright yellow amorphous powder. IR (NaCl): 3211, 2945, 1697, 1628, 1513, 1451, 1348, 1247, 1148, 1120, 1082, 744. ^1H NMR (600 MHz, CDCl_3) δ 7.29 (m, 2H), 6.30 (d, $J = 8.1$ Hz, 1H), 6.15 (d, $J = 8.0$ Hz, 1H), 5.12 (s, 1H), 3.67 (s, 3H), 3.63 (s, 3H), 3.17 – 3.11 (m, 2H), 2.83 – 2.75 (m, 1H), 2.36 – 2.29 (m, 2H), 2.26 – 2.20 (m, 2H), 2.14 (dd, $J = 11.2, 2.8$ Hz, 1H), 1.98 – 1.90 (m, 1H), 1.69 – 1.57 (m, 2H), 1.51 (d, $J = 11.2$ Hz, 1H), 1.24 – 1.16 (m, 1H), 1.14 – 1.08 (m, 1H), 0.86 (t, $J = 7.4$ Hz, 3H). ^{13}C NMR (151 MHz, CDCl_3) δ 203.60, 169.03, 160.52, 160.05, 157.31, 140.24, 111.63, 108.97, 103.68, 102.27, 75.49, 72.99, 61.69, 54.93, 53.32, 51.41, 40.21, 38.63, 34.79, 23.97, 19.43, 13.07. HRMS calcd for $\text{C}_{22}\text{H}_{28}\text{N}_2\text{O}_5$ (MH^+), 401.2076; found 401.2068.

(E)-Methyl-2-((1'S,6'S,7'S)-6'-ethyl-3-oxo-4-(trifluoromethylsulfonyloxy)-3',5',6',7',8',8a'-hexahydro-2'H-spiro[indoline-2,1'-indolizine]-7'-yl)-3-methoxyacrylate (9-O-trifluoromethane sulfonyl corynantheidine pseudoindoxyl, 5): **4** (200 mg, 0.5 mmol) was dissolved in dry DCM (15 mL) and pyridine (647 μL , 16 equiv) was added. Then the solution was cooled to -40°C on a dry ice acetone bath, and the following solution was slowly added over 2-3 minutes: 5 mL DCM and triflic anhydride (340 μL , 4 equiv). The reaction was stirred for 1h at -40°C . After warming up to rt, the solution was purified using flash column chromatography without immediately (gradient: 20-75 % EtOAc in hexanes) to yield 233 mg (83%) of **5** as a brown

amorphous solid. ^1H NMR (600 MHz, CDCl_3) δ 7.41 (t, $J = 8.1$ Hz, 1H), 7.29 (s, 1H), 6.84 (d, $J = 8.3$ Hz, 1H), 6.52 (d, $J = 7.9$ Hz, 1H), 5.58 (s, 1H), 3.68 (s, 3H), 3.63 (s, 3H), 3.15 (d, $J = 9.7$ Hz, 2H), 2.80 (dd, $J = 12.5, 3.5$ Hz, 1H), 2.35 (d, $J = 7.4$ Hz, 2H), 2.27 (d, $J = 8.8$ Hz, 2H), 2.18 – 2.13 (m, 1H), 1.66 – 1.58 (m, 1H), 1.52 (d, $J = 10.9$ Hz, 1H), 1.22 – 1.16 (m, 2H), 0.85 (t, $J = 7.3$ Hz, 4H). ^{13}C NMR (151 MHz, CDCl_3) δ 198.15, 171.40, 168.92, 161.46, 160.55, 145.34, 138.13, 111.73, 110.01, 73.58, 61.69, 60.62, 54.84, 53.65, 53.33, 51.38, 40.14, 38.33, 35.12, 23.84, 21.28, 19.41, 14.41, 13.00. HRMS calcd for $\text{C}_{23}\text{H}_{27}\text{F}_3\text{N}_2\text{O}_7\text{S}$ (MH⁺), 533.1569; found 533.1547.

(E)-Methyl-2-((1'S,6'S,7'S)-6'-ethyl-3-oxo-3',5',6',7',8',8a'-hexahydro-2'H-spiro[indoline-2,1'-indolizin]-7'-yl)-3-methoxyacrylate (Corynantheidine pseudoindoxyl, 6): **5** (10 mg, 0.019 mmol) was dissolved in dry DMF (500 μL) in a sealed tube and the following reagents were added: $\text{Pd}(\text{OAc})_2$ (1.4 mg, 0.3 equiv), dppp (4 mg, 0.5 equiv), triethylamine (52.4 μL , 20 equiv) and formic acid (1 μL , 1.8 equiv). The mixture was stirred at 60 $^\circ\text{C}$ for 1h. The reaction mixture was diluted with EtOAc (10 mL) and washed with brine (5 mL) 5 times. The organic layer was dried over Na_2SO_4 and evaporated under reduced pressure. The residue was redissolved in DCM and purified using preparative TLC (75% EtOAc in hexanes) to yield 4 mg (57%) of **6** as an amorphous solid. IR (NaCl): 3286, 2927, 2360, 1676, 1620, 1437, 1248, 1200, 1138, 755. ^1H NMR (600 MHz, CDCl_3) δ 7.55 (d, $J = 7.7$ Hz, 1H), 7.43 – 7.38 (m, 1H), 7.27 (d, $J = 6.7$ Hz, 1H), 6.85 (d, $J = 8.3$ Hz, 1H), 6.77 – 6.72 (m, 1H), 5.20 (s, 1H), 3.66 (s, 3H), 3.62 (s, 3H), 3.20 – 3.12 (m, 2H), 2.82 – 2.76 (m, 1H), 2.38 – 2.29 (m, 2H), 2.25 (s, 2H), 2.20 – 2.13 (m, 1H), 1.97 – 1.89 (m, 1H), 1.69 – 1.61 (m, 1H), 1.55 – 1.48 (m, 1H), 1.23 – 1.17 (m, 1H), 1.05 (s, 1H), 0.86 (t, $J = 7.4$ Hz, 3H). ^{13}C NMR (151 MHz, CDCl_3) δ 202.60, 169.02, 160.95, 160.51, 137.30, 124.61, 118.30, 111.85, 111.68, 75.22, 73.52, 61.67, 54.94, 53.42, 51.40, 40.20, 38.59,

35.13, 23.92, 21.29, 19.41, 13.06. HRMS calcd for $C_{22}H_{28}N_2O_4$ (MH⁺), 385.2127; found 385.2120.

(E)-Methyl-2-((1'S,6'S,7'S)-4-cyano-6'-ethyl-3-oxo-3',5',6',7',8',8a'-hexahydro-2'H-spiro[indoline-2,1'-indolizine]-7'-yl)-3-methoxyacrylate (9-Cyano corynantheidine pseudoindoxyl, 7): **5** (20 mg, 0.038 mmol) was dissolved in dry DMF (500 μ L) in a sealed tube, Pd(PPh₃)₄ (4.3 mg, 0.1 equiv) and Zn(CN)₂ (8.8 mg, 2 equiv) were added. The reaction mixture was stirred at 80 °C for 3h. After 3h, the reaction was diluted with EtOAc (20 mL) and washed with brine 5 times. The organic layer was dried over Na₂SO₄ and evaporated under reduced pressure. The residue was redissolved in DCM and purified using flash column chromatography (gradient: 20-80% EtOAc in hexanes) to yield 10 mg (65%) of **7** as an amorphous brown solid. IR (NaCl): 3355, 2940, 2233, 1699, 1606, 1501, 1438, 1242, 993, 859, 1082, 776. ¹H NMR (600 MHz, CDCl₃) δ 7.47 – 7.43 (m, 1H), 7.28 (s, 1H), 7.07 (d, *J* = 8.4 Hz, 1H), 7.05 (d, *J* = 7.2 Hz, 1H), 5.43 (s, 1H), 3.68 (s, 3H), 3.62 (s, 3H), 3.15 (t, *J* = 10.1 Hz, 2H), 2.79 (dt, *J* = 12.8, 3.5 Hz, 1H), 2.40 – 2.33 (m, 2H), 2.31 – 2.27 (m, 1H), 2.23 (t, *J* = 12.1 Hz, 1H), 2.19 – 2.13 (m, 1H), 1.96 – 1.90 (m, 1H), 1.66 – 1.59 (m, 1H), 1.51 (d, *J* = 11.2 Hz, 1H), 1.23 – 1.16 (m, 1H), 1.07 (d, *J* = 12.7 Hz, 1H), 0.85 (t, *J* = 7.4 Hz, 3H). ¹³C NMR (151 MHz, CDCl₃) δ 201.47, 169.06, 162.01, 160.51, 142.35, 137.85, 136.72, 129.43, 128.16, 127.88, 120.06, 111.79, 110.73, 75.10, 73.89, 61.70, 55.07, 53.59, 51.40, 40.28, 38.57, 36.87, 36.86, 35.53, 24.92, 24.08, 19.48, 13.08. HRMS calcd for $C_{23}H_{27}N_3O_4$ (MH⁺), 410.2080; found 410.2068.

(E)-Methyl-2-((1'S,6'S,7'S)-6'-ethyl-3-oxo-4-phenyl-3',5',6',7',8',8a'-hexahydro-2'H-spiro[indoline-2,1'-indolizine]-7'-yl)-3-methoxyacrylate (9-Phenyl corynantheidine pseudoindoxyl, 8): **5** (75 mg, 0.14 mmol) was dissolved in dry toluene (0.5 mL) and the solvent was removed under reduced pressure to ensure azeotropic removal of water residues. Dry

methanol (1 mL) and dry toluene (1.5 mL) were added. To the resulting solution were added phenylboronic acid (19 mg, 1.1 equiv), K₂CO₃ (38.9 mg, 2 equiv) and Pd(PPh₃)₄ (8.1 mg, 0.05 equiv). The mixture was stirred at 80 °C for 2 hrs. The solvent was evaporated under reduced pressure and the residue suspended in DCM, rinsed with water and brine (20 mL), then the organic layer was dried over Na₂SO₄ and was evaporated. Purified using flash column chromatography (gradient: 20-50% EtOAc in hexanes) to yield: 21 mg (32%) of **8** as a yellow amorphous solid. IR (NaCl): 3364, 2936, 2360, 1698, 1600, 1483, 1436, 1233, 1150, 759. ¹H NMR (600 MHz, CDCl₃) δ 7.52 – 7.49 (m, 2H), 7.43 – 7.34 (m, 4H), 7.30 (s, 1H), 6.82 (d, *J* = 8.2 Hz, 1H), 6.68 (d, *J* = 7.3 Hz, 1H), 5.26 (s, 1H), 3.69 (s, 3H), 3.63 (s, 3H), 3.15 (d, *J* = 10.0 Hz, 2H), 2.79 (dt, *J* = 13.0, 3.3 Hz, 1H), 2.34 – 2.25 (m, 3H), 2.21 (s, 1H), 2.17 – 2.10 (m, 1H), 1.95 – 1.88 (m, 1H), 1.69 – 1.63 (m, 1H), 1.53 – 1.47 (m, 1H), 1.24 – 1.18 (m, 1H), 1.14 – 1.09 (m, 1H), 0.86 (t, *J* = 7.4 Hz, 3H). ¹³C NMR (151 MHz, CDCl₃) δ 201.47, 169.06, 162.01, 160.51, 142.35, 137.85, 136.72, 129.43, 128.16, 127.88, 120.06, 111.79, 110.73, 75.10, 73.89, 61.70, 55.07, 53.59, 51.40, 40.28, 38.57, 36.87, 36.86, 35.53, 24.92, 24.08, 19.48, 13.08. HRMS calcd for C₂₈H₃₂N₂O₄ (MH⁺), 461.2440; found 461.2422.

(E)-Methyl-2-((1'S,6'S,7'S)-6'-ethyl-4-(furan-3-yl)-3-oxo-3',5',6',7',8',8a'-hexahydro-2'H-spiro[indoline-2,1'-indolizine]-7'-yl)-3-methoxyacrylate (9-Furyl corynantheidine pseudoindoxyl, 9): The procedure described for the synthesis of **8** was used. Instead of phenylboronic acid, (furan-3-yl)boronic acid was employed. Yield: 81%. Compound **9** is a bright yellow amorphous powder. IR (NaCl): 3358, 2954, 2795, 2360, 2341, 1691, 1604, 1437, 1316, 1238, 1152, 796, 772. ¹H NMR (600 MHz, Chloroform-*d*) δ 8.49 (d, *J* = 1.4 Hz, 1H), 7.48 (d, *J* = 1.7 Hz, 1H), 7.40 (t, *J* = 7.8 Hz, 1H), 7.31 (s, 1H), 6.87 (d, *J* = 7.5 Hz, 1H), 6.84 (dd, *J* = 1.9, 0.9 Hz, 1H), 6.77 (d, *J* = 8.1 Hz, 1H), 5.22 (s, 1H), 3.70 (s, 3H), 3.65 (s, 3H), 3.22 – 3.14 (m, 2H),

2.81 (dt, $J = 12.6, 3.7$ Hz, 1H), 2.40 – 2.32 (m, 2H), 2.32 – 2.22 (m, 2H), 2.17 (dd, $J = 11.5, 3.2$ Hz, 1H), 2.00 – 1.92 (m, 1H), 1.75 – 1.65 (m, 1H), 1.57 – 1.51 (m, 1H), 1.23 (dtd, $J = 15.1, 7.4, 2.7$ Hz, 1H), 1.15 – 1.10 (m, 1H), 0.89 (t, $J = 7.4$ Hz, 3H). ^{13}C NMR (151 MHz, CDCl_3) δ 201.73, 169.03, 162.40, 160.45, 143.85, 142.48, 137.00, 132.42, 122.74, 118.06, 116.66, 111.77, 110.60, 110.18, 75.11, 74.00, 61.65, 55.07, 53.54, 51.38, 40.28, 38.61, 35.68, 24.01, 19.50, 13.11. HRMS calcd for $\text{C}_{26}\text{H}_{30}\text{N}_2\text{O}_5$ (MH⁺), 451.2433; found 451.2215.

(E)-Methyl-2-((1'S,6'S,7'S)-4-acetoxy-6'-ethyl-3-oxo-3',5',6',7',8',8a'-hexahydro-2'H-spiro[indoline-2,1'-indolizine]-7'-yl)-3-methoxyacrylate (9-O-Acetyl corynantheidine pseudoindoxyl, 10): **4** (20 mg, 0.05 mmol) was dissolved in pyridine (0.5 mL) and acetic anhydride (80 μL) was added. The mixture was stirred at rt for 2h. The solution was poured into sat. aqueous NaHCO_3 solution and extracted with DCM (30 mL). The organic layer was separated, rinsed with brine (10 mL), dried over Na_2SO_4 and evaporated under reduced pressure the residue was redissolved in DCM and purified using flash column chromatography (gradient: 0-5% MeOH in DCM) to yield 13 mg (59%) of **10** as a bright yellow amorphous solid. IR (NaCl): 3393, 2956, 2874, 2787, 1761, 1688, 1628, 1503, 1239, 1217, 910, 764. ^1H NMR (500 MHz, Chloroform- d) δ 7.38 (t, $J = 8.0$ Hz, 1H), 7.28 (s, 1H), 6.71 (d, $J = 8.2$ Hz, 1H), 6.36 (d, $J = 7.6$ Hz, 1H), 5.73 (s, 1H), 3.68 (s, 3H), 3.62 (s, 3H), 3.20 (dd, $J = 8.6, 3.0$ Hz, 2H), 2.79 (dd, $J = 12.6, 3.9$ Hz, 1H), 2.36 (s, 3H), 2.33 – 2.22 (m, 3H), 2.22 – 2.14 (m, 1H), 1.95 (dt, $J = 15.8, 6.3$ Hz, 1H), 1.63 (dt, $J = 13.0, 6.1$ Hz, 1H), 1.53 (d, $J = 11.6$ Hz, 1H), 1.30 – 1.16 (m, 2H), 1.16 – 1.09 (m, 1H), 0.86 (t, $J = 7.3$ Hz, 3H). ^{13}C NMR (151 MHz, CDCl_3) δ 199.57, 169.14, 168.98, 161.57, 160.47, 148.07, 138.28, 113.07, 111.70, 110.70, 109.35, 75.71, 73.70, 61.67, 55.02, 53.49, 51.38, 40.25, 38.51, 35.16, 23.91, 21.01, 19.49, 13.08. HRMS calcd for $\text{C}_{24}\text{H}_{30}\text{N}_2\text{O}_6$ (MH⁺), 443.2182; found 443.2174.

(E)-Methyl 2-((1'S,6'S,7'S)-1-benzyl-6'-ethyl-4-methoxy-3-oxo-3',5',6',7',8',8a'-hexahydro-2'H-spiro[indoline-2,1'-indolizine]-7'-yl)-3-methoxyacrylate (N-Benzyl mitragynine pseudoindoxyl, 11): Compound **3** was dissolved in dry acetonitrile (0.5 mL) and NaH (6 mg, 10 equiv) was added. The resulting suspension was stirred at rt for 30 minutes, during which its color turned red. Benzyl bromide (7.2 uL, 2.5 equiv) was added and the mixture stirred for 2h at rt. The red color disappeared promptly after the addition of benzyl bromide. After the reaction time, the mixture was carefully poured into cold water (20 mL) and extracted with DCM (30 mL). The organic layer was rinsed with brine (5 mL) separated, dried over Na₂SO₄ and evaporated under reduced pressure. The residue was redissolved in DCM and purified using flash column chromatography (gradient: 0-5 % MeOH in DCM) to yield: 7.8 mg (64 %) of **11** as a bright yellow amorphous solid. IR (NaCl): 2940, 2794, 1694, 1610, 1497, 1337, 1265, 1239, 1078, 732. ¹H NMR (500 MHz, Chloroform-*d*) δ 7.42 – 7.28 (multiple overlapping peaks, 5H), 7.25 – 7.17 (multiple overlapping peaks, 2H), 6.08 (d, *J* = 8.1 Hz, 1H), 6.04 (d, *J* = 8.3 Hz, 1H), 5.35 (d, *J* = 17.4 Hz, 1H), 4.73 (d, *J* = 17.3 Hz, 1H), 3.90 (s, 3H), 3.70 (s, 3H), 3.64 (s, 3H), 3.13 (dd, *J* = 11.1, 2.2 Hz, 1H), 3.02 (t, *J* = 8.1 Hz, 1H), 2.78 – 2.72 (m, 1H), 2.33 – 2.18 (m, 4H), 2.09 – 2.02 (m, 1H), 1.94 (dt, *J* = 13.7, 8.5 Hz, 1H), 1.66 (dt, *J* = 19.2, 6.8 Hz, 1H), 1.50 (d, *J* = 10.9 Hz, 1H), 1.13 (d, *J* = 11.8 Hz, 1H), 0.86 (t, *J* = 7.4 Hz, 3H). ¹³C NMR (151 MHz, CDCl₃) δ 199.55, 169.06, 161.60, 160.40, 158.81, 138.88, 138.82, 129.25, 129.02, 128.76, 128.64, 126.95, 126.51, 111.99, 108.49, 101.82, 98.07, 61.63, 55.89, 55.23, 53.92, 51.38, 47.61, 40.46, 38.67, 33.80, 32.06, 24.18, 19.82, 13.13. HRMS calcd for C₃₀H₃₆N₂O₅ (MH⁺), 505.2702; found 505.2726.

(E)-Methyl 2-((1'S,6'S,7'S)-6'-ethyl-4-methoxy-1-methyl-3-oxo-3',5',6',7',8',8a'-hexahydro-2'H-spiro[indoline-2,1'-indolizine]-7'-yl)-3-methoxyacrylate (N-Methyl mitragynine

pseudoindoxyl, 12): The procedure described for the synthesis of **11** was used. Instead of benzyl bromide, iodomethane was employed. Yield: 58%. Compound **12** is a bright yellow amorphous powder. IR (NaCl): 2947, 2778, 2361, 1687, 1611, 1500, 1337, 1273. ¹H NMR (600 MHz, Chloroform-*d*) δ 7.34 (t, *J* = 8.1 Hz, 1H), 7.28 (s, 1H), 6.25 (d, *J* = 8.2 Hz, 1H), 6.05 (d, *J* = 8.0 Hz, 1H), 3.88 (s, 3H), 3.69 (s, 3H), 3.62 (s, 3H), 3.17 (s, 3H), 3.16 – 3.10 (m, 2H), 2.74 (dt, *J* = 13.0, 3.7 Hz, 1H), 2.34 (q, *J* = 8.9 Hz, 1H), 2.26 (dd, *J* = 11.3, 2.6 Hz, 1H), 2.21 – 2.12 (m, 2H), 2.08 – 2.03 (m, 1H), 1.92 (dt, *J* = 13.8, 8.7 Hz, 1H), 1.69 – 1.62 (m, 1H), 1.48 (dt, *J* = 11.1, 3.1 Hz, 1H), 1.28 – 1.23 (m, 1H), 1.20 (ddd, *J* = 13.3, 7.6, 2.8 Hz, 1H), 1.08 (dt, *J* = 13.0, 3.1 Hz, 1H), 0.84 (t, *J* = 7.4 Hz, 3H). ¹³C NMR (151 MHz, CDCl₃) δ 199.87, 169.08, 162.01, 160.36, 158.90, 139.01, 112.04, 108.08, 100.37, 97.46, 78.27, 74.62, 61.59, 55.86, 55.14, 53.88, 51.34, 40.54, 38.72, 31.99, 30.02, 24.29, 19.71, 13.17. HRMS calcd for C₂₄H₃₂N₂O₅ (MH⁺), 429.2389; found 429.2393.

ACKNOWLEDGEMENTS

This work was supported by research grants from the National Institute on Drug Abuse (DA034106) to SM, The Harrington Discovery Institute Scholar Award and a grant from the Peter F. McManus Charitable Trust to GWP, DA013997 and DA02944 to YXP, a National Science Foundation Graduate Research Fellowship (DGE-1257284) to GFM. The State of Florida, Executive Office of the Governor's Office of Tourism, Trade, and Economic Development to JPM. This research was funded in part through the NIH/NCI Cancer Center Support Grant P30 CA008748 to MSKCC. AB was supported by the János Bolyai Research Fellowship. The authors would like to thank Professor Bryan L. Roth and the National Institute of Mental Health's Psychoactive Drug Screening Program for their assistance in determining binding affinities at non-opioid CNS targets. The authors wish to thank Dr. Antara Majumdar of

Non-clinical Biostatistics division at Bristol-Meyers Squibb, NJ for performing the statistical analysis for this manuscript and George Sukenick and Rong Wang of the NMR Analytical Core Facility at MSKCC for their assistance with NMR and MS instruments and experiments.

Corresponding Author Information

Susruta Majumdar, PhD

Tel: 646-888-3669

Email: majumdas@mskcc.org

Abbreviations Used:

%MPE – percent maximal effect; 6TM – six transmembrane; 7TM – seven transmembrane; CHO – Chinese hamster ovary; CPA – conditioned place aversion; CPP – conditioned place preference; DAMGO – [D-Ala², N-MePhe⁴, Gly-ol⁵]-enkephalin; DCM – dichloromethane; DMF – *N,N*-dimethylformamide; DOR-1 – a delta opioid receptor (*Oprd1*) clone; DPDPE – [D-Pen², D-Pen⁵]Enkephalin; dppp - 1,3-bis(diphenylphosphino)propane; GDP – guanosine diphosphate; KO – knockout; KOR-1 – a kappa opioid receptor (*Oprk1*) clone; MOR-1 – a mu opioid receptor (*Oprm1*) clone; norBNI – norbinaltorphimine; NTI – naltrindole; PIFA – [bis(trifluoroacetoxy)iodo]benzene.

REFERENCES

1. Minami, M.; Satoh, M. Molecular biology of the opioid receptors: structures, functions and distributions. *Neurosci. Res.* **1995**, *23*, 121-145.
2. Eguchi, M. Recent advances in selective opioid receptor agonists and antagonists. *Med. Res. Rev.* **2004**, *24*, 182-212.
3. Trescot, A. M.; Datta, S.; Lee, M.; Hans, H. Opioid pharmacology. *Pain Physician* **2008**, *11*, S133-S153.
4. Pasternak, G. W. Pharmacological mechanisms of opioid analgesics. *Clin. Neuropharmacol.* **1993**, *16*, 1-18.
5. Ananthan, S. Opioid ligands with mixed mu/delta opioid receptor interactions: an emerging approach to novel analgesics. *AAPS J.* **2006**, *8*, E118-125.
6. Mosberg, H. I.; Yeomans, L.; Anand, J. P.; Porter, V.; Sobczyk-Kojiro, K.; Traynor, J. R.; Jutkiewicz, E. M. Development of a bioavailable mu opioid receptor (MOPr) agonist, delta opioid receptor (DOPr) antagonist peptide that evokes antinociception without development of acute tolerance. *J. Med. Chem.* **2014**, *57*, 3148-3153.
7. Váradi, A.; Hosztafi, S.; Le Rouzic, V.; Tóth, G.; Urai, A.; Noszal, B.; Pasternak, G. W.; Grinnell, S. G.; Majumdar, S. Novel 6beta-acylaminomorphinans with analgesic activity. *Eur. J. Med. Chem.* **2013**, *69C*, 786-789.
8. Váradi, A.; Marrone, G. F.; Eans, S. O.; Ganno, M. L.; Subrath, J. J.; Le Rouzic, V.; Hunkele, A.; Pasternak, G. W.; McLaughlin, J. P.; Majumdar, S. Synthesis and characterization of a dual kappa-delta opioid receptor agonist analgesic blocking cocaine reward behavior. *ACS Chem. Neurosci.* **2015**, *6*, 1813-1824.

9. Váradi, A.; Palmer, T. C.; Haselton, N.; Afonin, D.; Subrath, J. J.; Le Rouzic, V.; Hunkele, A.; Pasternak, G. W.; Marrone, G. F.; Borics, A.; Majumdar, S. Synthesis of carfentanil amide opioids using the ugi multicomponent reaction. *ACS Chem. Neurosci.* **2015**, *6*, 1570-1577.
10. Law, P. Y.; Reggio, P. H.; Loh, H. H. Opioid receptors: toward separation of analgesic from undesirable effects. *Trends Biochem. Sci.* **2013**, *38*, 275-282.
11. Pradhan, A. A.; Smith, M. L.; Kieffer, B. L.; Evans, C. J. Ligand-directed signalling within the opioid receptor family. *Br. J. Pharmacol.* **2012**, *167*, 960-969.
12. Raehal, K. M.; Bohn, L. M. The role of beta-arrestin2 in the severity of antinociceptive tolerance and physical dependence induced by different opioid pain therapeutics. *Neuropharmacology* **2011**, *60*, 58-65.
13. Bohn, L. M.; Lefkowitz, R. J.; Caron, M. G. Differential mechanisms of morphine antinociceptive tolerance revealed in (beta)arrestin-2 knock-out mice. *J. Neurosci.* **2002**, *22*, 10494-10500.
14. Dewire, S. M.; Yamashita, D. S.; Rominger, D. H.; Liu, G.; Cowan, C. L.; Graczyk, T. M.; Chen, X. T.; Pitis, P. M.; Gotchev, D.; Yuan, C.; Koblish, M.; Lark, M. W.; Violin, J. D. A G protein-biased ligand at the mu-opioid receptor is potently analgesic with reduced gastrointestinal and respiratory dysfunction compared with morphine. *J. Pharmacol. Exp. Ther.* **2013**, *344*, 708-717.
15. Soergel, D. G.; Subach, R. A.; Burnham, N.; Lark, M. W.; James, I. E.; Sadler, B. M.; Skobieranda, F.; Violin, J. D.; Webster, L. R. Biased agonism of the mu-opioid receptor by TRV130 increases analgesia and reduces on-target adverse effects versus morphine: A

- randomized, double-blind, placebo-controlled, crossover study in healthy volunteers. *Pain* **2014**, *155*, 1829-1835.
16. Rives, M. L.; Rossillo, M.; Liu-Chen, L. Y.; Javitch, J. A. 6'-Guanidinonaltrindole (6'-GNTI) is a G protein-biased kappa-opioid receptor agonist that inhibits arrestin recruitment. *J. Biol. Chem.* **2012**, *287*, 27050-27054.
17. White, K. L.; Robinson, J. E.; Zhu, H.; DiBerto, J. F.; Polepally, P. R.; Zjawiony, J. K.; Nichols, D. E.; Malanga, C. J.; Roth, B. L. The G protein-biased κ -opioid receptor agonist RB-64 is analgesic with a unique spectrum of activities in vivo. *J. Pharmacol. Exp. Ther.* **2015**, *352*, 98-109.
18. Lovell, K. M.; Frankowski, K. J.; Stahl, E. L.; Slauson, S. R.; Yoo, E.; Prisinzano, T. E.; Aubé, J.; Bohn, L. M. structure-activity relationship studies of functionally selective kappa opioid receptor agonists that modulate ERK 1/2 phosphorylation while preserving G protein over β arrestin2 signaling bias. *ACS Chem. Neurosci.* **2015**, *6*, 1411-1419.
19. Zhou, L.; Lovell, K. M.; Frankowski, K. J.; Slauson, S. R.; Phillips, A. M.; Streicher, J. M.; Stahl, E.; Schmid, C. L.; Hodder, P.; Madoux, F.; Cameron, M. D.; Prisinzano, T. E.; Aubé, J.; Bohn, L. M. Development of functionally selective, small molecule agonists at kappa opioid receptors. *J. Biol. Chem.* **2013**, *288*, 36703-36716.
20. Li, J. W.; Vederas, J. C. Drug discovery and natural products: end of an era or an endless frontier? *Science* **2009**, *325*, 161-165.
21. Prisinzano, T. E. Psychopharmacology of the hallucinogenic sage *Salvia divinorum*. *Life Sci.* **2005**, *78*, 527-531.
22. Prisinzano, T. E. Natural products as tools for neuroscience: discovery and development of novel agents to treat drug abuse. *J. Nat. Prod.* **2009**, *72*, 581-587.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
23. Tidgewell, K.; Harding, W. W.; Lozama, A.; Cobb, H.; Shah, K.; Kannan, P.; Dersch, C. M.; Parrish, D.; Deschamps, J. R.; Rothman, R. B.; Prisinzano, T. E. Synthesis of salvinorin A analogues as opioid receptor probes. *J. Nat. Prod.* **2006**, *69*, 914-918.
24. Tidgewell, K.; Groer, C. E.; Harding, W. W.; Lozama, A.; Schmidt, M.; Marquam, A.; Hiemstra, J.; Partilla, J. S.; Dersch, C. M.; Rothman, R. B.; Bohn, L. M.; Prisinzano, T. E. Herkinorin analogues with differential beta-arrestin-2 interactions. *J. Med. Chem.* **2008**, *51*, 2421-2431.
25. Aldrich, J. V.; Kulkarni, S. S.; Senadheera, S. N.; Ross, N. C.; Reilley, K. J.; Eans, S. O.; Ganno, M. L.; Murray, T. F.; McLaughlin, J. P. Unexpected opioid activity profiles of analogues of the novel peptide kappa opioid receptor ligand CJ-15,208. *ChemMedChem* **2011**, *6*, 1739-1745.
26. Jansen, K. L.; Prast, C. J. Ethnopharmacology of kratom and the Mitragyna alkaloids. *J. Ethnopharmacol.* **1988**, *23*, 115-119.
27. Karinen, R.; Fosen, J. T.; Rogde, S.; Vindenes, V. An accidental poisoning with mitragynine. *Forensic Sci. Int.* **2014**, *245C*, e29-e32.
28. Kronstrand, R.; Roman, M.; Thelander, G.; Eriksson, A. Unintentional fatal intoxications with mitragynine and O-desmethyltramadol from the herbal blend krypton. *J. Anal. Toxicol.* **2011**, *35*, 242-247.
29. McIntyre, I. M.; Trochta, A.; Stolberg, S.; Campman, S. C. Mitragynine 'kratom' related fatality: a case report with postmortem concentrations. *J. Anal. Toxicol.* **2015**, *39*, 152-155.
30. Neerman, M. F.; Frost, R. E.; Deking, J. A drug fatality involving Kratom. *J. Forensic Sci.* **2013**, *58* Suppl 1, S278-279.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
31. Macko, E.; Weisbach, J. A.; Douglas, B. Some observations on the pharmacology of mitragynine. *Arch. Int. Pharmacodyn. Ther.* **1972**, *198*, 145-161.
32. Matsumoto, K.; Hatori, Y.; Murayama, T.; Tashima, K.; Wongseripipatana, S.; Misawa, K.; Kitajima, M.; Takayama, H.; Horie, S. Involvement of mu-opioid receptors in antinociception and inhibition of gastrointestinal transit induced by 7-hydroxymitragynine, isolated from Thai herbal medicine *Mitragyna speciosa*. *Eur. J. Pharmacol.* **2006**, *549*, 63-70.
33. Matsumoto, K.; Horie, S.; Ishikawa, H.; Takayama, H.; Aimi, N.; Ponglux, D.; Watanabe, K. Antinociceptive effect of 7-hydroxymitragynine in mice: Discovery of an orally active opioid analgesic from the Thai medicinal herb *Mitragyna speciosa*. *Life Sci.* **2004**, *74*, 2143-2155.
34. Matsumoto, K.; Horie, S.; Takayama, H.; Ishikawa, H.; Aimi, N.; Ponglux, D.; Murayama, T.; Watanabe, K. Antinociception, tolerance and withdrawal symptoms induced by 7-hydroxymitragynine, an alkaloid from the Thai medicinal herb *Mitragyna speciosa*. *Life Sci.* **2005**, *78*, 2-7.
35. Matsumoto, K.; Mizowaki, M.; Suchitra, T.; Murakami, Y.; Takayama, H.; Sakai, S.-i.; Aimi, N.; Watanabe, H. Central antinociceptive effects of mitragynine in mice: contribution of descending noradrenergic and serotonergic systems. *Eur. J. Pharmacol.* **1996**, *317*, 75-81.
36. Matsumoto, K.; Mizowaki, M.; Suchitra, T.; Takayama, H.; Sakai, S.; Aimi, N.; Watanabe, H. Antinociceptive action of mitragynine in mice: evidence for the involvement of supraspinal opioid receptors. *Life Sci.* **1996**, *59*, 1149-1155.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
37. Matsumoto, K.; Yamamoto, L. T.; Watanabe, K.; Yano, S.; Shan, J.; Pang, P. K.; Ponglux, D.; Takayama, H.; Horie, S. Inhibitory effect of mitragynine, an analgesic alkaloid from Thai herbal medicine, on neurogenic contraction of the vas deferens. *Life Sci.* **2005**, *78*, 187-194.
38. Takayama, H.; Ishikawa, H.; Kitajima, M.; Aimi, N. Formation of an unusual dimeric compound by lead tetraacetate oxidation of a corynanthe-type indole alkaloid, mitragynine. *Chem. Pharm. Bull.* **2002**, *50*, 960-963.
39. Takayama, H.; Ishikawa, H.; Kurihara, M.; Kitajima, M.; Aimi, N.; Ponglux, D.; Koyama, F.; Matsumoto, K.; Moriyama, T.; Yamamoto, L. T.; Watanabe, K.; Murayama, T.; Horie, S. Studies on the synthesis and opioid agonistic activities of mitragynine-related indole alkaloids: discovery of opioid agonists structurally different from other opioid ligands. *J. Med. Chem.* **2002**, *45*, 1949-1956.
40. Zarembo, J. E.; Douglas, B.; Valenta, J.; Weisbach, J. A. Metabolites of mitragynine. *J. Pharm. Sci.* **1974**, *63*, 1407-1415.
41. Yamamoto, L. T.; Horie, S.; Takayama, H.; Aimi, N.; Sakai, S.; Yano, S.; Shan, J.; Pang, P. K.; Ponglux, D.; Watanabe, K. Opioid receptor agonistic characteristics of mitragynine pseudoindoxyl in comparison with mitragynine derived from Thai medicinal plant *Mitragyna speciosa*. *Gen. Pharmacol.* **1999**, *33*, 73-81.
42. Ponglux, D.; Wongseripipatana, S.; Takayama, H.; Kikuchi, M.; Kurihara, M.; Kitajima, M.; Aimi, N.; Sakai, S. A New Indole Alkaloid, 7 α -Hydroxy-7H-mitragynine, from *Mitragyna speciosa* in Thailand. *Planta Med.* **1994**, *60*, 580-581.
43. Mercado-Marin, E. V.; Garcia-Reynaga, P.; Romminger, S.; Pimenta, E. F.; Romney, D. K.; Lodewyk, M. W.; Williams, D. E.; Andersen, R. J.; Miller, S. J.; Tantillo, D. J.;

- Berlinck, R. G. S.; Sarpong, R. Total synthesis and isolation of citrinalin and cyclopiamine congeners. *Nature* **2014**, *509*, 318-324.
44. Váradi, A.; Marrone, G. F.; Palmer, T. C.; Hunkele, A.; Le Rouzic, V.; Pasternak, G. W.; Pan, Y.-X.; McLaughlin, J. P.; Majumdar, S. Mitragyna alkaloids revisited: pharmacology of mitragynine and related natural products. In *Poster Presentation, International Narcotics Research Conference 2015*, Phoenix, AZ, June 15-19, **2015**.
45. Besnard, J.; Ruda, G. F.; Setola, V.; Abecassis, K.; Rodriguiz, R. M.; Huang, X. P.; Norval, S.; Sassano, M. F.; Shin, A. I.; Webster, L. A.; Simeons, F. R.; Stojanovski, L.; Prat, A.; Seidah, N. G.; Constan, D. B.; Bickerton, G. R.; Read, K. D.; Wetsel, W. C.; Gilbert, I. H.; Roth, B. L.; Hopkins, A. L. Automated design of ligands to polypharmacological profiles. *Nature* **2012**, *492*, 215-220.
46. Rossi, G. C.; Pan, Y.-X.; Brown, G. P.; Pasternak, G. W. Antisense mapping the MOR-1 opioid receptor: Evidence for alternative splicing and a novel morphine-6b-glucuronide receptor. *FEBS Lett.* **1995**, *369*, 192-196.
47. Rossi, G. C.; Pan, Y.-X.; Cheng, J.; Pasternak, G. W. Blockade of morphine analgesia by an antisense oligodeoxynucleotide against the mu receptor. *Life Sci.* **1994**, *54*, L375-L379.
48. Standifer, K. M.; Chien, C.-C.; Wahlestedt, C.; Brown, G. P.; Pasternak, G. W. Selective loss of δ opioid analgesia and binding by antisense oligodeoxynucleotides to a δ opioid receptor. *Neuron* **1994**, *12*, 805-810.
49. Pasternak, K. R.; Rossi, G. C.; Zuckerman, A.; Pasternak, G. W. Antisense mapping KOR-1: evidence for multiple kappa analgesic mechanisms. *Brain Res.* **1999**, *826*, 289-292.
50. Pan, Y.-X.; Pasternak, G. W. Molecular Biology of Mu Opioid Receptors. In *The Opiate Receptors*, 2nd ed.; Pasternak, G. W., Ed. Springer: New York, 2011; pp 121-160.

- 1
2
3 51. Pasternak, G. W.; Pan, Y.-X. Mu opioids and their receptors: Evolution of a concept.
4
5 *Pharmacol. Rev.* **2013**, *65*, 1257-1317.
6
7
8 52. Pan, Y.-X.; Xu, J.; Mahurter, L.; Bolan, E. A.; Xu, M. M.; Pasternak, G. W. Generation of
9
10 the mu opioid receptor (MOR-1) protein by three new splice variants of the *Oprm* gene.
11
12 *Proc. Natl. Acad. Sci. U.S.A.* **2001**, *98*, 14084-14089.
13
14
15 53. Xu, J.; Xu, M.; Hurd, Y. L.; Pasternak, G. W.; Pan, Y. X. Isolation and characterization of
16
17 new exon 11-associated N-terminal splice variants of the human mu opioid receptor gene.
18
19 *J. Neurochem.* **2009**, *108*, 962-972.
20
21
22 54. Pan, Y. X.; Xu, J.; Xu, M.; Rossi, G. C.; Matulonis, J. E.; Pasternak, G. W. Involvement of
23
24 exon 11-associated variants of the mu opioid receptor MOR-1 in heroin, but not morphine,
25
26 actions. *Proc. Natl. Acad. Sci. U.S.A.* **2009**, *106*, 4917-4922.
27
28
29 55. Lu, Z.; Xu, J.; Rossi, G. C.; Majumdar, S.; Pasternak, G. W.; Pan, Y.-X. Mediation of
30
31 opioid analgesia by a truncated 6-transmembrane GPCR. *J. Clin. Invest.* **2015**, *125*, 2626-
32
33 2630.
34
35
36 56. Tryba, M.; Gehling, M. Clonidine--a potent analgesic adjuvant. *Curr. Opin. Anaesthesiol.*
37
38 **2002**, *15*, 511-517.
39
40
41 57. Schiller, P. W.; Fundytus, M. E.; Merovitz, L.; Weltrowska, G.; Nguyen, T. M. D.;
42
43 Lemieux, C.; Chung, N. N.; Coderre, T. J. The opioid μ agonist/ δ antagonist DIPP-NH₂[Ψ]
44
45 produces a potent analgesic effect, no physical dependence, and less tolerance than
46
47 morphine in rats. *J. Med. Chem.* **1999**, *42*, 3520-3526.
48
49
50 58. Ananthan, S.; Saini, S. K.; Dersch, C. M.; Xu, H.; McGlinchey, N.; Giuvelis, D.; Bilsky, E.
51
52 J.; Rothman, R. B. 14-Alkoxy- and 14-acyloxypyridomorphinans: μ agonist/ δ antagonist
53
54
55
56
57
58
59
60

- opioid analgesics with diminished tolerance and dependence side effects. *J. Med. Chem.* **2012**, *55*, 8350-8363.
59. Healy, J. R.; Bezawada, P.; Shim, J.; Jones, J. W.; Kane, M. A.; MacKerell, A. D.; Coop, A.; Matsumoto, R. R. Synthesis, modeling, and pharmacological evaluation of UMB 425, a mixed μ agonist/ δ antagonist opioid analgesic with reduced tolerance liabilities. *ACS Chem. Neurosci.* **2013**, *4*, 1256-1266.
60. Schiller, P. W.; Weltrowska, G.; Berezowska, I.; Nguyen, T. M. D.; Wilkes, B. C.; Lemieux, C.; Chung, N. N. The TIPP opioid peptide family: Development of δ antagonists, δ agonists, and mixed m agonist/ δ antagonists. *Biopolymers* **1999**, *51*, 411-425.
61. Abdelhamid, E. E.; Sultana, M.; Portoghese, P. S.; Takemori, A. E. Selective blockage of delta opioid receptors prevents the development of morphine tolerance and dependence in mice. *J. Pharmacol. Exp. Ther.* **1991**, *258*, 299-303.
62. Hepburn, M. J.; Little, P. J.; Gingras, J.; Kuhn, C. M. Differential effects of naltrindole on morphine-induced tolerance and physical dependence in rats. *J. Pharmacol. Exp. Ther.* **1997**, *281*, 1350-1356.
63. Kest, B.; Lee, C. E.; McLemore, G. L.; Inturrisi, C. E. An antisense oligodeoxynucleotide to the delta opioid receptor (DOR-1) inhibits morphine tolerance and acute dependence in mice. *Brain Res. Bull.* **1996**, *39*, 185-188.
64. Zhu, Y.; King, M. A.; Schuller, A. G.; Nitsche, J. F.; Reidl, M.; Elde, R. P.; Unterwald, E.; Pasternak, G. W.; Pintar, J. E. Retention of supraspinal delta-like analgesia and loss of morphine tolerance in delta opioid receptor knockout mice. *Neuron* **1999**, *24*, 243-252.

65. Charbogne, P.; Kieffer, B. L.; Befort, K. 15 years of genetic approaches in vivo for addiction research: Opioid receptor and peptide gene knockout in mouse models of drug abuse. *Neuropharmacology* **2014**, *76*, Part B, 204-217.
66. Bohn, L. M.; Gainetdinov, R. R.; Sotnikova, T. D.; Medvedev, I. O.; Lefkowitz, R. J.; Dykstra, L. A.; Caron, M. G. Enhanced rewarding properties of morphine, but not cocaine, in beta(arrestin)-2 knock-out mice. *J. Neurosci.* **2003**, *23*, 10265-10273.
67. Rankovic, Z.; Brust, T. F.; Bohn, L. M. Biased agonism: An emerging paradigm in GPCR drug discovery. *Bioorg. Med. Chem. Lett.* **2016**, *26*, 241-250.
68. Lefkowitz, R. J.; Shenoy, S. K. Transduction of receptor signals by beta-arrestins. *Science* **2005**, *308*, 512-517.
69. Correll, C. C.; McKittrick, B. A. Biased ligand modulation of seven transmembrane receptors (7TMRs): Functional implications for drug discovery. *J. Med. Chem.* **2014**, *57*, 6887-6896.
70. Bohn, L. M.; Lefkowitz, R. J.; Gainetdinov, R. R.; Peppel, K.; Caron, M. G.; Lin, F. T. Enhanced morphine analgesia in mice lacking beta-arrestin 2. *Science* **1999**, *286*, 2495-2498.
71. Bohn, L. M.; Gainetdinov, R. R.; Lin, F. T.; Lefkowitz, R. J.; Caron, M. G. Mu-opioid receptor desensitization by beta-arrestin-2 determines morphine tolerance but not dependence. *Nature* **2000**, *408*, 720-723.
72. Raehal, K. M.; Walker, J. K.; Bohn, L. M. Morphine side effects in beta-arrestin 2 knockout mice. *J. Pharmacol. Exp. Ther.* **2005**, *314*, 1195-1201.
73. Takayama, H. Chemistry and pharmacology of analgesic indole alkaloids from the rubiaceous plant, *Mitragyna speciosa*. *Chem. Pharm. Bull.* **2004**, *52*, 916-928.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
74. Granier, S.; Manglik, A.; Kruse, A. C.; Kobilka, T. S.; Thian, F. S.; Weis, W. I.; Kobilka, B. K. Structure of the delta-opioid receptor bound to naltrindole. *Nature* **2012**, *485*, 400-404.
75. Manglik, A.; Kruse, A. C.; Kobilka, T. S.; Thian, F. S.; Mathiesen, J. M.; Sunahara, R. K.; Pardo, L.; Weis, W. I.; Kobilka, B. K.; Granier, S. Crystal structure of the mu-opioid receptor bound to a morphinan antagonist. *Nature* **2012**, *485*, 321-326.
76. Huang, N.; Shoichet, B. K.; Irwin, J. J. Benchmarking sets for molecular docking. *J. Med. Chem.* **2006**, *49*, 6789-6801.
77. Huang, W.; Manglik, A.; Venkatakrishnan, A. J.; Laeremans, T.; Feinberg, E. N.; Sanborn, A. L.; Kato, H. E.; Livingston, K. E.; Thorsen, T. S.; Kling, R. C.; Granier, S.; Gmeiner, P.; Husbands, S. M.; Traynor, J. R.; Weis, W. I.; Steyaert, J.; Dror, R. O.; Kobilka, B. K. Structural insights into [micro]-opioid receptor activation. *Nature* **2015**, *524*, 315-321.
78. Majumdar, S.; Burgman, M.; Haselton, N.; Grinnell, S.; Ocampo, J.; Pasternak, A. R.; Pasternak, G. W. Generation of novel radiolabeled opiates through site-selective iodination. *Bioorg. Med. Chem. Lett.* **2011**, *21*, 4001-4004.
79. Majumdar, S.; Grinnell, S.; Le Rouzic, V.; Burgman, M.; Polikar, L.; Ansonoff, M.; Pintar, J.; Pan, Y. X.; Pasternak, G. W. Truncated G protein-coupled mu opioid receptor MOR-1 splice variants are targets for highly potent opioid analgesics lacking side effects. *Proc. Natl. Acad. Sci. U.S.A.* **2011**, *108*, 19776-19783.
80. Pickett, J. E.; Váradi, A.; Palmer, T. C.; Grinnell, S. G.; Schrock, J. M.; Pasternak, G. W.; Karimov, R. R.; Majumdar, S. Mild, Pd-catalyzed stannylation of radioiodination targets. *Bioorg. Med. Chem. Lett.* **2015**, *25*, 1761-1764.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
81. Lowry, O. H.; Rosebrough, N. J.; Farr, A. L.; Randall, R. J. Protein measurement with the Folin phenol reagent. *J. Biol. Chem.* **1951**, *193*, 265-275.
82. Bolan, E. A.; Pan, Y. X.; Pasternak, G. W. Functional analysis of MOR-1 splice variants of the mouse mu opioid receptor gene Oprm. *Synapse* **2004**, *51*, 11-8.
83. Haley, T. J.; McCormick, W. G. Pharmacological effects produced by intracerebral injection of drugs in the conscious mouse. *Br. J. Pharmacol. Chemother.* **1957**, *12*, 12-15.
84. Rossi, G. C.; Brown, G. P.; Leventhal, L.; Yang, K.; Pasternak, G. W. Novel receptor mechanisms for heroin and morphine-6b -glucuronide analgesia. *Neurosci. Lett.* **1996**, *216*, 1-4.
85. Rossi, G. C.; Standifer, K. M.; Pasternak, G. W. Differential blockade of morphine and morphine-6b-glucuronide analgesia by antisense oligodeoxynucleotides directed against MOR-1 and G-protein α subunits in rats. *Neurosci. Lett.* **1995**, *198*, 99-102.
86. Rossi, G. C.; Su, W.; Leventhal, L.; Su, H.; Pasternak, G. W. Antisense mapping DOR-1 in mice: further support for delta receptor subtypes. *Brain Res.* **1997**, *753*, 176-179.
87. Paul, D.; Pasternak, G. W. Differential blockade by naloxonazine of two μ opiate actions: analgesia and inhibition of gastrointestinal transit. *Eur. J. Pharmacol.* **1988**, *149*, 403-404.
88. Wu, H.; Wacker, D.; Mileni, M.; Katritch, V.; Han, G. W.; Vardy, E.; Liu, W.; Thompson, A. A.; Huang, X. P.; Carroll, F. I.; Mascarella, S. W.; Westkaemper, R. B.; Mosier, P. D.; Roth, B. L.; Cherezov, V.; Stevens, R. C. Structure of the human kappa-opioid receptor in complex with JDTic. *Nature* **2012**, *485*, 327-332.

TABLES

Table 1. Receptor affinities and subcutaneous antinociception in mice.

Compound	Affinity (K _i nM) ^a			Antinociception
	MOR-1	KOR-1	DOR-1	sc ^b
				mg/kg (CI)
1	230 ± 47	231 ± 21	1011 ± 49	166 (101-283)
2	37 ± 4	132 ± 7	91 ± 8	0.46 (0.39-0.71)
3	0.8 ± 0.2	24 ± 0.9	3.0 ± 1.3	0.76 (0.56-0.83)
4	1.4 ± 0.2	170 ± 61	6.1 ± 1.1	0.18 (0.16-0.20)
6	0.46 ± 0.01	19 ± 4.7	2.9 ± 0.29	0.24 (0.20-0.26)
7	0.5 ± 0.01	47 ± 3.3	2.4 ± 0.3	0.32 (0.26-0.41)
8	0.91 ± 0.06	51 ± 9.7	0.8 ± 0.13	1.0 (0.70-1.35)
9	0.94 ± 0.02	39 ± 11	1.5 ± 0.37	1.1 (0.83-1.38)
10	2.5 ± 0.6	31 ± 14	20 ± 1	0.38 (0.29-0.49)
11	249 ± 41	136 ± 8	258 ± 34	-
12	375 ± 138	>1000	>1000	-
DAMGO	3.3 ± 0.43 ^c	-	-	-
U50,488H	-	0.73 ± 0.32 ^c	-	-
DPDPE	-	-	1.39 ± 0.67 ^c	-
NTI	-	-	0.46 ± 0.32 ^c	-
norBNI	-	0.23 ± 0.03 ^c	-	-
morphine	4.6 ± 1.8 ^c	-	-	2.5(1.8, 3.4)

^aCompetition studies were performed with the indicated compounds against ¹²⁵I-IBNtxA (0.1 nM) in membranes from CHO cells stably expressing the indicated cloned mouse opioid receptors. Results are presented as nM ± SEM from 3 independent experiments performed in triplicate. ^bCumulative dose–response curves were carried out on groups of CD1 mice (n = 10) using radiant heat tail-flick assays with indicated compound at the indicated doses (sc), and antinociception was tested 15 min later at peak effect. Results from two independent experiments are shown as mean (95% CI). ^cValues from the literature.⁷⁸ "-" Denotes not determined or not applicable.

Table 2. [³⁵S]GTPγS functional assays in transfected cell lines.

Compound	[³⁵ S]GTPγS functional assays ^a					
	MOR-1		KOR-1		DOR-1	
	EC ₅₀ (nM)	E _{max} (%)	EC ₅₀ (nM)	IC ₅₀ (nM)	EC ₅₀ (nM)	IC ₅₀ (nM)
1	203 ± 13	65 ± 2.8	-	>4 uM	-	>4 uM
2	53 ± 4	77 ± 5	-	2524 ± 552	-	691 ± 434
3	1.7 ± 0.1	84 ± 5	-	31 ± 3	-	61 ± 6
4	2.0 ± 0.1	124 ± 2	-	-	-	293 ± 129
6	1.4 ± 0.03	116 ± 2	-	252 ± 48	-	193 ± 48
7	0.7 ± 0.2	122 ± 2	-	721 ± 58	-	73 ± 3
8	1.4 ± 0.25	123 ± 5	-	171 ± 10	0.83 ± 0.37 (89 ± 3) ^b	-
9	1.5 ± 0.3	100 ± 1	-	202 ± 14	-	39 ± 5
10	3.9 ± 0.5	120 ± 6	-	-	-	-
11	-	-	-	-	-	-
12	-	-	-	-	-	-
DAMGO	19 ± 7	-	-	-	-	-
U50,488H	-	-	17 ± 6.1	-	-	-
DPDPE	-	-	-	-	10 ± 2.2	-
NTI	-	-	-	-	-	0.72 ± 0.11
norBNI	-	-	-	2.9 ± 0.14	-	-

^aEfficacy data were obtained using agonist induced stimulation of [³⁵S]GTPγS binding assay. Efficacy is represented as EC₅₀ (nM) and percent maximal stimulation (E_{max}) relative to standard agonist DAMGO (MOR-1), DPDPE (DOR-1), or U50,488H (KOR-1) at 1000 nM. To determine the antagonist properties of a compound, membranes were incubated with 100 nM of the appropriate agonist in the presence of varying concentrations of the compound ^bCompound **8** is an agonist at DOR-1. Results are presented as nM ± SEM from 3 independent experiments performed in triplicate. "-" Denotes not determined or not applicable.

Table 3. *In vitro* and *in silico* binding affinities and specific contacts of mitragynine and derivatives.

	Cmpd	In vitro K _i (nM)	in silico K _i (nM) non-specific ^a	in silico K _i (nM) specific ^b	receptor side chains in polar contact with the ligand	receptor side chains in non-polar contact with the ligand
Mu	1	230 ± 47	23	569	Gln ¹²⁴ , Tyr ¹²⁸ , Asp ¹⁴⁷ , Tyr ³²⁶	Met ¹⁵¹ , Trp ²⁹³ , Ile ²⁹⁶ , Ile ³²²
	2	37 ± 4	4.0	55	Asp ¹⁴⁷ , Lys ²³³ , Trp ²⁹³	Tyr ¹⁴⁸ , Met ¹⁵¹ , Leu ²¹⁹ , Leu ²³² , Ile ²⁹⁶
	3	0.75 ± 0.18	2.5	2.5	Asp ¹⁴⁷ , Tyr ¹⁴⁸ , Lys ²³³	Ile ¹⁴⁴ , Leu ²¹⁹ , Leu ²³² , Trp ²⁹³ , Ile ²⁹⁶ , His ²⁹⁷
Delta	1	1011 ± 49	159	1565	Gln ¹⁰⁵ , Lys ¹⁰⁸ , Asp ¹²⁸ , Tyr ¹²⁹ , Tyr ¹²⁹	Leu ¹²⁸ , Lys ²¹⁴ (aliphatic chain), Val ²¹⁷
	2	90 ± 8	19	19	Asp ¹²⁸ , Tyr ¹²⁹	Met ¹³² , Lys ²¹⁴ (aliphatic chain), Val ²¹⁷ , Trp ²⁷⁴ , Ile ²⁷⁷ , Ile ³⁰⁴
	3	3 ± 1.3	17	17	Asp ¹²⁸ , Tyr ¹²⁹	Met ¹³² , Lys ²¹⁴ (aliphatic chain), Val ²¹⁷ , Trp ²⁷⁴ , Ile ²⁷⁷ , Val ²⁸¹
Kappa	1	231 ± 21	17	69	Asp ¹³⁸ , Tyr ¹³⁹ , Ser ²¹¹	Trp ¹²⁴ , Val ¹³⁴ , Leu ¹³⁵
	2	131 ± 7	8.3	53	Thr ¹¹¹ , Asp ¹³⁸ , Tyr ¹³⁹ , Lys ²²⁷ , Tyr ³¹² , Tyr ³²⁰	Gln ¹¹⁵ , Leu ¹³⁵ , Ile ²⁹⁴
	3	24 ± 9	3.2	23	Gln ¹¹⁵ , Asp ¹³⁸ , Tyr ¹³⁹ , Ser ²¹¹ , Tyr ³¹²	Phe ¹¹⁴ , Val ¹¹⁸ , Ile ²⁹⁴ , Ile ³¹⁶

^aInhibitory constants calculated for the lowest energy docked complexes including non-specifically bound poses. ^bInhibitory constants calculated for the lowest energy docked complexes excluding low energy poses which were regarded as false positives.

FIGURE CAPTIONS

Figure 1. Structure of several important opioid natural product analogs.

Figure 2. Structure of the studied mitragynine analogs.

Scheme 1. Synthesis of mitragynine derivatives **2-12**.

Reagents and conditions: (a) PIFA, H₂O, acetonitrile, 0 °C, 1h; (b) Zn(OTf)₂, toluene, 110 °C, 2h; (c) AlCl₃, EtSH, DCM, 0 °C, 5h; (d) Tf₂O, pyridine, DCM, -40 °C, 1h; (e, yielding **6**) Pd(OAc)₂, dppp, HCOOH, DMF, 60 °C, 1h; (f, yielding **7**) Zn(CN)₂, Pd(PPh₃)₄, DMF, 80 °C, 2h; (g, yielding **8** and **9**) phenylboronic acid (**8**) or 3-furanylboronic acid (**9**), Pd(PPh₃)₄, K₂CO₃, MeOH, toluene, 80 °C, 2h; (h) Ac₂O, pyridine, rt, 1h; (i, yielding **11** and **12**) benzyl bromide (**11**) or iodomethane (**12**), NaH, acetonitrile, rt, 2h.

Figure 3. β -arrestin-2 recruitment and antagonism of β -arrestin-2 recruitment.

A) β -arrestin-2 recruitment: β -arrestin-2 recruitment was determined using the DiscoverX PathHunter enzyme complementation assay using modified MOR-1 in CHO cells. Compounds were found to be completely G-protein biased. B) Antagonism of β -arrestin-2 recruitment: The same cells were incubated with the antagonist (**2**, **3**) for 30 minutes at 37 °C prior to the addition of agonist (10 μ M DAMGO) at MOR-1. Compounds **2** and **3** were able to antagonize β -arrestin-2 recruitment by DAMGO. IC₅₀ values: **2**: 725 \pm 292 nM; **3**: 34 \pm 2 nM

Figure 4. Antinociception of compound **3** upon intracerebroventricular, subcutaneous, and oral administration.

A) Dose-response curves of antinociception of **3** and morphine given supraspinally in CD1 mice. Two independent determinations of the cumulative dose-response curves were performed on groups of mice (n = 5) for antinociception in the tail flick assay with **3** intracerebroventricularly.

Animals were tested 15 min later at peak effect to generate the analgesic dose-response curve. Each point represents mean \pm SEM for 10 mice. ED₅₀ values (and 95% CI) were 0.38 (0.18-0.81) μ g for **3**. B) Dose-response curves of antinociception of **3** given subcutaneously in CD1 mice. Three independent determinations of the cumulative dose-response curves were performed on groups of mice (n = 10) for antinociception in the tail flick assay, 30 mice in total. ED₅₀ (and 95% CI) = 0.76 (0.56-0.83) mg/kg. C) Time course of tail flick antinociception of **3** given orally in CD1 male mice. Groups of mice (n=10) were given different doses of **3** orally by gavage and tested for analgesic response at the indicated time points. ED₅₀ (and 95% CI) = 7.5 (4.3-13) mg/kg. D) Hot plate. Groups of CD1 mice (n = 10) were assessed for antinociception of **3** at peak effect in two independent experiments (n = 20 total) in a cumulative dose-response paradigm. Analgesia was determined using a 55 °C hot plate where the latency to respond with a hind paw lick or shake/flutter, whichever came first, was recorded. ED₅₀ (and 95% CI) = 0.99 mg/kg (0.75–1.3).

Figure 5. Pharmacological and genetic reversal of antinociception of **3**.

A) Reversal of antinociception by selective antagonists. Groups of CD1 mice (n = 10) received **3** (1.5 mg/kg sc) and the indicated antagonist. β -Funaltrexamine (β -FNA; 40 mg/kg sc) and norbinaltorphimine (norBNI; 10 mg/kg sc) were administered 24 hours before agonist testing. Naltrindole (NTI; 0.5 mg/kg sc), naloxone (1 mg/kg), and yohimbine (10 mg/kg) were administered 15 minutes before **3**. All antinociception testing was performed 15 min after the administration of **3**. Similar results were observed in two independent replications. **3** antinociception is insensitive to NTI, norBNI, and yohimbine, whereas antinociception is antagonized by β -FNA and naloxone (two-way ANOVA followed by Bonferroni post hoc comparisons test, $p < 0.05$). All values are expressed as the mean \pm SEM. B) Antisense

oligodeoxynucleotide injection: Groups of mice (n=15) received the stated antisense (5-10 μ g) or mismatch (5-10 μ g) oligodeoxynucleotide icv under light isoflurane anesthesia on days 1, 3 and 5. Tail flick antinociception was tested on day 6. Control groups received no injection prior to testing. On test day, mice received **3** (1.5 mg/kg, sc). All experiments were performed 3 times with similar results observed with each determination. Analgesic response of **3** was only affected in mu receptor downregulated mice (MOR-1 AN)(one-way ANOVA followed by Bonferroni post hoc comparisons test). *Significantly different from control ($p < 0.05$). Data for agonist controls is shown in the Supporting Material (Figure S6) and sequences of AN and MIS oligos are shown in table 4 (experimental section). C) Antinociception of **3** in wild-type, exon 11 KO, and exon 1/exon 11 double KO C57 mice. Two independent determinations of the cumulative dose-response curves were performed on groups of mice (n = 5) for antinociception in the tail flick assay with **3** given subcutaneously. Compound **3** displayed similar antinociceptive effects in wild-type ($ED_{50} = 0.83$ mg/kg (0.37-1.9)) and exon 11 KO mice ($ED_{50} = 1.4$ mg/kg (0.34-5.8)), however, no antinociception was observed in exon 1/exon 11 double KO mice suggesting that the antinociceptive effect of **3** is mediated by the E1 MOR-1 variants.

Figure 6. Side effect studies with compound **3**.

A) Antinociceptive tolerance: Mice were dosed 2x daily with 2x antinociceptive ED_{50} with either morphine or **3** until they showed complete analgesic tolerance. **3** showed very slow onset of tolerance compared with morphine. *Significantly different from morphine ($p < 0.05$). The experiment was replicated at least twice with similar results. B) Physical dependence: Groups of mice were dosed 2x daily with 2x antinociceptive ED_{50} with morphine or **3**. Separate groups of mice were used for the 5, 22 and 29-day treatment with **3** and the animals within each group were sacrificed following the experiment with naloxone. Animals were challenged with naloxone


(1 mg/kg) on day 5 of the morphine group, and days 5, 22, and 29 of the **3** groups. Number of jumps was counted over a 15 min period post-injection. The response of mice treated with **3** on either day was not significantly greater than that of mice treated with saline. *Significantly different from saline. (One-way ANOVA followed by Dunnett's multiple comparison test, $p < 0.05$). C) Gastrointestinal transit. Groups of mice ($n = 10$) received saline, morphine (5 mg/kg), or **3** (1.5 and 4 mg/kg) before receiving an oral dose of 0.2 mL of charcoal meal by gavage. Animals were sacrificed 30 min later, and the distance traveled by charcoal was measured. **3** lowered transit significantly compared with saline ($P < 0.05$) but less than morphine at both doses ($P < 0.05$) as determined by ANOVA followed by Tukey's multiple-comparison test. The inhibition of gastrointestinal transit seems to plateau even at doses ~five times higher than the antinociceptive ED_{50} . D) Respiratory rate. Animals were randomly assigned to receive saline ($n = 5$), **3** (1.2 and 3 mg/kg, sc, $n = 5$ at each dose), or morphine (5 and 10 mg/kg, $n = 5$ at each dose). While morphine caused respiratory depression at both 2x and 5x antinociceptive ED_{50} dose (5 and 10 mg/kg, respectively), **3** did not depress respiratory rate at ~2x antinociceptive ED_{50} dose (1.2 mg/kg) and was not significantly different from saline at any time point, whereas morphine (5 mg/kg) decreased respiratory depression in comparison with **3** ($p < 0.05$) as determined by repeated-measures ANOVA followed by Tukey's multiple-comparison test. However, at ~5x antinociceptive ED_{50} dose (3 mg/kg), **3** showed signs of respiratory depression albeit significantly less than the equianalgesic dose of morphine (10 mg/kg) at any given time point. (repeated-measures ANOVA followed by Tukey's multiple-comparison test). E) Conditioned place-preference and aversion. Compound **3** alone did not produce conditioned place preference or aversion. After determination of initial preconditioning preferences, mice were place conditioned daily for 2 days with morphine (10 mg/kg/d, ip), U50,488 (30 mg/kg/d,

ip), cocaine (10 mg/kg/d, ip) or **3** (1.3 mg/kg/d, ip and 3.2 mg/kg/d, ip). Mean difference in time spent on the drug-paired side \pm SEM is presented (n=17-21). *Significantly different from matching preconditioning preference ($p < 0.05$); + significantly different from cocaine, morphine and U50,488 preference (two-way repeated measures ANOVA with Sidak's *post hoc* test).

Figure 7. Lowest energy docked complexes of specifically bound mitragynine derivatives (**1-3**) to opioid receptors.

Binding pocket side chains which were identified in the crystal structures to take part in receptor ligand interactions and were kept flexible during docking are shown. Side chains which are in direct contact with the ligands are depicted in orange. Non-polar hydrogens are omitted for clarity.

Figure 1.



The figure displays 11 chemical structures of indole alkaloids, numbered 1 through 11. Structures 1-11 are indole alkaloids, while structures 12-14 are their precursors. The structures are arranged in three rows: the first row contains structures 1, 2, 3, and 4; the second row contains structures 6, 7, 8, and 9; and the third row contains structures 10, 11, and 12. Each structure is a complex polycyclic molecule featuring an indole ring system fused to a piperidine ring, which is further substituted with a methoxy group, a methyl group, and a methoxycarbonyl group. The structures are labeled with their respective numbers (1, 2, 3, 4, 6, 7, 8, 9, 10, 11, 12) and chemical groups (MeO, HO, Me, OMe, COOMe, CN, Ph, AcO, TZ, Me).

Scheme 1.




Figure 3.




Figure 4.




Figure 5.




Figure 6.




Figure 7.




Table of Contents Graphic

