

## Some characterizations of two-sided regular rings

By S. LAJOS and F. SZÁSZ in Budapest

*To Professor L. Rédei on his seventieth birthday*

Throughout this paper by a ring we mean a not necessarily commutative but associative ring and by the radical of the ring we mean the Jacobson radical [6]. Following J. VON NEUMANN [14] we shall say that the ring  $A$  is *regular* if for every element  $a$  of  $A$  there exists an element  $x$  in  $A$  such that  $axa = a$ . It is well known that the class of regular rings plays a very important rôle in the abstract algebra, in the theory of Banach algebras (cf. C. E. RICKART [17]) and in the continuous geometry [15]. An interesting result is that the ring of all linear transformations of a vector space over a division ring is a regular ring. Some ideal-theoretical characterizations of regular rings were obtained by L. KOVÁCS [8] and J. LUH [12]. The regularity criterion of KOVÁCS reads as follows. An associative ring  $A$  is regular if and only if the relation

$$(1) \quad R \cap L = RL$$

holds for every left ideal  $L$  and for every right ideal  $R$  of  $A$ .

Following E. HILLE [5] a ring  $A$  is called a *two-sided* ring if every one-sided (left or right) ideal of  $A$  is a two-sided ideal of  $A$ . Clearly every division ring is a two-sided ring, and so is every commutative ring. It is easy to see that there exists a two-sided ring which is neither commutative nor a division ring. Two-sided rings, called *duo rings*, were investigated by E. H. FELLER [3] and G. THIERRIN [22]. Thierrin proved, using the classical method of N. H. MCCOY [13], that every two-sided ring can be represented as a subdirect sum of subdirectly irreducible two-sided rings.

A. FORSYTHE and N. H. MCCOY [4] proved the assertion that a nonzero regular ring  $A$  is a subdirect sum of division rings if and only if the ring  $A$  does not contain nonzero nilpotent elements. Their proof uses among others the following lemmas: (1) If a nonzero idempotent element  $e$  of a subdirectly irreducible ring  $A$  lies in the center of  $A$ , then  $e$  is the identity element of  $A$ . (2) If a nonzero subdirectly irreducible regular ring does not contain nonzero nilpotent elements, then it is a division ring.

A ring  $A$  is called *strongly regular* (see R. F. ARENS and I. KAPLANSKY [2])

if to every element  $a$  of  $A$  there exists at least one element  $x$  of  $A$  such that  $a = a^2x$ . It can be seen that every strongly regular rings is regular (see T. KANDÔ [7]), and in a strongly regular ring  $a = a^2x$  if and only if  $a = xa^2$ .

In a paper of the second author [18] it was proved that a ring with minimum condition on principal right ideals is a discrete direct sum of division rings if and only if the ring has no nonzero nilpotent elements. It is clear that this class of rings contains only regular two-sided rings.

The first named author has recently obtained ideal-theoretical characterizations of two-sided regular rings which are analogous to his characterizations of semi-lattices of groups [9], [10], [11]. His earlier criteria are also contained in the following result.

**Theorem.** *For an associative ring  $A$  the following conditions are mutually equivalent:*

- (I)  $A$  is a two-sided regular ring.
- (II)  $L \cap R = LR$  for every left ideal  $L$  and for every right ideal  $R$  of  $A$ .
- (III) The intersection of any two left ideals is equal to their product and the same holds for right ideals too.
- (IV)  $L \cap I = LI$  and  $R \cap I = IR$  for every left ideal  $L$ , for every right ideal  $R$  and, for every two-sided ideal  $I$  of  $A$ .
- (V)  $A$  is regular and a subdirect sum of division rings.
- (VI)  $A$  is a regular ring with no nonzero nilpotent elements.
- (VII)  $A$  is strongly regular.
- (VIII) The intersection of any two left ideals coincides with their product.
- (IX) The intersection of any two right ideals coincides with their product.
- (X)  $L \cap I = LI$  holds for every left ideal  $L$  and for every two-sided ideal  $I$  of  $A$ .
- (XI)  $R \cap I = IR$  holds for every right ideal  $R$  and for every two-sided ideal  $I$  of  $A$ .

**Proof.** (I) $\Rightarrow$ (II). Let  $A$  be a two-sided regular rings. Then  $A$  satisfies the relation

$$(2) \quad L \cap R = RL$$

for every left ideal  $L$  and for every right ideal  $R$  of  $A$  by the regularity criterion of KOVÁCS. In case of two-sided rings this is equivalent to condition (II).

(II) $\Rightarrow$ (I). Let  $A$  be an associative ring having the property (II). In the case of  $R=A$  the condition (II) implies

$$(3) \quad A \cap L = LA,$$

that is, every left ideal  $L$  of  $A$  is also a right ideal of  $A$ . Similarly in case  $L=A$  relation (II) implies

$$(4) \quad A \cap R = AR,$$

thus the right ideal  $R$  of  $A$  is a two-sided ideal of  $A$ . Therefore  $A$  is a two-sided ring. Finally (II) implies relation (2) which is equivalent to the regularity of  $A$ .

(I) $\Leftrightarrow$ (III). The proof is similar to the above proof of the equivalence (I) $\Leftrightarrow$ (II).

(I) $\Rightarrow$ (IV). The proof is analogous to that of (I) $\Rightarrow$ (II).

(IV) $\Rightarrow$ (I). Let  $A$  be a ring with property (IV). In case  $I=A$  we have

$$(5) \quad A \cap L = LA.$$

This means that any left ideal  $L$  of  $A$  is also a right ideal of  $A$ . Consequently the intersection of any two left ideals is equal to their product by (IV). Similarly it can be proved that every right ideal is a two-sided ideal of  $A$  and, the intersection of any two right ideals coincides with their product. Therefore (IV) implies (III), and we have already proved the implication (III) $\Rightarrow$ (I), thus (IV) implies (I).

(I) $\Rightarrow$ (V). Let  $A$  be an arbitrary regular two-sided ring. By the regularity of  $A$  the Jacobson radical  $J$  of  $A$  coincides with the ideal (0). Suppose that  $J \neq (0)$ . Then every nonzero principal right ideal of  $A$  contains a nonzero idempotent element  $e$  and the quasi-regularity condition

$$(6) \quad e + x - ex = 0$$

multiplied on the left by  $e$  yields

$$(7) \quad e = 0,$$

which is a contradiction to the supposition  $e \neq 0$ . Therefore we have  $J = (0)$ . Hence the intersection of all modular maximal right ideals  $I_\alpha$  of  $A$  equals the ideal (0), that is

$$(8) \quad \bigcap_{\alpha} I_{\alpha} = (0).$$

Since  $A$  is a two-sided ring, every right ideal  $I_\alpha$  is two-sided, hence the factor ring  $A/I_\alpha$  has no nontrivial right ideals. By the modularity of  $I_\alpha$  the factor ring  $A/I_\alpha$  is a division ring and, the relation (8) implies the condition (V).

(V) $\Rightarrow$ (VI). The proof is almost trivial, and we omit it.

(VI) $\Rightarrow$ (VII). Let  $A$  be an arbitrary regular ring with no nonzero nilpotent elements. By the mentioned paper of FORSYTHE and MCCOY every idempotent element of  $A$  belongs to the center of  $A$ . Suppose that  $a = axa$  for  $a \in A$ ,  $x \in A$ . Then the idempotent element  $e = ax$  commutes with the element  $a \in A$ , therefore  $a = a^2x$ . Similarly the idempotent element  $f = xa$  also commutes with  $a$  consequently  $a = xa^2$ , that is,  $A$  is strongly regular.

(VII) $\Rightarrow$ (I). Let  $A$  be an arbitrary strongly regular ring. Then the relation  $a \in a^2A$  for every  $a \in A$  implies the fact that  $A$  has no nonzero nilpotent elements because in case  $a^n = 0$  one can conclude

$$(9) \quad a \in a^2A \subseteq (a^3)_r \subseteq a^2 \cdot a^2A \subseteq (a^5)_r \subseteq a^6A \subseteq \dots,$$

whence  $a = 0$ .

We must yet prove that  $A$  is a two-sided ring. To this purpose it is enough to show that every principal right ideal of  $A$  is a two-sided ideal. By the regularity of  $A$  every principal right ideal of  $A$  can be generated by an idempotent element  $e$  of  $A$ . Let now  $a$  be an arbitrary element of the principal right ideal  $(e)$  generated by  $e$ . Then we have  $e^2 = e$  and  $a = ea$  which imply

$$(10) \quad (ae - a)^2 = 0.$$

Since  $A$  has no nonzero nilpotent element, we have  $ae = a$  and hence  $a \in (e)_l$  which implies the inclusion  $(e)_l \subseteq (e)_r$ . The converse inclusion  $(e)_l \subseteq (e)_r$  can be proved similarly; consequently  $(e)_r = (e)_l$ , which means that  $A$  is indeed a two-sided regular ring. Therefore condition (I) holds.

(VII)  $\Leftrightarrow$  (IX). This result was proved by V. A. ANDRUNAKIEVIČ [1].

(VIII)  $\Leftrightarrow$  (IX). By a left-right duality and by the mentioned result of ANDRUNAKIEVIČ it is sufficient to prove that the condition  $a \in Aa^2$  for every element  $a \in A$  is equivalent to one of  $a \in Aa^2$  for every element  $a$  of  $A$ . It was proved in the part (VII)  $\Rightarrow$  (I) that in the case  $a \in a^2A$  ( $\forall a \in A$ ) the ring  $A$  has no nonzero nilpotent elements and, hence every idempotent element lies in the center of  $A$  by FORSYTHE and MCCOY. Therefore  $a = a^2x$  implies  $a = axa$  and  $a = xa^2$ . The proof of the converse statement is similar.

(I)  $\Rightarrow$  (X). The proof is similar to that of (I)  $\Rightarrow$  (II).

(X)  $\Rightarrow$  (I). First in case  $I = A$  condition (X) implies that every left ideal  $L$  of  $A$  is a two-sided ideal. Therefore assertion (X) implies (VIII), which is equivalent to (I).

(I)  $\Rightarrow$  (XI). The proof is the same as in the case (I)  $\Rightarrow$  (II).

(XI)  $\Rightarrow$  (I). The proof is similar to that of (X)  $\Rightarrow$  (I).

The proof of our Theorem is complete.

Remark 1. If the condition

$$(11) \quad \bigcap_{\alpha} (R + I_{\alpha}) \subseteq R + \bigcap_{\alpha} I_{\alpha}$$

holds for every right (and left) ideal  $R$  and for any system of two-sided ideals  $I_{\alpha}$  of a ring  $A$  and  $A$  is a subdirect sum of division rings, then it can be proved by another method that  $A$  is a two-sided ring. Namely let us suppose that

$$(12) \quad \bigcap_{\alpha} I_{\alpha} = (0)$$

holds for the two-sided ideals  $I_{\alpha}$  of  $A$ , where the factor rings  $A/I_{\alpha}$  are division rings. Then the images of the arbitrary right ideal  $R$  of  $A$  are two-sided ideals in the rings  $A/I_{\alpha}$ . Furthermore the complete inverse images  $R + I_{\alpha}$  of  $R$  are two-sided ideals

in  $A$  by the first isomorphism theorem (see e.g. L. RÉDEI [16]). Then the condition (11) together with (12) implies

$$(13) \quad \bigcap_{\alpha} (R + I_{\alpha}) \subseteq R.$$

But conversely we trivially have

$$(14) \quad R \subseteq \bigcap_{\alpha} (R + I_{\alpha}),$$

whence

$$(15) \quad R = \bigcap_{\alpha} (R + I_{\alpha}).$$

Here the intersection  $\bigcap_{\alpha} (R + I_{\alpha})$  is a two-sided ideal of  $A$ , and thus  $R$  is also a two-sided ideal. Therefore  $A$  is a two-sided regular ring. Condition (11) seems to be very similar to the modularity condition of a lattice (see G. SZÁSZ [21]).

**Remark 2.** We mention a nontrivial example for a two-sided regular ring which is neither a commutative nor a division ring. Let  $A$  be the direct sum of two non-commutative division rings. Then  $A$  has obviously the wished properties.

### Bibliography

- [1] B. A. Андрунакиевич, О строго регулярных кольцах, *Изв. Акад. Наук Молдавской ССР*, **11** (1963), 75—77.
- [2] R. F. ARENS and I. KAPLANSKY, Topological representation of algebras, *Trans. Amer. Math. Soc.*, **63** (1948), 457—481.
- [3] E. H. FELLER, Properties of primary noncommutative rings, *Trans. Amer. Math. Soc.*, **89** (1958), 79—91.
- [4] A. FORSYTHE and N. H. MCCOY, On the commutativity of certain rings, *Bull. Amer. Math. Soc.*, **52** (1946), 523—526.
- [5] E. HILLE, *Functional analysis and semi-groups* (New York, 1948).
- [6] N. JACOBSON, *Structure of rings* (Providence, 1956).
- [7] T. KANDÔ, Strong regularity of arbitrary rings, *Nagoya Math. J.*, **4** (1952), 51—53.
- [8] L. KOVÁCS, A note on regular rings, *Publ. Math. Debrecen*, **4** (1955—56), 465—468.
- [9] S. LAJOS, On regular duo rings, *Proc. Japan Acad.*, **45** (1969), 157—158.
- [10] S. LAJOS, Characterization of regular duo rings, to appear.
- [11] S. LAJOS, On semilattices of groups, *Proc. Japan Acad.*, **45** (1969), to appear.
- [12] J. LUH, A characterization of regular rings, *Proc. Japan Acad.*, **39** (1963), 741—742.
- [13] N. H. MCCOY, Subdirectly irreducible commutative rings, *Duke Math. J.*, **12** (1945), 381—387.
- [14] J. v. NEUMANN, On regular rings, *Proc. Nat. Acad. Sci. U.S.A.*, **22** (1936), 707—713.
- [15] J. v. NEUMANN, *Continuous geometry* (Princeton, 1937).
- [16] L. RÉDEI, *Algebra*, I (Budapest, 1967).
- [17] C. E. RICKART, *General theory of Banach algebras* (Princeton, Toronto, London, New York, 1960).

- [18] F. Szász, Über Ringe mit Minimalbedingung für Hauptideale. I, *Publ. Math. Debrecen*, **7** (1960), 54—64.
- [19] F. Szász, Über Ringe mit Minimalbedingung für Hauptideale. II, *Acta Math. Acad. Sci. Hung.*, **12** (1961), 417—439.
- [20] F. Szász, Über Ringe mit Minimalbedingung für Hauptideale. III, *Acta Math. Acad. Sci. Hung.*, **14** (1963), 447—461.
- [21] G. Szász, *Introduction to lattice theory* (Budapest, 1963).
- [22] G. THIERRIN, On duo rings, *Canad. Math. Bull.*, **3** (1960), 167—172.

(Received May 15, 1969)