

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24

# Is subterranean lifestyle reversible? Independent and recent large-scale dispersal into surface waters by two species of the groundwater amphipod genus *Niphargus*

Denis Copilaş-Ciocianu<sup>1\*</sup>, Cene Fišer<sup>2</sup>, Péter Borza<sup>3</sup> and Adam Petrušek<sup>1</sup>

<sup>1</sup>Charles University, Faculty of Science, Department of Ecology, Viničná 7, CZ-12844, Prague, Czech Republic

<sup>2</sup>Department of Biology, Biotechnical Faculty, University of Ljubljana, Večna pot 111, SI-1000 Ljubljana, Slovenia

<sup>3</sup>Danube Research Institute, MTA Centre for Ecological Research, Karolina út 29-31, H-1113 Budapest, Hungary

**Keywords** dispersal, ecological shift, morphology, phylogeography, reversibility, stygobitism

\*Correspondence: Denis Copilaş-Ciocianu, Charles University, Faculty of Science, Department of Ecology, Viničná 7, 12844, Prague, Czech Republic. Fax: +420 - 221 951 673. E-mail: [copilas.denis@gmail.com](mailto:copilas.denis@gmail.com).

## **Abstract**

Groundwater is an extreme environment due to its absence of light, resource scarcity and highly fragmentary nature. Successful groundwater colonizers underwent major

25 evolutionary changes and exhibit eye and pigment loss (troglomorphies). Consequently,  
26 their chances of dispersal and survival in the well-connected surface waters are greatly  
27 decreased, resulting in significant endemism. The West Palaearctic subterranean amphipod  
28 genus *Niphargus* comprises hundreds of narrowly endemic and troglomorphic species.  
29 Nevertheless, a few are known to occur in surface waters, two of which, *N. hrabei* and *N.*  
30 *valachicus*, have extremely large ranges that even exceed those of many surface-water  
31 amphipods. We studied whether this pattern results from a secondary colonization of the  
32 relatively well-connected epigeal environment, and that this ecological shift promoted the  
33 large-scale dispersal of these species. Results showed that despite their ecological and  
34 zoogeographic similarities, *N. hrabei* and *N. valachicus* are not closely related and  
35 independently colonized surface waters. Their phylogeographic patterns indicate Middle to  
36 Late Pleistocene dispersal episodes throughout the Danube lowlands, and relatively modest  
37 yet significant genetic differentiation among populations. Clustering based on morphology  
38 revealed that the two species are phenotypically closer to each other than they are to most  
39 other epigeal congeners. We presume that the ecological shift to surface environments was  
40 facilitated by their ability to thrive in hypoxic waters where rheophilic competitors from  
41 the family Gammaridae cannot survive. In conclusion, our results indicate that adaptation  
42 to groundwater is not a one-way evolutionary path and that troglomorphic species can  
43 occasionally recolonize and widely disperse in surface waters.

44

## 45 **1. Introduction**

46 Groundwater macrofaunal species represent a substantial part of freshwater diversity in  
47 Europe (Zagmajster et al., 2014). Among the most remarkable features of groundwater  
48 fauna is high endemism (Trontelj et al., 2009; Eme et al., 2017). The key mechanism  
49 underlying narrow endemism is weak dispersal, presumably reflecting the physical and

50 ecological properties of groundwater habitats that are fragmented and poorly connected  
51 (Strayer, 1994; Lefébure et al., 2006, 2007; Eme et al., 2013). There are only a few widely  
52 distributed groundwater taxa, usually meiofaunal species (< 1 mm), living in better  
53 connected environments such as hyporheic alluvial habitats along rivers (Ward and Palmer,  
54 1994). Furthermore, groundwater is a challenging environment due to its permanent  
55 darkness and resource scarcity (Gilbert et al., 1994; Hüppop, 2000). Consequently,  
56 groundwater species display a suite of convergent adaptations (troglo-morphies) such as eye  
57 loss, depigmentation, body and appendage elongation, low metabolic rates, and resistance  
58 to hypoxia (Malard and Hervant, 1999; Hüppop, 2000). The apparent cost of their  
59 specialization is a lower ability to cope with the ecological conditions of photic  
60 environments. These are stressful for groundwater inhabitants, due to, e.g., damage from  
61 ultraviolet light because of depigmentation (Ginet, 1960; Maguire 1960; Langecker, 2000)  
62 or strong interspecific competition from well adapted and more prolific surface-water  
63 relatives (Fišer et al., 2007; Sket, 2008; Luštrik et al., 2011). Therefore, subterranean  
64 species seem to be restricted to the fragmented subsurface, their dispersal through the  
65 better connected surface waters is limited, and species ranges greater than 200 km are  
66 exceptional (Trontelj et al., 2009). Thus, these patterns from groundwater are an excellent  
67 case illustrating how evolutionary processes can shape macroecological patterns (discussed  
68 by Weber et al., 2017).

69 *Niphargus* is the most diverse genus of freshwater amphipods, comprising over 400  
70 species distributed in West Palaearctic groundwaters (Väinölä et al., 2008; Esmaili-Rineh  
71 et al., 2015; Horton et al., 2017). *Niphargus* species are ecologically diverse, inhabiting  
72 almost all types of aquatic subterranean habitats, from deep cave lakes to small pores in the  
73 epikarst (Fišer, 2012; Fišer et al., 2006, 2014). Several species are not strictly limited to  
74 groundwater and, in addition to permanent subterranean populations they also have stable

75 populations in surface or ecotonal habitats (Karaman, 1977; Fišer et al., 2006, 2014;  
76 Copilaş-Ciocianu et al., 2017a). The great majority of *Niphargus* species are narrow-range  
77 endemics, most being known only from their type localities (Fišer et al., 2008; Eme et al.,  
78 2017). Genetic evidence suggests that most of the presumably widespread taxa within  
79 *Niphargus* are actually complexes of endemic cryptic species (Lefébure et al., 2006, 2007;  
80 Delić et al., 2017).

81 Two similar species from the middle and lower Danube lowlands (Southeast  
82 Europe) remarkably break this general pattern: *N. hrabei* Karaman, 1932 and *N. valachicus*  
83 Dobreanu and Manolache, 1933 (Copilaş-Ciocianu et al., 2017a). They inhabit the muddy,  
84 dimly lit and densely vegetated substrate near the shores of slow-flowing or stagnant water  
85 bodies and have rarely been reported from groundwater habitats (Mejering et al., 1995;  
86 Copilaş-Ciocianu et al., 2017a and references therein). This, coupled with the seasonal life-  
87 cycle of *N. valachicus* (Copilaş-Ciocianu and Boroş, 2016), implies they are more tied to  
88 the surface than to the subterranean environment. The two species have the widest known  
89 geographical ranges of any niphargid (>1300 km), rivalling those of many epigean  
90 amphipods (Neseman et al., 1995; Borza et al., 2015; Copilaş-Ciocianu et al., 2017a). They  
91 exhibit a shallow genetic divergence among distant populations (Copilaş-Ciocianu et al.,  
92 2017a) in comparison with other amphipods from the same region (Meleg et al., 2013;  
93 Copilaş-Ciocianu and Petrussek 2015, 2017), suggesting relatively efficient dispersal in the  
94 recent past, though details about their phylogeographic histories and dispersal mechanism  
95 are unknown.

96 Based on previous phylogenetic studies (e.g. McInerney et al., 2014; Esmaeili-  
97 Rineh et al., 2015; Delić et al., 2016), it appears that surface-water affinity in *Niphargus*  
98 might not be the ancestral condition, indicating that surface dwelling species could be  
99 derived from subterranean ancestors. However, this assumption has neither been postulated

100 nor tested. The epigean lifestyle of *N. hrabei* and *N. valachicus* coupled with their  
101 troglomorphic phenotypes further point out that they secondarily colonized surface-waters.  
102 This secondary transition to ecotonal / epigean habitats might explain the large geographic  
103 ranges of both species. Therefore, the first aim of our study was to investigate if indeed  
104 surface-water *Niphargus* species are derived from groundwater ancestors and if so, to infer  
105 how many surface colonization events have occurred during the evolutionary history of the  
106 genus. Second, we examined the phylogeographic consequences of surface colonization by  
107 inferring the spatio-temporal dispersal patterns of *N. hrabei* and *N. valachicus*. Finally, we  
108 tested whether the similar ecology of these species is reflected in their morphological  
109 similarity.

110

## 111 **2. Material and methods**

### 112 *2.1. Sampling, laboratory protocols, sequence alignment and assembly of datasets*

113 Specimens were collected throughout the distribution range of both species between 2009  
114 and 2016 by sweeping a hand net through the dense riparian vegetation of various water  
115 bodies. After collection, animals were fixed in 95% ethanol. Depending on sample size,  
116 between one and six individuals per sampling locality were molecularly analysed. A total  
117 of 19 and 38 localities for *N. hrabei* (54 individuals) and *N. valachicus* (111 individuals),  
118 respectively, were investigated (Fig. 1, Table S1).

119 Genomic DNA was extracted using the Genomic DNA Mini Kit for tissue (Geneaid  
120 Biotech Ltd, Taipei). For phylogeographic purposes we used fast evolving mitochondrial  
121 and nuclear markers as they can provide phylogenetic resolution at fine spatio-temporal  
122 scales. As such, we sequenced a part of the mitochondrial gene for cytochrome c oxidase  
123 subunit I (COI) and a substantial fraction of the nuclear internal transcribed spacer (ITS1,  
124 5.8S rRNA and ITS2). These markers proved useful in a preliminary study of the genetic

125 variation in the two focal species (Copilaş-Ciocianu et al., 2017a). Amplification of the  
126 COI and ITS fragments followed protocols of Copilaş-Ciocianu et al. (2017a) and Flot et  
127 al. (2010a), respectively. For phylogenetic purposes we additionally sequenced two parts  
128 of the large ribosomal subunit (28S) and the histone H3 gene (H3), following the protocols  
129 in Fišer et al. (2013). These nuclear markers are more conserved and provided sufficient  
130 resolution for uncovering the niphargid phylogenetic relationships (e.g. Trontelj et al.  
131 2012, Fišer et al., 2013). Details about primers are provided in Table S3. The length of  
132 amplified fragments, and numbers of variable and parsimony informative sites are  
133 presented in Table S4.

134         The protein-coding COI and H3 sequences were aligned with MUSCLE (Edgar,  
135 2004) in MEGA 6 (Tamura et al., 2013) and checked for possible evidence of pseudogenes  
136 (i.e., presence of stop codons or reading frame shifts) by subsequent amino acid  
137 translation. The ITS and 28S fragments were aligned with MAFFT (Katoh and Standley,  
138 2013) with the Q-INS-i option (Katoh and Toh, 2008). Indels and regions of questionable  
139 homology in the 28S marker were identified and removed with GBLOCKS 0.9 (Talavera  
140 and Castresana, 2007). Double peaks in the ITS chromatograms (indicating heterozygosity  
141 or multiple gene copies) were coded according to the IUPAC ambiguity codes and  
142 haplotypes were phased with SeqPHASE (Flot, 2010) and PHASE (Stephens et al., 2001).  
143 Contigs were assembled using DNA Baser 4 (Heracle BioSoft 2013; www.DnaBaser.com).

144         For the analyses of phylogeography, we complemented the newly obtained dataset  
145 of both species (41 and 100 individuals of *N. hrabei* and *N. valachicus*, respectively) with  
146 additional COI and ITS data from previous studies (Flot et al., 2014; Copilaş-Ciocianu et  
147 al., 2017a) (see Table S1). For phylogenetic analyses, we gathered a large dataset  
148 comprising 157 ingroup taxa from 21 previous studies (see Table S2). We used the family

149 Pseudoniphargidae as an outgroup since it is a sister clade to niphargids (Jurado-Rivera et  
150 al., unpublished). The concatenated supermatrix contained 28% missing data.

151

## 152 *2.2. Phylogeny, topology tests and ancestral state reconstruction*

153 Phylogenetic analyses were carried out to investigate the phylogenetic position of  
154 *N. hrabei* and *N. valachicus* within the genus and to infer whether their presence in surface  
155 waters is due to a secondary colonization from subterranean habitats. Potential loss of  
156 phylogenetic signal due to substitution saturation at the COI marker was inspected using  
157 the test of Xia et al. (2003) implemented in DAMBE 5.3 (Xia and Xie, 2003).

158 PartitionFinder 1.1.1 (Lanfear et al., 2012) was used to determine the best fitting  
159 evolutionary models and partitioning schemes by employing the greedy algorithm and the  
160 Bayesian information criterion. Models and partitions are shown in Table S4.

161 We used Bayesian inference (BI) and maximum-likelihood (ML) approaches to  
162 reconstruct phylogenetic relationships within *Niphargus* using the concatenated  
163 supermatrix approach in BEAST 1.8.0 (Drummond et al., 2012) and RAxML-HPC 8.2.9  
164 (Stamatakis, 2014). For the BEAST analysis we used the initial alignment from which  
165 poorly alignable regions in the 28S marker were removed (see previous section). For  
166 RAxML we used an alignment which kept these regions and was produced with SATé  
167 2.2.7 (Liu et al., 2009). Further details on analysis settings and evolutionary models are  
168 provided in Supplementary Information.

169 To test whether the two focal taxa that show many ecological, morphological and  
170 biogeographic similarities are sister species that represent a single surface water  
171 colonization event, an alternative topology where they were constrained to monophyly was  
172 compared with the unconstrained phylogeny by applying the Shimodaira-Hasegawa (SH;

173 Shimodaira and Hasegawa, 1999) and the approximately unbiased tests (AU; Shimodaira,  
174 2002) implemented in the software Treefinder (Jobb, 2011).

175 We evaluated if, and how many times, groundwater *Niphargus* species secondarily  
176 colonized the epigeal / shallow subterranean habitat by mapping species habitat onto a  
177 phylogenetic tree. The habitat preferences were inferred from available publications and  
178 unpublished field data; we treated them as a binary character, simplified to “subterranean”  
179 and “surface”. The state “subterranean” refers to an exclusively stygobiotic lifestyle, while  
180 “surface” encompasses a broad array of habitats at the boundary between the surface and  
181 subterranean environments, including roots of submerged plants, forest ditches, *Sphagnum*  
182 moss, springs, and shallow subterranean habitats (hypotelminorheic). Ancestral states were  
183 inferred with likelihood and Bayesian methods using 1000 post burn-in trees from the  
184 BEAST analysis to account for phylogenetic uncertainty. Likelihood mapping using  
185 Markov k-state 1 parameter model was performed in Mesquite 3.04 (Maddison and  
186 Maddison, 2015). The character state at the root was estimated from the model (Mesquite  
187 default setting) and not constrained as in the original method of likelihood reconstruction  
188 (Schluter et al., 1997). Bayesian mapping was performed using Bayes Traits v. 3 (Meade  
189 and Pagel, 2016). Priors for evolutionary rates from “surface” to “subterranean” and from  
190 “subterranean” to “surface” were drawn from a uniform distribution between 0 and 100.  
191 The selection of the optimal model of the evolutionary rates was made by comparison of  
192 Bayes factors (function *stepping stone*). The best performing models allowed for traits to  
193 vary their rate of evolution within and between branches (function *covarion*, adds one  
194 additional parameter to the model) (Table S5). Among these models, there was no  
195 significant difference in marginal likelihoods if the transition rates from “surface” to  
196 “subterranean” and *vice versa* are identical or not. For this reason, we selected a simpler  
197 model (function *restrict*, one parameter less, i.e., forward evolutionary rate is equal to

198 backward rate); the final model had two parameters (a single evolutionary rate modified by  
199 covarion). We ran 1 010 000 iterations, which were sampled every 1000<sup>th</sup> generation with  
200 burn-in of 10 000. The analyses were repeated four times to check for the consistency of  
201 the results. Acceptance rate between 20-40% was achieved by an automatic tuning method  
202 implemented in Bayes Traits v.3 as a default.

203

### 204 2.3. Phylogeographic analyses

205 The number of haplotypes (H), segregating sites (S), haplotype (Hd) and nucleotide  
206 diversity ( $\pi$ ) and mean number of pairwise nucleotide differences (K) were calculated for  
207 *N. hrabei* and *N. valachicus* and their intraspecific clades (see Results) with DnaSP 5.1  
208 (Librado and Rozas, 2009) using both the COI and ITS markers.

209 In order to explore intraspecific patterns of genetic diversity, we constructed  
210 haplotype networks and time-calibrated phylogenetic trees. The models and partitions were  
211 estimated with PartitionFinder 1.1.1 and are shown in Table S4. Haplotype networks were  
212 constructed for both COI and phased ITS sequences with Haploviewer (Salzburger et al.,  
213 2011); maximum likelihood trees inferred with MEGA 6 with the models presented in  
214 Table S4 were used as input. Unique haplotypes were selected with the online tool FaBox  
215 (Villesen, 2007; <http://www.birc.au.dk/software/fabox>) and were used to build time-  
216 calibrated COI gene trees for both species using BEAST 1.8.0. The best fitting coalescent  
217 (constant size, logistic, expansion and exponential growth) and clock models (strict and  
218 relaxed) were selected using the modified Akaike information criterion (AICM) with  
219 moment's estimator (Baele et al., 2012) in TRACER 1.6 (Drummond and Rambaut, 2007)  
220 with 1000 bootstrap replicates. The best clock and coalescent models are shown in Table  
221 S6. The MCMC chain was run for 20 million generations and sampled every 1000

222 generations. Effective sample size of parameters was checked using TRACER 1.6; values  
223 of at least 200 were considered appropriate. The first 20% of trees were discarded as burn-  
224 in and the maximum clade credibility tree was built using TreeAnnotator 1.8.0 (Drummond  
225 and Rambaut, 2007).

226         The main goal of the dating analysis was to provide an approximate estimation of  
227 the time frame of dispersal (i.e. Pleistocene vs. Pliocene or Miocene) and not to pinpoint  
228 specific historical factors responsible for the dispersal events. Because the rates of  
229 molecular evolution seem to be time-dependent, intraspecific rates can evolve faster than  
230 interspecific ones (Ho et al., 2005, 2011; but see Emerson and Hickerson, 2015). This  
231 phenomenon has been observed in malacostracans as well, where the intraspecific clock  
232 rates vary from 6.58% Ma<sup>-1</sup> (mantis shrimp *Haptosquilla oulchella*, Crandall et al., 2012)  
233 to very fast post-glacial rates of up to 27% Ma<sup>-1</sup> (mysid *Mysis salemaai*, Audzijonyte and  
234 Väinölä, 2006). However, because we do not have any strong *a priori* assumption of fast  
235 post-glacial rates, we prefer the 6.58% Ma<sup>-1</sup> rate as it is intermediary between the  
236 interspecific rate of 2.3% Ma<sup>-1</sup>, commonly used in dating amphipod divergence (e.g.  
237 Lefébure et al., 2006; Copilaş-Ciocianu and Petrussek 2015), and the other extreme of very  
238 fast intraspecific post-glacial rate of 27% Ma<sup>-1</sup>. Furthermore, the magnitude of divergence  
239 and the time scale of our study are also intermediary (see Results).

240         The following landscape genetic analyses were performed on the COI marker  
241 because it had a greater variability than ITS and sufficient sample size. To test for a pattern  
242 of isolation by distance (IBD), we performed a Mantel test in the software Alleles In Space  
243 (Miller 2005) using pairwise p-distance values as a measure of genetic distance and 1000  
244 replicates. The genetic population structure was examined using an analysis of molecular  
245 variance (AMOVA; Excoffier et al., 1992) in ARLEQUIN 3.5 (Excoffier and Lischer,  
246 2010) by grouping the sampling sites according to the main river drainages. Tests were

247 performed with Tamura-Nei distances and significance was assessed using 10 000  
248 permutations. Population differentiation among drainages was investigated using pairwise  
249 fixation indices ( $\Phi_{ST}$ ; Weir and Cockerham, 1984) computed with the Tamura-Nei model  
250 in ARLEQUIN. Significance levels were assessed using 10 000 permutations. Correction  
251 for multiple testing was performed with the Benjamini-Hochberg procedure (Benjamini  
252 and Hochberg, 1995) in the software SGoF+ (Carvajal-Rodriguez and de Uña-Alvarez,  
253 2011). In the case of *N. hrabei*, four individuals were removed from the AMOVA and  $\Phi_{ST}$   
254 analyses as they were the only representatives of their respective drainages.

255         The historical demographic patterns of both species were explored using three  
256 approaches based on the COI data. First, to examine if a demographic expansion took place  
257 recently, we employed three neutrality tests: Tajima's  $D$  (Tajima, 1989) and Fu's  $F_s$  (Fu,  
258 1997), implemented in ARLEQUIN, and  $R_2$  (Ramos-Onsins and Rozas, 2002) in DnaSP  
259 5.1. Their statistical significance was evaluated using 10 000 simulated samples. As a  
260 second test for demographic expansion, we calculated mismatch distributions (Rogers and  
261 Harpending, 1992) under a sudden-expansion model in ARLEQUIN with 1000 bootstrap  
262 replicates. The validity of this model was assessed using the sum of squared deviations  
263 (SSD) and Harpending's raggedness statistic (Hri; Harpending, 1994). Third, Bayesian  
264 skyline plot analyses (BSP; Drummond et al., 2005) were employed in BEAST 1.8.0 in  
265 order to visualize demographic changes through time. The same clock rate and  
266 evolutionary models were used as for the time calibrated phylogenetic analyses (see  
267 above). The MCMC chain was run for 50 million generations, sampled every 1000  
268 generations and the first 10% of trees were discarded as burn-in. The analyses were  
269 repeated three times in order to ensure convergence on the same result. Effective sample  
270 sizes were checked with TRACER 1.6.

271 To explore the origin and dispersal history of each species while accounting for  
272 phylogenetic uncertainty, we used continuous Bayesian phylogeographic diffusion models  
273 (Lemey et al., 2010) implemented in BEAST 1.8.0. The analysis was based on the COI  
274 marker and we used the same settings as for the time-calibrated phylogenetic analyses.  
275 Best fitting diffusion models are presented in Table S7 and further details are provided in  
276 the Supplementary Information.

277

#### 278 *2.4. Detection of cryptic lineages*

279 Considering the wide and fragmented range of the studied species, we used the COI  
280 and ITS markers to investigate if they are composed of independently evolving cryptic  
281 lineages. For the COI dataset (encompassing all individuals of both species) we used a  
282 Bayesian implementation of the Poisson tree process on the bPTP server ([http://species.h-](http://species.h-its.org/ptp)  
283 [its.org/ptp](http://species.h-its.org/ptp)) (Zhang et al., 2013). The input phylogenetic tree was generated with PhyML  
284 3.0 (Guindon et al., 2010) using the evolutionary models in Table S4. We ran the analysis  
285 for 300 000 MCMC iterations with a thinning of 100 and 20% burn-in. Due to the fact that  
286 putative heterozygous individuals were present in the ITS dataset (see Results), potential  
287 cryptic lineages were investigated using Doyle's concept of species as fields for  
288 recombination (FFRs; Doyle 1995), i.e. assuming that species are characterized by mutual  
289 allelic exclusivity. For this purpose, we transformed the haplotype network of phased ITS  
290 sequences (obtained as indicated above) into a haploweb by connecting the haplotypes  
291 which were co-occurring in heterozygous individuals (Flot et al., 2010b). Interconnection  
292 among all haplotypes indicates a common gene pool, and therefore conspecificity under  
293 the FFR concept, while groups of haplotypes which are not interconnected might indicate  
294 cryptic species or cessation of gene-flow among geographically isolated populations.

295

## 296 2.5. Morphological analyses

297 In order to test the hypothesis that similar ecology leads to morphological similarity of the  
298 two focal species, we analyzed a dataset of 81 *Niphargus* species, of which 15 live in  
299 similar habitats as the two focal ones (Tables S2, S8). We measured between 1 and 10  
300 individuals per species. For the analyses we used only adult males; females were  
301 considered only in those species that show no sexual dimorphism and only when males  
302 were not available. We analyzed 35 traits describing in detail the body shape and size,  
303 appendage length, and spine patterns (Fišer et al., 2009) (Table S8). We calculated mean  
304 values for the traits for the species and prior to the analysis transformed the data as  
305 follows: 1) the number of spines and the body lengths were log-transformed, and 2) in  
306 order to remove the impact of body size, all length-measures were regressed onto body  
307 lengths, and standardized residuals were calculated. We inferred morphological similarity  
308 from cluster analyses using squared Euclidean distances and Ward's agglomeration  
309 method. All analyses were run using SPSS ver. 20 (IBM Corp 2011).

310

## 311 3. Results

### 312 3.1. Phylogeny, topology tests and ancestral state reconstruction

313 The substitution saturation test indicated no significant saturation at the COI marker  
314 ( $p < 0.0001$ ). Both ML and BI analyses on the concatenated dataset revealed similar results.  
315 The removal or retention of poorly alignable regions in the 28S marker did not influence  
316 the overall phylogenetic pattern (Fig. S1). The topology of the phylogeny is largely  
317 congruent with the recent phylogenetic reconstructions of the genus by recovering the  
318 same major clades (e.g. McInerney et al., 2014; Esmaili-Rineh et al., 2015; Delić et al.,  
319 2016). All the species inhabiting surface-water habitats, including *N. hrabei* and *N.*  
320 *valachicus*, were recovered in the same major clade; however, the two focal species are not

321 in a sister relationship (Figs 1, S1). The position of *N. valachicus* is recovered at the base  
322 of a clade that mostly contains surface-water species, though the support for the clade is  
323 not high in the ML analysis. In contrast, *N. hrabei* is clustered together with strong support  
324 with two subterranean species, *N. plateaui* and *N. puteanus* (Figs 1, S1). The sister relation  
325 of both species (monophyly) was rejected by both the SH and AU tests at  $p = 0.036$  and  $p$   
326  $= 0.023$ , respectively (likelihood unconstrained = -60408.49, AICc = 121554; likelihood  
327 constrained = -60430.26, AICc = 121598).

328 All ancestral state reconstruction methods support the hypothesis that the ancestor  
329 of Niphargidae and Pseudoniphargidae was a subterranean species (Fig. 1, Table S9). Both  
330 families apparently diversified in the subterranean environment. Both likelihood and  
331 Bayesian mapping indicate that the probability of ancestors having lived in a subterranean  
332 environment is above 0.95 across all basal splits. Bayesian mapping suggests that the  
333 surface and/or shallow subterranean environments were colonized at least four times  
334 independently (Fig. 1; nodes 7, 9, 11 and 13), when the probability for a subterranean  
335 ancestor abruptly fell below 0.58. However, the likelihood analysis was less conservative  
336 and indicated that ecological change took place along terminal branches, i.e. there might  
337 have been even more transitions to surface / shallow subterranean environments (Table  
338 S9). In the case of *N. hrabei*, the transition occurred along the terminal branch of the tree,  
339 while in the case of *N. valachicus*, the transition probably took place in the common  
340 ancestor of clade 11. In any case, these results coupled with the topology of the tree  
341 indicate that both focal species colonized surface habitats independently from each other.

342

### 343 3.2. Phylogeography

344 The COI marker was represented by 12 and 42 haplotypes in *N. hrabei* and *N. valachicus*,  
345 respectively, and the ITS marker by 6 and 23 haplotypes (Table 1). Eight out of the 22

346 analysed individuals (36%) were heterozygous at the ITS locus in *N. hrabei*, and 26 out of  
347 40 individuals (65%) were heterozygous in *N. valachicus* (Table 1). The geographical  
348 distribution of haplotypes is shown in Fig. 2. Haplotype and nucleotide diversity, and mean  
349 number of pairwise nucleotide differences were higher in *N. valachicus* than in *N. hrabei* at  
350 both markers (Table 1).

351 The haplotype networks based on COI indicate that both species have a  
352 geographically structured genetic variation, most haplotypes apparently being endemic in  
353 relatively narrow parts of the species' ranges (Fig 2A, B). This pattern was less  
354 pronounced at the ITS marker where a single haplotype was widespread across the entire  
355 range in each of the species (nH1 in *H. hrabei* and nV1 in *N. valachicus*). However, groups  
356 of locally restricted haplotypes could also be observed (Fig. 2C, D). The intraspecific COI  
357 time-calibrated trees (assuming the COI rate of  $6.58 \text{ Ma}^{-1}$ ) indicate that *N. hrabei* has a  
358 shorter mean coalescence time (103 ka, 95% HPD: 56–152 ka) than *N. valachicus* (353 ka,  
359 95% HPD: 214–504 ka) (Fig. 3A). No strongly supported intraspecific lineages could be  
360 observed in *N. hrabei*; however, *N. valachicus* was composed of two distinct clades: clade  
361 A distributed in the Pannonian lowlands and clade B distributed in the SE parts of the  
362 Pannonian lowlands (where it partly overlaps with clade A), Wallachian Plain, Danube  
363 Delta and northern Turkey (Fig 1B).

364 The IBD test revealed a highly significant relationship between geographic and  
365 genetic distances in both species (*N. hrabei*,  $r^2=0.43$ ; *N. valachicus*  $r^2=0.37$ ;  $p<0.0001$  in  
366 both species), indicating that dispersal is limited (Fig. S2). The AMOVA analysis indicated  
367 that most of the observed variation is explained by differences among drainages (*N. hrabei*:  
368 80.05%,  $\Phi_{CT} = 0.80$ ,  $p=0.007$ ; *N. valachicus*: 58.5%,  $\Phi_{CT} = 0.58$ ,  $p<0.0001$ ) (Table 2).  
369 Drainages are listed in Tables S10 and S11. Pairwise  $\Phi_{ST}$  distances indicated a significant  
370 genetic differentiation at COI among most of the main drainages, with 90% and 82% of all

371 pairwise comparisons being statistically significant in *N. hrabei* and *N. valachicus*,  
372 respectively (Tables S10 and S11).

373 The three demographic tests (Tajima's  $D$ , Fu's  $F_s$  and  $R_2$ ) were applied to each  
374 species and to each of the two clades of *N. valachicus*. In the case of *N. hrabei*, only Fu's  
375  $F_s$  was statistically negatively significant ( $p = 0.031$ ), however, the other two tests were  
376 close to the significance threshold (Tajima's  $D$   $p = 0.056$ ;  $R_2$   $p = 0.083$ ), indicating recent  
377 population expansion (Table 1). In *N. valachicus* overall and in its clade A none of the tests  
378 were significant, suggesting a stable populations size, but population expansion was  
379 indicated in clade B as Fu's  $F_s$  was significantly negative ( $p < 0.0001$ ), while the other two  
380 tests were close to statistical significance (Tajima's  $D$   $p = 0.061$ ;  $R_2$   $p = 0.06$ ) (Table 1).  
381 The mismatch distribution of haplotype pairwise differences was unimodal in *N. hrabei*  
382 and the SSD and Hri tests indicate no significant departure from the assumption of rapid  
383 population expansion (Fig. 3B, Table 1). In the case of *N. valachicus*, the overall mismatch  
384 distribution was bimodal with the SSD and Hri having no statistically significant values  
385 (Fig. 3B, Table 1). Clade A had a multimodal mismatch distribution while clade B was  
386 unimodal (Fig. 3B). The SSD and Hri tests were not significant in the former, but the Hri  
387 was significant in the latter (Table 1).

388 Taken together the evidence points to a relatively recent and rapid population  
389 expansion in *N. hrabei* and *N. valachicus* clade B and a stable population size in clade A.  
390 The BSP indicates a population growth in *N. hrabei* during the last 50 ka with an increase  
391 in the last 10-15 ka (Fig. 3B). In *N. valachicus* a population growth took place ca. 100 ka,  
392 followed by a decline during the Last Glacial Maximum (ca. 25 ka) and another growth  
393 during the last 5-10 ka (Fig. 3B).

394 The different Bayesian phylogeographic diffusion models produced congruent  
395 results regarding the presumed past dispersal patterns. The analyses suggest the origin of

396 *N. hrabei*'s dispersal is in the lower Danube lowlands, ca. 90 ka (Wallachian Plain, S  
397 Romania). From there, it has subsequently spread to the west and east multiple times along  
398 the Danube River and only recently, likely postglacially, arrived to the upper parts of the  
399 river basin (Fig. 4A). In contrast, the dispersal of *N. valachicus* apparently started earlier,  
400 in the southwest of the Pannonian lowlands in northern Croatia, ca. 350 ka. It spread to the  
401 east, along the Sava and Drava rivers, reaching the Danube and the Wallachian Plain  
402 somewhere between 100 and 200 ka. It reached northern Turkey and the north of the  
403 Pannonian Plain only in the last 100 ka, possibly postglacially (Fig. 4B). Although we  
404 acknowledge a substantial uncertainty of the absolute values, the results based on a  
405 conservative mutation rate strongly suggest that dispersal of these species is relatively  
406 recent (the second half of the Pleistocene) and that *N. valachicus* has occupied the Danube  
407 lowlands before *N. hrabei*.

408

### 409 *3.3. Absence of cryptic lineages*

410 Despite the fact that both species have wide and fragmented ranges, the COI and ITS data  
411 do not indicate the existence of reproductively separated lineages. The bPTP analysis  
412 based on COI supports the conspecificity of all analysed populations in both species  
413 (posterior probability of 0.5 and 0.8 in *N. hrabei* and *N. valachicus*, respectively) (Fig.  
414 S3A). Furthermore, the haploweb analysis based on ITS indicates that many of the  
415 analysed individuals share the same haplotype (nH1 in *H. hrabei* and nV1 in *N.*  
416 *valachicus*) and all haplotypes co-occurring in heterozygous individuals are interconnected  
417 (Fig. S3B), indicating a common gene pool.

418

### 419 *3.4. Morphological analyses*

420 The morphological analysis split all 81 analyzed species into two major clusters (Fig. 5).  
421 One cluster comprises species living in the stagnant water of permanently flooded parts of  
422 karstic massifs while the second includes species found in flowing water, interstitial and  
423 surface habitats. The first cluster is further split into three subclusters corresponding to  
424 lake, lake giant and daddy-longlegs ecomorphs (Trontelj et al., 2012). The second cluster  
425 has a more complex structure and its two principal subclades are split even further. Species  
426 occurring in surface habitats were clustered into four separate groups. *N. hrabei* and *N.*  
427 *valachicus* along with *N. elegans* from northern Italy were clustered together with small  
428 bodied, interstitial and epikarst taxa (small pore ecomorph *sensu* Trontelj et al., 2012).  
429 Apparently, they are not very similar to the other species occurring in surface habitats that  
430 were rather clustered together with cave stream species (cave stream ecomorph *sensu*  
431 Trontelj et al., 2012) (Fig. 5).

432

#### 433 **4. Discussion**

434 Our results indicate that, despite their ecological and morphological similarities, *N. hrabei*  
435 and *N. valachicus* are not closely related. Their phylogenetic position and reconstructed  
436 ancestral states imply independent colonization of surface waters from subterranean  
437 ancestors. Their phylogeographies reveal large-scale dispersal across the Danube lowlands  
438 throughout the Pleistocene, and their morphologies are more similar to each other than to  
439 most other congeners known to occur in surface-water habitats. These results indicate that  
440 the habitat shift from ground- to surface waters enabled the dispersal and range expansion  
441 of these species. Below we discuss the possible factors that have facilitated this ecological  
442 shift and examine the biogeographical histories of these two species.

443

##### 444 *4.1. Reversal to surface waters*

445 It appears that the ecological barrier between subsurface and surface is weaker than  
446 previously thought for troglomorphic species. Reversal to surface habitats from  
447 subterranean ancestors has been proposed so far in typhlochactid scorpions (Prendini et al.,  
448 2010) and in phalangopsid crickets (Desutter-Grandcolas, 1997), but both studies relied on  
449 morphology only, and did not test this phenomenon on molecular phylogenies. To our  
450 knowledge, our study is the first to use a molecular phylogeny to test whether  
451 troglomorphic subterranean species can recolonize surface habitats. In the case of  
452 *Niphargus*, all of the surface-water species belong to a large, morphologically and  
453 ecologically diverse clade.

454         Although most of the surface-water *Niphargus* species have known populations  
455 both in subterranean and surface / ecotonal habitats (Fišer C. et al., 2006; 2010b; 2014;  
456 Fišer Ž. et al., 2015), *N. hrabei* and *N. valachicus* are probably the most detached from the  
457 subterranean environment as they have much larger ranges and a far greater number of  
458 documented occurrences in surface versus groundwater (Copilaș-Ciocianu et al., 2017a).  
459 Therefore, it appears that there were at least two independent colonization events of truly  
460 surface-waters and multiple shifts to ecotonal habitats from subterranean ancestors during  
461 the evolutionary history of *Niphargus*.

462         According to the molecular dating analysis of McInerney et al. (2014), the clade  
463 that contains the surface-water species has radiated during the Late Eocene. This is in  
464 accordance with the amber fossil evidence which indicates that some species of *Niphargus*  
465 were already living in surface freshwaters during this time period (Coleman and Myers,  
466 2000; Jazdzewski and Kupryjanowicz, 2010). Thus, the evidence indicates that there were  
467 recurrent colonization events of surface waters during the evolutionary history of the  
468 genus. Moreover, we cannot rule out a scenario that the ancestors of some clades for a

469 certain period lived and dispersed in ecotonal habitats, leading to a secondary colonization  
470 of subterranean waters.

471 The occurrence of a high number of troglomorphic groundwater species has been  
472 long documented at the surface/subsurface boundary (shallow subterranean habitats;  
473 Culver and Pipan, 2014). It is generally thought that they are ancestral forms which  
474 represent the initial stages of colonization of the deeper subterranean realm (Culver and  
475 Pipan, 2009, 2014). However, our results add to the growing body of evidence which  
476 indicates that the opposite is also possible. Nevertheless, the factors that promoted the  
477 ecological shift to shallow subterranean and eventually surface habitats are unknown. The  
478 persistence of troglomorphic species in ecotonal habitats is probably determined by the  
479 greater availability of nutritional resources (assumed by Sket, 2008; Culver and Pipan,  
480 2009, 2014; Fišer et al., 2010a), but their presence in epigeal habitats can be realized in  
481 rare circumstances in which competitors are absent or scarce (Humphreys, 2000; Prendini  
482 et al., 2010). Indeed, the distribution of troglomorphic species in surface environments  
483 seems to be limited by competition, predation and risk of ultraviolet radiation exposure due  
484 to irreversible adaptations such as lack of eyes and pigment (reviewed in Fišer C. et al.,  
485 2014; Fišer Ž. et al., 2016).

486 *N. hrabei* and *N. valachicus* are bound to the muddy, dimly lit and densely  
487 vegetated bottom of stagnant or slow flowing lowland waters (Copilaș-Ciocianu et al.,  
488 2017a and references therein). These habitats are characterized by eutrophic conditions,  
489 high temperature and frequent hypoxia (Junk et al., 1989; Parr and Mason 2004; Graeber et  
490 al., 2013) and are avoided by the presumably ecologically closest competitors, the  
491 oxyphilic and mostly rheophilous gammarid amphipods (Mejering, 1991; Mejering et al.,  
492 1995; Henry and Danielopol, 1999; Copilaș-Ciocianu et al., 2014; Copilaș-Ciocianu and  
493 Boroș, 2016; Borza et al., 2017; Mauchart et al., 2017). Experimental evidence has shown

494 that niphargids can tolerate hypoxia and other adverse environmental conditions much  
495 better than surface-water gammarids (Danielopol et al., 1994; Hervant et al., 1995; Malard  
496 and Hervant, 1999; Coppellotti Krupa and Guidolin, 2003; Simčič et al., 2005, 2006; Flot  
497 et al., 2014). Therefore, it seems that *N. hrabei* and *N. valachicus* might have a competitive  
498 edge over gammarids in these habitats; however, experimental proof is needed in this  
499 particular case.

500 Furthermore, it appears that gammarids of the genus *Gammarus* survived  
501 Pleistocene glaciation episodes in montane refugia from where only a few species  
502 regionally dispersed to lower elevations (Copilaş-Ciocianu and Petrussek 2015, 2017;  
503 Copilaş-Ciocianu et al., 2017b). *G. roeselii* and some invasive Ponto-Caspian species,  
504 which are common in the Danube lowlands, have dispersed only postglacially or even in  
505 historical times from the Balkans or the Black Sea (Barnard and Barnard, 1983; de Vaate et  
506 al., 2002; Cristescu et al., 2004; Rewicz et al., 2015). We therefore hypothesise that the  
507 colonization of surface waters was possible because of the general and historical scarcity  
508 or even absence of gammarid competitors in the habitats in which *N. hrabei* and *N.*  
509 *valachicus* thrive.

510

#### 511 *4.2. Morphological evolution*

512 The similarity of species living at the boundary between the surface and subterranean  
513 ecosystems was noted earlier and some authors even considered them as members of own  
514 subgenus or species group (Karaman 1950; Sket 1958, Straškraba 1972a). Our results  
515 imply that this similarity is of convergent origin due to multiple independent colonization  
516 events of the epigeal / ecotonal environment. However, this ecological shift does not  
517 always lead to the same phenotype: secondary colonizers of the surface-subterranean  
518 boundary are grouped into two morphological clusters. Interestingly, the two focal species

519 cluster together with *N. elegans*, another nearly entirely surface-water species with a large  
520 range, widespread across the northern Italian lowlands (Karaman 1977). The astonishing  
521 similarities in their morphology, ecology, and range-size deserve further attention. A key  
522 question arising from this pattern is whether the large range size is a result of an enhanced  
523 possibility of passive dispersal connected with transition to better-connected habitats (see  
524 below), or whether some morphological traits may be (indirectly) linked with dispersal  
525 capacity. Thus, the functional links between morphological similarities, ecological  
526 conditions and dispersal ability remain yet to be explored.

527

#### 528 4.3. Phylogeography and dispersal

529 According to our phylogeny, *N. hrabei* is a sister species to *N. plateaui*, a taxon known  
530 from western France. Such a vast distance (ca. 1500 km) between these species might  
531 indicate a long-range dispersal in the past, or that additional related lineages remain  
532 undiscovered or went extinct. On the other hand, *N. valachicus* belongs to a clade of  
533 northern Dinaric species which are known to occur in surface-water habitats. This agrees  
534 with its SW Pannonian dispersal origin, which is in geographical proximity to its relatives.

535         If we consider the scenario of relatively conservative mutation rates, coalescence  
536 times for both species correspond with periods of warm interglacial stages. Haplotypes of  
537 *N. hrabei* coalesce ca. 100 ka (Eemian), roughly corresponding with the Marine Isotopic  
538 Stage 5d (ca. 109 ka), while coalescence time for *N. valachicus* is ca. 350 ka,  
539 corresponding with the Marine Isotopic Stage 9 (ca. 337 ka) (Lisiecki and Raymo, 2005),  
540 suggesting that their dispersal may have started during these warmer periods. In the case of  
541 *N. hrabei*, its initial Eemian expansion would coincide with the expansion of *Fagus*  
542 *sylvatica* in Central and Southern Europe (Magri et al., 2006). The presence of *N.*  
543 *valachicus* in isolated streams along the Black Sea coast is likely explained by the lower

544 water levels and freshwater conditions during the Last Glacial Maximum (Ryan et al.,  
545 1997; Bahr et al., 2006; Georgievski and Stanev, 2006). Our data reveal that *N. valachicus*  
546 has reached northern Turkey very recently, possibly during the Holocene, but definitely  
547 before the last connection with the saline Mediterranean ca. 7-9 ka (Federov, 1971; Ryan et  
548 al., 1997; Badertscher et al., 2011). During this time the shelf of the Black Sea was  
549 exposed, forming vast deltaic systems that probably facilitated coastal dispersal (Federov,  
550 1971; Ryan et al., 1997). The presence of this species along the Caspian Sea shores in Iran  
551 (Karaman, 1998) might be explained by the frequent Pleistocene connections with the  
552 Black Sea which have facilitated biotic interchange (Leonov et al., 2002; Grigorovich et  
553 al., 2003; Badertscher et al., 2011). However, it is not yet known if the Iranian populations  
554 are indeed conspecific with *N. valachicus*.

555         Although we urge caution in interpreting absolute dates, these results confidently  
556 refute previous hypotheses according to which *N. hrabei* and *N. valachicus* invaded  
557 freshwater from the brackish Paratethys Sea during the Late Miocene (Straškraba 1972b;  
558 Sket 1981) and are in accordance with the preliminary data presented in our previous study  
559 (Copilaș-Ciocianu et al., 2017a). A faster mutation rate would, of course, indicate an even  
560 more recent expansion, and even two to three times slower rate would not push  
561 coalescence times further back than the Pleistocene. The Pannonian clade of *N. valachicus*  
562 (clade A) seems to have had a stable demographic history throughout the Late Pleistocene  
563 as opposed to clade B and *N. hrabei* which exhibit substantial recent demographic growth.  
564 The stable demography of clade A is in agreement with the emerging view that the  
565 Pannonian Basin functioned as a glacial refugium for a wide array of taxa, ranging from  
566 aquatic invertebrates and vertebrates to terrestrial plants and mammals (Neumann et al.,  
567 2005; Verovnik et al., 2005; Fussi et al., 2010; Antal et al., 2016; Vörös et al., 2016).

568           The significant correlation between genetic and geographic distances and genetic  
569 differentiation among drainages indicates that despite the wide ranges of the focal  
570 *Niphargus* species, their dispersal is limited. This apparent contradiction might be  
571 explained by the fact that their dispersal is passive and happened throughout relatively long  
572 periods of time (ca. 100 ka in *N. hrabei* and 350 ka in *N. valachicus*), although in some  
573 parts of their ranges they have very likely experienced postglacial demographic expansion.  
574 This indicates that chances for dispersal are not equal in time and the highest probability  
575 for spreading could be restricted to favourable climatic periods or linked to rare long-  
576 distance dispersal events. Given that freshwater amphipods are usually poor and passive  
577 dispersers, their dispersal among different water bodies could be achieved either by animal  
578 vectors – especially waterfowl and aquatic mammals (Peck, 1975; Swanson, 1984;  
579 Rachalewski et al., 2013) – or during flooding episodes (e.g. Van Leeuwen et al., 2013). In  
580 any case, the large expanses of flat and homogeneous relief with interconnected water  
581 bodies of the Danube floodplains seem to have facilitated the range expansion of these  
582 species. The same pattern might be true for *N. elegans*, a species which is widespread  
583 throughout the floodplains of the Po River in northern Italy (Karaman, 1977) and also  
584 exhibits low intraspecific genetic divergence (Fabio Stoch, pers. comm.).

585           Considering that groundwater habitats are poorly connected in comparison to  
586 surface ones, the occurrence of at least two independent large-scale dispersal events of  
587 troglomorphic species in surface waters indicates that habitat connectivity might play a  
588 greater role in limiting dispersal than species' ecology or biology.

589

## 590 **5. Conclusion**

591           Our results strongly indicate that more than one invasion of surface waters and even  
592 more shifts to the surface/subsurface boundary from subterranean ancestors have occurred

593 during the evolutionary history of *Niphargus*. Furthermore, we show for the first time that  
594 troglomorphic species are capable of relatively rapid and large-scale dispersal in surface  
595 waters. This indicates that adaptation to groundwater might not be a one-way evolutionary  
596 path and subterranean troglomorphic species can occasionally recolonize and widely  
597 disperse in surface waters.

598

### 599 **Acknowledgements**

600 We thank Murat Sezgin, Monika Hess and Wolfram Graf for providing material, Marius G.  
601 Berchi for his help during the fieldwork, José A. Jurado-Rivera and Damià Jaume for  
602 kindly providing *Pseudoniphargus* sequences, and two anonymous reviewers for their  
603 helpful suggestions. Andrew Meade kindly changed the code in Bayes Traits such that we  
604 could reconstruct ancestral states under the *covarion* model. This study was supported by  
605 the Charles University Grant Agency, project no. 1398214, and Slovenian Research  
606 Agency, program P1-0184.

607

### 608 **References**

- 609 Antal, L., László, B., Kotlík, P., Mozsár, A., Czeglédi, I., Oldal, M., Kemenesi, G., Jakab,  
610 F., Nagy, S.A., 2016. Phylogenetic evidence for a new species of *Barbus* in the  
611 Danube River basin. *Mol. Phylogenet. Evol.* 96, 187–194. doi:  
612 10.1016/j.ympev.2015.11.023
- 613 Audzijonyte, A., Väinölä, R., 2006. Phylogeographic analyses of a circumarctic coastal  
614 and a boreal lacustrine mysid crustacean, and evidence of fast postglacial mtDNA  
615 rates. *Mol. Ecol.* 15, 3287–3301. doi: 10.1111/j.1365-294X.2006.02998.x
- 616 Baele, G., Lemey, P., Bedford, T., Rambaut, A., Suchard, M.A., Alekseyenko, A.V., 2012.  
617 Improving the accuracy of demographic and molecular clock model comparison

618 while accommodating phylogenetic uncertainty. *Mol. Biol. Evol.* 29, 2157–2167.  
619 doi: 10.1093/molbev/mss084

620 Badertscher, S., Fleitmann, D., Cheng, H., Edwards, R.L., Göktürk, O.M., Zumbühl, A.,  
621 Leuenberger, M., Tüysüz, O., 2011. Pleistocene water intrusions from the  
622 Mediterranean and Caspian seas into the Black Sea. *Nat. Geosci.* 4, 236–239.  
623 doi:10.1038/ngeo1106

624 Bahr, A., Arz, H.W., Lamy, F., Wefer, G., 2006. Late glacial to Holocene  
625 paleoenvironmental evolution of the Black Sea, reconstructed with stable oxygen  
626 isotope records obtained on ostracod shells. *Earth Planet. Sci. Lett.* 241, 863–875.  
627 doi.org/10.1016/j.epsl.2005.10.036

628 Barnard, J.L., Barnard, C.M., 1983. *Freshwater Amphipoda of the World*. Hayfield  
629 Associates, Mt. Vernon, Virginia.

630 Benjamini, Y., Hochberg, Y., 1995. Controlling the false discovery rate: a practical and  
631 powerful approach to multiple testing. *J. R. Stat. Soc.* 57, 289–300.

632 Borza, P., Huber, T., Leitner, P., Remund, N., Graf, W., 2017. Current velocity shapes co-  
633 existence patterns among invasive *Dikerogammarus* species. *Freshwater Biol.* 62,  
634 317–328. doi: 10.1111/fwb.12869

635 Borza, P., Csányi, B., Huber, T., Leitner, P., Paunović, M., Remund, N., Szekeres, J., Graf,  
636 W., 2015 Longitudinal distributional patterns of Peracarida (Crustacea,  
637 Malacostraca) in the River Danube. *Fundam. Appl. Limnol.* 187, 113–126. doi:  
638 10.1127/fal/2015/0769

639 Carvajal-Rodriguez, A., de Uña-Alvarez, J., 2011. Assessing significance in high  
640 throughput experiments by sequential goodness of fit and q-value estimation. *PLoS*  
641 *ONE*, 6, e24700. doi: 10.1371/journal.pone.0024700

642 Coleman, C.O., Myers, A.A., 2000. New Amphipoda from Baltic amber. Pol. Arch.  
643 Hydrobiol. 47, 457–464.

644 Coppellotti Krupa, O., Guidolin, L., 2003. Responses of *Niphargus montellianus* and  
645 *Gammarus balcanicus* (Crustacea, Amphipoda) from karst waters to heavy metal  
646 exposure. J. Phys. IV 107, 323–326. doi: 10.1051/jp4:20030307

647 Copilaş-Ciocianu, D., Grabowski, M., Pârvulescu, L., Petrusek, A., 2014. Zoogeography  
648 of epigean freshwater Amphipoda (Crustacea) in Romania: fragmented distributions  
649 and wide altitudinal variability. Zootaxa 3893, 243–260. doi:  
650 10.11646/zootaxa.3893.2.5

651 Copilaş-Ciocianu, D., Petrusek, A., 2015. The southwestern Carpathians as an ancient  
652 centre of diversity of freshwater gammarid amphipods: insights from the *Gammarus*  
653 *fossarum* species complex. Mol. Ecol. 24, 3980–3992. doi: 10.1111/mec.13286

654 Copilaş-Ciocianu, D., Boroş, B.V., 2016. Contrasting life history strategies in a  
655 phylogenetically diverse community of freshwater amphipods (Crustacea:  
656 Malacostraca). Zoology 119, 21–29. doi: 10.1016/j.zool.2015.11.001

657 Copilaş-Ciocianu, D., Petrusek, A., 2017. Phylogeography of a freshwater crustacean  
658 species complex reflects a long-gone archipelago. J. Biogeogr. 44, 421–432.  
659 doi:10.1111/jbi.12853

660 Copilaş-Ciocianu, D., Fišer, C., Borza, P., Balázs, G., Angyal, D., Petrusek, A., 2017a.  
661 Low intraspecific genetic divergence and weak niche differentiation despite wide  
662 ranges and extensive sympatry in two epigean *Niphargus* species (Crustacea:  
663 Amphipoda). Zool. J. Linn. Soc. doi: 10.1093/zoolinnean/zlw031

664 Copilaş-Ciocianu, D., Rutová, T., Pařil, P., Petrusek, A., 2017b. Epigean gammarids  
665 survived millions of years of severe climatic fluctuations in high latitude refugia

666 throughout the Western Carpathians. *Mol. Phylogenet. Evol.* 112, 218–229. doi:  
667 10.1016/j.ympev.2017.04.027

668 Crandall, E.D., Sbrocco, E.J., DeBoer, T.S., Barber, P.H., Carpenter, K.E., 2012.  
669 Expansion dating: calibrating molecular clocks in marine species from expansions  
670 onto the Sunda Shelf following the Last Glacial Maximum. *Mol. Biol. Evol.* 29,  
671 707–719. doi: 10.1093/molbev/msr227

672 Cristescu, M.E.A., Witt, J.D.S., Grigorovich, I.A., Hebert, P.D.N., MacIsaac, H.J., 2004.  
673 Dispersal of the Ponto-Caspian amphipod *Echinogammarus ischnus*: invasion waves  
674 from the Pleistocene to the present. *Heredity* 92, 197–203. doi:  
675 10.1038/sj.hdy.6800395

676 Culver, D.C., Pipan, T., 2009. Superficial subterranean habitats – gateway to the  
677 subterranean realm? *Cave Karst Sci.* 35, 5–12.

678 Culver, D.C., Pipan T., 2014. *Shallow Subterranean Habitats. Ecology, Evolution, and*  
679 *Conservation.* New York, Oxford University Press.

680 Danielopol, D.L., Creuzé des Châtelliers, M., Mösslacher, F., Pospisil, P., Popa, R., 1994.  
681 Adaptation of crustacea to interstitial habitats: A practical agenda for ecological  
682 studies. In: Gilbert, J., Danielopol, D.L., Stanford, J.A. (Eds.), *Groundwater Ecology.*  
683 *Academic Press, New York*, pp. 217–243.

684 Delić, T., Trontelj, P., Zakšek, V., Fišer, C., 2016. Biotic and abiotic determinants of  
685 appendage length evolution in a cave amphipod. *J. Zool.* 299, 42–50. doi:  
686 10.1111/jzo.12318

687 Delić, T., Trontelj, P., Rendoš, M., Fišer, C., 2017. The importance of naming cryptic  
688 species and the conservation of endemic subterranean amphipods. *Sci. Rep.* 7, 3391.  
689 doi:10.1038/s41598-017-02938-z

690 Desutter-Grandcolas, L., 1997. Studies in cave life evolution: a rationale for future  
691 theoretical developments using phylogenetic inference. *J. Zool. Syst. Evol. Res.* 35,  
692 23–32. doi: 10.1111/j.1439-0469.1997.tb00400.x

693 Doyle, J.J., 1995. The irrelevance of allele tree topologies for species delimitation, and a  
694 non-topological alternative. *Syst. Bot.* 20, 574–588. doi: 10.2307/2419811

695 Drummond, A.J., Rambaut, A., 2007. BEAST: Bayesian evolutionary analysis by sampling  
696 trees. *BMC Evol. Biol.* 7, 214. doi: 10.1186/1471-2148-7-214

697 Drummond, A.J., Rambaut, A., Shapiro, B., Pybus, O.G., 2005. Bayesian coalescent  
698 inference of past population dynamics from molecular sequences. *Mol. Bio. Evol.*  
699 22, 1185–1192. doi: 10.1093/molbev/msi103

700 Drummond, A.J., Suchard, M.A., Xie, D., Rambaut, A., 2012. Bayesian phylogenetics with  
701 BEAUti and the BEAST 1.7. *Mol. Biol. Evol.* 29, 1969–1973. doi:  
702 10.1093/molbev/mss075

703 Eme, D., Malard, F., Konecny-Dupré, L., Lefébure, T., Douady, C.J., 2013. Bayesian  
704 phylogeographic inferences reveal contrasting colonization dynamics among  
705 European groundwater isopods. *Mol. Ecol.* 22, 5685–5699. doi: 10.1111/mec.12520

706 Eme, D., Zagamajster, M., Delić, T., Fišer, C., Flot, J.-F., Konecny-Dupré, L., Pálsson, S.,  
707 Stoch, F., Zakšek, V., Douady, C.J., Malard, F., 2017. Do cryptic species matter in  
708 macroecology? Sequencing European groundwater crustaceans yields smaller ranges  
709 but does not challenge biodiversity determinants. *Ecography*, doi:  
710 10.1111/ecog.02683.

711 Emerson, B.C., Hickerson, M.J., 2015. Lack of support for the time-dependent molecular  
712 evolution hypothesis. *Mol. Ecol.* 24, 702–709. doi: 10.1111/mec.13070

713 Esmaeili-Rineh, S., Sari, A., Delić, T., Moškrič, A., Fišer, C., 2015. Molecular phylogeny  
714 of the subterranean genus *Niphargus* (Crustacea: Amphipoda) in the Middle East: a

715 comparison with European Niphargids. Zool. J. Linn. Soc. 175, 812–826. doi:  
716 10.1111/zoj.12296

717 Excoffier, L., Smouse, P.E., Quattro, J.M., 1992. Analysis of molecular variance inferred  
718 from metric distances among DNA haplotypes: application to human mitochondrial  
719 DNA restriction data. Genetics 131, 479–491.

720 Excoffier, L., Lischer, H.E.L., 2010. Arlequin suite ver 3.5: A new series of programs to  
721 perform population genetics analyses under Linux and Windows. Mol. Ecol. Resour.  
722 10, 564-567.doi: 10.1111/j.1755-0998.2010.02847.x

723 Federov, P.V., 1971. Postglacial transgression of the Black Sea. Int. Geol. Rev. 14, 160–  
724 164. doi: 10.1080/00206817209475678

725 Fišer, C., 2012. *Niphargus*: a model system for evolution and ecology. In: Culver, D.C.,  
726 White, W.B. (Eds.), Encyclopedia of caves. Academic Press, New York, pp. 555–  
727 564.

728 Fišer, C., Sket, B., Stoch, F., 2006. Distribution of four narrowly endemic *Niphargus*  
729 species (Crustacea: Amphipoda) in the western Dinaric region with description of a  
730 new species. Zool. Anz. 245, 77–94. doi: 10.1016/j.jcz.2006.05.003

731 Fišer, C., Keber, R., Kereži, V., Moškrič, A., Palandančić, A., Petkovska, V., Potočnik, H.,  
732 Sket, B., 2007. Coexistence of species of two amphipod genera: *Niphargus timavi*  
733 (*Niphargidae*) and *Gammarus fossarum* (*Gammaridae*). J. Nat. Hist. 41, 2641–2651.  
734 doi: 10.1080/00222930701661225

735 Fišer, C., Sket, B., Trontelj, P., 2008. A phylogenetic perspective on 160 years of troubled  
736 taxonomy of *Niphargus* (Crustacea: Amphipoda). Zool. Scr. 37, 665-680. doi:  
737 10.1111/j.1463-6409.2008.00347.x

738 Fišer, C., Trontelj, P., Luštrik, R., Sket, B., 2009. Toward a unified taxonomy of  
739 *Niphargus* (Crustacea: Amphipoda): A review of morphological variability. *Zootaxa*  
740 22, 1–22.

741 Fišer, C., Kovačec, Ž., Pustovrh, M., Trontelj, P., 2010a. The role of predation in the diet  
742 of *Niphargus* (Amphipoda: Niphargidae). *Speleobiology notes* 2, 4–6. doi:  
743 10.5563/spbn.v2i0.17

744 Fišer, C., Konec, M., Kobe, Z., Osanič, M., Gruden, P., Potočnik, H., 2010b. Conservation  
745 problems with hypothelminorheic *Niphargus* species (Amphipoda: Niphargidae).  
746 *Aquat. Conserv.* 20, 602–604. doi: 10.1002/aqc.1119

747 Fišer, C., Zagmajster, M., Zakšek, V. 2013., Coevolution of life history traits and  
748 morphology in female subterranean amphipods. *Oikos* 122, 770–778. doi:  
749 10.1111/j.1600-0706.2012.20644.x

750 Fišer, C., Pipan, T., Culver, D.C., 2014. The vertical extent of groundwater metazoans: an  
751 ecological and evolutionary perspective. *BioScience* 64, 971–979. doi:  
752 10.1093/biosci/biu148

753 Fišer, Ž., Altermatt, F., Zakšek, V., Knapič, T., Fišer, C., 2015. Morphologically cryptic  
754 amphipod species are “ecological clones” at regional but not at local scale. *PLoS*  
755 *ONE*, 10, e0134384. doi: 10.1371/journal.pone.0134384

756 Fišer, Ž., Novak, L., Luštrik, R., Fišer, C., 2016. Light triggers habitat choice of eyeless  
757 subterranean but not of eyed surface amphipods. *Sci. Nat.* 103, 1–12. doi:  
758 10.1007/s00114-015-1329-9

759 Flot, J.-F., 2010. SeqPHASE: a web tool for interconverting PHASE input/output files and  
760 FASTA sequence alignments. *Mol. Ecol. Resour.* 10, 162–166. doi: 10.1111/j.1755-  
761 0998.2009.02732.x

762 Flot, J.-F., Couloux, A., Tillier, S., 2010b. Haplowebs as a graphical tool for delimiting  
763 species: a revival of Doyle’s “field for recombination” approach and its application  
764 to the coral genus *Pocillopora* in Clipperton. BMC Evol. Biol. 10, 372. doi:  
765 10.1186/1471-2148-10-372

766 Flot, J.-F., Wörheide, G., Dattagupta, S., 2010a. Unsuspected diversity of *Niphargus*  
767 amphipods in the chemoautotrophic cave ecosystem of Frasassi, central Italy. BMC  
768 Evol. Biol. 10, 171. doi: 10.1186/1471-2148-10-171

769 Flot, J.-F., Bauermeister, J., Brad, T., Hillebrand-Voiculescu, A., Sarbu, S.M., Dattagupta,  
770 S., 2014. *Niphargus–Thiothrix* associations may be widespread in sulphidic  
771 groundwater ecosystems: evidence from southeastern Romania. Mol. Ecol. 23, 1405–  
772 1417. doi: 10.1111/mec.12461

773 Fu, Y.X., 1997. Statistical tests of neutrality of mutations against population growth,  
774 hitchhiking and background selection. Genetics 147, 915–925.

775 Fussi, B., Lexer, C., Heinze, B., 2010. Phylogeography of *Populus alba* (L.) and *Populus*  
776 *tremula* (L.) in Central Europe: secondary contact and hybridisation during  
777 recolonisation from disconnected refugia. Tree Genet. Genomes 6, 439–450. doi:  
778 10.1007/s11295-009-0262-5

779 Georgievski, G., Stanev, E.V., 2006. Paleo-evolution of the Black Sea watershed: sea level  
780 and water transport through the Bosphorus Straits as an indicator of the Lateglacial–  
781 Holocene transition. Clim. Dynam. 26, 631–644. doi: 10.1007/s00382-006-0123-y

782 Gilbert, J., Danielopol, D.L., Stanford, J.A., 1994. Groundwater Ecology. Academic Press,  
783 New York.

784 Ginet, R., 1960. Écologie, éthologie et biologie de *Niphargus* (Amphipodes Gammaridés  
785 hypogés). Ann. Spéléol. 15, 127–376.

786 Graeber, D., Pusch, M.T., Lorenz, S., Brauns, M., 2013. Cascading effects of flow  
787 reduction on the benthic invertebrate community in a lowland river. *Hydrobiologia*  
788 717, 147–159. doi: 10.1007/s10750-013-1570-1

789 Grigorovich, I.A., Therriault, T.W., MacIsaac, H.C., 2003. History of aquatic invertebrate  
790 invasions in the Caspian Sea. *Biol. Invasions* 5, 103–115. doi:  
791 10.1023/A:1024050824073

792 Guindon, S., Dufayard, J.F., Lefort, V., Anisimova, M., Hordijk, W., Gascuel, O., 2010.  
793 New algorithms and methods to estimate maximum-likelihood phylogenies:  
794 assessing the performance of PhyML 3.0. *Syst. Biol.* 59, 307–321. doi:  
795 10.1093/sysbio/syq010

796 Harpending, H.C., 1994. Signature of ancient population growth in a low-resolution  
797 mitochondrial DNA mismatch distribution. *Hum. Biol.* 66, 591–600.

798 Henry, K.S., Danielopol, D.L., 1999. Oxygen dependent habitat selection in surface and  
799 hyporheic environments by *Gammarus roeseli* Gervais (Crustacea, Amphipoda):  
800 experimental evidence. *Hydrobiologia* 390, 51–60. doi: 10.1023/A:1003541300460

801 Hervant, F., Mathieu, J., Garin, D., Fréminet, A., 1995. Behavioral, ventilatory, and  
802 metabolic responses to severe hypoxia and subsequent recovery of the hypogean  
803 *Niphargus rhenorhodanensis* and the epigeal *Gammarus fossarum* (Crustacea:  
804 Amphipoda). *Physiol. Zool.* 68, 223–244. doi: 10.1086/physzool.68.2.30166501

805 Horton, T., Lowry, J., De Broyer, C., Bellan-Santini, D., Coleman, C.O., Daneliya, M.,  
806 Dauvin, J-C., Fišer, C., Gasca, R., Grabowski, M., Guerra-García, J. M., Hendrycks,  
807 E., Holsinger, J., Hughes, L., Jaume, D., Jazdzewski, K., Just, J., Kamaltynov, R.M.,  
808 Kim, Y.-H., King, R., Krapp-Schickel, T., LeCroy, S., Lörz, A.-N., Senna, A.R.,  
809 Serejo, C., Sket, B., Tandberg, A.H., Thomas, J., Thurston, M., Vader, W., Väinölä,  
810 R., Vonk, R., White, K., Zeidler, W., 2017. World Amphipoda Database. Accessed at

811 <http://www.marinespecies.org/amphipoda./aphia.php?p=taxdetails&id=545672> on  
812 2017-06-08.

813 Humphreys, W.F., 2000. Background and glossary. In: Wilkens, H., Culver, D.C.,  
814 Humphreys, W.F. (Eds), *Ecosystems of the World*, vol. 30: Subterranean  
815 Ecosystems. Elsevier, Amsterdam, pp. 3–14.

816 Hüppop, K., 2000. How do cave animals cope with the food scarcity in caves? In: Wilkens,  
817 H., Culver, D.C., Humphreys, W.F. (Eds.), *Ecosystems of the World*, vol. 30:  
818 Subterranean ecosystems. Elsevier, Amsterdam, pp. 189–210.

819 Ho, S.Y.W., Phillips, M.J., Cooper, A., Drummond, A.J., 2005., Time dependency of  
820 molecular rate estimates and systematic overestimation of recent divergence times.  
821 *Mol. Biol. Evol.* 22, 1561–1568. doi: 10.1093/molbev/msi145

822 Ho, S.Y.W., Lanfear, R., Bromham, L., Phillips, M.J., Soubrier, J., Rodrigo, A.G., Cooper,  
823 A., 2011. Time-dependent rates of molecular evolution. *Mol. Ecol.* 20, 3087–3101.  
824 doi: 10.1111/j.1365-294X.2011.05178.x

825 Jazdzewski, K., Kupryjanowicz, J., 2010. One more fossil niphargid (Malacostraca:  
826 Amphipoda) from Baltic amber. *J. Crustacean Biol.* 30, 413–416. doi: 10.1651/09-  
827 3259.1

828 Jobb, G., 2011. TREEFINDER Version of March 2011. Munich, Germany. (computer  
829 program). Available at: [www.treefinder.de](http://www.treefinder.de)

830 Junk, W.J., Bayley, P.B., Sparks, R.E., 1989. The flood pulse concept in river-flood- plain  
831 systems. *Can. Spec. Publ. Fish. Aquat. Sci.* 106, 110–27.

832 Karaman, S., 1950a. *Niphargus ilidzensis* Schäferna i njegovi srodnici u Jugoslaviji. O  
833 nekim amfipodima — izopodima Balkana i o njihovoj sistematici. *Poseb. Izd.* 163,  
834 51–86.

835 Karaman, G.S., 1977. Contribution to the knowledge of the Amphipoda 78: *Niphargus*  
836 *elegans* Garbini, 1894, in Italy. Crustaceana Suppl. 4, 177–187.

837 Karaman, G.S., 1998. First discovery of the family Niphargidae (Gammaridea) in Iran.  
838 Glas. Od. Prirod. Nauk. – Crnog. Akad. Nauka umet. 12, 9–22.

839 Katoh K., Standley D.M., 2013. MAFFT Multiple Sequence Alignment Software Version  
840 7: Improvements in performance and usability. Mol. Biol. Evol. 30, 772–780.doi:  
841 10.1093/molbev/mst010

842 Katoh K., Toh H., 2008. Improved accuracy of multiple ncRNA alignment by  
843 incorporating structural information into a MAFFT-based framework. BMC  
844 Bioinformatics 9, 212. doi: 10.1186/1471-2105-9-212

845 Lanfear, R., Calcott, B., Ho, S.Y.W. Guindon, S., 2012. PartitionFinder: combined  
846 selection of partitioning schemes and substitution models for phylogenetic analyses.  
847 Mol. Biol. Evol. 29, 1695–1701. doi: 10.1093/molbev/mss020

848 Langecker, T.G., 2000. The effects of continuous darkness on cave ecology and  
849 cavernicolous evolution. In: Wilkens, H., Culver, D.C., Humphreys, W.F. (Eds),  
850 Ecosystems of the World, vol. 30: Subterranean Ecosystems. Elsevier, Amsterdam,  
851 pp. 135–157.

852 van Leeuwen, C.H.A., Huig, N., van der Velde, G., van Alen, T.A., Wagemaker, C.A.M.,  
853 Sherman, C.D.H., Klaassen, M., Figuerola, J., 2013. How did this snail get here?  
854 Several dispersal vectors inferred for an aquatic invasive species. Freshwater Biol.  
855 58, 88–99. doi: 10.1111/fwb.12041

856 Lefébure, T., Douady, C.J., Gouy, M., Trontelj, P., Briolay, J., Gibert, J., 2006.  
857 Phylogeography of a subterranean amphipod reveals cryptic diversity and dynamic  
858 evolution in extreme environments. Mol. Ecol. 15, 1797–1806. doi: 10.1111/j.1365-  
859 294X.2006.02888.x

860 Lefébure, T., Douady, C.J., Malard, F., Gibert, J., 2007. Testing dispersal and cryptic  
861 diversity in a widely distributed groundwater amphipod (*Niphargus*  
862 *rhenorhodanensis*). Mol. Phylogenet. Evol. 42, 676–686. doi:  
863 10.1016/j.ympev.2006.08.020

864 Lemey, P., Rambaut, A., Welch, J.J., Suchard, M.A., 2010. Phylogeography takes a  
865 relaxed random walk in continuous space and time. Mol. Biol. Evol. 27, 1877–1885.  
866 doi: 10.1093/molbev/msq067

867 Leonov, Y.G., Lavrushin, Y.A., Antipov, M.P., Sprididonova, Y.A., Kuzmin, Y.V., Jull,  
868 E.J.T., Burr, S., Jelinowska, A., Chalif, F., 2002. New age data on sediments of the  
869 transgressive phase of the Early Khvalyn transgression of the Caspian Sea. Dokl.  
870 Earth Sci. 386, 748–751.

871 Librado, P., Rozas, J., 2009. DnaSP v5: a software for comprehensive analysis of DNA  
872 polymorphism data. Bioinformatics 25, 1451–1452. doi:  
873 10.1093/bioinformatics/btp187

874 Lisiecki, L.E., Raymo, M.E., 2005. A Pliocene-Pleistocene stack of 57 globally distributed  
875 benthic  $\delta^{18}\text{O}$  records. Paleoceanography 20, PA1003. doi: 10.1029/2004PA001071

876 Liu, K., Raghavan, S., Nelesen, S., Linder, C.R., Warnow, T., 2009. Rapid and accurate  
877 large scale coestimation of sequence alignments and phylogenetic trees. Science 324,  
878 1561–1564. doi: 10.1126/science.1171243

879 Luštrik, R., Turjak, M., Kralj-Fišer, S., Fišer, C., 2011. Coexistence of surface and cave  
880 amphipods in an ecotone environment. Contrib. Zool. 80, 133–141.

881 Maddison, W.P., Maddison, D.R., 2015. Mesquite: a modular system for evolutionary  
882 analysis. Retrieved from <https://mesquiteproject.wikispaces.com>

883 Magri, D., Vendramin, G.G., Comps, B., Dupanloup, I., Geburek, T., Gömöry, D.,  
884 Latalowa, M., Litt, T., Paule, L., Roure, J.M., Tantau, I., van der Knaap, W.O., Petit,

885 R.J., de Beaulieu, J.L., 2006. A new scenario for the Quaternary history of European  
886 beech populations: Palaeobotanical evidence and genetic consequences. *New Phytol.*  
887 171, 199–221. doi: 10.1111/j.1469-8137.2006.01740.x

888 Maguire, B., 1960. Lethal effect of visible light on cavernicolous ostracods. *Science* 132,  
889 226–227. doi: 10.1126/science.132.3421.226

890 Malard, F., Hervant, F., 1999. Oxygen supply and the adaptations of animals in  
891 groundwater. *Freshwater Biol.* 41, 1–30. doi: 10.1046/j.1365-2427.1999.00379.x

892 Mauchart, P., Czirok, A., Horvai, V., Herczeg, R., Móra, A., Csabai, Z., 2017. Effects of  
893 meso- and microhabitat characteristics on the coexistence of two native gammarid  
894 species (Crustacea, Gammaridae). *Int. Rev. Hydrobiol.* 102, 38–46. doi:  
895 10.1002/iroh.201601855

896 McInerney, C.I., Maurice, L., Robertson, A.L., Knight, L.R.F.D., Arnscheidt, J., Venditti,  
897 C., Dooley, J.S.G., Mathers, T., Matthijs, S., Eriksson, K., Proudlove, G.S., Hänfling,  
898 B., 2014. The Ancient Britons: Groundwater fauna survived extreme climate changes  
899 over tens of millions of years across NW Europe. *Mol. Ecol.* 23, 1153–1166. doi:  
900 10.1111/mec.12664

901 Meade, A., Pagel, M., 2016. Bayes Traits v.3. Retrieved from  
902 <http://www.evolution.rdg.ac.uk/BayesTraitsV3/BayesTraitsV3.html>

903 Meleg, I.N., Zakšek, V., Fišer, C., Kelemen, B.S., Moldovan, O.T., 2013. Can environment  
904 predict cryptic diversity? The case of *Niphargus* inhabiting western Carpathian  
905 groundwater. *PLoS ONE*, 8, e76760. doi: 10.1371/journal.pone.0076760

906 Meijering, M.P.D., 1991. Lack of oxygen and low pH as limiting factors for *Gammarus* in  
907 Hessian brooks and rivers. *Hydrobiologia* 223, 159–169. doi: 10.1007/978-94-011-  
908 3542-9\_15

909 Meijering, M.P.D., Jazdzewski, K., Kohn, J., 1995. Ecotypes of Amphipoda in Central  
910 European inland waters. *Pol. Arch. Hydrobiol.* 42, 527–536.

911 Miller, M.P., 2005. Alleles In Space (AIS): computer software for the joint analysis of  
912 interindividual spatial and genetic information. *J. Hered.* 96, 722–724. doi:  
913 10.1093/jhered/esi119

914 Nesemann, H., Pöckl, M., Wittmann, K.J., 1995. Distribution of epigean Malacostraca in  
915 the middle and upper Danube (Hungary, Austria, Germany). *Miscnea. Zool. Hung.*  
916 10, 49–68.

917 Neumann, K., Michaux, J.R., Maak, S., Jansman, H.A.H., Kayser, A., Mundt, G.,  
918 Gattermann, R., 2005. Genetic spatial structure of European common hamsters  
919 (*Cricetus cricetus*) – a result of repeated range expansion and demographic  
920 bottlenecks. *Mol. Ecol.* 14, 1473–1483. doi: 10.1111/j.1365-294X.2005.02519.x

921 Parr, L.B., Mason, C.F., 2004. Causes of low oxygen in a lowland, regulated eutrophic  
922 river in Eastern England. *Sci. Total Environ.* 321, 273–286. doi:  
923 10.1016/j.scitotenv.2003.09.004

924 Peck, S.B., 1975. Amphipod dispersal in the fur of aquatic mammals. *Can. Field Nat.* 89,  
925 181–82.

926 Prendini, L., Francke, O.F., Vignoli, V., 2010. Troglomorphism, trichobothriotaxy and  
927 typhlochactid phylogeny (Scorpiones, Chactoidea): more evidence that troglobitism  
928 is not an evolutionary dead-end. *Cladistics* 26, 117–142. doi: 10.1111/j.1096-  
929 0031.2009.00277.x

930 Rachalewski, M., Banha, F., Grabowski, M., Anastacio, P.M., 2013. Ectozoochory as a  
931 possible vector enhancing the spread of an alien amphipod *Crangonyx*  
932 *pseudogracilis*. *Hydrobiologia* 717, 109–117. doi:10.1007/s10750-013-1577-7

933 Ramos-Onsins, S.E., Rozas, J., 2002. Statistical properties of new neutrality tests against  
934 population growth. *Mol. Biol. Evol.* 19, 2092–2100. doi:  
935 10.1093/oxfordjournals.molbev.a004034

936 Rewicz, T., Wattier, R., Grabowski, M., Rigaud, T., Bazcela-Spychalska, K., 2015. Out of  
937 the Black Sea: phylogeography of the invasive killer shrimp *Dikerogammarus*  
938 *villosus* across Europe. *PLoS ONE*, 10, e0118121. doi:  
939 10.1371/journal.pone.0118121

940 Rogers, A.R., Harpending, H., 1992. Population growth makes waves in the distribution of  
941 pairwise genetic differences. *Mol. Biol. Evol.* 9, 552–569. doi:  
942 10.1093/oxfordjournals.molbev.a040727

943 Ryan, W.B.F., Pitman III, W.C., Major, C.O., Shimkus, K., Moskalenko, V., Jones, G.A.,  
944 Dimitrov, P., Gorür, N., Sakinç, M., Yüce, H., 1997. An abrupt drowning of the  
945 Black Sea shelf. *Mar. Geol.* 138, 119–126. doi: 10.1016/S0025-3227(97)00007-8

946 Salzburger, W., Ewing, G., von Haeseler, A., 2011. The performance of phylogenetic  
947 algorithms in estimating haplotype genealogies with migration. *Mol. Ecol.* 20, 1952–  
948 1963. doi: 10.1111/j.1365-294X.2011.05066.x

949 Schluter, D., Price, T., Mooers, A.O., Ludwig D., 1997. Likelihood of ancestor states in  
950 adaptive radiation. *Evolution* 51, 1699-1711. doi: 10.2307/2410994

951 Shimodaira, H., 2002. An approximately unbiased test of phylogenetic tree selection. *Syst.*  
952 *Biol.* 51, 492–508. doi: 10.1080/10635150290069913

953 Shimodaira, H., Hasegawa, M., 1999. Multiple comparisons of log-likelihoods with  
954 applications to phylogenetic inference. *Mol. Biol. Evol.* 16, 1114–1116. doi:  
955 10.1093/oxfordjournals.molbev.a026201

956 Simčič, T., Brancelj, A., 2006. Effects of pH on electron transport system (ETS) activity  
957 and oxygen consumption in *Gammarus fossarum*, *Asellus aquaticus* and *Niphargus*  
958 *sphagnicolus*. *Freshwater Biol.* 51, 686–694. doi: 10.1111/j.1365-2427.2006.01522.x

959 Simčič, T., Lukančič, S., Brancelj, A., 2005. Comparative study of electron transport  
960 system activity and oxygen consumption of amphipods from caves and surface  
961 habitats. *Freshwater Biol.* 50, 494–501. doi: 10.1111/j.1365-2427.2005.01339.x

962 Sket, B., 1958. Prispevek k poznavanju nasih amfipodov Is, II. *Biol. Vestn.* 6, 67–75.

963 Sket, B., 1981. Distribution, ecological character and phylogenetic importance of  
964 *Niphargus valachicus* (Amphipoda, Gammaridae). *Biol. Vestn.* 29, 87–103.

965 Sket, B., 2008. Can we agree on an ecological classification of subterranean animals? *J.*  
966 *Nat. Hist.* 42, 1549–1563. doi: 10.1080/00222930801995762

967 Stamatakis, A., 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis  
968 of large phylogenies. *Bioinformatics* 30, 1312–1313. doi:  
969 10.1093/bioinformatics/btu033

970 Stephens, M., Smith, N., Donnelly, P., 2001. A new statistical method for haplotype  
971 reconstruction from population data. *Am. J. Hu. Genet.* 68, 978–989. doi:  
972 10.1086/319501

973 Straškraba, M., 1972a. Les groupments des especes du genre *Niphargus* (sensu lato). Actes  
974 du Ier Colloque International sur le genre *Niphargus*. Museo Civico di Storia  
975 Naturale di Verona Memorie fuori Serie 5, 85–90.

976 Straškraba, M., 1972b. L'état actuel de nos connaissances sur le genre *Niphargus* en  
977 Tchecoslovaquie et dans les pays voisins. Actes du Ier Colloque International sur le  
978 genre *Niphargus*. Museo Civico di Storia Naturale di Verona Memorie fuori Serie 5,  
979 35–45.

980 Strayer, D.L., 1994. Limits to biological distributions in groundwater. In: Gilbert, J.,  
981 Danielopol, D.L., Stanford, J.A. (Eds.), *Groundwater Ecology*. Academic Press, New  
982 York, pp. 287–310.

983 Swanson, G.A., 1984. Dissemination of amphipods by waterfowl. *J. Wildl. Manag.* 48,  
984 988–991. doi: 10.2307/3801453

985 Tajima, F., 1989. Statistical method for testing the neutral mutation hypothesis by DNA  
986 polymorphism. *Genetics* 123, 585–595.

987 Talavera, G., Castresana, J., 2007. Improvement of phylogenies after removing divergent  
988 and ambiguously aligned blocks from protein sequence alignments. *Syst. Biol.* 56,  
989 564–577. doi: 10.1080/10635150701472164

990 Tamura, K., Stecher, G., Peterson, D., Filipiński, A., Kumar, S., 2013. MEGA6: Molecular  
991 Evolutionary Genetics Analysis version 6.0. *Mol. Biol. Evol.* 30, 2725–2729. doi:  
992 10.1093/molbev/mst197

993 Trontelj, P., Blejec, A., Fišer, C., 2012. Ecomorphological convergence of cave  
994 communities. *Evolution* 66, 3852–3865. doi: 10.1111/j.1558-5646.2012.01734.x

995 Trontelj, P., Douady, C., Fišer, C., Gibert, J., Gorički, Š., Lefébure, T., Sket, B., Zakšek,  
996 V., 2009. A molecular test for hidden biodiversity in groundwater: how large are the  
997 ranges of macro-stygobionts? *Freshwater Biol.* 54, 727–744. doi: 10.1111/j.1365-  
998 2427.2007.01877.x

999 de Vaate, A.B., Jazdzewski, K., Ketelaars, H.A.M., Gollasch, S., Van der Velde, G., 2002.  
1000 Geographical patterns in range extension of Ponto-Caspian macroinvertebrate  
1001 species in Europe. *Can. J. Fish. Aquat. Sci.* 59, 1159–1174. doi: 10.1139/f02-098

1002 Väinölä, R., Witt, J.D.S., Grabowski, M., Bradbury, J.H., Jazdzewski, K., Sket, B., 2008.  
1003 Global diversity of amphipods (Amphipoda; Crustacea) in freshwater. *Hydrobiologia*  
1004 595, 241–255. doi: 10.1007/978-1-4020-8259-7\_27

1005 Verovnik, R., Sket, B., Trontelj, P., 2005. The colonization of Europe by the freshwater  
1006 crustacean *Asellus aquaticus* (Crustacea: Isopoda) proceeded from ancient refugia  
1007 and was directed by habitat connectivity. *Mol. Ecol.* 14, 4355–4369. doi:  
1008 10.1111/j.1365-294X.2005.02745.x

1009 Villesen, P., 2007. FaBox: an online toolbox for fasta sequences. *Mol. Ecol. Notes* 7, 965–  
1010 968. doi: 10.1111/j.1471-8286.2007.01821.x

1011 Vörös, J., Mikulíček, P., Major, Á., Recuero, E., Arntzen, J.W., 2016. Phylogeographic  
1012 analysis reveals northerly refugia for the riverine amphibian *Triturus dobrogicus*  
1013 (Caudata: Salamandridae). *Biol. J. Linn. Soc.* 119, 974–991. doi: 10.1111/bij.12866

1014 Ward, J.V., Palmer, M.A., 1994. Distribution patterns of interstitial freshwater meiofauna  
1015 over a range of spatial scales, with emphasis on alluvial river-aquifer systems  
1016 *Hydrobiologia* 287, 147–156. doi: 10.1007/BF00006903

1017 Weber, M.G., Wagner, C.E., Best, R.J., Harmon, L.J., Matthews, B., 2017. Evolution in a  
1018 Community Context: On Integrating Ecological Interactions and Macroevolution.  
1019 *Trends Ecol. Evol.* 32, 291–304. doi: 10.1016/j.tree.2017.01.003

1020 Weir, B.S., Cockerham, C.C., 1984. Estimating F-Statistics for the analysis of population-  
1021 structure. *Evolution* 38, 1358–1370. doi: 10.2307/2408641

1022 Zgmajster, M., Eme, D., Fišer, C., Galassi, D., Marmonier, P., Stoch, F., Cornu, J.F.,  
1023 Malard, F., 2014. Geographic variation in range size and beta diversity of  
1024 groundwater crustaceans: insights from habitats with low thermal seasonality. *Global*  
1025 *Ecol. Biogeogr.* 23, 1135–1145. doi: 10.1111/geb.12200

1026 Zhang, J., Kapli, P., Pavlidis, P., Stamatakis, A., 2013. A general species delimitation  
1027 method with applications to phylogenetic placements. *Bioinformatics* 29, 2869–  
1028 2876. doi: 10.1093/bioinformatics/btt499

1029

1030

1031 **Supporting information**

1032 **Tables S1-S11** Details about samples used in the phylogeographic and phylogenetic  
1033 analyses, GenBank accession numbers, geographic coordinates, PCR primers,  
1034 evolutionary, coalescent and clock models, morphological data, ancestral states, and  
1035 population differentiation based on  $\Phi$  statistics.

1036 **Figures S1-S3** Fully annotated BI and ML phylogenies, isolation by distance plots and  
1037 cryptic lineage delimitation results.

1038

1039

1040

1041

1042

1043

1044 Table 1. Genetic polymorphism (COI and ITS) and historical demographic results (COI)

1045 for the studied species and clades. For the neutrality tests, p-values are provided in

1046 parentheses and significant values are shown in bold.

| Species/clade                   | <i>N</i>    | <i>S</i> | <i>H</i> | <i>Hd</i> (SD)   | $\pi$ (SD)           | <i>K</i> | <i>D</i> ( <i>p</i> ) | <i>F<sub>s</sub></i> ( <i>p</i> ) | <i>R<sub>2</sub></i> ( <i>p</i> ) | $\tau$ | $\theta_0$ | $\theta_1$ | <i>SSD</i> ( <i>p</i> ) | <i>Hri</i> ( <i>p</i> ) |
|---------------------------------|-------------|----------|----------|------------------|----------------------|----------|-----------------------|-----------------------------------|-----------------------------------|--------|------------|------------|-------------------------|-------------------------|
| <b>COI</b>                      |             |          |          |                  |                      |          |                       |                                   |                                   |        |            |            |                         |                         |
| <i>N. hrabei</i>                | 54          | 15       | 12       | 0.768<br>(0.046) | 0.0028<br>(0.0004)   | 1.76     | -1.376<br>(0.056)     | -4.160<br><b>(0.031)</b>          | 0.108<br>(0.083)                  | 0.68   | 1.23       | 3451.34    | 0.003<br>(0.522)        | 0.034<br>(0.275)        |
| <i>N. valachicus</i><br>overall | 111         | 42       | 42       | 0.962<br>(0.007) | 0.0157<br>(0.0004)   | 7.13     | -0.413<br>(0.406)     | -5.742<br>(0.117)                 | 0.092<br>(0.402)                  | 12.26  | 0.00       | 16.48      | 0.009<br>(0.470)        | 0.008<br>(0.719)        |
| <i>N. valachicus</i> A          | 49          | 19       | 14       | 0.877<br>(0.030) | 0.0079<br>(0.0006)   | 3.56     | -0.724<br>(0.252)     | 3.044<br>(0.880)                  | 0.107<br>(0.385)                  | 3.21   | 1.54       | 8.32       | 0.004<br>(0.766)        | 0.015<br>(0.867)        |
| <i>N. valachicus</i> B          | 62          | 36       | 28       | 0.956<br>(0.011) | 0.0071<br>(0.0006)   | 4.42     | -1.385<br>(0.061)     | -14.238<br><b>(0.000)</b>         | 0.102<br>(0.06)                   | 3.00   | 1.90       | 3414.97    | 0.006<br>(0.078)        | 0.020<br><b>(0.028)</b> |
| <b>ITS</b>                      |             |          |          |                  |                      |          |                       |                                   |                                   |        |            |            |                         |                         |
| <i>N. hrabei</i>                | 30<br>(22)* | 7        | 6        | 0.736<br>(0.056) | 0.00047<br>(0.00008) | 1.26     |                       |                                   |                                   |        |            |            |                         |                         |
| <i>N. valachicus</i>            | 66<br>(40)* | 28       | 23       | 0.73<br>(0.060)  | 0.00339<br>(0.00045) | 4.96     |                       |                                   |                                   |        |            |            |                         |                         |

1047 *N* – sample size (no. of sequences); *S* – number of variable sites; *H* – number of haplotypes; *Hd* – haplotype

1048 diversity;  $\pi$  – nucleotide diversity; *K* – mean number of pairwise nucleotide differences; *D* – Tajima’s *D*

1049 statistics; *F<sub>s</sub>* – Fu’s *F<sub>s</sub>* statistics; *R<sub>2</sub>* – Ramos-Onsins and Rozas’s *R<sub>2</sub>* statistics;  $\tau$  – coalescence time in

1050 mutational units;  $\theta_0$ ,  $\theta_1$  – effective population size at the start and the end of the expansion; *SSD* – sum of

1051 squared deviations; *Hri* – Harpending’s raggedness index; \* – parentheses refer to no. of analysed individuals

1052

1053 Table 2. Analysis of molecular variance (AMOVA) in the studied species based on COI  
 1054 with populations grouped according to the major river drainages (listed in Tables S10 and  
 1055 S11). Significant p-values are shown in bold for the fixation indices ( $\Phi$ -statistics).

1056

| 1057 | <b>Species</b>       | <b>Source of variation</b>         | <b>d.f.</b> | <b>Sum of squares</b> | <b>Percentage of variation</b> | <b><math>\Phi</math>-statistics</b> | <b>P</b>          |
|------|----------------------|------------------------------------|-------------|-----------------------|--------------------------------|-------------------------------------|-------------------|
| 1058 | <i>N. hrabei</i>     | Among drainages                    | 4           | 31.28                 | 80.05                          | $\Phi_{ct} = 0.80$                  | <b>0.007</b>      |
| 1059 |                      | Among populations within drainages | 10          | 4.03                  | 11.19                          | $\Phi_{sc} = 0.56$                  | <b>0.002</b>      |
| 1060 |                      | Within populations                 | 35          | 3.4                   | 8.75                           | $\Phi_{st} = 0.91$                  | <b>&lt;0.0001</b> |
| 1061 |                      |                                    |             |                       |                                |                                     |                   |
| 1062 | <i>N. valachicus</i> | Among drainages                    | 12          | 381.68                | 58.5                           | $\Phi_{ct} = 0.58$                  | <b>&lt;0.0001</b> |
| 1063 |                      | Among populations within drainages | 25          | 131.3                 | 33.1                           | $\Phi_{sc} = 0.79$                  | <b>&lt;0.0001</b> |
| 1064 |                      | Within populations                 | 73          | 32.75                 | 8.4                            | $\Phi_{st} = 0.91$                  | <b>&lt;0.0001</b> |
| 1065 |                      |                                    |             |                       |                                |                                     |                   |
| 1066 |                      |                                    |             |                       |                                |                                     |                   |
| 1067 |                      |                                    |             |                       |                                |                                     |                   |

1068

1069

1070

1071

1072

1073

1074

1075

1076

1077

1078

1079

1080 Figure captions

1081

1082 Fig. 1 Bayesian phylogeny of *Niphargus* and ancestral state reconstruction of the species'

1083 habitat. Species that occur in surface habitats are highlighted with red, dashed branches.

1084 The widely-dispersed focal species, *N. hrabei* and *N. valachicus* are indicated with larger

1085 font size. Circles at nodes denote posterior probability for clade support (black  $\geq 0.95$ , grey

1086 = 0.90-0.94 and white = 0.70-0.89). The pie charts along the numbered nodes of the tree

1087 indicate the probability of subterranean (grey) and surface (red) ecology of the ancestors.

1088 The numbers at nodes correspond with those in Table S9. Inset image depicts a male *N.*

1089 *valachicus* (Photograph: Denis Copilaş-Ciocianu).

1090

1091 Fig. 2 Geographical distribution and haplotype networks of COI and ITS in *N. hrabei* (A,

1092 C) and *N. valachicus* (B, D). Colours indicate middle (red) and lower (yellow) Danube and

1093 Black Sea (green) populations. Distribution of clades A and B of *N. valachicus* are

1094 indicated by dotted and dashed lines, respectively. The size of the circles in the haplotype

1095 networks is proportional to the observed frequency of the corresponding haplotype. Each

1096 segment indicates one substitution. Relevant countries are indicated by corresponding 2-

1097 letter ISO codes: AT—Austria, HU—Hungary, HR—Croatia, RO—Romania, RS—Serbia

1098 and TR—Turkey.

1099

1100 Fig. 3 Coalescence and demography of *N. hrabei* (upper panel) and *N. valachicus* (lower

1101 panel). A) Time-calibrated COI trees. Posterior probability at nodes is indicated by circles

1102 (black  $\geq 0.95$ , dark grey = 0.85–0.94, light grey = 0.70–0.84, and white = 0.50–0.69). Blue

1103 bars at nodes denote the 95% HPD intervals of clade age. B) Bayesian skyline plots (BSP)

1104 and mismatch distribution histograms. Mean populations size through time is shown by

1105 thick black lines and 95% confidence intervals with grey in the BSP. Mismatch  
1106 distributions were also calculated separately for each of the two clades (A and B) of *N.*  
1107 *valachicus*. Continuous lines indicate the observed frequency of pairwise differences and  
1108 dotted lines indicate the expected frequency under a model of sudden demographic  
1109 expansion.

1110

1111 Fig. 4 Dispersal of *N. hrabei* (A) and *N. valachicus* (B) inferred from Bayesian  
1112 phylogeographic diffusion models. The putative origin of dispersal is shown with a dashed,  
1113 white line. Dispersal routes at different time intervals are indicated by different line  
1114 shadings. Country ISO codes are the same as in Fig. 2.

1115

1116 Fig. 5 Clustering of 81 *Niphargus* taxa based on 35 morphological traits. Surface-water  
1117 species are indicated with red font and grey shading. The focal *N. hrabei* and *N. valachicus*  
1118 are shown with larger font size. Ecomorph names follow Trontelj et al., (2012).


1119





1120

1121

1122




- PP ≥ 0.95
- PP = 0.90-0.94
- PP = 0.70-0.89







**A**

- PP ≥ 0.95
- PP = 0.85-0.94
- PP = 0.70-0.84
- PP = 0.50-0.69

**B***N. hrabei***A***N. valachicus*



0 5 10 15 20 25

small pore  
ecomorph

*N. kochianus*  
*N. labacensis*  
*N. glenniei*  
*N. serbicus*  
*N. foreli*  
*N. longicaudatus* Gargano  
*N. fongi*  
*N. pretneri*  
*N. puteanus*  
*N. carniolicus*  
*N. jovanovici*  
*N. multipennatus*  
*N. grandii*  
*N. hvarensis*  
*N. miljeticus*  
*N. elegans*  
*N. hrabei*  
*N. danconai*  
*N. skopljensis*  
*N. hebereri*  
*N. dimorphopus*  
*N. pectinicauda*  
*N. stefanelli*  
*N. lunaris*  
*N. valachicus*

cave stream  
ecomorph

*N. karamani*  
*N. likanus*  
*N. podpecanus*  
*N. timavi*  
*N. novomestanus*  
*N. sphagnicolus*  
*N. longicaudatus*  
*N. pasquinii*  
*N. schellenbergi*  
*N. scopicauda*  
*N. aitulosi*  
*N. aquilex*  
*N. rhenorhodanensis*  
*N. stygius*  
*N. kenki*  
*N. costozzae*  
*N. sanctinaumi*  
*N. illidzensis*  
*N. zagrebensis*  
*N. dalmatinus*  
*N. krameri*  
*N. vinodolensis*  
*N. hadziji*  
*N. spoeckeri*  
*N. spinulifemur*  
*N. brachytelson*  
*N. slovenicus*

daddy-  
longlegs  
ecomorph

*N. longicaudatus* Cres  
*N. tatrensis*

lake  
ecomorph

*N. balcanicus*  
*N. croaticus*  
*N. liburnicus*  
*N. steueri*  
*N. bilecanus*  
*N. subtypicus*  
*N. polymorphus*  
*N. trullipes*

lake giant  
ecomorph

*N. salonitanus*  
*N. stenopus*  
*N. orcinus*  
*N. vjetrnicensis*  
*N. dabarensis*  
*N. lourensis*  
*N. pachytelson*  
*N. arbiter*  
*N. jadranko*  
*N. rejici*

## Supplementary Information

### **Is subterranean lifestyle reversible? Independent and recent large-scale dispersal into surface waters by two species of the groundwater amphipod genus *Niphargus***

Denis Copilaş-Ciocianu, Cene Fişer, Péter Borza and Adam Petrusek

#### **Supplementary Methods**

##### *Phylogenetic analyses*

Bayesian inference (BI) and maximum-likelihood (ML) approaches were used to reconstruct phylogenetic relationships and examine the position of *N. hrabei* and *N. valachicus* within the genus. Both approaches were applied to the concatenated supermatrix. BI was carried out in BEAST 1.8.0 (Drummond et al. 2012). We employed the evolutionary models in Table S4 for each partition. Random starting topologies were used for each run and speciation was modelled using a Yule prior. As we were not interested in absolute divergence times, we used a relaxed molecular clock with a lognormal distribution and the rate was left at the default value of 1. Clock models and trees were linked across partitions. The MCMC chain was run for 100 million generations with a sampling frequency of 1000 and 30% of the trees were discarded as burn-in. Convergence and effective sample size was assessed using TRACER 1.6 and the maximum clade credibility (MCC) tree was produced with TreeAnnotator 1.8. The alignment used in the BEAST analysis had regions of questionable homology in the 28S marker removed with GBLOCKS 0.9 (Talavera and Castresana, 2007). To evaluate the effect of removal of these regions on the topology of the tree, we used a ML method that simultaneously estimates the sequence alignment and phylogenetic tree in SATé 2.2.7 (Liu et al. 2009), thus retaining the poorly alignable regions. MAFFT 6.7 (Katoh et al. 2005) was used as the initial aligner and OPAL 1.0.3 (Wheeler & Kececioglu 2007) was used to merge the alignment of subproblems into the final alignment. The tree was estimated with RAxML 7.2.6 (Stamatakis 2006) and the GTR+ $\Gamma$  model. The cycle of alignment and tree estimation was iterated ten times. Because SATé does not calculate bootstrap support on RAxML produced trees and does not handle codon partitions, the alignment with the best likelihood was used for the final tree estimation and bootstrapping in RAxML-HPC 8.2.9 (Stamatakis 2014). A thorough ML tree search was performed with the GTR+ $\Gamma$  model assigned to each partition and 1000 fast bootstrap iterations. We conducted two independent runs of both ML and BI. Analyses were carried out on the CIPRES Science Gateway (Miller et al. 2010).

##### *Bayesian phylogeographical diffusion models*

To explore the origin and dispersal history of each species while accounting for phylogenetic uncertainty, we used Bayesian phylogeographic diffusion models (Lemey et al. 2010) implemented in BEAST 1.8.0. The analysis was based on the COI marker and we used the same settings as for the time-calibrated phylogenetic analyses. The coordinates of each locality were used as a quantitative trait and individuals possessing the same haplotype but collected from different locations were retained because their corresponding coordinates contained spatial information. We compared four diffusion models available in BEAST: a

random walk model following a homogeneous Brownian diffusion (BD), and three relaxed random walk models (RRW) using gamma, lognormal and Cauchy distributions (Lemey et al. 2010). For each spatial diffusion model and species, we ran two independent runs of 30 million generations which were sampled every 1000 steps with 10% burn-in and convergence was evaluated with TRACER. The best-fit diffusion models were selected by calculating Bayes factors based on marginal likelihoods estimated with path and stepping stone sampling (Baele et al. 2012) and are presented in Table S6. The phylogeographic history of each species was visualized in Google Earth Pro 7.1.5 (<https://www.google.com/earth/>) by producing and input Keyhole Markup Language (kml) file with SPREAD 1.0.7 (Bielejec et al. 2011).

### *Analysed morphological traits*

A total of 34 quantitative and one qualitative trait were analysed. Landmarks and variation of these traits were presented in a previous study (Fišer et al. 2009). Many of the measured traits are presumably linked to the species ecology (see Trontelj et al. 2012; Fišer et al. 2015; Copilas-Ciocianu et al. 2017). *Body size* is related to the trophic ecology, but also to the pore size of microhabitats the species lives in (Trontelj et al. 2012). *Body shape* relates to hydrodynamic properties of the body (Dahl 1977) and reproductive biology (Fišer et al. 2013). It can be inferred from coxal plates II and III and bases of pereopods V-VII (all measured as width and length). *Appendage length* is a tradeoff between extra-optic sensory capacity and resilience to water flow (Pipan & Culver 2012; Trontelj et al. 2012, Delić et al. 2016). We measured lengths of antennae I-II and pereopods V-VII. *Shape and size of gnathopods I-II* is likely involved into feeding biology. We measured lengths of carpus and propodus, propodus palm and the length of propodus diagonal (Copilas-Ciocianu et al. 2017). *Uropods I and III* are sexually dimorphic in some species. In these species, males have either elongated inner ramus of uropod I, or distal article of exopodite of uropod III, or both. In addition, some males have a strange, flap-like appendix on a base of uropod I. We measured lengths of both rami of uropod I, and both articles of exopodite of uropod III; the flap-like appendix was treated as present-absent. Spines on dactyls of pereopods III-VII, and spines on urosomites I-II, the function of which is not known, vary in number between one and nine, were counted.

### **References**

- Bielejec F, Rambaut A, Suchard MA, Lemey P (2011) SPREAD: spatial phylogenetic reconstruction of evolutionary dynamics. *Bioinformatics*, 27, 2910–2912.
- Copilas-Ciocianu D, Fišer C, Borza P, Balázs G, Angyal D, Petrusek A (2017) Low intraspecific genetic divergence and weak niche differentiation despite wide ranges and extensive sympatry in two epigeal *Niphargus* species (Crustacea: Amphipoda). *Zoological Journal of the Linnean Society*, doi: 10.1093/zoolinnean/zw031
- Dahl E (1977) The amphipod functional model and its bearing upon systematics and phylogeny. *Zoologica Scripta*, 6, 221–228.
- Drummond AJ, Suchard MA, Xie D, Rambaut A (2012) Bayesian phylogenetics with BEAUti and the BEAST 1.7. *Molecular Biology and Evolution*, 29, 1969–1973.
- Fišer C, Zigmajster M, Zakšek V (2013) Coevolution of life history traits and morphology in female subterranean amphipods. *Oikos*, 122, 770–778.


- Fišer C, Trontelj P, Luštrik R, Sket B (2009) Toward a unified taxonomy of *Niphargus* (Crustacea: Amphipoda): A review of morphological variability. *Zootaxa*, 22, 1–22.
- Fišer C, Luštrik R, Sarbu S, Flot J-F, Trontelj P (2015) Morphological evolution of coexisting amphipod species pairs from sulfidic caves suggests competitive interactions and character displacement, but no environmental filtering and convergence. *PLoS ONE*, 10, e0123535.
- Katoh K, Kuma K-I, Toh H, Miyata T (2005) MAFFT version 5: improvement in accuracy of multiple sequence alignment. *Nucleic Acids Research*, 33, 511–8.
- Lemey P, Rambaut A, Welch JJ, Suchard MA (2010) Phylogeography takes a relaxed random walk in continuous space and time. *Molecular Biology and Evolution*, 27, 1877–1885.
- Liu K, Raghavan S, Nelesen S, Linder CR, Warnow T (2009) Rapid and accurate large scale coestimation of sequence alignments and phylogenetic trees. *Science*, 324, 1561–1564.
- Miller MA, Pfeiffer W, Schwartz T (2010) Creating the CIPRES Science Gateway for inference of large phylogenetic trees. In: Proceedings of the Gateway Computing Environments Workshop (GCE), pp 1–8. 14 Nov. 2010, New Orleans, Louisiana.
- Pipán T, Culver DC (2012) Convergence and divergence in the subterranean realm: a reassessment. *Biological Journal of the Linnean Society*, 101, 1–14.
- Stamatakis A (2006) RAxML-VI-HPC: maximum likelihood based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics*, 22, 2688–2690.
- Talavera G, Castresana J (2007) Improvement of phylogenies after removing divergent and ambiguously aligned blocks from protein sequence alignments. *Systematic Biology*, 56, 564–577.
- Trontelj P, Blejec A, Fišer C (2012) Ecomorphological convergence of cave communities. *Evolution*, 66, 3852–3865.
- Wheeler TJ, Kececioglu JD (2007) Multiple alignment by aligning alignments. Proceedings of the 15th ISCB Conference on Intelligent Systems for Molecular Biology, *Bioinformatics*, 23, 559–568.



Fig. S1. Maximum-likelihood (left) and Bayesian phylogenies (right) obtained from the concatenated dataset of COI, H3 and 28S sequences. The ML tree was obtained from an alignment produced with SATé and contained poorly alignable regions in the 28S marker, while the alignment for the BI tree had these regions removed and was obtained with MAFFT. Numbers above nodes are bootstrap percentages and posterior probabilities, respectively. Only values above 40% and 0.70 are shown. Surface-water species are shown with red branches. *N. hrabei* and *N. valachicus* are indicated with larger font size.

*N. hrabei*

$r = 0.4317, p = 0.00001$



*N. valachicus*

$r = 0.3737, p = 0.00001$




Fig. S2. Correlation between geographic and genetic (based on COI) distance.




Fig. S3. Cryptic lineage delimitation results based on COI and ITS in *N. hrabei* (above) and *N. valachicus* (below). A) Results from the bPTP analysis based on COI. Clustered red branches indicate putative lineages which correspond to the two focal species, i.e. no independent intraspecific lineages were detected. Numbers above branches indicate the posterior probability for species assignment. B) Haplowebs based on phased ITS sequences. Haplotypes that co-occur in the same individual are connected by thin, dotted lines. All haplotypes are interconnected, indicating a common gene pool.