

Integrating plant science and crop modelling: Assessment of the impact of climate change on soybean and maize production

Nándor Fodor^{1,2}, Andrew Challinor³, Ioannis Droutsas³, Julian Ramirez-Villegas^{4,5}, Florian Zabel⁶, Ann-Kristin Koehler³, Christine H. Foyer^{1*}

1 Centre for Plant Sciences, School of Biology, Faculty of Biological Sciences, University of Leeds, LS2 9JT Leeds, UK

2 Centre for Agricultural Research, Hungarian Academy of Sciences, 2462 Martonvásár
Brunszvik u. 2., Hungary

3 Institute for Climate and Atmospheric Science, School of Earth and Environment,
University of Leeds, LS2 9JT Leeds, UK

4 International Center for Tropical Agriculture (CIAT), km 17 recta Cali-Palmira, Cali,
Colombia

5 CGIAR Research Program on Climate Change, Agriculture and Food Security (CCAFS),
c/o CIAT, km 17 recta Cali-Palmira, Cali, Colombia

6 Ludwig-Maximilians-Universität München, Luisenstrasse 37, 80333 Munich, Germany

*Corresponding author e-mail: c.foyer@leeds.ac.uk

Running title: Climate change impacts on soybean and maize production

Key words: high CO₂, photosynthesis; crop production; land use; climate change modelling

ABSTRACT

Increasing global CO₂ emissions have profound consequences for plant biology, not least because of direct influences on carbon gain. However, much remains uncertain regarding how our major crops will respond to a future high CO₂ world. Crop models inter-comparison studies have identified large uncertainties and biases associated with climate change. The need to quantify uncertainty has drawn the fields of plant molecular physiology, crop breeding and biology and climate change modelling closer together. Comparing data from different models that have been used to assess the potential climate change impacts on soybean and maize production, future yield losses have been predicted for both major crops. However, when CO₂ fertilisation effects are taken into account significant yield gains are predicted for soybean, together with a shift in global production from the Southern to the Northern hemisphere. Maize production is also forecast to shift northwards. However, unless plant breeders are able to produce new hybrids with improved traits, the forecasted yield losses for maize will only be mitigated by agro-management adaptations. In addition, the increasing demands of a growing world population will require larger areas of marginal land to be used for maize and soybean production. We summarise the outputs of crop models, together with mitigation options for decreasing the negative impacts of climate on the global maize and soybean production, providing an overview of projected land-use change as a major determining factor for future global crop production.

INTRODUCTION

Atmospheric CO₂ concentrations have risen from about 280 μLL^{-1} in pre-industrial times to 400 μLL^{-1} at present (IPCC 2013). The increasing concentration rate has accelerated in recent years to the extent that [CO₂] may reach between 530 and 970 μLL^{-1} by the end of the 21st century, leading to significant global warming (IPCC 2013). Higher temperatures and high [CO₂] can be both beneficial and detrimental to plants, leading to changes in the global agricultural landscape. Average global temperatures have increased by 0.76 °C over the last 150 years and are likely to increase by at least another 1.7 °C by the end of this century. It is generally assumed that most plants are adapted to atmospheric [CO₂] below 300 μLL^{-1} and that they will be slow to adapt to the ongoing rapid increases (Ort et al. 2015).

Since high [CO₂] will favor photosynthetic carbon assimilation and depress photorespiration in plants with the C₃ pathway of photosynthesis, it is generally assumed that C₃ plants will benefit from increased carbon gain that will translate into increased biomass and yield. However, many aspects of plant metabolism, molecular physiology, structure and development are modified by growth under high atmospheric [CO₂], not least because the assimilation of carbon is tightly linked to primary nitrogen assimilation. Moreover, increased [CO₂] reduces the density of stomata and also aperture of the stomatal resulting in decreased evapotranspiration (Mansfield et al. 1990, Kim et al. 2010, Vavasseur and Raghavendra 2005). Stomatal development is also controlled by both [CO₂] and the phytohormone abscisic acid (ABA, Woodward 1987, Woodward and Kelly 1995, Tanaka et al. 2013). Several components have been identified in the signaling pathway that reduces stomatal apertures in response to elevated [CO₂] including β -carbonic anhydrases (Hu et al. 2010), the HT1 protein kinase, the RHC1 MATE transporter and the NtMPK4 protein kinase (Hashimoto et al. 2006, Marten et al. 2008, Tian et al. 2015). However, the generation of reactive oxygen species (ROS) is involved in both high [CO₂]-induced decreases in stomatal density and stoma, requiring the presence of ABA, PYR/RCAR and ABA receptors (Chater et al. 2015). Despite extensive research efforts over the last 50 years, the complex interplay between metabolic and environmental signals that determine the plant response to high CO₂ is far from resolved, particularly at the whole plant level. Much of our current understanding of the responses of crop growth to high atmospheric [CO₂] has come from either studies in free air CO₂ enrichment (FACE) sites or chamber (closed or open-top) experiments. However, such studies have not always yielded consistent results. CO₂ enrichment does not necessarily enhance plant growth or yield and differences in the responses of these

traits have been reported even within the same species (Ainsworth and Long 2005, Luo et al. 2006, Leakey et al. 2009a,b, Hasegawa et al. 2013, Bishop et al. 2015). These studies provide the essential foundation data underpinning crop models, predicting future changes in crop production and their implications for food security.

Crop models have a central role in informing agro-industry and policymakers about the risks and potential of adaptation strategies to counter climate change, as well as directing plant scientists and breeders towards the required traits in improved varieties and cropping systems' management practices to mitigate global climate change impacts. Crop model inter-comparison studies have identified large uncertainties and biases (e.g. Asseng et al. 2013, 2014, Bassu et al. 2014), and unfortunately they do not often incorporate current knowledge of plant responses to growth under high atmospheric [CO₂] (Durand et al. 2017). This review summarises current crop models and the complexity of analysis, within the context of our current knowledge on the impacts of a high [CO₂] on the C3 crop plant soybean (*Glycine max*), and the C4 crop maize (*Zea mays*), which has an internal CO₂ concentrating mechanism. Maize and soybean are used to produce a wide range of food and non-food products including pharmaceuticals and biofuels, as well as important sources of livestock feed. Here, we consider the projected impacts of a future high CO₂ world on the global production of maize and soybean. Currently, maize is the most important grain crop and soybean the fourth most important in terms of global production. Since 1960, soybean and maize grain yields increased 7.6 and 2.6 times, respectively. Together, the USA, Brazil and Asia (mainly China and India) account for respectively 92% and 84% of the world soybean and maize production. However, while the land area on which grain legumes, such as soybean are grown has gradually increased over the past 50 years, this is still only a quarter of that planted to cereals, such as maize (Foyer et al. 2016). In addition, while increases in cereal production over this period have been predominantly due to increases in yield, driven by the introduction of new varieties and improvements in agronomic practices, whereas increases in grain legume production are due to both increases in land area planted and grain yield (Foyer et al. 2016). For soybean in particular, grain yields have increased in proportion to the land area planted. Moreover, year-on-year increases in soybean yields are slowing while area planted is increasing, suggesting that more marginal land is being planted.

In this review, we will provide a brief overview of our current understanding of the molecular, metabolic and physiological responses of plants to increasing atmospheric [CO₂] and briefly summarise the history and types of crop models that are currently available. We then

specifically address the question of how increasing atmospheric [CO₂] will alter global soybean and maize production patterns. Using 118 peer-reviewed publications (31 for soybean and 87 for maize), we review the main issues that should be taken into account when modelling of these two important crops, namely model inputs, the roles of [CO₂] adaptation, mitigation, and modelling uncertainties. Finally, we discuss projected land-use change as a major determining factor for future global crop production.

THE PLASTICITY OF PLANT RESPONSES TO HIGH CO₂

There is now an extensive literature on the responses of plant biology to growth under high [CO₂] conditions, with reviews ranging from the control of photosynthetic electron transport and re-programming of photosynthetic gene expression that accompanies the suppression of photorespiration (Foyer et al. 2012) to effects on abiotic stress tolerance (AbdElgawad et al. 2016). It is not our intention therefore to describe the complex and many-faceted responses of plants to CO₂ enrichment but rather to highlight a few of the salient points that form the basis for current assumptions made in crop models.

Current atmospheres have a CO₂:O₂ ratio of 0.0018 but this may increase to values as high as 0.0047 by the end of this century (IPCC 2013), because CO₂ is currently increasing with an annual rate average of 2.1 μLL^{-1} (Dlugokencky and Tans 2017). This will benefit plants such as soybean that rely on C₃ photosynthesis. High atmospheric [CO₂] in FACE experiments resulted in increased soybean photosynthesis rates of up to 46 % (Leakey et al. 2009a). This enhancement is possible because the current atmospheric [CO₂] of 400 μLL^{-1} is insufficient to saturate the enzyme responsible for photosynthetic carbon assimilation, ribulose-1,5-bisphosphate carboxylase-oxygenase (Rubisco). Gaseous CO₂ is much more soluble in water than O₂, and thus the local CO₂:O₂ ratio in the chloroplast environment is currently about 0.026 at 25 °C. Rubisco has about a 100-fold greater affinity for CO₂ than O₂ in higher plants, dictating that this enzyme catalyses 2/3 cycles of carboxylation for every cycle of oxidation. In this way, carbon is partitioned between the assimilatory C₃ cycle and the photorespiratory pathways. Hence, higher CO₂:O₂ ratios will competitively inhibit the oxygenase activity of Rubisco and C₃ carbon fixation will be favoured over photorespiration. However, the potential benefits offered by increased carbon gain are often not fully realized because of insufficient sink capacity when C₃ plants are grown at elevated [CO₂] (Paul and Foyer 2001, Bernacchi et al. 2005). This results in carbohydrate accumulation in source leaves, a signal that causes

repression of genes encoding photosynthetic proteins leading to a down-regulation of photosynthesis and a decrease in leaf nitrogen (N) content (Leakey et al. 2009a). Limitations in soil nitrate availability can also lead to down-regulation of photosynthesis in plants grown at elevated [CO₂]. The “progressive N limitation” hypothesis suggests that under CO₂ enrichment, plant N uptake from soils fails to keep pace with photosynthesis and shoot carbohydrate accumulation (Foyer et al. 2009). A decline in soil quality is accompanied by increases in microbial immobilization because of high C/N ratios, a decline in soil N availability so that plants become increasingly N-limited, and as a result there is a decrease in photosynthesis (Foyer et al. 2009). It has also been argued that photorespiration plays an important role in providing the reductant required to drive the assimilation of nitrate into ammonium (Rachmilevitch et al. 2004). Hence, that increasing [CO₂] will favor C3 plants particularly in environments where NH₄⁺ is available as a nitrogen source.

The effects of increasing [CO₂] on plant architecture and partitioning of biomass between roots and shoots remains uncertain. Much depends on the C/N balance in roots and shoots. N-availability signals in the shoot influence the root system. The shoot promotes root growth in proportion to total N-demand. Plant architecture responses to increasing [CO₂] are likely to involve complex pathways of root-to-shoot and shoot-to-root signaling. Signaling molecules include the small C-TERMINALLY ENCODED PEPTIDE (CEP) family peptides, which control root system architecture in a non-cell-autonomous manner (Mohd-Radzman et al. 2015). In N-deprived roots CEP peptides are produced and transported to the shoot, where they induce expression of ‘CEP-DOWNSTREAM’ peptides that are transported back to the root to increase the expression of N-uptake transporters. There is a paucity of literature to date concerning how high [CO₂] influences whole plant signaling.

One particularly important result of the growth of C3 plants under elevated CO₂ is the priming of pathogen defenses (Mhamdi and Noctor 2016). Multiple pathogen defense pathways are activated when C3 plants are grown with atmospheric CO₂ enrichment, leading to increased resistance to bacterial and fungal pathogens. This high [CO₂]-dependent priming of pathogen defenses is linked to metabolic adjustments involving redox signaling (Mhamdi and Noctor 2016). While growth under elevated [CO₂] may enhance the resistance/resilience of C3 plants to pests and pathogens, a FACE study showed no effects on aphid performance (Mondor et al. 2005).

C4 plants such as maize are able to concentrate CO₂ in the Rubisco-containing photosynthetic cells of the bundle sheath. The CO₂-concentrating mechanism allows high rates of photosynthesis to occur even when stomata are closed while limiting flux through the photorespiratory pathway. Hence the C4 pathway of photosynthesis provides a competitive advantage under growth conditions that promote carbon loss through photorespiration, such as high temperatures or decreased water availability (Lopes and Foyer 2011). The transpiration rates and water status of maize leaves, particularly the older leaf ranks, are changed under conditions of atmospheric CO₂ enrichment even when plants are maintained under well-watered conditions (Prins et al. 2010). Under well-watered conditions, elevated CO₂ has little effect on the photosynthesis or growth of C4 plants in controlled environment (Soares et al. 2007, Prins et al. 2010) or in the FACE studies (Leakey et al. 2009a,b, Manderscheid et al. 2014). Moreover, the negative impact of drought on yield is attenuated at high CO₂ because of stomatal closure (Lopes et al. 2011, Manderscheid et al. 2014). Such observations indicate that maize should perform better under drought stress conditions when plants are grown at high [CO₂]. While higher temperatures should favor C4 plants over C3 plants (Long and Ort 2010), a negative response of global yields has been projected for maize as well as wheat and barley as a result of increased temperatures (Tatsumi et al. 2011, Asseng et al. 2014). Elevated temperatures have been reported to exert a negative influence on a range of plant processes such as photosynthesis through decreased activation of Rubisco, stomatal closure, flower development, pollen viability and hence fertility, and fruit ripening but in many cases the precise mechanisms remain to be characterised.

THE RISE OF CROP MODELLING

Crop models are designed to calculate crop yield (and other important parameters of the soil-plant system) as a function of weather and soil conditions, plant-specific characteristics as well as a choice of agricultural management practices. Models of cropping systems were first conceived of in the 1960s (Jones et al. 2016). Although it is fundamentally a curiosity-driven activity, the development of crop models received major boosts from various economic, technological and political events. During the Cold War, fueled by the unexpected large volume purchase of wheat by the Soviet Union in 1972, another type of curiosity played an important role in the development of key components of the DSSAT model suite (Jones et al. 2003) enabling the USA to predict the yield of major crops produced and traded worldwide, especially

in the COMECON (Council for Mutual Economic Assistance) countries (Ritchie 2000). Governmental funds helped experts from different disciplines to develop crop models with new capabilities: EPIC (Williams et al. 1989) with a soil erosion module, APSIM (Keating et al. 2003) able to simulate large number of different crops including trees and weeds. The release of the first personal computers in the early 1980s revolutionized not only the use and development of crop models but it led to many innovations in other fields (computer graphics, statistical analysis, GIS, etc.) that have contributed to the modeling of agricultural systems (Jones et al. 2016).

Crop modelling has been used for various applications over the past few decades. Field-scale applications for decision support have a long history (Hoogenboom et al. 1994) that in turn enabled work with seasonal weather forecasting (Hansen 2005), frameworks to link crop and climate models (Challinor et al. 2003), or integrated assessments within watersheds or across multiple sectors (Warszawski et al. 2014, Wriedt et al. 2009). Crop models have been used to develop adaptation options (Webber et al. 2014, Challinor 2009) and there is now recognition of the need for combined assessments of adaptation and mitigation, in support of achieving emissions targets (Jarvis et al. 2011, Shirsath et al. 2017). The need to quantify uncertainty (Challinor et al. 2013) and to improve models has led to an increasing number of international collaborations across modelling groups (Rosenzweig et al. 2013), and to work supporting the use of crop models with climate model ensembles (Ramirez-Villegas et al. 2013). Recognition of the importance of vulnerability and agricultural management in determining impacts and adaptation options has led to work across the natural-social science interface (Simelton et al. 2012). For a detailed history of crop models see the comprehensive work of Jones et al. (2016).

MAJOR TYPES OF CROP MODELS

Approaches used to assess the impacts of climate change on agriculture include four major types. 1) Climate or more generally, environmental index-based methods (Olesen et al. 2011) utilize a multidimensional scoring system of production determining factors to provide a quasi-quantitative assessment of the vulnerability of the investigated agricultural system or area. 2) Statistical models express the relationship between yield or yield components and weather parameters in a form of regression equations (Lobell and Burke 2010) or other type of more “black-box” models (Delerce et al. 2016) which are calibrated by using corresponding observed yield and weather data varying in time or space or in both domains. 3) Niche-based models

describe the geographical distribution of a crop species using either a set of explicit fuzzy-logic equations that describe the crop's response to various environmental factors (Zabel et al. 2014) or a statistical model fitted with presences and absences (Estes et al. 2013). 4) Process-based models (Rosenzweig et al. 2014, Ewert et al. 2015, Müller et al. 2017) are the mathematical (and nowadays usually computer-based) representation of the most important processes of the soil-plant system consisting of a set of ordinary or partial differential equations and empirical equations organised into procedures or modules where the outputs of one procedure can serve as input to other procedures and the model as a whole is able to describe the temporal pattern of the key system parameters. That is why these models are also called crop simulation models. Each type of model has advantages and disadvantages as well as limitations. However, all are useful tools in considerations of the potential impacts of climate change. Researchers select the model that best suits the application. From the point of view of the present question, the major limitations of the first three approaches are that they cannot capture future climate-soil-crop relationships, adaptation and carbon dioxide fertilization effect, though there are techniques to estimate the latter in statistical methods (McGrath and Lobell 2011). Probably this is the main reason why process-based crop models are the most commonly used tools for climate impact assessments (White et al. 2011).

STATE OF THE ART OF CROP MODELLING

The capabilities of crop models depend in large part on the observed data used for developing and testing the model, and on modelling the crop at a degree of complexity that is appropriate to the aims of the study (Sinclair and Seligman 2000). The results of any one particular study are highly dependent upon input data quality and adequate quantification of uncertainty, though synthesis across many studies helps achieving consensus (Challinor et al. 2014b). Crop model ensembles should represent the underlying distribution of probabilities, which is not straightforward (Wallach et al. 2016). Attention should be paid to bias correction of climate data where necessary (Hawkins et al. 2013). The assumptions underlying the results of the study should be made explicit, for example using a common uncertainty reporting format (Wesselink et al. 2015). For adaptation, there are number issues that need attention when formulating a study (see Lobell 2014).

Whilst the spread of results produced by crop models has increased over time, robust conclusions can still result from analysis of outputs (Challinor et al. 2014b). Crop models are

increasingly used for global assessments (Rosenzweig et al. 2014). There are currently two large modelling initiatives, AgMIP (agmip.org) and Modelling European Agriculture for Climate Change (MACSUR: macsur.eu). These networking hubs coordinate and support crop model development, together with crop model based studies and impact assessments, providing information for producers, policy-makers and the public in the area of integrated climate change risk assessment for global agriculture and food security. The [projections described for maize and soybean below are the derived from integrated MACSUR model development and application.

Understanding the influence of land use on crop production is an important challenge for such studies (Challinor et al. 2015). Effective use of crop models within integrated assessment models is another important challenge (Ewert et al. 2015). Coupling crop models with general equilibrium models to bring demand and supply together and consider global trade as a major driver of future land use change is another step forward in the evolution of crop models (Mauser et al. 2015). These challenges for the use of crop models do nothing to detract from the need for continued model improvement and representation of processes (Hollaway et al. 2012, Challinor et al. 2014a), particularly where experimental limitations occur (Reich and Hobbie 2013).

PROJECTIONS FOR THE FUTURE OF C3 AND C4 CROPS, FOCUSING ON SOYBEAN AND MAIZE

Crop models have been widely used to estimate the potential impacts of climate change on future agricultural productivity. The protocols of the assessments vary to such an extent that they impose serious limitations to cross-study syntheses and increase the potential for bias in projected impacts (White et al. 2011). Despite this fact, the available results allow us to draw some robust conclusions that are outlined below. With the help of the SCOPUS database, we reviewed 118 peer-reviewed publications (31 for soybean and 87 for maize) that used crop models to investigate the impact of climate change on the production of maize and soybean worldwide in the second half of the 21st century. These modelling studies covered all the most important production areas in America, Asia, Europe and Africa. Using these studies, we draw conclusions on model inputs, consideration of [CO₂] response, adaptation and mitigation for both crops.

MODELS AND KEY MODEL INPUTS IN THE SOYBEAN STUDIES

Fifteen different models were used to assess the potential climate change impacts on soybean. However, only two models were used in more than 2 studies. CROPGRO and EPIC model results were reported in 15 and 4 papers, respectively. 17 studies investigated more than one location (from 2 to 100) within the study area (point-based studies) and 11 studies used the gridded modelling approach covering the total investigated area with a specific spatial resolution. No studies used gridded and point-based estimates jointly. Regarding uncertainty quantification, only 2 papers used more than one crop model, though this technique helps avoiding model related biases in the climate change impact projections. Conversely, with the exception of two studies, all used several (2-72) future climate projections to assess (or show) the uncertainty arising from different climate model- and/or climate change scenario-related issues. The projected temperature rise used in the climate projections (compared to the baseline) varied between 0.9 and 9 °C, but the majority of the studies examined the effect of a 2-4 °C temperature rise. These temperature changes were associated with an increase in the atmospheric [CO₂] from 450-700 μLL^{-1} , although the majority of the papers postulated a [CO₂] of 550-650 μLL^{-1} for the future.

THE EFFECT OF HIGH [CO₂] AND ADAPTATION OPTIONS ON FUTURE SOYBEAN PRODUCTION

Of the literature use in this analysis, six studies failed to consider the direct effect of high [CO₂] on soybeans. All studies projected yield losses for soybean, which might be mitigated by agricultural management adaptations such as changing the planting date (do Rio et al. 2016), change of cultivars (Battisti et al. 2017) or introducing double-cropping systems (Lant et al. 2016). The global study of Tatsumi et al. (2011) projected yield decrease for all the major soybean producing areas. However, this study applied several significant simplifications such as use of monthly step climatic data, ignoring CO₂ fertilisation effects and the water holding capacity of soils. Twenty seven studies that took into account CO₂ fertilisation effects projected significant yield gains. Of these, only one global assessment that took into account the added carbon gain arising from future high atmospheric [CO₂], projected moderate (5-15%) yield losses and this was only for regions in US and Latin-America (Deryng et al. 2014). The same study did not investigate the potential of management adaptation options. In relation to adaptation, in fact, we find that some 16 out of the 37 studies investigated adaptation options. These studies suggest that management adaptation options can have a significant effect in

counterbalancing the negative effects of climate change (Tubiello et al. 2000, Challinor et al. 2014b). According to some simulations, some crop management options (e.g. winter rye cover) have no effect on future yields but they have the potential to reduce soil erosion and nitrous oxide emissions significantly (Basche et al. 2016).

THE ROLE OF CLIMATE CHANGE MITIGATION POLICIES IN FUTURE SOYBEAN PRODUCTION

Representative Concentration Pathways (RCPs) are four greenhouse gas concentration trajectories (IPCC, 2013), all of which are plausible depending on how much greenhouse gases (GHG) are emitted in the years to come. The four RCPs, RCP2.6, RCP4.5, RCP6, and RCP8.5, are named after the prospective radiative forcing values in the year 2100 relative to pre-industrial values (+2.6, +4.5, +6.0, and +8.5 Wm⁻², respectively). The ultimate aim of climate change mitigation policies is to reduce emissions consistent with specific targets, thus helping to avoid high-end emissions scenarios such as RCP8.5. The Paris Agreement (2015), for example, aims at maintaining global average temperature well below 2 °C above pre-industrial levels; this has been reported to significantly reduce the risks and impacts of climate change (Schleussner et al. 2016). This aim could be achieved in many ways including the use of low-carbon technologies, renewable energy sources, transportation optimization, as well as promoting individual-lifestyle changes (cycling instead of driving, alternative diets, etc.). In the agricultural sector, climate change mitigation policies may be implemented via promoting reforestation, low input soil management, resource efficient farm management, more sustainable fertiliser subsidy provision, and improving knowledge and transfer mechanisms all aiming at increasing carbon sequestration and/or decreasing GHG emissions. Climate mitigation policies have an important role in ensuring the implementation of technologies, meeting mitigation targets, ultimately helping avoid global yield losses.

Across the soybean studies reviewed here, mitigation policies are typically addressed by modelling crop yields for different RCPs. Comparison between different RCPs allows determination of the likely benefits of climate change mitigation. For example, the yield reduction reported by Deryng et al. (2014) was the result of using the most extreme RCP8.5 based climate projections which is in fine agreement with the findings of Bhattarai et al. (2017) who, on the other hand, used not only RCP8.5 but RCP2.6 and RCP4.5 based projections resulting in marginal yield losses (-2%) for RCP8.5 and yield gains (11 and 13%) for RCP2.6

and RCP4.5, respectively. The studies reviewed thus indicate that successful climate change mitigation policies that secure the future [CO₂] pathway below RCP4.5, will allow future resolution of soybean production problems.

Another important aspect of future crop production is the extent to which areas where crops are grown may shift as conditions change. Some studies have shown that land that is suitable for soybean production displays a large northward shift (Lant et al. 2016). This shift incorporates significant areas of the Northern hemisphere reaching as far as Ireland (Holden and Brereton 2003). Soybeans are already grown in Canada and varieties are already being trialled for production in the UK. Thus, due to the projected future yield and sowing area gains an expansion of soya production could be expected worldwide, although as with projected yield changes, these shifts in production areas could change depending upon the emissions pathway.

MODELS AND KEY MODEL INPUTS IN THE MAIZE STUDIES

Twenty one different models were used for assessing the potential climate change impacts on maize. The two most frequently used models were the CERES and EPIC that were used in 45 and 8 studies, respectively. About a third (23) of the assessments were based on data of only one particular site of the study area and/or applied only one climate projection for the future. The projected temperature rise and the associated atmospheric carbon dioxide increase of the climate projections of the maize studies were similar to those of the soybean studies. Regarding crop model uncertainty, twenty two studies used the gridded modelling approach and five papers used more than one crop model for the impact assessments. The most comprehensive of these was the study of Bassu et al. (2014), which evaluated 23 maize simulation models for four locations representing a wide range of maize production conditions in the world. They found that only an ensemble of models (a minimum of about 8 to 10 needed) was able to simulate absolute yields accurately and that there was a large uncertainty in the yield response to [CO₂] among models. The uncertainty envelope is mainly due to inconsistency in the way models simulate assimilation, as well as in whether or not models simulate enhanced [CO₂] effects on transpiration.

MODEL AND SCALE RELATED UNCERTAINTY IN THE MAIZE STUDIES

In a global study, Blanc and Sultan (2015) showed that the projected changes for maize production were highly model-dependent, ranging from a 15% decrease to a 20% increase in yield in the Corn Belt. However, large scale investigations may contradict local (country scale) studies even if the same model was used. For example, Supit et al. (2012) projected a yield increase for Turkey as a result of climate change while Sen et al. (2012) predicted that yields will decrease in this region. One reason for this kind of discrepancy could be the lack of use or quality in the soil data used for yield projections (Tatsumi et al. 2011). The impact of climate change on specific regions could vary significantly because of differences in soil characteristics (Chipanshi et al. 2003). Surprisingly, no local model-based impact studies were found for France, Indonesia, Ukraine or South-Africa, although these countries are among the top 10 global maize producers.

PROSPECTS FOR FUTURE MAIZE PRODUCTION

While a number of studies have predicted increases in maize yields in the major corn-producing areas of the world such as the USA (Tubiello et al. 2002), China (Guo et al. 2010) and Argentina (Travasso et al. 2009), most studies have projected global decreases in maize yields (Lin et al. 2015, Supit et al. 2012, Byjesh et al. 2010, Schlenker and Roberts 2009; Deryng et al. 2014), even in studies that took the beneficial effect of CO₂ fertilisation into account. Many studies accounted the predicted yield reduction by one or more of the three main reasons: 1) Increasing frequency and severity of drought; 2) Increasing risk of heat waves around flowering; 3) Shortening of the vegetation period. However, it may also be the case that current models fail to account for the water saving mechanisms afforded by C₄ metabolism and physiology appropriately. Higher water use efficiencies would be expected in maize under high [CO₂]. Thus, models failing to take this feature into account might underestimate biomass and yield gains under high [CO₂]. Durand et al. (2017) assessed the accuracy of maize crop models in simulating the interactions of changes at high atmospheric [CO₂]. Under well-watered conditions the models were able to reproduce the absence of yield response to elevated [CO₂]. However, under water deficit conditions the models failed to capture the extent of the [CO₂] response that was observed in the field.

Regional gridded modelling studies are particularly important in maize yield projections because they are able to distinguish between sub-regions that may be positively or negatively affected by climate change. The currently high yielding sub-regions of China may face yield

decreases while the current low yielding sub-regions may expect yield increase (Xiong et al. 2007). Current high yielding sub-regions are near-optimum zones providing very favourable conditions for maize production. Almost any environmental change in these areas could only be negative as it would distance the system from its near-optimum state. On the other hand, marginal areas (far from the optimum) most likely benefit from the environmental changes, by getting closer to the optimum state of the system. However, yield losses per unit area do not necessarily translate into overall productivity for a given region, because the projected area of cultivated land used for multiple-cropping systems may be significantly increased as a result of climate change (Yang et al. 2015). Moreover, the indirect effects of climate change can become important for example the projected increases in insect pests as a result of increased winter survival (Diffenbaugh et al. 2008). Such factors could significantly alter the pest management landscape of North American maize production, leading to substantial economic impacts through increased seed and insecticide costs, as well as decreased yields.

ROLES FOR ADAPTATION OPTIONS AND CLIMATE CHANGE MITIGATION POLICY IN FUTURE MAIZE PRODUCTION

Modelling studies do not depict a clear positive or negative picture for future global maize production but they clearly emphasize the need for explicit adaptation actions such as breeding of heat/drought tolerant hybrids. The majority of the studies (13 out of 20) that assessed certain adaptation options concluded that a shift in planting date, together with the use of longer maturing hybrids and alternative soil and nitrogen management practices will be insufficient to counter negative impacts of climate change (Tubiello et al. 2000, Ko et al. 2012, Moradi et al. 2013). Studies also agree that the more extreme the scenario (RCP8.5 or similar scenarios from the earlier IPCC reports) the more severe the yield losses that could be expected. This highlights the necessity and opportunities for joint mitigation-adaptation efforts. A global study suggest that the drastic climate mitigation policy of RCP2.6 could avoid more than 80% of the projected global average yield losses (USA: -20%, Brazil: -50%, Argentina: -40%) that are otherwise projected by the 2080s under RCP8.5 (Deryng et al. 2014).

PROJECTED LAND USE CHANGES FOR MAIZE AND SOYBEAN PRODUCTION BY 2100

Coupling land use (Monfreda et al. 2008) and baseline and future land suitability data (Zabel et al. 2014) with future diet (Tilman and Clark 2014) and GHG emission (Smith et al. 2008) scenarios we projected future of global maize (Fig. 1) and soybean (Fig. 2) production areas. a Baseline (1981-2010) and future (2071-2100) land suitability determinations for each grid cell were made using the methodology of Zabel et al. (2014) by incorporating data on local climate, soil and topography. The ‘No Change’ scenario is the extrapolation of the current trends i.e. assuming that no major GHG emission reductions will be achieved by the introduction of mitigation policies or enhanced climate-smart agro-technologies. Moreover, the scenario predicts that increases in income and urbanization will drive a global dietary transition that involves increasingly higher consumption of refined sugars, fats, oils and meats (Tilman and Clark 2014). Together, these features will result in increased demands for maize and soybean production. In contrast, the ‘Major Change’ scenario envisages successful and effective GHG mitigation policies, together with the instigation of new GHG emission reducing agricultural practices. Together with significant health-driven changes in diets and adoption of alternative diets such as Mediterranean, pescetarian or vegetarian diets that are characterised by higher consumption of fruits, vegetables and pulses and a lower meat consumption (Tilman and Clark 2014), these will result in an decreasing demand for maize and soybean. Using these scenarios global crop production area maps were created in a 10 km (5 arc minute) spatial resolution. According to current land use (LU) given by (Monfreda et al. 2008) each grid cell can have two states: used (harvested area fraction of the crop is at least 1% of the grid cell area) and not used.

The crop production scenarios reported here predict significantly different demands for land use for maize (Fig. 1) and soybean (Fig. 2) production. Both the used and the not used cells may remain in the same land use category or may be changed in the future providing four options that can be defined by certain rules for both scenarios (Table 1). If land is ‘used’ today according to the definitions used above, we assume that these areas will be unaltered in the future (2071-2100) in the ‘Major Change’ scenario, if the suitability increases by at least 10 %. If suitability increases less than 10 % or decreases until 2071-2100, we assume that these areas will be abandoned and not be used in the future.

Crucially, areas that are currently not used for maize (Fig. 1B) and soybean (Fig. 2B) production will probably be added if future land suitability is higher than the 67th percentile of today’s global suitability on used areas. Conversely, areas that are currently not used will also not be used in the future if suitability is lower than the 67th percentile. Since demands for soybean and

maize production are higher in the 'No Change' scenarios than in the 'Major Change' scenarios, more areas will be required for the production of these crops. Accordingly, we assume lower thresholds for future land suitability, as well as a lower percentiles of suitability on today's production areas for maize and soybean respectively. Hence, greater areas of marginal land will have to be used for the cultivation maize and soybean in order to fulfil the increasing demands.

CONCLUSIONS AND PERSPECTIVES

Future land use maps were created for maize and soybean using the basic rules outlined in Table 1 (Fig. 1). Major changes in policy, agricultural practice and diet imply that major shifts will occur in the area used for maize and soya production. Our assessment of modelling outputs predicts that large portions of current areas of significant maize and soya production may be abandoned from in the future. On the other hand, large new areas will become available in the future (Table 2) in order to meet the increasing demands on maize and soya production, particularly if no significant policy, agro-technological and diet-related changes take place in the future. According to the projections Europe will face major challenges in both production scenarios, especially in case of maize. Aligned to other studies (Ruiz-Ramos and Mínguez 2010, Supit et al. 2012, Fodor et al. 2014, Mihailović et al. 2015) a stern warning sign could be given to the European Union that effective adaptation actions are required to mitigate the harmful impacts of climate change across the continent. At the other end of the spectrum is Africa, where climate change may allow a massive increase in soybean production no matter which production scenario becomes a reality in the future. This it is not surpassing that soybean is called Africa's Cinderella crop (Kolapo, 2011). The studies that were assessed here predict a more promising future for soybean, particularly in terms of production areas, gained and abandoned (Table 2). These crop models provide essential underpinning information to farmers, agro-industries and policymakers, so that appropriate cropping systems and/or management practices can be put in place to counter global climate change.

Crop models also have an important role to play in informing plant scientists and breeders of essential traits that must be developed in future crop varieties. However, many current models are not based on current knowledge of plant responses to elevated atmospheric [CO₂] and they do not incorporate the latest findings about how crops respond to a changing climate. There is therefore an urgent need for a new interface of information exchange between crop modellers

and plant scientists highlighting weaknesses and overlooked processes, and to influence how models are built, to include how recent changes in our understanding of [CO₂]-mediated effects on plants might be formalised and incorporated into models. It is thus timely to renew discussions in order to remove the large uncertainties and biases in some current crop models, as well as informing plant scientists of the essential underpinning traits that will ensure food security over the next 50 years. Current crop varieties are not well suited to future unpredictable weather patterns caused by climate change. Modern breeding programs have selected for dwarf shoot systems, minimizing the production of vegetative tissues. Moreover, elite crop varieties are developed and bred under ideal growth conditions so the selective pressure for plant performance under sub-optimal conditions has largely been removed. This has favoured small root systems, a trait that may have inadvertently decreased the resilience of plants to both abiotic and biotic stresses, which are likely to increase as a result of climate change.

Finally, plant physiologists should be aware of areas where collaboration and data generation would greatly assist crop modellers:

1. Grain quality aspects: While FACE experiments clearly indicate that CO₂ enrichment affects grain quality characteristics that are important for consumer nutrition and health, and for industrial processing and marketing (Högy et al. 2009), CO₂ enrichment effects on grain quality traits remain poorly characterised in terms of metabolite, proteome and transcript profiles. Some field-scale crop models already include yield quality related outputs, including sugar and acid concentrations (Bindi and Maselli, 2001), grain protein (Asseng et al. 2002) and grain protein composition (Martre et al. 2006) protein composition. The embedded yield quality calculation methods are not thoroughly tested, especially not by using data from elevated CO₂ experiments. While manipulation of some of the enzymes of primary carbon assimilation was found to protect soybean seed yields against the negative effects of elevated temperature on plants grown at high CO₂ (Köhler et al. 2016), there are no comparable studies in the literature on effects on grain quality.
2. More accurate vegetation-related to CO₂ fluxes: An important aspect of the crop simulation models typically used for climate change impacts assessments is that they harness important, widely validated knowledge on crop responses to biotic and abiotic factors (Boote et al. 2013). Recent progress in crop, ecosystem, and climate modelling has led to integration of these disciplines in support of integrated assessments of agro-ecosystems at the global or regional level (e.g. Osborne et al. 2007, 2015, Wang et al. 2005). In these cases, crop models may provide the underlying information, parameters

and mathematical formulations that underpin the vegetation models used. Nevertheless, much work remains to be done in crop simulation models if these are to be fully integrated within vegetation models. Foremost, adequately simulating vegetation within complex agro-ecosystems requires detailed consideration of CO₂ uptake for gross primary productivity and CO₂ release through respiration (Cramer et al. 2001). While progress has been made in developing and testing leaf-to-canopy assimilation in some crop models, only a handful of models for the major crops, including maize and soybean, include detailed photosynthesis-respiration routines for both assimilation and CO₂ fertilisation (Bassu et al. 2014, Li et al. 2015). Moreover, respiration costs for production of new or maintenance of existing tissue (growth and maintenance respiration) are either highly uncertain or not estimated or reported in crop simulation studies. Furthermore, testing of CO₂ fluxes or canopy assimilation using eddies of air, although feasible, is rarely if at all conducted for crop simulation models (Hollinger et al. 2005, Paul et al. 1999). Finally, appropriate consideration and validation of CO₂ fluxes in crop models will also help improving water fluxes and evapotranspiration, which is a key source of uncertainty in crop simulation (Liu et al. 2016).

3. Canopy temperature and evapotranspiration: The importance of models predicting global warming effects on crop yield to include canopy temperature instead of using air temperature was demonstrated by Julia and Dingkuhn (2013). They found that rice panicle temperature varied between 9.5 below and 2 °C above air temperature at 2 m depending on the microclimate and therefore heat stress causing sterility was more likely to occur in warm-humid than hot-arid environments due to humidity effects on transpiration cooling. Even though some crop models calculate canopy from air temperature, which is then used on some but not necessarily all temperature-related processes in the crop model, Webber et al. (2015) found that this did not necessarily improve yield simulations. The study compared nine process based crop models that used three different approaches of simulating canopy temperature (empirical, energy balance assuming neutral atmospheric stability, and energy balance correcting for the atmospheric stability conditions) in their ability to simulate heat stress in irrigated wheat in a semi-arid environment. They found that for all models the reduction in RMSE was larger if canopy temperature was only used for the processes simulating heat stress but that using canopy temperatures for all processes did not necessarily improved yield simulations. Models that performed well in simulating yield under heat stress had varying skill in simulating canopy temperature (the method energy balance assuming

neutral atmospheric stability performed worst). Models differ in parameter values which might be able to somewhat alleviate the impact from using air temperature. Unfortunately the models could not be tested with observed canopy temperature as it was not measured continuously throughout the growing season. Webber et al concluded that a more systematically understanding of heat stress events and how to model them is needed.

4. Effects of high ozone concentrations: Ozone is highly phytotoxic and can cause significant damage to vegetation and crops even at current concentrations in many parts of the world (Mills et al. 2011, Booker et al. 2009, Hollaway et al. 2012, Wang and Mauzerall 2004). Both maize and soybean are sensitive to ozone (McGrath et al. 2015), with predicted global yield losses ranging from 2.5 - 8% for maize and 9.5 – 15% for soybean for the year 2030 (Avnery et al. 2011). However, the negative effects of ozone are included only in a few crop models. For example, the WOFOST model accounts for ozone damage to crops by using a flux-based approach in which the ozone flux inside the plant is regulated by the stomatal conductance (Cappelli et al. 2016). The model shows that for wheat there are large yield losses under high ozone exposure (i.e. up to 30% loss for ozone concentration of 60 ppb; Cappelli et al. 2016). While the effects of ozone on plant biology have been extensively studied, the effect of pollution on crop productivity and quality is an important area for future work, particularly as global ozone concentrations are projected to remain at high levels (Fowler et al. 2008). The responses of plants to atmospheric ozone should be assessed in combination with other stresses to address current as well as the future responses under climate change.
5. Acclimation to elevated CO₂: current knowledge of how plants sense and signal changes in atmospheric [CO₂] over and above effects on photosynthesis, is limited. Moreover, much remains uncertain concerning the mechanisms that define many of the observed plant responses to increased atmospheric [CO₂] or how these mechanisms will influence biotic and abiotic stress responses under field conditions. In particular, relatively little is known about how high [CO₂] will influence the soil microbiome or plant interactions with beneficial fungi and bacteria.

Table 1. Rules of projections of future of crop production areas. LSt: Land Suitability today (1981-2010); LSf: Land Suitability in the future (2071-2100); PERC33(LSt) and PERC67(LSt): 33rd and 67th percentile of the distribution of the LSt values of the grid cells used for maize/soya production over the global grid. LU denotes Land Use.

LU today	used	used	not used	not used
LU change	unaltered	abandoned	added	unaltered
LU in the future	used	not used	used	not used
Scenario	No Change			
Rule	if $LSf > 0.9 \times LSt$	if $LSf \leq 0.9 \times LSt$	if $LSf > PERC33(LSt)$	if $LSf \leq PERC33(LSt)$
Acronym	NoCh_Used	NoCh_Aband	NoCh_Added	NoCh_Notused
Scenario	Major Change			
Rule	if $LSf > 1.1 \times LSt$	if $LSf \leq 1.1 \times LSt$	if $LSf > PERC67(LSt)$	if $LSf \leq PERC67(LSt)$
Acronym	MaCh_Used	MaCh_Aband	MaCh_Added	MaCh_Notused

Table 2 Predicted global gains and abandoned areas of maize and soya production. . The ‘no change’ scenario is the extrapolation of the current trends with no major GHG emission reductions or no major changes in dietary trends that would result in an increasing need for maize or soybean production. The ‘Major Change’ scenario will be attained if successful GHG mitigation policies are enforced and significant health-driven changes in diets occur that result in a decreasing need for maize or soybean production.

Scenario	Transition	Acronym (see Fig. 1)	maize [km ²]	soya [km ²]
No change	Abandoned	NoCh_Aband	3 364 115	299 005
	Added	NoCh_Added	27 740 977	30 524 853
Major change	Abandoned	MaCh_Aband	13 287 592	6 506 380
	Added	MaCh_Added	10 137 774	6 547 211

Figure 1. Current maize growing areas (blue), together with predicted abandoned (red) and added (green) maize growing areas by 2100. The ‘no change’ scenario (A) is the extrapolation of the current trends with no major GHG emission reductions or no major changes in dietary trends that would result in an increasing need for maize production. The ‘Major Change’ scenario (B) will be attained if successful GHG mitigation policies are enforced and significant health-driven changes in diets occur that result in a decreasing need for maize production.





Figure. 2 Current soybean growing areas (blue), together with predicted abandoned (red) and added (green) soybean growing areas by 2100. The ‘no change’ scenario (A) is the extrapolation of the current trends with no major GHG emission reductions or no major changes in dietary trends that would result in an increasing need for soybean production. The ‘Major Change’ scenario (B) will be attained if successful GHG mitigation policies are enforced and significant health-driven changes in diets occur that result in a decreasing need for soybean production.



ACKNOWLEDGEMENTS

NF, AC and CF thank BBSRC for financial support (BB/N004914/1). ID thanks ‘Oatley PhD Scholarship’ for financial support.. JRV and AJC are supported by the CGIAR Research Program on Climate Change, Agriculture and Food Security (CCAFS). CCAFS is carried out with support from CGIAR Fund Donors and through bilateral funding agreements. For details please visit <https://ccafs.cgiar.org/donors>. The views expressed in this paper cannot be taken to reflect the official opinions of these organizations. The results were obtained within an international research project named “FACCE MACSUR – Modelling European Agriculture with Climate Change for Food Security, a FACCE JPI knowledge hub” and acknowledge the respective national or regional funding organizations. The research was supported by the Széchenyi 2020 programme, the European Regional Development Fund - "Investing in your future" and the Hungarian Government (GINOP-2.3.2-15-2016-00028).

REFERENCES

- AbdElgawad, H., Zinta, G., Beemster, G.T.S., Janssens I.A. and Asard H. (2016) Future Climate CO₂ Levels Mitigate Stress Impact on Plants: Increased Defense or Decreased Challenge? *Frontiers in Plant Science*. 7: 1–7.
- Ainsworth, E.A. and Long, S.T. (2005) What have we learned from 15 years of free-air CO₂ enrichment (FACE)? A meta-analytic review of the responses of photosynthesis, canopy properties and plant production to rising CO₂. *New Phytologist*. 165: 351–372.
- Asseng, S., Bar-Tal, A., Bowden, J. W., Keating, B.A., Van Herwaarden, A., Palta, J.A., Huth, N.I. and Probert, M.E. (2002) Simulation of grain protein content with APSIM-Nwheat. *European Journal of Agronomy*. 16: 25–42.
- Asseng, S., Ewert, F., Rosenzweig, C., Jones, J.W., Hatfield, J.L., Ruane, A.C., Boote, K.J., Thorburn, P.J., Rötter, R.P., Cammarano, D., Brisson, N., Basso, B., Martre, P., Aggarwal, P.K., Angulo, C., Bertuzzi, P., Biernath, C., Challinor, A.J., Doltra, J., Gayler, S., Goldberg, R., Grant, R., Heng, L., Hooker, J., Hunt, L.A., Ingwersen, J., Izaurralde, R.C., Kersebaum, K.C., Müller, C., Naresh Kumar, S., Nendel, C., O’Leary, G., Olesen, J.E., Osborne, T.M., Palosuo, T., Priesack, E., Ripoche, D., Semenov, M.A., Shcherbak, I., Steduto, P., Stöckle, C., Stratonovitch, P., Streck, T., Supit, I., Tao, F., Travasso, M., Waha, K., Wallach, D., White, J.W., Williams, J.R. and Wolf, J. (2013) Uncertainty in simulating wheat yields under climate change. *Nature Climate Change*. 3: 827–832.
- Asseng, S., Ewert, F., Martre, P., Rötter, R.P., Lobell, D.B., Cammarano, D., Kimball, B.A., Ottman, M.J., Wall, G.W., White, J.W., Reynolds, M.P., Alderman, P.D., Prasad, P.V.V., Aggarwal, P.K., Anothai, J., Basso, B., Biernath, C., Challinor, A.J., De Sanctis, G., Doltra, J., Fereres, E., Garcia-Vila, M., Gayler, S., Hoogenboom, G., Hunt, L.A., Izaurralde, R.C., Jabloun, M., Jones, C.D., Kersebaum, K.C., Koehler, A.-K., Müller, C., Naresh Kumar, S., Nendel, C., O’Leary, G., Olesen, J.E., Palosuo, T., Priesack, E., Eyshi Rezaei, E., Ruane, A.C., Semenov, M.A., Shcherbak, I., Stöckle, C., Stratonovitch, P., Streck, T., Supit, I., Tao, F., Thorburn, P.J., Waha, K., Wang, E., Wallach, D., Wolf, J., Zhao, Z. and Zhu, Y. (2014) Rising temperatures reduce global wheat production. *Nature Climate Change*. 5: 143–147.

Avnery, S., Mauzerall, D.L., Liu, J. and Horowitz, L.W. (2011) Global crop yield reductions due to surface ozone exposure: 2. Year 2030 potential crop production losses and economic damage under two scenarios of O₃ pollution. *Atmospheric Environment*. 45: 2297–2309.

Basche, A.D., Archontoulis, S.V., Kaspar, T.C., Jaynes, D.B., Parkin, T.B. and Miguez, F.E. (2016) Simulating long-term impacts of cover crops and climate change on crop production and environmental outcomes in the Midwestern United States. *Agriculture, Ecosystems and Environment*. 218: 95–106.

Bassu, S., Brisson, N., Durand, J.L., Boote, K., Lizaso, J., Jones, J.W., Rosenzweig, C., Ruane, A.C., Adam, M., Baron, C., Basso, B., Biernath, C., Boogaard, H., Conijn, S., Corbeels, M., Deryng, D., De Sanctis, G., Gayler, S., Grassini, P., Hatfield, J., Hoek, S., Izaurralde, C., Jongschaap, R., Kemanian, A.R., Kersebaum, K.C., Kim, S-H. Kumar, N.S., Mikowski, D., Müller, C., Nendel, C., Priesack, E., Pravia, M.V., Sau, F., Shcherbak, I., Tao, F., Teixeira, E., Timlin, D. and Waha, K. (2014) How do various maize crop models vary in their responses to climate change factors? *Global Change Biology*. 20: 2301–2320.

Battisti, R., Sentelhas, P.C., Boote, K.J., de S. Câmara, G.M., Farias, J.R.B. and Basso, C.J. (2017) Assessment of soybean yield with altered water-related genetic improvement traits under climate change in Southern Brazil. *European Journal of Agronomy*. 83: 1–14.

Bernacchi, C.J., Morgan, P.B., Ort, D.R. and Long, S.P. (2005) The growth of soybean under free air [CO₂] enrichment (FACE) stimulates photosynthesis while decreasing in vivo Rubisco capacity. *Planta*. 220: 434–446.

Bhattarai, M.D., Secchi, S. and Schoof, J. (2017) Projecting corn and soybeans yields under climate change in a Corn Belt watershed. *Agricultural Systems*. 152: 90–99.

Bindi, M., and Maselli, F. (2001) Extension of crop model outputs over the land surface by the application of statistical and neural network techniques to topographical and satellite data. *Climate Research*. 16: 237–246.

Bishop, K.A., Betzelberger, A.M., Long, S.P. and Ainsworth, E.A. (2015) Is there potential to adapt soybean (*Glycine max* Merr.) to future [CO₂]? An analysis of the yield response of 18 genotypes in free-air CO₂ enrichment. *Plant, Cell & Environment*. 38: 1765–1774.

Blanc, E. and Sultan, B. (2015) Emulating maize yields from global gridded crop models using statistical estimates. *Agricultural and Forest Meteorology*. 214–215: 134–147.

Booker, F., Muntifering, R., McGrath, M., Burkey, K., Decoteau, D., Fiscus, E., Manning, W., Krupa, S., Chappelka, A., and Grantz, D. (2009) The ozone component of global change: potential effects on agricultural and horticultural plant yield, product quality and interactions with invasive species. *Journal of Integrative Plant Biology*. 51: 337–351.

Boote, K.J., Jones, J.W., White, J.W., Asseng, S. and Lizaso, J.I. (2013) Putting mechanisms into crop production models. *Plant Cell & Environment*. 36: 1658–72.

Byjesh, K., Kumar, S.N. and Aggarwal, P. K. (2010) Simulating impacts, potential adaptation and vulnerability of maize to climate change in India. *Mitigation and Adaptation Strategies for Global Change*. 15: 413–431.

Cappelli, G., Confalonieri, R., Van Den Berg, M. and Dentener, F. (2016) Modelling inclusion, testing and benchmarking of the impacts of ozone pollution on crop yields at regional level. EUR 28395 EN. *Luxembourg: Publications Office of the European Union*. doi:10.2788/68501.

Challinor, A. (2009) Towards the development of adaptation options using climate and crop yield forecasting at seasonal to multi-decadal timescales. *Environmental Science & Policy*. 12: 453–465.

Challinor, A., Martre, P., Asseng, S., Thornton, P. and Ewert, F. (2014a) Making the most of climate impacts ensembles. *Nature Climate Change*. 4: 77–80.

Challinor, A.J., Watson, J., Lobell, D.B., Howden, S.M., Smith, D.R. and Chhetri, N. (2014b) A meta-analysis of crop yield under climate change and adaptation. *Nature Climate Change*. 4: 287–291.

Challinor, A., Slingo, J., Wheeler, T., Craufurd, P. and Grimes, D. (2003) Toward a combined seasonal weather and crop productivity forecasting system: Determination of the working spatial scale. *Journal of Applied Meteorology and Climatology*. 42: 175–192.

Challinor, A.J., Parkes, B. and Ramirez-Villegas, J. (2015) Crop yield response to climate change varies with cropping intensity. *Global Change Biology*. 21: 1679–1688.

Challinor, A.J., Smith, M.S. and Thornton, P. (2013) Use of agro-climate ensembles for quantifying uncertainty and informing adaptation. *Agricultural and Forest Meteorology*. 170: 2–7.

Chipanshi, A.C., Chanda, R. and Totolo, O. (2003) Vulnerability assessment of the maize and sorghum crops to climate change in Botswana. *Climatic Change*. 61: 339–360.

Chater, C., Peng, K., Movahedi, M., Dunn, J.A., Walker, H.J., Liang, Y.-K., McLachlan, D.H., Casson, S., Isner, J.C., Wilson, I., Neill, S.N., Hedrich, R., Gray, J.E. and Hetherington, A.M. (2015) Elevated CO₂-induced responses in stomata require ABA and ABA signaling. *Current Biology*. 25: 2709–2716.

Cramer, W., Bondeau, A., Woodward, F.I., Prentice, I.C., Betts, R.A., Brovkin, V., Cox, P.M., Fisher, V., Foley, J.A., Friend, A.D., Kucharik, C., Lomas, M.R., Ramankutty, N., Sitch, S., Smith, B., White, A. and Young-Molling, C. (2001) Global response of terrestrial ecosystem structure and function to CO₂ and climate change: results from six dynamic global vegetation models. *Global Change Biology*. 7: 357–373.

Delerce, S., Dorado, H., Grillon, A., Rebolledo, M.C., Prager, S.D., Patiño, V.H., Garcés Varón, G. and Jiménez, D. (2016) Assessing Weather-Yield Relationships in Rice at Local Scale Using Data Mining Approaches. *PLoS ONE*. 11: e0161620.

Deryng, D., Conway, D., Ramankutty, N., Price, J. and Warren, R. (2014) Global crop yield response to extreme heat stress under multiple climate change futures. *Environmental Research Letters*. 9: 34011 (13pp).

Diffenbaugh, N.S., Krupke, C.H., White, M.A. and Alexander, C.E. (2008) Global warming presents new challenges for maize pest management. *Environmental Research Letters*. 3: 44007 (9pp).

do Rio, A., Sentelhas, P.C., Farias, J.R.B., Sibaldelli, R.N.R. and Ferreira, R.C. (2016) Alternative sowing dates as a mitigation measure to reduce climate change impacts on soybean yields in southern Brazil. *International Journal of Climatology*. 36: 3664–3672.

Dlugokencky, E. and Tans, P. (2017) Trends in Atmospheric Carbon Dioxide. National Oceanic and Atmospheric Administration (NOAA/ESRL). <http://www.esrl.noaa.gov/gmd/ccgg/trends/global.html#global>. (Accessed April 2017)

Durand, J., Delusca, K., Boote, K., Lizaso, J., Manderscheid, R., Weigel, J.H., Ruane, A.C., Rosenzweig, C., Jones, J., Ahuja, L., Anapalli, S., Basso, B., Baron, C., Bertuzzi, P., Biernath, C., Deryng, D., Ewert, F., Gaiser, T., Gayler, S., Heinlein, F., Kersebaum, K.C., Kim, S.-H., Müller, C., Nendel, C., Olinaro, A., Priesack, E., Ramirez-Villegas, J., Ripoche, D., Rötter, R.P., Seidel, S.I., Srivastava, A., Tao, F., Timlin, D., Twine, T., Wang, E., Webber, H. and Zhao, Z. (2017) How accurately do maize crop models simulate the interactions of atmospheric CO₂ concentration levels with limited water supply on water use and yield? *European Journal of Agronomy*. (in press)

Ewert, F., Rötter, R.P., Bindi, M., Webber, H., Trnka, M., Kersebaum, K.C., Olesen, J.E., van Ittersum, M.K., Janssen, S., Rivington, M., Semenov, M.A., Wallach, D., Porter, J.R., Stewart, D., Verhagen, J., Gaiser, T., Palosuo, T., Tao, F., Nendel, C., Roggero, P.P., Bartosova, L. and Asseng, S. (2015) Crop modelling for integrated assessment of risk to food production from climate change. *Environmental Modelling & Software*. 72: 287–303.

Estes, L.D., Bradley, B.A., Beukes, H., Hole, D.G., Lau, M., Oppenheimer, M.G., Schulze, R., Tadross, M.A. and Turner, W.R. (2013) Comparing mechanistic and empirical model projections of crop suitability and productivity: implications for ecological forecasting. *Global Ecology and Biogeography*. 22: 1007–1018.

Fodor, N., Pásztor, L. and Németh, T. (2014) Coupling the 4M crop model with national geo-databases for assessing the effects of climate change on agro-ecological characteristics of Hungary. *International Journal of Digital Earth*. 7: 391–410.

Fowler, D., Amann, M., Anderson, F., Ashmore, M., Cox, P., Depledge, M., Derwent, D., Grennfelt, P., Hewitt, N., Hov, O. and Jenkin, M. (2008) Ground-level ozone in the 21st century: future trends, impacts and policy implications. *Royal Society Science Policy Report* 15.

Foyer, C.H., Bloom, A., Queval, G. and Noctor, G. (2009) Photorespiratory metabolism: genes, mutants, energetics, and redox signaling. *Annual Reviews of Plant Biology*. 60: 455–484.

Foyer, C.H., Lam, H-M., Nguyen, H.T., Siddique, K.H.M., Varshney, R.K., Colmer, T.D., Cowling, W., Bramley, H., Mori, T.A., Hodgson, J.M., Cooper, J.W., Miller, T., Kunert, K., Vorster, J., Cullis, C., Ozga, J.A., Wahlqvist, M.L., Liang, Y., Shou, H., Shi, K., Yu, J., Fodor, N., Kaiser, B.N., Wong, F-L., Valliyodan, B. and Considine, M.J. (2016) Neglecting legumes has compromised human health and sustainable food production. *Nature Plants*. 2: Article: 16112.

Foyer, C.H., Neukermans, J., Queval, G., Noctor, G. and Harbinson, J. (2012) Photosynthetic control of electron transport and the regulation of gene expression. *The Journal of Experimental Botany*. 63: 1637–1661.

Guo, R., Lin, Z., Mo, X. and Yang, C. (2010) Responses of crop yield and water use efficiency to climate change in the North China Plain. *Agricultural Water Management*. 97: 1185–1194.

Hansen, J.W. (2005) Integrating seasonal climate prediction and agricultural models for insights into agricultural practice. *Philosophical Transactions of the Royal Society B*. 360: 2037–2047.

Hasegawa, T., Sakai, H., Tokida, T., Nakamura, H., Zhu, C., Usui, Y., Yoshimoto, M., Fukuoka, M., Wakatsuki, H., Katayanagi, N., Matsunami, T., Kaneta, Y., Sato, T., Takakai, F., Sameshima, R., Okada, M., Mae, T. and Amane Makino, A. (2013) Rice cultivar response to elevated CO₂ at two free-air CO₂ enrichment (FACE) sites in Japan. *Functional Plant Biology*. 40: 148–159.

Hashimoto, M., Negi, J., Young, J., Israelsson, M., Schroeder, J.I. and Iba, K. (2006) Arabidopsis HT1 kinase controls stomatal movements in response to CO₂. *Nature Cell Biology*. 8: 391–397.

Hawkins, E., Osborne, T.M., Ho, C.K. and Challinor, A.J. (2013) Calibration and bias correction of climate projections for crop modelling: An idealised case study over Europe. *Agricultural and Forest Meteorology*. 170: 19–31.

Holden, N.M. and Brereton, A.J. (2003) Potential impacts of climate change on maize production and the introduction of soybean in Ireland. *Irish Journal of Agricultural and Food Research*. 42: 1–15.

Hollaway, M.J., Arnold, S.R., Challinor, A.J. and Emberson, L.D. (2012) Intercontinental trans-boundary contributions to ozone-induced crop yield losses in the Northern Hemisphere. *Biogeosciences*. 9: 271–292.

Hollinger, D.Y. and Richardson, A.D. (2005) Uncertainty in eddy covariance measurements and its application to physiological models. *Tree Physiology*. 25: 873–885.

Hoogenboom, G., Jones, J.W., Wilkens, P.W., Batchelor, W.D., Bowen, W.T., Hunt, L.A., Pickering, N.B., Singh, U., Godwin, D.C., Baer, B., Boote, K.J., Ritchie, J.T. and White, J.W. (1994) *Crop Models*. University of Hawaii, Department of Agronomy and Soil Science, Honolulu, Hawaii, USA.

Högy, P., Wieser, H., Köhler, P., Schwadorf, K., Breuer, J., Franzaring, J., Muntiferung, R. and Fangmeier, A. (2009) Effects of elevated CO₂ on grain yield and quality of wheat: Results from a 3-year free-air CO₂ enrichment experiment. *Plant Biology*. 11: 60–69.

Hu, H., Boisson-Dernier, A., Israelsson-Nordström, M., Böhmer, M., Xue, S., Ries, A., Godoski, J., Kuhn, J.M. and Schroeder, J.I. (2010) Carbonic anhydrases are upstream regulators of CO₂-controlled stomatal movements in guard cells. *Nature Cell Biology*. 12: 87–93.

IPCC (2013) Summary for Policymakers. In *Climate Change 2013: The Physical Science Basis*, Edited by Stocker, T.F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S.K., Boschung, J. et al., pp.1–27, Cambridge Univ. Press, Cambridge.

Jarvis, A., Lau, C., Cook, S., Wollenberg, E., Hansen, J., Bonilla, O. and Challinor, A. (2011) An Integrated Adaptation and Mitigation Framework for Developing Agricultural Research: Synergies and Trade-offs. *Experimental Agriculture*. 47: 185–203.

Jones, J.W., Hoogenboom, G., Porter, C.H., Boote, K.J., Batchelor, W.D., Hunt, L.A., Wilkens, P.W., Singh, U., Gijsman, A.J. and Ritchie, J.T. (2003) The DSSAT cropping system model. *European Journal of Agronomy*. 18: 235–265.

Jones, J.W., Antle, J.M., Basso, B., Boote, K.J., Conant, R.T., Foster, I., Godfray, H.C.J., Herrero, M., Howitt, R.E., Janssen, S., Keating, B.A., Munoz-Carpena, R., Porter, C.H., Rosenzweig, C. and Wheeler, T.R. (2016) Brief history of agricultural systems modeling. *Agricultural Systems*. (in press)

Julia, C. and Dingkuhn, M. (2013) Predicting temperature induced sterility of rice spikelets requires simulation of crop-generated microclimate. *European Journal of Agronomy*. 49: 50–60.

Keating, B.A., Carberry, P.S., Hammer, G.L., Probert, M.E., Robertson, M.J., Holzworth, D., Huth, N.I., Hargreaves, J.N.G., Meinke, H., Hochman, Z., McLean, G., Verburg, K., Snow, V., Dimes, J.P., Silburn, M., Wang, E., Brown, S., Bristow, K.L., Asseng, S., Chapman, S., McCown, R.L., Freebairn, D.M. and Smith, C.J. (2003) An overview of APSIM, a model designed for farming systems simulation. *European Journal of Agronomy*. 18: 267–288.

Kim, T.-H., Böhmer, M., Hu, H., Nishimura, N. and Schroeder, J.I. (2010) Guard cell signal transduction network: advances in understanding abscisic acid, CO₂, and Ca²⁺ signaling. *Annual Review of Plant Biology*. 61: 561–591.

Ko, J., Ahuja, L.R., Saseendran, S.A., Green, T.R., Ma, L., Nielsen, D.C. and Walthall, C.L. (2012) Climate change impacts on dryland cropping systems in the Central Great Plains, USA. *Climatic Change*. 111: 445–472.

Kolapo, A. (2011) Soybean: Africa's Potential Cinderella Food Crop. INTECH Open Access Publisher.

Köhler, I.H., Ruiz-Vera, U.M., VanLoocke, A., Thomey, M.L., Clemente, T., Long, S.P., Ort, D.R. and Bernacchi, C.J. (2016) Expression of cyanobacterial FBP/SBPase in soybean prevents yield depression under future climate conditions. *Journal of Experimental Botany*. 68: 715–726.

Lant, C., Stoebner, T.J., Schoof, J.T. and Crabb, B. (2016) The effect of climate change on rural land cover patterns in the Central United States. *Climatic Change*. 138: 585–602.

Leakey, A.D.B., Ainsworth, E.A., Bernacchi, C.J., Rogers, A., Long, S.P. and Ort, D.R. (2009a) Elevated CO₂ effects on plant carbon, nitrogen, and water relations: six important lessons from FACE. *Journal of Experimental Botany*. 60: 2859–2876.

Leakey, A.D.B., Ainsworth, E.A., Bernard, S.M., Markelz, C., Ort, D.R., Placella, S.A., Rogers, A., Smith, M.D., Sudderth, E.A., Weston, D.J., Wullschleger, S.D. and Yuan, S. (2009b) Gene expression profiling—opening the black box of plant ecosystem responses to global change. *Global Change Biology*. 15: 1201–1213.

Li, T., Hasegawa, T., Yin, X., Zhu, Y., Boote, K., Adam, M., Bregaglio, S., Buis, S., Confalonieri, R., Fumoto, T., Gaydon, D., Marcaida, M., Nakagawa, H., Oriol, P., Ruane, A.C., Ruget, F., Singh, B., Singh, U., Tang, L., Tao, F., Wilkens, P., Yoshida, H., Zhang, Z. and Bouman, B. (2015) Uncertainties in predicting rice yield by current crop models under a wide range of climatic conditions. *Global Change Biology*. 21: 1328–1341.

Lin, Y., Wu, W. and Ge, Q. (2015) CERES-Maize model-based simulation of climate change impacts on maize yields and potential adaptive measures in Heilongjiang Province, China. *Journal of the Science of Food and Agriculture*. 95: 2838–2849.

Liu, W., Yang, H., Folberth, C., Wang, X., Luo, Q. and Schulin, R. (2016) Global investigation of impacts of PET methods on simulating crop-water relations for maize. *Agricultural and Forest Meteorology*. 211: 164–175.

- Lobell, D.B. and Burke, M.B. (2010) On the use of statistical models to predict crop yield responses to climate change. *Agricultural and Forest Meteorology*. 150: 1443–1452.
- Lobell, D.B. (2014) Climate change adaptation in crop production: Beware of illusions. *Global Food Security*. 3: 72–76.
- Long, S.P. and Ort, D.R. (2010) More than taking the heat: crops and global change. *Current Opinion Plant Biology*. 13: 241–248.
- Lopes, M.S. and Foyer, C.H. (2011) The impact of high CO₂ on plant abiotic stress tolerance. Chapter: 6. In: *Crop Stress Management and Global Climate Change*, CABI, (Araus, J.L. and Slafer, G. eds). 85–104.
- Lopes, M.S., Araus, J.L., van Heerden, P.D.R. and Foyer, C.H. (2011) Enhancing drought tolerance in C₄ crops. *The Journal of Experimental Botany*. 62: 3135–3153.
- Luo, Y., Hui, D. and Zhang, D. (2006) Elevated CO₂ stimulated net accumulation of carbon and nitrogen in land ecosystems: a meta-analysis. *Ecology*. 87: 53–63.
- Manderscheid, R., Erbs, M. and Weigel, H.-J. (2014) Interactive effects of free-air CO₂ enrichment and drought stress on maize growth. *European Journal of Agronomy*. 52: 11–21.
- Mansfield, T.A., Hetherington, A.M. and Atkinson, C.J. (1990) Some current aspects of stomatal physiology. *Annual review of plant physiology and plant molecular biology*. 41: 55–75.
- Marten, H., Hyun, T., Gomi, K., Seo, S., Hedrich, R. and Roelfsema M.R.G. (2008) Silencing of NtMPK4 impairs CO₂-induced stomatal closure, activation of anion channels and cytosolic Ca signals in *Nicotiana tabacum* guard cells. *Plant Journal*. 55: 698–708.
- Martre, P., Jamieson, P.D., Semenov, M.A., Zyskowski, R.F., Porter, J.R. and Triboi, E. (2006) Modelling protein content and composition in relation to crop nitrogen dynamics for wheat. *European Journal of Agronomy*. 25: 138–154.

Mauser, W., Klepper, G., Zabel, F., Delzeit, R., Hank, T., Putzenlechner, B. and Calzadilla, A. (2015) Global biomass production potentials exceed expected future demand without the need for cropland expansion. *Nature Communications*. 6: 8946.

McGrath, J.M. and Lobell, D.B. (2011) An independent method of deriving the carbon dioxide fertilization effect in dry conditions using historical yield data from wet and dry years. *Global Change Biology*. 17: 2689–2696.

McGrath, J.M., Betzelberger, A.M., Wang, S., Shook, E., Zhu, X.G., Long, S.P. and Ainsworth, E.A. (2015) An analysis of ozone damage to historical maize and soybean yields in the United States. *Proceedings of the National Academy of Sciences*. 112: 14390–14395.

Mhamdi, A. and Noctor, G. (2016) High CO₂ primes plant biotic stress defences through redox-linked pathways. *Plant Physiology*. 172: 929–942.

Mihailović, D.T., Lalić, B., Drešković, N., Mimić, G., Djurdjević, V. and Jančić, M. (2015) Climate change effects on crop yields in Serbia and related shifts of Köppen climate zones under the SRES-A1B and SRES-A2. *International Journal of Climatology*. 35: 3320–3334.

Mills, G., Hayes, F., Simpson, D., Emberson, L., Norris, D., Harmens, H. and Büker, P. (2011) Evidence of widespread effects of ozone on crops and (semi-) natural vegetation in Europe (1990–2006) in relation to AOT40- and flux-based risk maps. *Global Change Biology*. 17: 592–613.

Mohd-Radzman, N.A., Binos, S., Truong, T.T., Imin, N., Mariani, M. and Djordjevic, M.A. (2015) Novel MtCEP1 peptides produced in vivo differentially regulate root development in *Medicago truncatula*. *Journal of Experimental Botany*. 66: 5289–5300.

Mondor, E.B., Tremblay, M.N., Awmack, C.S. and Lindroth, R.L. (2005) Altered genotypic and phenotypic frequencies of aphid populations under enriched CO₂ and O₃ atmospheres. *Global Change Biology*. 11: 1990–1996.

Monfreda, C., Ramankutty, N. and Foley, J.A. (2008) Farming the planet. Part 2: Geographic distribution of crop areas, yields, physiological types, and net primary production in the year 2000. *Global Biogeochemical Cycles*. 22: GB1022.

Moradi, R., Koocheki, A., Nassiri Mahallati, M. and Mansoori, H. (2013) Adaptation strategies for maize cultivation under climate change in Iran: Irrigation and planting date management. *Mitigation and Adaptation Strategies for Global Change*. 18: 265–284.

Müller, C., Elliott, J., Chryssanthacopoulos, J., Arneth, A., Balkovic, J., Ciais, P., Deryng, D., Folberth, C., Glotter, M., Hoek, S., Iizumi, T., Izaurralde, R.C., Jones, C., Khabarov, N., Lawrence, P., Liu, W., Olin, S., Pugh, T.A.M., Ray, D.K., Reddy, A., Rosenzweig, C., Ruane, A.C., Sakurai, G., Schmid, E., Skalsky, R., Song, C.X., Wang, X., de Wit, A. and Yang, H. (2017) Global gridded crop model evaluation: benchmarking, skills, deficiencies and implications. *Geoscientific Model Development*. 10: 1403–1422.

Olesen, J.E., Trnka, M., Kersebaum, K.C., Skjelvåg, A.O., Seguin, B., Peltonen-Sainio, P., Rossi, F., Kozyra J. and Micale, F. (2011) Impacts and adaptation of European crop production systems to climate change. *European Journal of Agronomy*. 34: 96–112.

Osborne, T.M., Lawrence, D.M., Challinor, A.J., Slingo, J.M. and Wheeler, T.R. (2007) Development and assessment of a coupled crop–climate model. *Global Change Biology*. 13: 169–183.

Osborne, T., Gornall, J., Hooker, J., Williams, K., Wiltshire, A., Betts, R. and Wheeler, T. (2015) JULES-crop: a parametrisation of crops in the Joint UK Land Environment Simulator. *Geoscientific Model Development*. 8: 1139–1155.

Ort, D.R., Merchant, S.S., Alric, J., Barkan, A., Blankenship, R.E., Bock, R., Croce, R., Hanson, M.R., Hibberd, J.M., Long, S.P., Moore, T.A., Moroney, J., Niyogi, K.K., Parry, M.A.J., Peralta-Yahya, P.P., Prince, R.C., Redding, K.E., Spalding, M.H., van Wijk, K.J., Vermaas, W.F.J., von Caemmerer, S., Weber, A.P.M., Yeates, T.O., Yuan, J.S. and Zhu, X.G. (2015) Redesigning photosynthesis to sustainably meet global food and bioenergy demand. *Proceedings of the National Academy of Sciences*. 112: 8529–8536.

Paul, M.J. and Foyer, C.H. (2001) Sink regulation of photosynthesis. *Journal of Experimental Botany*. 52: 1383–1400.

Paul, E.A., Harris, D., Collins, H.P., Schulthess, U. and Robertson, G.P. (1999) Evolution of CO₂ and soil carbon dynamics in biologically managed, row-crop agroecosystems. *Applied Soil Ecology*. 11: 53–65.

Prins, A., Muchwesi Mukubi, J., Pellny, T.K., Verrier, P., Beyene, G., Sabino, S., Lopes, M., Emami, K., Treumann, A., Lelarge-Trouverie, C., Noctor, G., Kunert, K.J., Kerchev, P. and Foyer, C.H. (2010) Acclimation to high CO₂ in maize is dependent on water status and leaf rank. *Plant, Cell & Environment*. 34: 314–331.

Rachmilevitch, S., Cousins, A.B. and Bloom, A.J. (2004) Nitrate assimilation in plant shoots depends on photorespiration. *Proceedings of the National Academy of Sciences of the USA*. 101: 11506–11510.

Ramirez-Villegas, J., Challinor, A.J., Thornton, P.K. and Jarvis, A. (2013) Implications of regional improvement in global climate models for agricultural impact research. *Environmental Research Letters*. 8: 024018 (12pp).

Reich, P.B. and Hobbie, S.E. (2013) Decade-long soil nitrogen constraint on the CO₂ fertilization of plant biomass. *Nature Climate Change*. 3: 278–282.

Ritchie, J.T. (2000) Oral communication. At Nowlin Chair Conference on crop and soil modelling, 10-11 November 2000, Detroit, Michigan, USA.

Rosenzweig, C., Elliott, J., Deryng, D., Ruane, A.C., Müller, C., Arneth, A., Boote, K.J., Folberth, C., Glotter, M., Khabarov, N., Neumann, K., Piontek, F., Pugh, T.A.M., Schmid, E., Stehfest, E., Yang, H. and Jones, J.W. (2014) Assessing agricultural risks of climate change in the 21st century in a global gridded crop model intercomparison. *Proceedings of the National Academy of Sciences of the USA*. 111: 3268–3273.

Rosenzweig, C., Jones, J.W., Hatfield, J.L., Ruane, A.C., Boote, K.J., Thorburn, P., Antle, J.M., Nelson, G.C., Porter, C., Janssen, S., Asseng, S., Basso, B., Ewert, F., Wallach, D., Baigorría,

G. and Winter, J.M. (2013) The Agricultural Model Intercomparison and Improvement Project (AgMIP): Protocols and pilot studies. *Agricultural and Forest Meteorology*. 170: 166–182.

Ruiz-Ramos, M. and Mínguez, M.I. (2010) Evaluating uncertainty in climate change impacts on crop productivity in the Iberian Peninsula. *Climate Research*. 44: 69–82.

Sen, B., Topcu, S., Türkeş, M., Sen, B. and Warner, J.F. (2012) Projecting climate change, drought conditions and crop productivity in Turkey. *Climate Research*. 52: 175–191.

Shirsath, P.B., Aggarwal, P.K., Thornton, P.K. and Dunnett, A., (2017) Prioritizing climate-smart agricultural land use options at a regional scale. *Agricultural Systems*. 151: 174–183.

Schlenker, W. and Roberts, M.J. (2009) Nonlinear temperature effects indicate severe damages to U.S. crop yields under climate change. *Proceedings of the National Academy of Sciences*. 106: 15594–15598.

Schleussner, C.-F., Rogelj, J., Schaeffer, M., Lissner, T., Licker, R., Fischer, E.M., Knutti, R., Levermann, A., Frieler, K. and Hare, W. (2016) Science and policy characteristics of the Paris Agreement temperature goal. *Nature Climate Change*. 6: 827–835.

Sinclair, T.R. and Seligman, N.A. (2000) Criteria for publishing papers on crop modeling. *Field Crops Research*. 68: 165–172.

Simelton, E., Fraser, E.D.G., Termansen, M., Benton, T.G., Gosling, S.N., South, A., Arnell, A.W., Challinor, A.J., Dougill, A.J. and Forster, P.M. (2012) The socioeconomics of food crop production and climate change vulnerability: a global scale quantitative analysis of how grain crops are sensitive to drought. *Food Security*. 4: 163–179.

Smith, P., Martino, D., Cai, Z., Gwary, D., Janzen, H., Kumar, P., McCarl, B., Ogle, S., O'Mara, F., Rice, C., Scholes, B., Sirotenko, O., Howden, M., McAllister, T., Pan, G., Romanenkov, V., Schneider, U., Towprayoon, S., Wattenbach, M. and Smith J. (2008) Greenhouse gas mitigation in agriculture. *Philosophical Transactions of the Royal Society B: Biological Sciences*. 363: 789–813.

- Soares, A.S., Driscoll, S.P., Olmos, E., Harbinson, J., Arrabacxa, M.C. and Foyer, C.H. (2007) Adaxial/abaxial specification in the regulation of photosynthesis with respect to light orientation and growth with CO₂ enrichment in the C₄ species *Paspalum dilatatum*. *New Phytologist*. 177: 186–198.
- Supit, I., van Diepen, C.A., De Wit, A.J.W., Wolf, J., Kabat, P., Baruth, B. and Ludwig, F. (2012) Assessing climate change effects on European crop yields using the Crop Growth Monitoring System and a weather generator. *Agricultural and Forest Meteorology*. 16: 96–111.
- Tanaka, Y., Nose, T., Jikumaru, Y. and Kamiya, Y. (2013) ABA inhibits entry into stomatal-lineage development in *Arabidopsis* leaves. *The Plant Journal*. 74: 448–457.
- Tatsumi, K., Yamashiki, Y., Valmir da Silva, R., Takara, K., Matsuoka, Y., Takahashi, K., Maruyama, K. and Kawahara, N. (2011) Estimation of potential changes in cereals production under climate change scenarios. *Hydrological Processes*. 25: 2715–2725.
- Tian, W., Hou, C., Ren, Z., Pan, Y., Jia, J., Zhang, H., Bai, F., Zhang, P., Zhu, H. and He Y. (2015) A molecular pathway for CO₂ response in *Arabidopsis* guard cells. *Nature Communications*. 6: 6057.
- Tilman, D. and Clark, M. (2014) Global diets link environmental sustainability and human health. *Nature*. 515: 518–522.
- Travasso, M., Magrin, G.O., Rodriguez, G.R., Solman, S. and Nunez, M. (2009) Climate change impacts on regional maize yields and possible adaptation measures in Argentina. *International Journal of Global Warming*. 1: 201–213.
- Tubiello, F.N., Donatelli, M., Rosenzweig, C. and Stockle, C.O. (2000) Effects of climate change and elevated CO₂ on cropping systems: Model predictions at two Italian locations. *European Journal of Agronomy*. 13: 179–189.
- Tubiello, F.N., Rosenzweig, C., Goldberg, R.A., Jagtap, S. and Jones, J.W. (2002) Effects of climate change on US crop production: Simulation results using two different GCM scenarios. Part I: Wheat, potato, maize, and citrus. *Climate Research*. 20: 259–270.

Vavasseur, A. and Raghavendra, A.S. (2005) Guard cell metabolism and CO₂ sensing. *New Phytologist*. 165: 665–682.

Wallach, D., Thorburn, P., Asseng, S., Challinor, A.J., Ewert, F., Jones, J.W., Rotter, R. and Ruane, A.C. (2016) Estimating model prediction error: Should you treat predictions as fixed or random? *Environmental Modelling & Software*. 84: 529–539.

Wang, X. and Mauzerall, D.L. (2004) Characterizing distributions of surface ozone and its impact on grain production in China, Japan and South Korea: 1990 and 2020. *Atmospheric Environment*. 38: 4383-4402.

Wang, Q., Watanabe, M. and Ouyang, Z. (2005) Simulation of water and carbon fluxes using BIOME-BGC model over crops in China. *Agricultural and Forest Meteorology*. 131: 209–224.

Warszawski, L., Frieler, K., Huber, V., Piontek, F., Serdeczny, O. and Schewe, J. (2014) The Inter-Sectoral Impact Model Intercomparison Project (ISI-MIP): Project framework. *Proceedings of the National Academy of Sciences of the USA*. 111: 3228–3232.

Webber, H., Gaiser, T. and Ewert, F. (2014) What role can crop models play in supporting climate change adaptation decisions to enhance food security in Sub-Saharan Africa? *Agricultural Systems*. 127: 161–177.

Webber, H., Martre, P., Asseng, S., Kimball, B., White, J., Ottman, M., Wall, G.W., De Sanctis, G., Doltra, J., Grant, R., Kassie, B., Maiorano, A., Olesen, J.E., Ripoche, D., Rezaei, E.E., Semenov, M.A., Stratonovitch, P. and Ewert, F. (2015) Canopy temperature for simulation of heat stress in irrigated wheat in a semi-arid environment: A multi-model comparison. *Field Crops Research*. 20: 21–35.

Wesselink, A., Challinor, A., Watson, J., Beven, K., Allen, I., Hanlon, H., Lopez, A., Lorenz, S., Otto, F., Morse, A., Rye, C., Saux-Picard, S., Stainforth, D. and Suckling, E. (2015) Equipped to deal with uncertainty in climate and impacts predictions: lessons from internal peer review. *Climatic Change*. 132: 1–14.

Williams, J.R., Jones, C.A., Kiniry, J.R. and Spanel, D.A. (1989) EPIC crop growth model. *Transactions of the American Society of Agricultural Engineers*. 32: 497–511.

White, J.W., Hoogenboom, G., Kimball, B.A. and Wall, G.W. (2011) Methodologies for simulating impacts of climate change on crop production. *Field Crops Research*. 124: 357–368.

Woodward, F.I. and Kelly, C.K. (1995) The influence of CO₂ concentration on stomatal density. *New Phytologist*. 131: 311–327.

Woodward, F.I. (1987) Stomatal numbers are sensitive to increases in CO₂ from pre-industrial levels. *Nature*. 327: 617–618.

Wriedt, G., Van der Velde, M., Aloe, A. and Bouraoui, F. (2009) Estimating irrigation water requirements in Europe. *Journal of Hydrology*. 373: 527–544.

Xiong, W., Matthews, R., Holman, I., Lin, E. and Xu, Y. (2007) Modelling China's potential maize production at regional scale under climate change. *Climatic Change*. 85: 433–451.

Yang, X., Chen, F., Lin, X., Liu, Z., Zhang, H., Zhao, J., Li, K., Ye, Q., Li, Y., Lv, S., Yang, P., Wu, W., Li, Z., Lal, R. and Tang, H. (2015) Potential benefits of climate change for crop productivity in China. *Agricultural and Forest Meteorology*. 208: 76–84.

Zabel, F., Putzenlechner, B. and Mauser, W. (2014) Global Agricultural Land Resources – A High Resolution Suitability Evaluation and Its Perspectives until 2100 under Climate Change Conditions. *PLoS ONE*. 9: e107522.