

## RESOLVABILITY IN C.C.C. GENERIC EXTENSIONS

LAJOS SOUKUP AND ADRIENNE STANLEY

ABSTRACT. Every crowded space  $X$  is  $\omega$ -resolvable in the c.c.c generic extension  $V^{\text{Fn}(|X|,2)}$  of the ground model.

We investigate what we can say about  $\lambda$ -resolvability in c.c.c-generic extensions for  $\lambda > \omega$ ?

A topological space is *monotonically  $\omega_1$ -resolvable* if there is a function  $f : X \rightarrow \omega_1$  such that

$$\{x \in X : f(x) \geq \alpha\} \subset^{dense} X$$

for each  $\alpha < \omega_1$ .

We show that given a  $T_1$  space  $X$  the following statements are equivalent:

- (1)  $X$  is  $\omega_1$ -resolvable in some c.c.c-generic extension,
- (2)  $X$  is monotonically  $\omega_1$ -resolvable.
- (3)  $X$  is  $\omega_1$ -resolvable in the Cohen-generic extension  $V^{\text{Fn}(\omega_1,2)}$ .

We investigate which spaces are monotonically  $\omega_1$ -resolvable. We show that if a topological space  $X$  is c.c.c, and  $\omega_1 \leq \Delta(X) \leq |X| < \omega_\omega$ , where  $\Delta(X) = \min\{|G| : G \neq \emptyset \text{ open}\}$ , then  $X$  is monotonically  $\omega_1$ -resolvable.

On the other hand, it is also consistent, modulo the existence of a measurable cardinal, that there is a space  $Y$  with  $|Y| = \Delta(Y) = \aleph_\omega$  which is not monotonically  $\omega_1$ -resolvable.

The characterization of  $\omega_1$ -resolvability in c.c.c generic extension raises the following question: is it true that crowded spaces from the ground model are  $\omega$ -resolvable in  $V^{\text{Fn}(\omega,2)}$ ?

We show that (i) if  $V = L$  then every crowded c.c.c. space  $X$  is  $\omega$ -resolvable in  $V^{\text{Fn}(\omega,2)}$ , (ii) if there is no weakly inaccessible cardinals, then every crowded space  $X$  is  $\omega$ -resolvable in  $V^{\text{Fn}(\omega_1,2)}$ .

On the other hand, it is also consistent, modulo a measurable cardinals, that there is a crowded space  $X$  with  $|X| = \Delta(X) = \omega_1$  such that  $X$  remains irresolvable after adding a single Cohen real.

---

*Date:* February 2, 2017.

*2010 Mathematics Subject Classification.* 54A35, 03E35, 54A25.

*Key words and phrases.* resolvable, monotonically  $\omega_1$ -resolvable, measurable cardinal.

The second author was supported by Fulbright Scholar Program.

The preparation of this paper was supported by OTKA grant no. K113047.

## 1. INTRODUCTION

Notion of resolvability was introduced and studied first by E. Hewitt, [4], in 1943. A topological space  $X$  is  $\kappa$ -resolvable if it can be partitioned into  $\kappa$  many dense subspaces.  $X$  is *resolvable* iff it is 2-resolvable, and *irresolvable* otherwise. Irresolvable spaces with many interesting extra properties were constructed, but there are no “absolute” examples for crowded irresolvable spaces, because if  $X$  is a crowded space, then clearly

$$V^{\text{Fn}(|X|,2)} \models X \text{ is } \omega\text{-resolvable.}$$

In this paper we investigate what we can say about  $\lambda$ -resolvability in c.c.c-generic extensions for  $\lambda > \omega$ ?

To characterize spaces which are  $\omega_1$ -resolvable in some c.c.c-generic extension we introduce the notion of monotonically  $\kappa$ -resolvable.

**Definition 1.1.** Let  $\kappa$  be an infinite cardinal. A topological space  $X$  is *monotonically  $\kappa$ -resolvable*<sup>†</sup> if there is a function  $f : X \rightarrow \kappa$  such that

$$\{x \in X : f(x) \geq \alpha\} \subset^{dense} X$$

for each  $\alpha < \kappa$ . We will say that  $f$  *witnesses* that  $X$  is monotonically  $\kappa$ -resolvable.

Clearly a space  $X$  is monotonically  $\kappa$ -resolvable iff  $X$  has a partition  $\{X_\zeta : \zeta < \kappa\}$  of  $X$  such that

$$\text{int} \left( \bigcup \{X_\zeta : \zeta < \xi\} \right) = \emptyset$$

for all  $\xi < \kappa$ .

**Theorem 1.2.** *Let  $X$  be a  $T_1$  topological space. The following statements are equivalent:*

- (1)  $X$  is  $\omega_1$ -resolvable in some c.c.c-generic extension,
- (2)  $X$  is monotonically  $\omega_1$ -resolvable,
- (3)  $X$  is  $\omega_1$ -resolvable in the Cohen generic extension  $V^{\text{Fn}(\omega_1,2)}$ .

Which spaces are monotonically  $\omega_1$ -resolvable?

**Theorem 1.3.** *If a topological space  $X$  is c.c.c, and  $\omega_1 \leq \Delta(X) \leq |X| < \omega_\omega$ , then  $X$  is monotonically  $\omega_1$ -resolvable.*

**Theorem 1.4.** *If  $\kappa$  is a measurable cardinal, then there is a space  $X$  with  $|X| = \Delta(X) = \kappa$  which is not monotonically  $\omega_1$ -resolvable.*

---

<sup>†</sup>In [13] a “monotonically  $\omega$ -resolvable” space is called “almost- $\omega$ -resolvable”. However, in [12] a space  $X$  is *almost- $\kappa$ -resolvable* if it contains a family of  $\kappa$  dense sets with pairwise nowhere dense intersections.

What about spaces of cardinality  $\omega_\omega$ ?

**Theorem 1.5.** *It is consistent, modulo the existence of a measurable cardinal, that there is a space  $X$  with  $|X| = \Delta(X) = \omega_\omega$  which is not monotonically  $\omega_1$ -resolvable.*

Do we really need to add  $|X|$ -many Cohen reals to make  $X$  resolvable?

**Theorem 1.6.** (1) *It is consistent, modulo a measurable cardinal, that there is a crowded space  $X$  with  $|X| = \Delta(X) = \omega_1$  (so  $X$  is monotonically  $\omega_1$ -resolvable) such that*

$$V^{\text{Fn}(\omega, 2)} \models "X \text{ is irresolvable}."$$

- (2) *If  $V = L$ , then every crowded space with  $|X| = \Delta(X) = \text{cf}(|X|)$  is monotonically  $\omega$ -resolvable, and so it is  $\omega$ -resolvable in  $V^{\text{Fn}(\omega, 2)}$ .*  
(3) *If the cardinality of a crowded c.c.c space  $X$  is less than the first weakly inaccessible cardinal, then  $X$  is  $\omega$ -resolvable in  $V^{\text{Fn}(\omega_1, 2)}$  §.*

The almost resolvability of c.c.c spaces was investigated by Pavlov in [11]: on page 53 Pavlov writes that – mimicked Malykhin’s method by using Ulam matrices – he showed that every crowded ccc space of cardinality  $\omega_1$  is almost resolvable. In [3, Theorem 2.22] a stronger result was proved: a crowded c.c.c. space is almost resolvable, if its cardinality is less than the first weakly inaccessible cardinal. Theorem 1.6(2) is a further improvement of this result because monotonically  $\omega$ -resolvability implies almost resolvability.

In [1, 3.12 Problem (2)] the authors ask *if every space with countable cellularity and cardinality less than the first inaccessible non-countable cardinal almost- $\omega$ -resolvable?* As we will see Theorem 1.6 (3) gives a positive answer to a weakening of this question.

## 2. CHARACTERIZATION OF $\omega_1$ -RESOLVABILITY IN C.C.C EXTENSIONS.

Instead of Theorem 1.2 we prove the following stronger result.

**Theorem 2.1.** *Assume that  $X$  is a crowded topological space and  $\kappa$  is an infinite cardinal. If  $\kappa = \text{cf}([\kappa]^\omega, \subset)$  then following statements are equivalent.*

- (1)  *$X$  is  $\kappa$ -resolvable in some c.c.c-generic extension,*
- (2) *there is a function  $h : X \rightarrow [\kappa]^\omega$  such that  $\bigcup h''U = \kappa$  for each non-empty open  $U \subset X$ .*
- (3)  *$X$  is  $\kappa$ -resolvable in the Cohen-generic extension  $V^{\text{Fn}(\kappa, 2)}$ .*

---

§ $\omega_1$  is not a misprint here

We say that a function  $g : X \rightarrow \kappa$  *witnesses* that  $X$  is  $\kappa$ -resolvable if

$$\{x \in X : g(x) = \alpha\} \subset^{dense} X$$

for each  $\alpha < \kappa$ .

*Proof.* First we show that (1)  $\rightarrow$  (2). Assume that  $\mathbb{P}$  is a c.c.c. poset such that there is a function  $g \in V^{\mathbb{P}}$  witnessing the  $\kappa$ -resolvability of  $X$ .

For each  $x \in X$  define

$$h(x) = \{\alpha < \kappa : \exists p_\alpha^x \in \mathbb{P}(p_\alpha^x \Vdash \dot{g}(\check{x}) = \check{\alpha})\}.$$

Since the conditions  $\{p_\alpha^x : \alpha \in h(x)\}$  are pairwise incomparable and  $\mathbb{P}$  is c.c.c., the set  $h(x)$  is countable.

We now show that the function  $h$  defined above satisfies (2). Fix  $\alpha < \kappa$  and  $U$  an open subset of  $X$ . We need to show that there exists  $x \in U$  such that  $\alpha \in h(x)$ . Since

$$V^{\mathbb{P}} \models g^{-1}(\{\alpha\}) \subset^{dense} X$$

it follows that there is  $x \in U$  such that

$$V^{\mathbb{P}} \models g(x) = \alpha.$$

Thus, there exists  $p \in \mathbb{P}$  such that

$$p \Vdash \text{"}\dot{g}(\check{x}) = \check{\alpha}\text{"}$$

Then  $\alpha \in h(x)$ .

Next we now show that (2)  $\rightarrow$  (3). Let  $\mathcal{A}$  be a cofinal subset of  $[\kappa]^\omega$  with  $|\mathcal{A}| = \kappa$ .

Let  $\{A_\alpha : \alpha < \kappa\}$  be an enumeration of  $\mathcal{A}$ , and for each  $x \in X$  pick

$$h^*(x) \in \mathcal{A} \text{ such that } h^*(x) \supset \bigcup_{\alpha \in h(x)} A_\alpha.$$

Then for all non-empty open  $U$

$$\{h^*(x) : x \in U\} \text{ is cofinal in } [\kappa]^\omega. \quad (+)$$

Next we note that forcing with  $\text{Fn}(\kappa, 2)$  is the same as forcing with  $\text{Fn}(\kappa, \omega)$ . Further,  $\text{Fn}(\kappa, \omega)$  is isomorphic to

$$\mathbb{P} = \{p \in \text{Fn}(\mathcal{A}, \kappa) : \forall A \in \text{dom}(p) \ p(A) \in A\}.$$

Indeed, for each  $A \in \mathcal{A}$  fix a bijection  $\rho_A : \omega \rightarrow A$ , and then for  $q \in \text{Fn}(\kappa, \omega)$  define  $\varphi(q) \in \mathbb{P}$  as follows:

- (i)  $\text{dom}(\varphi(q)) = \{A_\alpha : \alpha \in \text{dom}(q)\}$ , and
- (ii)  $\varphi(q)(A_\alpha) = \rho_{A_\alpha}(q(\alpha))$  for  $A_\alpha \in \text{dom}(\varphi(q))$ .

Then  $\varphi$  is clearly an isomorphism between  $\text{Fn}(\kappa, \omega)$  and  $\mathbb{P}$ .

We will proceed using  $\mathbb{P}$ .

Let  $G$  be a  $\mathbb{P}$ -generic filter, and let  $g = \bigcup G$ . Then  $g \in V^{\mathbb{P}}$  and  $g : \mathcal{A} \rightarrow \kappa$  such that  $g(A) \in A$ .

We claim that  $f = g \circ h^*$  witnesses that  $X$  is  $\kappa$ -resolvable.

Fix  $\alpha < \kappa$  and an open  $U \subset X$ .

Let  $q \in \mathbb{P}$  be arbitrary. Then, by  $(+)$ , there is  $x \in U$  such that

$$\{\alpha\} \cup \bigcup \text{dom}(q) \subsetneq h^*(x).$$

Then  $h^*(x) \notin \text{dom}(q)$ , and  $\alpha \in h^*(x)$ , so

$$p = q \cup \{\langle h^*(x), \alpha \rangle\} \in \mathbb{P}_1,$$

and

$$p \Vdash (g \circ h^*)(\check{x}) = \check{\alpha}.$$

Thus, by genericity, there is  $p \in G$  and  $x \in U$  such that

$$p \Vdash (g \circ h^*)(\check{x}) = \check{\alpha}.$$

Hence

$$V^{\mathbb{P}} \models X \text{ is } \kappa\text{-resolvable}.$$

Finally  $(3) \rightarrow (1)$  is trivial.  $\square$

**Problem 2.2.** *Can we drop the assumption  $\kappa = \text{cf}([\kappa]^\omega, \subset)$  from Theorem 2.1?*

### 3. ON MONOTONICALLY $\omega_1$ -RESOLVABILITY OF C.C.C SPACES

We start with an easy observation.

**Lemma 3.1.** *Let  $X$  be a topological space and  $\mathcal{B} \subset \mathcal{P}(X)$ . If every  $B \in \mathcal{B}$  is monotonically  $\kappa$ -resolvable, then so is  $\overline{\bigcup \mathcal{B}}$ . So every space contains a greatest monotonically  $\kappa$ -resolvable subspace (that subspace can be empty, of course).*

**Corollary 3.2.** *Let  $X$  be a topological space. Let  $Z$  be a dense subset of  $X$ . If  $Z$  is monotonically  $\kappa$ -resolvable, then  $X$  is also monotonically  $\kappa$ -resolvable.*

Before proving Theorem 1.3 we prove the following “stepping-down” theorem.

**Theorem 3.3.** *If  $X$  is a  $\kappa$ -c.c., monotonically  $\kappa^+$ -resolvable space, then  $X$  is monotonically  $\kappa$ -resolvable as well.*

The proof uses ideas from [8].

*Proof.* Since an open subspace of a  $\kappa$ -c.c., monotonically  $\kappa^+$ -resolvable space is also  $\kappa$ -c.c. and monotonically  $\kappa^+$ -resolvable, by Lemma 3.1 it is enough to show that

- (\*) every  $\kappa$ -c.c., monotonically  $\kappa^+$ -resolvable space  $X$  has a monotonically  $\kappa$ -resolvable non-empty open subset.

Ulam [14] proved that there is a “matrix”

$$\langle M_{\alpha,\zeta} : \alpha < \kappa^+, \zeta < \kappa \rangle \subset \mathcal{P}(\kappa^+)$$

such that

- (i)  $M_{\alpha,\xi} \cap M_{\beta,\xi} = \emptyset$  for  $\{\alpha, \beta\} \in [\kappa^+]^2$  and  $\xi \in \kappa$ ,
- (ii)  $M_{\alpha,\xi} \cap M_{\alpha,\zeta} = \emptyset$  for  $\alpha \in \kappa^+$  and  $\{\xi, \zeta\} \in [\kappa]^2$ ,
- (iii) and  $|M_\alpha^-| \leq \kappa$ , where  $M_\alpha^- = \kappa^+ \setminus \bigcup_{\zeta < \kappa} M_{\alpha,\zeta}$  for  $\alpha < \kappa^+$ .

Fix a partition  $\{Y_\eta : \eta < \kappa^+\}$  witnessing that  $X$  is monotonically  $\kappa^+$ -resolvable.

Let

$$Z_{\alpha,\zeta} = \bigcup \{Y_\eta : \eta \in M_{\alpha,\zeta}\}$$

for  $\alpha < \kappa^+$  and  $\zeta < \kappa$ , and let

$$Z_\alpha = \bigcup_{\zeta < \kappa} Z_{\alpha,\zeta}.$$

Since  $Z_\alpha = \bigcup \{Y_\eta : \eta \in \kappa^+ \setminus M_\alpha^-\}$ , assumption (iii) implies that every  $Z_\alpha$  is dense in  $X$ .

**Case 1.** There is  $\alpha < \kappa^+$  such that for all  $\zeta < \kappa$

$$\bigcup_{\zeta \leq \xi} Z_{\alpha,\zeta} \subset^{dense} Z_\alpha.$$

Then  $(Z_{\alpha,\zeta})_{\zeta < \kappa}$  witnesses  $Z_\alpha$  is monotonically  $\kappa$ -resolvable and so by corollary 3.2,  $X$  is also monotonically  $\kappa$ -resolvable.

**Case 2.** For all  $\alpha < \kappa^+$  there is  $\zeta_\alpha < \kappa$  and there is a non-empty open set  $U_\alpha \in \tau_X$  such that

$$\bigcup_{\zeta_\alpha \leq \xi} Z_{\alpha,\zeta} \cap U_\alpha = \emptyset. \quad (\dagger)$$

Then there is a set  $I \in [\kappa^+]^{\kappa^+}$  and there is an ordinal  $\zeta < \kappa$  such that  $\zeta_\alpha = \zeta$  for all  $\alpha \in I$ .

Fix an arbitrary  $K \in [I]^\kappa$ . By (iii) we can find  $\rho < \kappa^+$  such that

$$\bigcup_{\alpha \in K} M_\alpha^- \subset \rho.$$

Let  $Z = \bigcup_{\rho < \eta} Y_\eta$ . Then  $Z \subset^{dense} X$  and  $Z \subset Z_\alpha$  for all  $\alpha \in K$ .

**Claim.** If  $L \in [K]^\kappa$  then

$$\bigcap_{\alpha \in L} U_\alpha \cap Z = \emptyset.$$

*Proof of the Claim.* Assume on the contrary that  $z \in \bigcap_{\alpha \in L} U_\alpha \cap Z$ . Then  $z \in Y_\eta$  for some  $\rho < \eta$ .

Let  $\alpha \in L$ . Then  $\eta \in \kappa^+ \setminus \rho \subset \bigcup_{\xi < \kappa} M_{\alpha, \xi}$ . Pick  $\xi_\alpha < \kappa$  with  $\eta \in M_{\alpha, \xi_\alpha}$ . Then  $Y_\eta \subset Z_{\alpha, \xi_\alpha}$ , so  $Z_{\alpha, \xi_\alpha} \cap U_\alpha \neq \emptyset$ , so  $\xi_\alpha < \zeta_\alpha = \zeta$  by  $(\dagger)$ .

Since  $\zeta < \kappa = |L|$ , there are  $\alpha \neq \beta \in [L]^2$  such that  $\xi_\alpha = \xi_\beta$ . Thus  $\eta \in M_{\alpha, \xi_\alpha} \cap M_{\beta, \xi_\beta}$  which contradicts (i) because  $\xi_\alpha = \xi_\beta$ .  $\square$

Fix an enumeration  $K = \{\chi_\xi : \xi < \kappa\}$ , and let  $V_\zeta = \bigcup_{\xi < \zeta} U_{\chi_\xi}$ . Then the sequence  $\langle V_\zeta : \zeta < \kappa \rangle$  is decreasing and

$$\bigcap_{\zeta < \kappa} V_\zeta \cap Z = \emptyset$$

by the Claim.

Since  $X$  is  $\kappa$ -c.c. there is  $\xi < \kappa$  such that  $\overline{V_\zeta} = \overline{V_\xi}$  for all  $\xi < \zeta < \kappa$ .

We can assume that  $\xi = 0$ . Let

$$T_\zeta = \begin{cases} V_0 \setminus Z & \text{if } \zeta = 0, \\ ((\bigcap_{\xi < \zeta} V_\xi) \setminus V_\zeta) \cap Z & \text{if } \zeta > 0. \end{cases}$$

Then

$$\bigcup_{\xi < \zeta} T_\zeta \supset V_\xi \cap Z \subset^{dense} V.$$

thus the partition  $\{T_\zeta : \zeta < \kappa\}$  witnesses that  $V$  is monotonically  $\kappa$ -resolvable.  $\square$

*Proof of Theorem 1.3.* Let  $\mathcal{Y} = \{Y \in \tau_X : |Y| = \Delta(Y)\}$ .

Then  $\bigcup \mathcal{Y}$  is dense in  $X$ , and every open subset of every  $Y \in \mathcal{Y}$  is also in  $\mathcal{Y}$ . Thus by lemma 3.1 it is enough to prove that a c.c.c. space  $Y$  with  $\omega_1 \leq |Y| = \Delta(Y) < \omega_\omega$  is monotonically  $\omega_1$ -resolvable.

Let  $Y \in \mathcal{Y}$  such that  $\omega_n = |Y|$ . Clearly,  $Y$  is monotonically  $\omega_n$ -resolvable as  $|Y| = \Delta(Y) = \omega_n$ . Since  $Y$  is c.c.c. then  $Y$  is  $\omega_{n-1}$ -c.c.c.. By theorem 3.3,  $Y$  is monotonically  $\omega_{n-1}$ -resolvable. By continually applying theorem 3.3 we conclude that  $Y$  is monotonically  $\omega_1$ -resolvable.  $\square$

**Problem 3.4.** Is it true that every crowded c.c.c space with  $\Delta(X) \geq \omega_1$  is monotonically  $\omega_1$ -resolvable?

4. SPACES WHICH ARE NOT MONOTONICALLY  $\omega_1$ -RESOLVABLE.

If  $X$  is a topological space, and  $\mathcal{D} \subset \mathcal{P}(X)$ , write

$$\overline{\overline{\mathcal{D}}} = \{\overline{D} : D \in \mathcal{D}\}$$

**Lemma 4.1.** *Let  $X$  be a topological space. Assume that  $\overline{\overline{\mathcal{D}}}$  is point-countable for each point-countable family  $\mathcal{D} \subset \mathcal{P}(X)$ . Then  $X$  is not contain any monotonically  $\omega_1$ -resolvable subspace  $Y$ .*

*Proof.* Assume that  $\{Y_\zeta : \zeta < \omega_1\}$  is a partition of  $Y$ . Let  $D_\xi = \bigcup\{Y_\zeta : \xi < \zeta\}$  for  $\xi < \omega_1$ . Then the family  $\mathcal{D} = \{D_\xi : \xi < \omega_1\}$  is point-countable. So  $\overline{\overline{\mathcal{D}}}$  is also point-countable. So  $D_\xi$  is not dense in  $Y$  for all but countably many  $\xi$ . So the partition  $\{Y_\zeta : \zeta < \omega_1\}$  does not witness that  $Y$  is monotonically  $\omega_1$ -resolvable.  $\square$

To prove Theorems 1.4 and 1.5 we should recall some definitions and results from [6] and [5].

**Definition 4.2** ([6, Definition 3.1]). Let  $\kappa$  be an infinite cardinal, and let  $\mathcal{F}$  be a filter on  $\kappa$ . Let  $T$  be the tree  $\kappa^{<\omega}$ . A topology  $\tau_{\mathcal{F}}$  is defined on  $T$  by

$$\tau_{\mathcal{F}} = \{V \subset T : \forall t \in V \{\alpha \in \kappa : t \frown \alpha \in V\} \in \mathcal{F}\},$$

and the space  $\langle T, \tau_{\mathcal{F}} \rangle$  is denoted by  $X(\mathcal{F})$ .

*Proof of Theorem 1.4.* Let  $\mathcal{U}$  be a  $\kappa$ -complete non-principal ultrafilter on  $\kappa$ .

The space  $X = X(\mathcal{U})$  is monotonically normal by [6, Theorem 3.1].

An ultrafilter  $\mathcal{U}$  is  $\lambda$ -descendingly complete if  $\bigcap\{U_\zeta : \zeta < \lambda\} \neq \emptyset$  for each decreasing sequence  $\{U_\zeta : \zeta < \lambda\} \subset \mathcal{U}$ .

A  $\sigma$ -complete ultrafilter is clearly  $\omega$ -descendingly-complete. In the proof of [6, Theorem 3.5] the authors prove Lemma 3.6 which claims that  $\overline{\overline{\mathcal{D}}}$  is point-countable for each point-countable family  $\mathcal{D} \subset \mathcal{P}(X(\mathcal{F}))$  provided that  $\mathcal{F}$  is a  $\omega$ -descendingly complete ultrafilter. So  $\overline{\overline{\mathcal{D}}}$  is point-countable for each point-countable family  $\mathcal{D} \subset \mathcal{P}(X)$ , and so  $X$  is not monotonically  $\omega_1$ -resolvable by Lemma 4.1.  $\square$

Instead of Theorem 1.5 we prove the following theorem which is a slight improvement of [5, Theorem 5].

**Theorem 4.3.** *If it is consistent that there is a measurable cardinal, then it is also consistent that there is an  $\omega$ -resolvable monotonically normal space  $X$  with  $|X| = \Delta(X) = \omega_\omega$  such that if a family  $\mathcal{D} \subset \mathcal{P}(X)$  is point-countable, then the family  $\overline{\overline{\mathcal{D}}} = \{\overline{D} : D \in \mathcal{D}\}$  is also point*



countable. Hence  $X$  does not contain any monotonically  $\omega_1$ -resolvable subspace.

*Proof.* In [5, page 665] the authors write that "starting from one measurable, Woodin ([15]) constructed a model in which  $\aleph_\omega$  carries an  $\omega_1$ -descendingly complete uniform ultrafilter. Woodin's model  $V_1$  can be embedded into a bigger ZFC model  $V_2$  so that the pair of models  $(V_1, V_2)$  with  $\kappa = \aleph_\omega$  satisfies the two models situation", i.e.

- (1)  $\omega_1^{V_1} = \omega_1^{V_2}$ ,
- (2) there is a countable subset  $A$  of  $\omega_\omega$  in  $V_2$  such that no  $B \in V_1$  of cardinality  $< \omega_\omega$  covers  $A$ ;
- (3) for the filter  $\mathcal{G}$  on  $\omega_\omega$  defined in  $V_2$  by  $B \in \mathcal{G}$  iff  $A - B$  is finite, we have  $\mathcal{G} \cap V_1 \in V_1$ .

(the "two model situation" is defined in [5, Theorem 4.5]).

Let  $\mathcal{F} = \mathcal{G} \cap V_1$  and consider the space  $X = X(\mathcal{F})$ . As it was observed in [6], spaces obtained as  $X(\mathcal{H})$  from some filter  $\mathcal{H}$  are monotonically normal and  $\omega$ -resolvable.

In [5, Theorem 4.1] Juhász and Magidor showed that the space  $X(\mathcal{F})$  is actually hereditarily  $\omega_1$ -irresolvable. They proved the following lemma:

**Lemma 4.2 from [5].** For any  $D \subset X(F)$  and  $t \in \overline{D}$  there is a finite sequence  $s$  of members of  $A$  such that  $t \frown s \in D$ .

Using this lemma we show that  $\overline{\overline{D}}$  is point-countable for each point-countable family  $\mathcal{D} \subset \mathcal{P}(X)$ , and so  $X$  is not monotonically  $\omega_1$ -resolvable by Lemma 4.1.

Indeed, let  $\mathcal{D} \subset \mathcal{P}(X)$  be an uncountable family such that  $t \in \bigcap_{D \in \mathcal{D}} \overline{D}$ . Then, by [5, Lemma 4.3], for each  $D \in \mathcal{D}$  we can pick a finite sequence  $s_D$  of members of  $A$  such that  $t \frown s_D \in D$ . Since there are only countable many finite sequences of elements of  $A$  there is  $s$  such that  $s_D = s$  for uncountably many  $D \in \mathcal{D}$ . Then  $t \frown s$  is in uncountably many elements of  $\mathcal{D}$ , so  $\mathcal{D}$  is not point-countable.

So we proved that no subspace of  $X$  is monotonically  $\omega_1$ -resolvable.  $\square$

## 5. $\omega$ -RESOLVABILITY AFTER ADDING A SINGLE COHEN REALS

Before proving Theorem 1.6 we need some preparation.

The notion of almost resolvability was introduced by Bolstein ([2]) in 1973: a topological space is *almost-resolvable* if it is a countable union of sets with empty interiors. The notion of monotonically  $\omega$ -resolvability was first considered in [13] under the name almost- $\omega$ -resolvability.

Clearly almost  $\omega$ -resolvable (i.e. monotonically  $\omega$ -resolvable) spaces are almost resolvable.

**Lemma 5.1.** *Let  $X$  be a crowded topological space.*

- (1) *If  $X$  is monotonically  $\omega$ -resolvable, then  $X$  is  $\omega$ -resolvable in  $V^{\text{Fn}(\omega, 2)}$ .*
- (2) *If  $X$  is resolvable in  $V^{\text{Fn}(\omega, 2)}$ , then  $X$  is almost-resolvable.*

*Proof of Lemma 5.1.* (1) Assume that the function  $f : X \rightarrow \omega$  witnesses the monotonically  $\omega$ -resolvability of  $X$ .

If  $\mathcal{G}$  is the  $V$ -generic filter in  $\text{Fn}(\omega, \omega)$ , and  $g = \bigcup \mathcal{G}$ , then the function  $h = g \circ f$  witnesses that  $X$  is  $\omega$ -resolvable.

We need to show that  $\{y \in X : (g \circ f)(y) = n\}$  is dense in  $X$

Indeed, let  $p \in \text{Fn}(\omega, \omega)$ ,  $\emptyset \neq U \in \tau_X$ . Since  $f : X \rightarrow \omega$  witnesses the monotonically  $\omega$ -resolvability of  $X$  there is  $y \in U$  such that

$$f(y) > \max \text{dom}(p).$$

Let

$$q = p \cup \{\langle f(y), n \rangle\}.$$

Then  $q \leq p$  and

$$g \Vdash (g \circ f)(y) = n.$$

So we proved that  $g \circ f$  witnesses that  $X$  is  $\omega$ -resolvable in the generic extension.

(2) Assume

$$V^{\text{Fn}(\omega, 2)} \models "X \text{ has a partition } \{D_0, D_1\} \text{ into dense subsets}."$$

For all  $p \in \text{Fn}(\omega, 2)$  and  $i < 2$  let

$$D_i^p = \{x \in X : p \Vdash x \in \dot{D}_i\}.$$

Then  $X = \bigcup \{D_i^p : p \in \text{Fn}(\omega, 2), i < 2\}$ , and we claim that  $\text{int } D_i^p = \emptyset$  for each  $p \in \text{Fn}(\omega, 2)$ , and  $i < 2$ .

Indeed, fix  $p$  and  $i$  and let  $U$  be an arbitrary non-empty open subset. Then  $p \Vdash U \cap \dot{D}_{1-i} \neq \emptyset$ , so there is  $q \leq p$  and  $y \in U$  such that  $q \Vdash y \in \dot{D}_{1-i}$ . Then  $q \Vdash y \notin \dot{D}_i$ , so  $p \nVdash y \in \dot{D}_i$ , and so  $y \notin D_i^p$ . Thus  $U \not\subset D_i^p$ . Since  $U$  was arbitrary, we proved  $\text{int } D_i^p = \emptyset$ .  $\square$

After this preparation we can prove Theorem 1.6.

*Proof of Theorem 1.6.* (1) Kunen [7] proved that it is consistent, modulo a measurable cardinal, that there is a maximal independent family  $\mathcal{A} \subset \mathcal{P}(\omega_1)$  which is also  $\sigma$ -independent.

In [9, Theorems 3.1 and 3.2] the authors proved that if there is a maximal independent family  $\mathcal{A} \subset \mathcal{P}(\omega_1)$  which is also  $\sigma$ -independent, then there is a Baire space  $X$  with  $|X| = \Delta(X) = \omega_1$  such that every open subspace of  $X$  is irresolvable, i.e. the space  $X$  is *OHI*.

It is well-known that a crowded OHI Baire space  $X$  is not almost resolvable: if  $X = \bigcup_{n \in \omega} X_n$ , then  $\text{int } X_n \neq \emptyset$  for some  $n \in \omega$ .

Indeed, if  $\text{int } X_n = \emptyset$ , then  $X \setminus X_n$  is dense, so  $U_n = \text{int}(X \setminus X_n)$  is dense in  $X$  because every open subset of  $X$  is irresolvable. Thus  $\bigcap_{n \in \omega} U_n \neq \emptyset$  because  $X$  is Baire. However

$$\bigcap_{n \in \omega} U_n \subset \bigcap_{n \in \omega} (X \setminus X_n) = X \setminus \bigcup_{n \in \omega} X_n = \emptyset,$$

which is a contradiction.

Thus  $X$  is not almost resolvable, so it is not  $\omega$ -resolvable in the model  $V^{\text{Fn}(\omega, 2)}$  by Lemma 5.1(2).

(2) In [10] the authors proved that if  $V = L$ , then there are no crowded Baire irresolvable spaces. Hence, by [13], if  $V = L$ , then every crowded space  $X$  is almost- $\omega$ -resolvable (i.e. monotonically  $\omega$ -resolvable).

So these spaces are  $\omega$ -resolvable in the model  $V^{\text{Fn}(\omega, 2)}$  by Lemma 5.1(1).  $\square$

*Proof of Theorem 1.6(3).* Let  $X$  be a crowded c.c.c space.

We can assume that  $|X| = |\Delta(X)|$ .

By induction we define a strictly decreasing sequence of cardinals:

$$\kappa_0, \kappa_1, \dots, \kappa_n \dots$$

as follows.

- (i)  $\kappa_0 = \Delta(X)$ ,
- (ii) if  $\kappa_i$  is singular, then  $\kappa_{i+1} = \text{cf}(\kappa_i)$ ,
- (iii) if  $\kappa_i > \omega$  is regular, then  $\kappa_i = \lambda^+$  (because  $|X|$  is below the first weakly inaccessible cardinal,) and let  $\kappa_{i+1} = \lambda$ ,
- (iv) if  $\kappa_i = \omega$  or  $\kappa_i = \omega_1$ , then we stop.

Assume that the construction stopped in the  $n$ th step.

Then we can prove, by finite induction, then  $X$  is monotonically  $\kappa_i$ -resolvable for all  $i \leq n$  by theorem 3.3. Thus  $X$  is monotonically  $\omega$ -resolvable or monotonically  $\omega_1$ -resolvable, and so either  $X$  is  $\omega$ -resolvable in  $V^{\text{Fn}(\omega, 2)}$  by Lemma 5.1(1), or  $X$  is  $\omega_1$ -resolvable in  $V^{\text{Fn}(\omega_1, 2)}$  by Theorem 2.1.  $\square$

**Problem 5.2** ([13, Questions 5.2.]). *Are almost resolvability and almost- $\omega$ -resolvability equivalent in the class of irresolvable spaces?*

**Problem 5.3.** *Is there, in ZFC, a crowded topological space  $X$  which is irresolvable in the Cohen generic extension  $V^{\text{Fn}(\omega, 2)}$ .*

## REFERENCES

- [1] J. Angoa; M. Ibarra; Angel Tamariz-Mascarúa *On  $\omega$ -resolvable and almost- $\omega$ -resolvable spaces* Commentationes Mathematicae Universitatis Carolinae, Vol. 49 (2008), No. 3, 485–508
- [2] Richard Bolstein *Sets of Points of Discontinuity* Proceedings of the American Mathematical society 38.1 (1973), 193-197
- [3] A. Dorantes-Aldama, *Baire irresolvable spaces with countable Souslin number*, Topology Appl. 188, (2015) 16–26.
- [4] Hewitt, E. *A problem of set theoretic topology*, Duke Math. J. 10 (1943) 309-333.
- [5] István Juhász, Menachem Magidor, *On the Maximal Resolvability of Monotonically Normal Spaces* Israel Journal Of Mathematics 192 (2012), 637–666.
- [6] Juhász, István; Soukup, Lajos; Szentmiklóssy, Zoltán *Resolvability and monotone normality* Israel Journal of Mathematics 166 (2008), 1–16.
- [7] Kunen, Kenneth *Maximal  $\sigma$ -independent families*. Fund. Math. 117 (1983), no. 1, 75–80.
- [8] Kenneth Kunen, Karel Prikry, *On descendingly incomplete ultrafilters*, The journal of symbolic logic volume 36.4 (1971), pp 650–652.
- [9] K. Kunen, A. Szymanski, F. Tall *Naire irresolvable spaces and ideal theory* Ann. Math Syleziana 2(14) (1986) 98.107
- [10] K. Kunen, F. Tall, *On the consistency of the non-existence of Baire irresolvable spaces*, Manuscript privately circulated, Topology Atlas, <http://at.yorku.ca/v/a/a/a/27.htm>, 1998.
- [11] O. Pavlov, *Problems on (ir)resolvability*, in: Open Problems in Topology II, Elsevier, 2007, pp. 51–59.
- [12] *Recent Progress in General Topology, III* ed. K.P. Hart, Jan van Mill, P Simon Springer Science & Business Media, 2013, pp 903.
- [13] A. Tamariz-Mascarúa, H. Villegas-Rodríguez *em Spaces of continuous functions, box products and almost- $\omega$ -resolvable spaces* Comment.Math.Univ.Carolin. 43,4 (2002)687–705
- [14] S. Ulam, *Zur Masstheorie in der allgemeinen Mengenlehre*, Fundamenta Mathematicae, vol. 16 (1930), pp. 140-150.
- [15] W. H. Woodin, *Descendingly complete ultrafilter on  $\aleph_\omega$* , Personal communication.

ALFRÉD RÉNYI INSTITUTE OF MATHEMATICS, HUNGARIAN ACADEMY OF SCIENCES

*E-mail address:* soukup@renyi.hu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NORTHERN IOWA

*E-mail address:* adrienne.stanley@uni.edu