

Pleistocene vertebrate faunas of the Süttő Travertine Complex (Hungary)

*Piroska Pazonyi; László Kordos; Enikő Magyari;
Elena Marinova; Levente Fükőh; Márton Venczel*

Abstract

Numerous fossil remains (vertebrates, molluscs and plants) were found in more than twenty sites of the Süttő Travertine Complex during the last 150 years. The majority of these remains were recovered from fissures of the travertine, but also from the travertine and an overlying loess-paleosol sequence. The aims of this study were to review the fossil content, to determine the stratigraphical positions of the various vertebrate faunas of Süttő and provide paleoecological interpretation of the periods on the basis of their faunas and floras. In addition, this paper describes new faunas and floras from the sites Süttő 16–20 and provides ^{14}C dates for Süttő 16.

On the basis of the new uranium series isotope and optical dating (OSL), the age of the travertine complex is Middle Pleistocene (235 ± 21 – 314 ± 45 ka, MIS 7–9), while the age of the loess-paleosol sequence in superposition of the travertine is Middle-Late Pleistocene (MIS 2–MIS 6). In contrast, the fossils of the travertine indicated an older, Pliocene–Early Pleistocene age. A fissure (Süttő 17) and a red clay layer (Süttő 19) contained mammal faunas of Early–Middle Pleistocene age. These results indicated the existence of older travertine in certain quarries (Hegyháti quarry, Cukor quarry).

Sedimentological and OSL data of well-dated layers of the loess-paleosol sequence (Süttő/LPS) at Süttő allowed a correlation with the layers of Süttő 6. The paleosol layer in the upper part of the sequence of Süttő 6, was correlated with a pedocomplex of the overlying loess-paleosol sequence, which was dated to MIS 5c (upper, dark soil) and MIS 5e (lower, reddish brown soil). The paleoecological analysis of the mammal and mollusc faunas supported the former interpretation of Novothny et al., inferring warm, dry climate during the sedimentation of the upper layers, and more humid climate for the lower layers). However, the fauna of the lower soil layer indicated cold climate, so an age of MIS 5d is suggested.

Dating of the fissure faunas is based on similarity studies. For some faunas, this method cannot be used, because of the low number of species. On the basis of the species compositions and former interpretations, these faunas originated mainly from sediments that were deposited under cold climatic conditions. Other fissure faunas

were dated by AMS ^{14}C (Süttő 16), or by correlation with soil layers of Süttö 6. According to these results, most of the fissure faunas can be correlated with different phases of MIS 5. However, there are a younger (MIS 2) and an older (Early–Middle Pleistocene) fissure fauna.

1. Introduction

The Süttö Travertine Complex (northern Hungary) was known in the geological literature as a series of paleontological sites for more than a century. The age of this complex was discussed by several authors since the middle of the 19th century (Hantken, 1861; Hauer, 1870; Hofmann, 1884; Liffa, 1907; Kormos, 1911, 1913, 1926; Schréter, 1953; Scheuer and Schweitzer, 1974; Jánossy, 1969, 1979; Brunnacker et al., 1980; Jánossy and Krolopp, 1981; Novothny et al., 2009, 2011; Kele, 2009; Sierralta et al., 2010).

To the authors' knowledge, Kormos (1926) was the first who identified vertebrate remains in the travertine and its fissures. His observation, that the age of the travertine's vertebrates is much older than the age of the fissure infill faunas, was a good take-off for later research. In addition to the vertebrate sites, he also described two archaeological sites from Süttö (Fleissig and Kormos, 1934). He suggested that the fauna of Süttö is not very similar to the mammalian faunas of Brasov and Püspökfürdő (Kormos, 1926), and that it is younger than those. He placed the fauna at the end of the preglacial interval characterised by Mediterranean climate (Kormos, 1926, Kretzoi, 1927). In contrast, Kretzoi (1927) suggested that the fauna of Süttö originated from the last interglacial fauna of Germany and Austria that migrated to this area when the ice front expanded southward in Scandinavia.

In the 1970s, Jánossy and Krolopp made excavations in this area and described several new faunas from travertine fissure infills and one unique fauna from the loess-paleosol sequence that covers it (Jánossy, 1969, 1979; Brunnacker et al., 1980; Jánossy and Krolopp, 1981). This site (Süttő 6) is the type locality of the Süttö Biochronological Phase (Kretzoi, 1953). In the 1980s, Kordos and Krolopp discovered and excavated some further sites in the quarries (Süttő 16–20), which are not published yet.

The aim of this study is to review the fossil content of the various sites of the Süttö Travertine Complex, their correlation and to describe the vertebrate and mollusc faunas, as well as the macrocharcoal-based floras of Süttö 16–20 and correlate them with the formerly discovered sites from Süttö and other Hungarian localities.

2. Geological setting

The town of Süttő is located in northern Hungary, about 60 km northwest of Budapest, close to the right bank of the river Danube (47°44.26' N, 18°26.87' E). The vertebrate sites of Süttő are located in the Diósvölgyi, Hegyháti, Új Haraszti and Páchl quarries on the northern slope of the Haraszt Hill in the Gerecse Hills, part of the West Hungarian Mountain Range (Fig. 1).

Süttő is one of the largest travertine deposits in the Gerecse Hills. The travertine covers an area of more than 1 km². The immediate bedrock of the travertine is Upper Pannonian gravel, sand and clay, which were correlated with the VIIth terrace of the river Danube by Pécsi et al. (1982). The minimum age of the terrace is Late Pliocene, based on the record of *Anancus arvernensis* and *Tapirus arvernensis* found in the travertine in superposition of the fluvial sediments (Jánossy and Krolopp, 1981). The eroded surface of the travertine is covered by loose Upper Pleistocene eolian sediments: Riss-Würmian loess and Holocene topsoil. In the lower part of the loess section, a reddish paleosol horizon can be found, that developed during the last interglacial (Pécsi et al., 1982; Novothny et al., 2009, 2011). Petrographic description and interpretation of a site from the loess-paleosol sequence (Süttő 6) are in Table 1 (Brunnacker et al., 1980).

The fissures' infills also contain loess and paleosol layers, but red clay layers also occur. For example, the sequence of Süttő 16 is the following: red sandy loess, which was divided to three sublayers by visible remains ("csontos" – the lower layer wherein there are numerous bones (6); "marhás" – the middle layer wherein there are Bovidae remains (5); "eszközös" – the upper layer wherein there are tools (4)); soil, above the fissure, which was also divided into three sublayers on the basis of their colour ("legalsó" – the lightest lower layer (3); "sötét alatt középső" – the darker middle layer (2); "sötét talaj felső" – the darkest upper layer (1)) (Fig. 2). However, the sequence of Süttő 17 varies from the former one: the lower layer is red clay (17/C), and the upper one is sandy loess (17/L) (Fig. 3).

3. Material and methods

3.1. Fossil sites at Süttő

The known 20 fossil sites at Süttő (Fig. 1) are grouped into: (1) sites from the travertine; (2) sites from the loess-paleosol sequence (Table 2); and (3) sites from the fissures in the

travertine (Table 3).

3.1.1. Non-numbered sites from the travertine

Fossils from the travertine have been found/recovered during mining, but they have never been collected systematically. These fossils, mainly vertebrates, often were found far from the quarry, mainly near renovation works of buildings, or in separated travertine blocks. Therefore, the exact location of these fossils is unknown. The exact provenance of fossils collected by Kormos or deposited earlier in his collection (see Kormos, 1926) is unknown. However, the localities of some vertebrate remains, which were found in the Hegyhát quarry (*Anancus arvernensis*, *Dicerorhinus* cf. *jeanvireti*, *Cervus* cf. *philisi*), and that of one fossil (*Cervus* cf. *ardei*), which is known from the top layer of Gazda quarry (Jánossy and Krolopp, 1981), are known.

The molluscan fauna of the travertines was identified exclusively on the basis of impressions and casts. Former lists of the malacofauna were revised by Krolopp (in Jánossy and Krolopp, 1981).

3.1.2. Sites from the loess-paleosol sequence (Süttő 5–6)

Süttő 6 is the most important site from the loess-paleosol sequence. It is situated in the northwestern corner of the Diósvölgyi quarry, about 210 m above sea level, in sandy loess (Fig. 1). The upper layers of this profile yielded fossils (vertebrates and molluscs, referred to site Süttő 5), but Jánossy started his excavation in 1973 from the paleosol horizon situated in the lower tier of the profile. He burrowed a 5 m deep, 1 m wide and 8 m long trench to the surface of the travertine. Except for the paleosol horizon, the loess seemed to be uniform sandy loess. He collected material per 20–50 cm and separated 13 layers (Jánossy, 1979; Brunnacker et al., 1980).

Süttő 6 is unique because, in contrast to other loess sites, of the co-occurrence of molluscs plus a significant volume of vertebrate remains in each layer (Table 2). This site is the type locality of the Süttő Biochronological Phase (Kretzoi, 1953; Jánossy, 1979).

3.1.3. Sites from fissures in the travertine

3.1.3.1. Kormos' sites (Süttő 1–2)

Kormos (1926) reported vertebrates from the fissures at first. However, the site of these vertebrates was not assigned exactly in his works. According to Jánossy (1969), Kormos

collected his material from two places: Diósvölgy quarry (Süttő 1), and Pachtl quarry (Süttő 2). In addition, he supposed Süttő 4/2 (Jánossy and Krolopp, 1981) is also related to one of Kormos' sites (Fig. 1, Table 3). Although the vertebrate fauna which was published by Kormos (1926) seems mixed, it is very important, because he noted small-sized carnivores, which have not been recorded from other sites of Süttő.

3.1.3.2. Jánossy's and Krolopp's sites (Süttő 3–4 and 7–15)

Jánossy and Krolopp found and excavated several sites between 1965 and 1974 (Jánossy, 1979; Jánossy and Krolopp, 1981). Some of them contain snail faunas only (Süttő 10, 11, 14 and 15), but in most cases they also found vertebrate remains (Fig. 1, Table 3).

3.1.3.3. Kordos' and Krolopp's sites (Süttő 16–20)

In the late eighties (1986–1988), Kordos and Krolopp collected five new faunas (sites 16–20) from the northern part of Hegyháti quarry (Müller quarry), Diósvölgyi quarry and New Haraszti quarry. These sites are mainly fissures (Süttő 16, 17, 18 and 20), but one (Süttő 19) is a fossiliferous, red clay layer in the travertine.

Süttő 16 is a fissure on the north wall of Hegyháti quarry (Fig. 1). Kordos and Krolopp found vertebrates, molluscs and charcoal in each layer (Table 4). To avoid referring to the original difficult names, the layers are numbered as shown in Figure 2, and henceforth this paper refers to these names (Süttő 16/1–6).

Süttő 17 is a fissure infill in Hegyháti quarry (Fig. 1), with two layers (Fig. 3, Table 4).

Süttő 18 is also a fissure infill in Hegyháti quarry (Fig. 1). Unfortunately from this fissure, only bone fragments were found.

Süttő 19 is a red clay layer in the southern wall of New Haraszti quarry (Fig. 1). This site has a poor vertebrate fauna (Table 4).

Süttő 20 is a fissure infill in Diósvölgyi quarry, near site Süttő 4 (Fig. 1). Some vertebrate remains and molluscs were found at this locality (Table 4).

Vertebrates (*Pliomys* sp., *Mimomys pusillus*) were found in the southern part of the area in Cukor quarry (Fig. 1). Cukor quarry is a small, disused quarry at higher elevation (260–270 m a.s.l.) than the other quarries. Its travertine is harder than other travertines from Süttő, with abundant calcite veins, dripstone fragments and reddish colour because of its red clay content. Probably this material originated from a cemented, red clay cave fill (Fig. 4).

3.2. Methods

3.2.1. Similarity study

For the correlation of the faunal assemblages, correlation measures were used. Percentage similarity is an index that measures the common part of two temporally consecutive or spatially adjacent communities and is calculated as follows (Krebs, 1989):

$$P = \sum \text{minimum}(p_{1i}, p_{2i});$$

where P is the percentage similarity between community 1 and community 2, p_{1i} is percentage of the i^{th} species of community 1, and p_{2i} is percentage of the i^{th} species of community 2.

3.2.2. AMS ^{14}C Dating

For this study, ^{14}C dates from two layers of Süttő 16 provide an independent age control. The measurements were done in ATOMKI (Debrecen, Hungary) by AMS.

3.2.3. Wood charcoal analysis

The material for wood charcoal analysis comes from handpicked visible concentrations of charcoal from layers 2-6 of Süttő 16 (Fig. 2). The wood charcoals were analysed under a reflected light microscope Zeiss Axiscope MAT, to 400X magnification. All fragments bigger than 5 mm (in total 46) were fractured in transversal, radial and tangential sections in order to observe the corresponding diagnostic anatomical features of the wood and to identify them. For identification, wood anatomical atlases (Greguss, 1955; Schweingruber, 1990) and a wood reference collection from Europe and West Asia were used. The determined charcoal specimens are given as absolute numbers in Table 4. Light microscope images made at Leuven University are displayed in Figure 6.

4. Results

4.1. Dating and correlation of faunas

4.1.1. Dating of the travertine

More than 50 travertine sites are known from the northern part of Gerecse Hill. These sites were found along a NE–SW line (Scheuer and Schweitzer, 1988). They were formed during the Pliocene (62%) and the Quaternary (32.8%), and the most dynamic interval of travertine formation within the Quaternary was in the Lower Pleistocene (Scheuer, 2002). However, the Pliocene travertines cannot be differentiated clearly from the Pleistocene

travertines by their morphology and elevations in Gerecse Hill. In general, the older occurrences (older than the lower limit of the uranium-series dating method) are located above 250 m a.s.l., while the Middle Pleistocene travertines are situated between 250 and 150 m a.s.l. (Kele, 2009). In Scheuer and Schweitzer's (1981) opinion, the Middle Gerecse was the main drainage area in the Lower Pleistocene, and the travertine at Süttö formed at this time.

At the same time, new research (Kele, 2009; Sierralta et al., 2010) indicated a far younger, Middle Pleistocene age in the case of travertine at Süttö. Sierralta et al. (2010) published uranium-series ($^{230}\text{Th}/^{234}\text{U}$) dates of the travertine from two sections. On the basis of their examinations, the age of the travertine is mid- Pleistocene, ranging from 235 ± 21 to 314 ± 45 ka (296 ± 22 ka at Új Haraszti quarry, 422 ± 21 ka at Diósvölgyi quarry and 273 ± 65 ka at Hegyháti quarry), that can be correlated with Marine Isotope Stages (MIS) 7–9. The age of the overlying loess-paleosol sequence was determined by OSL dating. Novothny et al. (2009, 2011) placed the formation of the loess sequence into MIS 2–6.

These results at first sight are in conflict with the paleontological data. Ranges of vertebrates from the travertine indicated a Pliocene–Lower Pleistocene age. However, Kele did not examine travertines from the northern part of Hegyháti quarry, from where fossils with exact location are known (see above). It is possible that the travertine of this part of Hegyháti quarry is older, as its elevation is not as high as in other parts.

The fauna of Süttö 17 also indicates an older age of the northern part of Hegyháti quarry. This fauna is very similar to that of the site Somssich-hegy 2 (Jánossy, 1983). Its age, based mainly on arvicolids (*Mimomys savini*, *Microtus pliocaenicus*, *Pliomys episcopalis*, *P. lenki*, *Microtus gregaloides*), is at the boundary of the Lower to the Middle Pleistocene (about 800 ka). The fauna of Süttö 19 indicated an even older age (Lower Pleistocene). The new finds from the travertine of Cukor quarry (*Mimomys pusillus*, *Microtus pliocaenicus*), its elevation and this travertine's differing appearance (pink, more compact and harder than the other travertines at Süttö) also suggests the existence of an older (Pliocene/Lower Pleistocene) travertine at Süttö.

4.1.2. Dating and correlation between the loess-paleosol sequence and Süttö 6

In previous years, several papers were published about a loess-paleosol sequence of Süttö (Süttö/LPS) (Novothny et al., 2009, 2011). The loess deposits are up to 20 m thick, and

contain two greyish, laminated horizons, three brownish steppe-like soils and a thick pedocomplex including a dark brown chernozem-like paleosol and a reddish-brown paleosol (Novothy et al., 2011). Sedimentological studies and luminescence ages, mainly from the lower tiers of the section (Unit 14–16), are of importance.

This part of the section contains the pedocomplex (Unit 14–15, 1.5 m) and the lowest loess layer (Unit 16, 4 m). In the former literature about Süttő 6 (Brunnacker et al., 1980; Jánossy, 1979), the similarity of the two sections is apparent. Both of them are about 5 m thick, their top layers are paleosols, and both are underlain by loess. On the basis of their sedimentological descriptions (and vertebrate remains in case of Süttő 6), the units of the loess-paleosol section were correlated with the layers of Süttő 6 (Fig. 5).

At Süttő 6, there are five paleosol layers at the top of the section underlain by sandy loess layers. Based on their sedimentology and vertebrate material, the paleosol layers were divided in two parts. Layers 6/2–3 consist of a dark brown chernozem (1 m). Layers 6/4–5 are light reddish brown, chalk enriched horizons (C_c -horizon) (Jánossy, 1979; Brunnacker et al., 1980). The proportion of the steppe species is higher (61%), while the proportion of the forest species is lower (25%) in layers 6/2–3 than in layers 6/4–5 (56% steppe and 33% forest species) (Table 5). The mollusc fauna of this site also showed significant difference between layers 6/2–3 and 6/4–5. The molluscs of layers 6/2–3 indicated warm and dry climate (proportion of xerotherm species was 18% and cold indicator species are missing). In layers 6/4–5, the mollusc fauna indicated a slightly more humid, but rather dry and cold climate (proportion of cold indicator species was 40%) (Table 5).

These results can be correlated with Units 14–15 of the Süttő loess sequence (Novothy et al., 2011) and their paleoclimatic and environmental interpretations (warm, dry climate during the sedimentation of the upper layers, and more humid climate for the lower layers). Units 14–15 were dated at 93.7 ± 21.1 ka and 106 ± 13 ka/ 137 ± 23 ka, respectively in the loess sequence. Layers 6/2–3 in Süttő 6 correspond with MIS 5c, while layers 6/4–5 correspond rather with MIS 5d than MIS 5e, because their fauna suggested rather cold climates. The lower loess layers were correlated with Unit 16 (MIS 6).

4.1.3. Dating and correlation of the fissures' faunas

Stratigraphic correlation of the fissure faunas is based on similarity studies. For some faunas, this method could not be used, because of the small number of species. On the

basis of their species composition and former interpretations, these faunas originated mainly from sediments, which were deposited under cold climatic conditions (Süttő 4/1, 8/1, 8/3 and 13). However, some small faunas deposited under warmer climate (Süttő 4/2, 8/3 and 16/1) yielded insufficient material for analysis. Fauna of Süttö 3 is also poor, but this fauna was fairly well correlated with other faunas based on the appearance of an index fossil (*Dama* sp.).

In the course of correlation studies, suitable faunas from the site Süttö 6 were compared with other mammalian faunas with similar ages (MIS 5), and with each other. Süttö 16 is, based on ^{14}C data, younger than the others, so this site was studied separately.

Similarity studies of small mammalian faunas showed strong connections between Süttö 6 and Süttö 3, 7/U and 9 (percentage similarity >60%). Between Süttö 6 and Süttö 7/L and 12A the percentage similarity is about 50%, whereas the site 12/B did not show correspondence with the site Süttö 6 (Table 6).

The mammalian faunas (including site 16) were divided into three groups: (1) proportion of steppe species is far higher (50–55%) than proportion of forest species (15–30%) (Süttő 9 and 16/2); (2) proportion of forest species is far higher (40–65%) than proportion of steppe species (16–40%) (Süttő 3 and 7/L) and (3) proportions of steppe and forest species are nearly equal (20–40%) (Süttő 7/U, 12/A, 16/3 and 16/5–6). These groups coincide with the results of comparison of the faunas. Within the first and second groups the percentage similarity values are about 60%, while in case of the third group, the values are about 50–55% (Table 7).

However, in the mollusc faunas there are slightly different groups. Most mollusc faunas indicated warm and dry climatic conditions (site 3, 9, 12/A and 12/B), whereas in sites 16 and 7/U they indicated cold and humid climates. The fauna of site 7/L, based on its mollusc fauna, arose in warm and humid circumstances (Table 7).

Based on these (mainly mollusc) results, faunas of Süttö 3, 9 and 12/A were correlated with Süttö 6/2–3 (although the site Süttö 9 also showed strong correlation with other faunas of the same region: Diósgyőr-Tapolca Caves (Hellebrandt et al, 1976), as well as Eger, Dobó-bástya (Kordos and Krolopp, 1980)). The result of this analysis showed cold and humid climatic conditions at Süttö 7/U, a site also well correlated with Süttö 6. Accordingly, this fauna was correlated with Süttö 6/4–5.

In the mammalian fauna of Süttö 7/L, the percentages of steppe and forest species are in contrast to values of Süttö 6. However, site 7/L was well correlated with Tarkő IV

(Jánossy, 1979) (MIS 5c). Beside this, the proportion of forest species is far higher (43%) than proportion of steppe species (17%) in the mammalian fauna of Tarkő IV, even as in the group 2 (Table 7). The mollusc fauna of Süttő 7/L also indicated a warm and humid climate. This result indicates that the age of these sites is probably not MIS 5c, which climate was warm and dry, but more likely MIS 5e.

Results of similarity study showed that mammalian fauna of site 12/B did not show strong correspondence with other faunas, although its similarity with the fauna of site 12/A is about 54%. However, its mollusc fauna indicated a more humid climate than site 12/A. The proportions of steppe (15%) and forest species (52%) in this fauna are similar to the Tarkő IV, Süttő 3 and 7/L. Although the connections between the mammalian fauna of 12/B and these sites are fairly weak, they were tentatively correlated. The correlation results showed that all the mammalian faunas, which were deposited in warmer climate conditions, might be correlated with MIS 5.

4.1.4. Dating of Süttő 16

In order to obtain numerical ages for the floras of Süttő 16/3 and 16/6, 2 charcoal pieces were selected for AMS ^{14}C dating. For Süttő 16/3, the measurement was made on *Pinus* sp., while the other was on *Betula* sp. The results are as follows:

Süttő 16/3 (DeA-1319) — 12499±54 conv. BP (1 σ); 15050–14590 cal BP (1 σ), 14430–14350 cal BP (1 σ);

Süttő 16/6 (DeA-1320) — 14079±63 conv. BP (1 σ); 17340–17030 cal BP (1 σ).

Three vertebrate (mainly mammal) cave localities (Peskő, Jankovich and Bivak Caves) are known from this period in Hungary. Comparing these with the mammalian fauna of Süttő 16, there is similarity with the sequence of Peskő Cave (Hír, 1991) only. This result was surprising, because the Bivak (Jánossy et al., 1957) and Jankovich Caves (Bácskay and Kordos, 1984) (Pilis and Gerecse Hills) are located very near to Süttő, whereas the Peskő Cave is in the Bükk Mountain. Based on Pazonyi (2006), the age of the top (yellow) layer of Bivak Cave is 15970±270 conv. BP, whereas the ages of Jankovich Cave are 11720±190 conv. BP (layer 7) and 12440±230 conv. BP (layer 8).

On the basis of the ^{14}C dates, the macrocharcoal assemblages of Süttő 16/4 postdate the last glacial maximum (LGM) and were likely deposited during the Oldest Dryas (late MIS 2). Overall, the layers of Süttő 16 are correlated with the top paleosol and loess layers of the loess- paleosol sequence (Unit 1–2) (Novothy et al., 2011) (Fig. 5).

4.2. *Paleoecological analysis*

According to their assumed ecological preferences, the mammalian community from the older (Pliocene/Lower Pleistocene) travertine, allude unequivocally to warm, humid climate and closed forest vegetation during its formation. Unfortunately, no vertebrate fossils have been recovered from the younger (Middle Pleistocene) travertine, but based on their Th/U ages and isotope geochemical ($\delta^{13}\text{C}$ and $\delta^{18}\text{O}$) results of Kele (2009), it was formed during the cold, but humid interval of MIS 7–9.

The mammalian and mollusc fauna of Süttő 6 and the sedimentological analysis of the loess-paleosol complex indicate warm/dry climate and steppe-like vegetation for layers 6/2–3 (MIS 5c), with colder and more humid climate during the formation of layers 6/4–5 (MIS 5d) (Table 5). The mammalian and mollusc faunas, in Süttő 3, 9 and 12/A indicate similar climatic conditions and vegetation as during the formation of Süttő 6/2–3. Süttő 7/U was formed under colder and humid circumstances, similar to Süttő 6/4–5 (Table 7).

Paleoecological analysis of the mammalian fauna of Süttő 16 also showed colder and humid climatic conditions (Table 8). The charcoal assemblages suggest the prevalence of boreal forest vegetation around the locality with birch (*Betula* sp.), pine (*Pinus* sp.), alder (*Alnus* sp.), spruce (*Picea* sp.) and larch (*Larix* sp.) (Fig. 6). Overall they suggest a cold continental climate during the terminal phase of MIS 2 (Oldest Dryas). The presence of spruce suggests that available moisture was relatively high (Rudner and Sümegei, 2001) in this period. Results of herpetological analysis of Süttő 16 also suggested cold and humid climate with parkland vegetation, in line with the results of the anthracological analysis. In contrast to this, the mammalian and mollusc fauna of Süttő 7/L indicated warm and humid climate and closed forest vegetation (Table 7).

In the mammalian fauna of site Süttő 17 (Early Biharian, MIS 22), the steppe species are dominant (60%), and the proportion of forest species is 40%. This result indicates probably steppe or parkland vegetation. This inference was supported by results of the herpetological analysis (dry, temperate climate, grassland vegetation).

On the basis of the results of the herpetological analysis, the fauna of site Süttő 19 (Late Villanyian–Early Biharian) was formed under warm (warmer than today), very dry (Mediterranean-like) climate and shrub-forest vegetation. The herpeto- and mollusc fauna of site Süttő 20 indicated temperate and humid climate and forest vegetation.

5. Conclusions

The data suggest the existence of older, Pliocene/Lower Pleistocene travertine deposits at Süttő in the north part of Hegyháti quarry and in Cukor quarry. These quarries have not yet been dated by U-Th. In addition to the vertebrates of the travertine, mammals of Süttő 17 and 19 also have signalled older ages.

The layers of Süttő 6 are correlated with parts of the loess-paleosol sequence (Units 14–16) (Novothny et al., 2011). Layers 6/2–3 are well correlated with Unit 14. In Novothny et al. and the authors' opinion, this unit corresponds to MIS 5c. This period was also correlated with some fissure faunas: Süttő 3, 9, 12A and probably 12/B (Fig. 7). According to Novothny et al.'s (2011) interpretation and based on the mammalian and snail faunas, it was a warm and dry period with steppe-like vegetation.

Other faunas showed strong connection to the site Süttő 9. Ringer and Moncel (2002) reviewed Taubachian tools from those layers of the Diósgyőr-Tapolca Cave with fossils (layers 4–5). The Taubachian culture is typical of MIS 5, so this data gave the age of the site. The mammalian fauna of Eger, Dobó-bástya is also correlated with MIS 5 (Pazonyi and Kordos, 2004).

Layers 6/4-5 of Süttő 6 are correlated with Unit 15 of the Süttő loess-paleosol complex. In Novothny et al.'s (2011) opinion, this unit corresponds to MIS 5e, but on the basis of the fauna it is suggested here to correspond to MIS 5d. In this period the climate was more humid than for Unit 14, but still rather dry and cold. The fissure fauna of Süttő 7/U is also correlated with this period. Layers 6/6–10 were correlated with Unit 16 (MIS 6) (Fig. 7).

Süttő 7/L, and the very similar Tarkó IV, are potential faunas of MIS 5e, which indicate warm, humid climate and closed forest vegetation.

Fauna of Süttő 17 is very similar to that of the site Somssich-hegy 2 (Jánossy, 1983), which is dated, mainly based on arviculids, to the Early/Middle Pleistocene boundary. Süttő 16 is a younger site, correlated with Unit 1–2 of the loess-paleosol sequence (Novothny et al., 2011). Its age, based on the ^{14}C dating, is 17290–14300 cal BP (Fig. 7). Paleoecological analysis of its fauna and flora showed cold, humid climate and boreal parkland vegetation.

Wood charcoal and macrofossil assemblages dating to the Oldest Dryas are relatively scarce in the Carpathian Basin and adjoining areas. The Süttő 16 wood assemblage is in agreement with the indication in other sites: a parkland boreal landscape with increasing

coverage of *Betula* in addition to *Pinus* species and *Picea abies* (e.g. Budapest-Csillaghegy, Balatonederics in Rudner and Sümegei 2001; Balatonederics in Juhász, 2007; Jakab, 2007), *Picea*, *Pinus cembra* and *Salix* (Poland, Damblon and Haesaerts, 1997; Haesaerts et al., 1998), and *Pinus cembra* and *Larix* (Slovak Tatra Mts, Jankovska and Pokorny, 2008; Kunes et al., 2008). Comparing the wood charcoal data with contemporaneous Transdanubian pollen records, the main difference concerns the scattered prevalence of relatively thermophilous temperate broad-leaved trees (*Corylus*, *Ulmus*, *Quercus*, *Fagus*) in several Oldest Dryas pollen records (e.g. Sárrét, Balatonederics, Mezölak, Willis et al. 2000; Juhász and Szegvári, 2007; Juhász 2007), which, however, have not yet been confirmed by wood charcoal or macrofossils. The Süttö wood charcoal flora holds no evidence for their local occurrence either. Comparing the full glacial (LGM) woody cover of the Carpathian Basin with the Oldest Dryas (OD), the main difference is the increasing coverage of *Betula* during the OD that is in line with its increasing pollen frequency in several pollen records (Járai-Komlódi, 2003, Feurdean et al., 2007; Juhász and Szegvári, 2007), but otherwise the wood assemblages are similar. Notable is the abundance of *Picea* in Süttö 16 that seems typical for MIS 3-2 wood charcoal assemblages in the hilly area of Northern Transdanubia (Rudner and Sümegei, 2001). Mixed boreo-nemoral parkland forests were likely widespread in this region.

Based on former investigations (Pazonyi, 2004, 2006, 2011), in the Oldest Dryas the composition and the paleoecological features of the mammalian fauna also changed in the Carpathian Basin. The composition of fauna indicates warmer and wetter climate with more closed (forest-steppe) vegetation, but the area of tundra vegetation was probably significant (Pazonyi, 2004, 2011). These observations are in agreement with the paleoecological results of the current studies of the Süttö 16 fauna.

Earlier pollen and plant macrofossil studies in the Süttö 6 profile provided a relatively detailed picture of the last interglacial vegetation (Brunnacker et al., 1980). Pollen assemblages were studied in the MIS 5c and MIS 5d paleosols and the underlying loess horizons of Süttö 6, and yielded mainly entomophily herb rich pollen floras dominated by the Compositae subfamily Tubuliflorae. Brown paleosols likely dating to MIS 5c were abundant in *Juglans* pollen and contained fruits of *Celtis australis* and *Crataegus* sp. (Brunnacker et al., 1980). *Juglans* was considered secondary in the pollen assemblages. However, the association of these trees and shrubs suggests warm Sub-Mediterranean climate and at least partially forested landscape that agrees well with the fauna-based

environmental inferences for this interglacial period. The reddish brown paleosol horizon was rich in *Vitis* seeds (both *V. vinifera* and *V. sylvestris*; Brunnacker et al., 1980) and inferred to liana rich forested environment probably near to a water body. These last interglacial botanical records are important additions to the Eemian floras of the Carpathian Basin, summarized by Járαι-Komlódi (2003). The richest pollen and macroflora was described from the travertine deposits of Tata (19 km SW from Süttő; Vértés, 1964). Here, *Celtis* was also recorded in the warmest phase of the interglacial together with Cupressaceae, *Biota*, *Corylus*, *Cornus* and *Rhamnus*. Similar to this study's interpretation, Járαι-Komlódi (2003) has inferred strong Mediterranean climatic influence in this period, but also emphasized the warm-microclimate of the area due to the presence of thermal springs. Brunnacker et al. (1980) made some important inferences regarding the MIS 6 (late Saalian glacial) steppe vegetation of Süttő as well. First of all, they argued that the absence of *Artemisia* and Chenopodiaceae pollen and the scarcity of Poaceae pollen in the loess deposit preclude the presence of periglacial sagebrush and grass steppes in this region unlike in Austria. Second, they argued that the high relative frequency of *Centranthus* sp., a typical Southern European drought-tolerant but warm-loving herb genus, suggests herb/forb rich warm steppe vegetation during MIS 6.

The composition of the Süttő 6 faunas and the current results is partly in agreement with these observations. All observations allude to warm and dry (Sub-Mediterranean) climate and steppe-like, but partially forested landscape, during MIS 5c (Süttő 6/2–3). Paleoecological interpretation of MIS 5d is unequivocal as well. There were more humid, but rather dry and cold climate and liana rich forested vegetation, probably near to a water body (Süttő 6/4–5). However, the current results do not support the former results. Mammalian and mollusc faunas of Süttő 6/6–10 (MIS 6) unequivocally indicate cold climate (*Microtus gregalis* was dominant in the mammalian fauna and *Dicrostonyx torquatus* also present; 68% of the mollusc fauna was oligothermic), so the former warm steppe hypothesis is untenable. Probably, there was cold, periglacial steppe in this region, but around the thermal springs, herb/forb rich warm steppe vegetation also was present.

Revision of the classical and studies of newly discovered vertebrate faunas from the Carpathian Basin allowed the reconstruction of an almost complete succession from the Late Saalian to the Early Weichselian (Pazonyi and Kordos, 2004):

Late Saalian–Eemian transition (MIS 6) — Type fauna Süttő 6, layers 6–10. The lowermost layer contains *Dicrostonyx* and *Lagopus muta* remains, while *Microtus*

gregalis decreases and is replaced in predominance by *M. arvalis*. The current results completed this idea. The periglacial steppe was changed to warmer steppe vegetation in the surroundings of the thermal springs.

Eemian (MIS 5e) — Formerly, Süttő 6 layers 2–5 was considered the type fauna of this period. On the strength of the current results, these layers do not represent MIS 5e, but MIS 5d and MIS 5c. Instead of Süttő 6, the localities from the last interglacial are the karst fissure faunas of Süttő 7/L and Tarkő IV. These faunas are dominated by *Apodemus sylvaticus*, *Myodes glareolus*, glirids and insectivores, without any significant steppe and cool elements. Another locality from the Carpathian Basin is Por-lyuk at Jósvalő (Jánossy et al., 1973) which has a similar faunal character with a series of forest elements (*Myodes glareolus*, murids and glirids) in combination with some steppe taxa (*Spermophilus*, *Sicista*, *Spalax*, *Ochotona*).

Eemian–Weichselian transition (MIS 5d, MIS 5c) — This transition period is characterised, on the basis of current results, by most of the faunas of Süttő (Süttő 6/4–5, 7/U (MIS 5d); Süttő 3, 6/2–3, 9, 12/A (MIS 5c)), and several newly discovered faunas, as Horváti-lik at Uppony (Füköh and Kordos, 1977, 1980; Pazonyi and Kordos, 2004), Eger Dobó-bástya (Kordos and Krolopp, 1980), Tatabánya Kálvária 4. (Kordos, 1994), Poros-lyuk at Répáshuta (Jánossy, 1979), and Bajót 3. rock-shelter 5a layer (Kordos, 1994). The faunas of these localities appear to fill the 25–30 ky gap between the end of the Eemian Süttő Phase (MIS 5e) and the Early Weichselian Varbó Phase (MIS 4) in Hungary.

Acknowledgements

We would like to thank to Mihály Molnár for the ^{14}C measurements. This is MTA-MTM Paleo contribution No.175.

References

- Bácskay, E., Kordos, L., 1984. Fontosabb szórványleletek a MÁFI gerinces-gyűjteményében. MÁFI Évi Jelentése az 1982. évről, Budapest, pp. 355–361.
- Brunnacker, K., Jánossy, D., Krolopp, E., Skoflek, I., Urban, B., 1980. Das jungmittelpleistozäne Profil von Süttő 6 (Westungarn). *Eiszeitalter und Gegenwart* 30, 1–18.

- Damblon, F., Haesaerts, P., 1997. Radiocarbon chronology of representative Upper Palaeolithic sites in the central European Plain: a contribution to the SC-004 project. *Préhistoire Européenne* 11, 255–276.
- Feurdean, A., Wohlfarth, B., Björkman, L., Tantau, I., Bennike, O., Willis, K.J., Farcas, S., Robertsson, A.M., 2007. The influence of refugial population on Late glacial and early Holocene vegetational changes in Romania. *Review of Palaeobotany and Palynology* 145, 305–320.
- Fleissig, J., Kormos, T., 1934. Die ältesten Menschenspuren in Ungarn. *Arbeiten des Archaeologischen Instituts der Kön. Ung. Franz-Josef Universität in Szeged* 9–10(1–2), 16–29.
- Fűköh, L., Kordos, L., 1977. Jelentés az Uppony Horváti-lik 1977. évi őslénytani ásatásáról [Bericht über die paläontologische Ausgrabung in Uppony Horváti-lik, im Jahre 1977]. *Egri Múzeum Évkönyve* 15, 21–32.
- Fűköh, L., Kordos, L., 1980. Jelentés az Uppony Horváti-lik 1978. évi őslénytani ásatásáról [Report on the paleontological excavation on the Horváti-hole (Uppony) in 1978]. *Egri Múzeum Évkönyve* 16–17, 21–43.
- Greguss, P., 1955. *Xylotomische Bestimmung der heute lebende Gymnospermen*. Akadémiai Kiadó, Budapest.
- Haesaerts, P., Borziak, I., Van der Plicht, J., Damblon, F., 1998. Climatic events and Upper Paleolithic chronology in the Dniester Basin: new ^{14}C results from Cosautsi. *Radiocarbon* 40, 649–657.
- Hantken, M., 1861. *Geológiai tanulmányok Buda és Tata között*. *Akad. Math. és Term.-tud. Közl.* 1, 213–279.
- Hauer, F., 1870. *Geologische Uebersichtskarte der österreichisch-ungarischen Monarchie*. Blatt VII. Ungarisches Tiefland. *Jahrb. k. k. Geol. Reichsanst.* 20(4), 463–500.
- Hellebrandt, M., Kordos, L., Tóth, L., 1976. A Diósgyőr-Tapolca-barlang ásatásának eredményei. *A Herman Ottó Múzeum Évkönyve*, pp. 7–36.
- Hír, J., 1991. Rétegazonosító ásatás a Pes-kő-barlangban. *Folia Historico-Naturalia Musei Matraensis* 16, 5–12.
- Hofmann, K., 1884. A Duna jobb partján Ó-Szőny és Piszke közt foganatosított földtani részletes felvételről. *Földtani Közlöny* 14, 174–190.
- Jakab, G. 2007. The macrobotanical remains from Balatonederics. In: Zatykó, Cs., I. Juhász and P. Sümegei (Eds.), *Environmental Archaeology in Transdanubia*. *Varia*

Archaeologica Hungarica, Budapest, pp. 63–67.

Jankovská, V., Pokorný, P., 2008. Forest vegetation of the last full-glacial period in West Carpathians, Slovakia and Czech Republic. *Preslia* 80, 307–324.

Jánossy, D., 1969. Stratigraphische Auswertung der europäischen mittelpleistozänen Wirbeltierfauna. Teile I.–II. *Berichte der Deutschen Gesellschaft für Geologische Wissenschaften. Reihe A, Geologie und Paläontologie* 14(4–5), 367–438, 573–643.

Jánossy, D., 1979. A magyarországi pleisztocén tagolása gerinces faunák alapján. Akadémiai Kiadó, Budapest, 207 pp.

Jánossy, D., 1983. Lemming-remain from the Older Pleistocene of Southern Hungary (Villány, Somssich-hegy 2). *Fragmenta Mineralogica et Palaeontologica* 11, 55–60.

Jánossy, D., Varrók, S., Herrmann, M., Vértes, L., 1957. Forschungen in der Bivakhöhle. *European Union of Geosciences* 8, 18–36.

Jánossy, D., Kordos, L., Krolopp, E., Topál, Gy., 1973. The Porlyuk Cave of Jósvafő. *Karszt- és Barlangkutatás* 7, 15–50.

Jánossy, D., Krolopp, E., 1981. Die pleistozänen Schnecken- und Vertebraten-Faunen von Süttő (Travertine, Deckschichten und Spalten). *Fragmenta Mineralogica et Palaeontologica* 10, 31–58.

Járai-Komlódi, M., 2003. Quaternary vegetation history in Hungary. *Theory–Methods–Practice* 59, Geographical Research Institute, Research Centre of Earth Sciences, Hungarian Academy of Sciences, Budapest.

Juhász, I., 2007. The pollen sequence from Balatonederics. In: Zatykó, Cs., I. Juhász and P. Sümegei (Eds.), *Environmental Archaeology in Transdanubia. Varia Archaeologica Hungarica*, Budapest, pp. 57–63.

Juhász, I., Szegvári, G., 2007. The pollen sequence from Mezőlak. In Zatykó, Cs., I. Juhász and P. Sümegei (Eds.), *Environmental Archaeology in Transdanubia. Varia Archaeologica Hungarica*, Budapest, pp. 314–316.

Kele, S., 2009. Édesvízi mészkövek vizsgálata a Kárpát-medencéből: paleoklimatológiai és szedimentológiai vizsgálatok (Study of freshwater limestones from the Carpathian basin: paleoclimatological and sedimentological implications). Ph.D. thesis, Eötvös University, Budapest (in Hungarian).

Kordos, L., 1994. A gerecsei barlangok ősgerinces kutatásának újabb eredményei (1970–1994), [New records of the palaeovertebrate research of the caves of Gerecse Mts., 1970–1994]. *Limes* 94(2), 93–112.

- Kordos, L., Krolopp, E., 1980. Felső-pleisztocén forrásmész-kő-üledék mollusca- és gerinces faunája az egri Dobó-bástya területéről. *Folia Historico-Naturalia Musei Matraensis* 6, 5–12.
- Kormos, T., 1911. Une nouvelle espèce de tortue (*Clemmys méhelyii* n. sp.) du pleistocène hongrois. *Földtani Közlöny* 41, 506–512.
- Kormos, T., 1913. Kleinere Mitteilungen aus dem ungarischen Pleistocän. *Centralbl. Min. Geol. Pal.*, 13–17.
- Kormos, T., 1926. Die Fauna des Quellenkalk-Komplexes von Süttő. *Állattani Közlemények* 22(3–4), 248–253.
- Krebs, J.R., 1989. *Ecological methodology*. Harper and Row Publishers, New York, 654 pp.
- Kretzoi, M., 1927. Kormos T.: A süttöi forrásmész-kő-komplexus faunája. Die Fauna des Quellenkalk-Komplexes von Süttő. (*Állattani Közl.* XXII. 159-175) [Review]. *Barlangkutatás* 14–15, 30–31 and 100–101.
- Kretzoi, M., 1953. A negyedkor taglalása gerinces faunák alapján. A MTA Műszaki Tud. Oszt. Alföldi Kongresszusa, Budapest, 89–99.
- Kuneš, P., Penánková, B., Chytrý, M., Jankovská, V., Pokorný, P., Petr, L., 2008. Interpretation of the last-glacial vegetation of eastern-central Europe using modern analogues from southern Siberia. *Journal of Biogeography* 35, 2223–2236.
- Liffa, A., 1907. Geologische Notizen aus der Umgebung von Nyergesujfalu und Neszmély. *Jahresbach Ung. Geologie Reichsanst. für 1907*, 168–192.
- Novothny, Á., Frechen, M., Horváth, E., Bradák, B., Oches, E.A., McCoy, W.D., Stevens, T., 2009. Luminescence and amino acid racemization chronology of the loess–paleosol sequence at Süttő, Hungary. *Quaternary International* 198, 62–76.
- Novothny, Á., Frechen, M., Horváth, E., Wacha, L., Rolf, C., 2011. Investigating the penultimate and last glacial cycles of the Sütto loess section (Hungary) using luminescence dating, high-resolution grain size, and magnetic susceptibility data. *Quaternary International* 234, 75–85.
- Pazonyi, P., 2004. Mammalian ecosystem dynamics in the Carpathian Basin during the last 27,000 years. *Palaeogeography, Palaeoclimatology, Palaeoecology* 212, 295–314.
- Pazonyi, P., 2006. A Kárpát-medence kvarter emlősfauna közösségeinek paleoökológiai és rétegtani vizsgálata (Palaeoecological and stratigraphical investigations of the Quaternary mammalian communities in the Carpathian Basin). Ph.D. thesis, Eötvös

University, Budapest (in Hungarian).

Pazonyi, P., 2011. Palaeoecology of Late Pliocene and Quaternary mammalian communities in the Carpathian Basin. *Acta zoologica cracoviensia* 54A(1–2), 1–29.

Pazonyi, P., Kordos, L., 2004. Late Eemian (Late Pleistocene) vertebrate fauna from the Horváti-lik (Uppony, NE Hungary). *Fragmenta Palaeontologica Hungarica* 22, 107–117.

Pécsi, M., Scheuer, Gy., Schweitzer, F., 1982. Geomorphological and chronological classification of Hungarian travertines. *Quaternary Studies in Hungary*. Geographical Research Institute of the Hungarian Academy of Sciences, Budapest. 113–117.

Ringer, Á., Moncel, M.–H., 2002. Le Taubachien dans la Grotte Diósgyőr-Tapolca (Montagne de Bükk, Hongrie du Nord-Est). *Praehistoria* 3, 177–201.

Rudner, Z.E., Sümegei, P. 2001. Recurring taiga forest-steppe habitats in the Carpathian Basin. *Quaternary International* 76-77, 177–189.

Scheuer, Gy., 2002. A nyugat-gerecsei pliocén és quarter mészképző hévforrások paleokarszthidrológiai vizsgálata. *Hidrológiai Közlöny* 2002(1), 7–14.

Scheuer, Gy., Schweitzer, F., 1974. New aspects in the formation of the fresh-water limestone series of the environs of Buda-Mountains. *Földrajzi Közlemények* 22(2), 113–134.

Scheuer, Gy., Schweitzer, F., 1981. A Gerecse-hegység paleokarszthidrológiai viszonyainak rekonstrukciója a felső-pannontól napjainkig. *Hidrológiai Közlöny* 61(8), 333–343.

Scheuer, Gy., Schweitzer, F., 1988. A Gerecse és a Budai-hegység édesvízi mészkőösszletei. *Földrajzi Tanulmányok* 20, Akadémiai Kiadó, Budapest. 131 p.

Schréter, Z., 1953. Les occurrences de calcaire d'eau douce des bords des montagnes de Buda et Gerecse. *Jahresb. Ung. Geol. Anst. für 1951*, 111–150.

Schweingruber, F., 1990. *Anatomie europäischer Hölzer*. Haupt, Stuttgart.

Sierralta, M., Kele, S., Melcher, F., Hambach, U., Reinders, J., van Geldern, R., Frechen, M., 2010. Characterisation and Uranium-series dating of Travertine from Süttő in Hungary. *Quaternary International* 222, 178–193.

Vértes, L. (Ed.), 1964. Tata, eine mittelpaläolithische Travertin-Siedlung in Ungarn. *Archaeologica Hungaria Serie Nova* 43, 1–284.

Willis, K.J., Rudner, E., Sümegei, P., 2000. The full-glacial forests of central and southeastern Europe: Evidence from Hungarian palaeoecological records. *Quaternary Research* 53, 203–213.

Figure captions

Fig. 1. Map showing location of Süttő in northern Hungary (Novothny et al., 2011) (left side), and the quarries of Süttő with numbers of vertebrate localities (right side). **A1** is an archaeological site (Süttő-Diószárok), **LPS** is the loess-paleosol sequence of Novothny et al. (2011).

Fig. 2. Section of site Süttő 16 with new layer numbers.

Fig. 3. Drawing of site Süttő 17. Fossils were found from the sand (17/S) and the red clay (17/C) layers.

Fig. 4. Travertine of the Cukor quarry with several vertebrate and mollusc fossils.

Fig. 5. Correlation of the loess profile at Süttő (Novothny et al., 2011) and the profile of site Süttő 6.

Fig. 6. Charcoals of site Süttő 16. **1.** *Betula* sp. ×5; **2.** *Betula* sp. ×10; **3.** *Picea/Larix* sp. ×20; **4.** *Picea/Larix* sp. ×20. Images were taken by Zeiss-InfinityX at Leuven University.

Fig. 7. Stratigraphic positions of sites of Süttő.

Table 1. Petrographic description and interpretation of site Süttő 6 (Brunnacker et al., 1980).

Table 2. Plants, vertebrate and mollusc faunas of site Süttő 6.

Table 3. Location, type of fossil deposit and paleontology of sites from the fissures in the travertine. **Dq** – Diósvölgyi quarry; **Pq** – Páchl quarry; **Hq** – Hegyháti quarry; **N-S** – orientation of the fissure.

Table 4. Fauna and flora of the new sites (Süttő 16–20).

Table 5. Results of the paleoecological analysis of site Süttő 6.

Table 6. Percentage similarity values of comparison between small mammal faunas of Süttő 6 and other fissures of travertine (percentage similarity values).

Table 7. Percentage of paleoecological groups of various sites from fissures in the travertine.

Table 8. Percentage of paleoecological groups of Süttő 16.

layers of Süttő 6	petrography	interpretation
1	brown friable clay (with some travertine rubbles)	Bv-horizon of an interglacial brown soil (approx. 0.5 m)
2-3	humus, dark brownish gray friable clay	chernozem (humus zone, with the transition to layer 4) (approx. 1 m)
4	thin humus, light brown clay	
5	light yellowish gray loess, with lime-pseudomycelium	chalk enriched horizon (C-horizon)
6	light yellowish gray loess, with a little lime-pseudomycelium	
7	light yellowish gray loess	bedrock
8-13	pale yellowish gray loess	

	Süttő 6/0	Süttő 6/1	Süttő 6/2-3	Süttő 6/4-5	Süttő 6/6-10	Süttő 6/12	Süttő 6/13
plant macrofossils							
<i>Celtis</i> sp.		+	+				
<i>Celtis</i> cf. <i>australis</i>			+	+			
<i>Vitis silvestris</i>			+				
<i>Vitis</i> cf. <i>vinifera</i>				+			
wood charcoal indet.			+				
pollen (%)*							
<i>Juglans</i>			70		3		
Compositae Subfam. Tubuliflorae Type A	70		10		15		5
Compositae Subfam. Tubuliflorae Type B	15		5		2		
Compositae Subfam. Tubuliflorae			3		24		5
<i>Artemisia</i>	2		3				
Compositae Subfam. Liguliflorae	8		3		60		10
<i>Centranthus</i> sp.			4		30	99	85
Chenopodiaceae			1		2		1
Caryophyllaceae			3				
Cruciferae	1						
Labiatae	1						
<i>Mentha</i> type	2						
Polemoniaceae							
Rosaceae			2				
Ranunculaceae			2				
Umbelliferae	1		2				1
<i>Plantago</i> sp.							
Poaceae			1		1		3
<i>Pinus</i>	0.1		0.6			0.3	
<i>Tilia</i>			0.2		0.3		
<i>Quercus</i>			1.3				
<i>Alnus</i>					0.3		
<i>Carpinus</i>					0.3	0.3	
mammals							
<i>Talpa europaea</i>			10				
<i>Sorex araneus</i>			6	2	2		
<i>Crocidura leucodon</i> group				1			
<i>Crocidura</i> cf. <i>suaveolens</i>			1				
<i>Sicista</i> cf. <i>subtilis</i>			2	2			
<i>Spalax leucodon</i>			3				
<i>Allocricetus bursae</i>					3		
<i>Spermophilus citelloides</i>			10	3			
<i>Glis glis</i>			2	1			

	Süttő 6/0	Süttő 6/1	Süttő 6/2-3	Süttő 6/4-5	Süttő 6/6-10	Süttő 6/12	Süttő 6/13
<i>Dryomys nitedula</i>			1				
<i>Mus</i> sp.			1				
<i>Apodemus sylvaticus</i>			13	6			
<i>Arvicola terrestris</i>			2		1		
<i>Microtus (Terricola) subterraneus</i>			3				
<i>Lagurus</i> cf. <i>lagurus</i>				1			
<i>Microtus arvalis</i>			57	17	9		
<i>Microtus gregalis</i>				1	50		
<i>Myodes</i> sp.			6	5	4		
<i>Dicrostonyx torquatus</i>					1		
<i>Mustela putorius</i>					1		
<i>Dama</i> sp.			1				
molluscs							
<i>Succinea oblonga</i>				96	318		
<i>Cochlicopa lubrica</i>				1			
<i>Columella columella</i>				16	329		
<i>Granaria frumentum</i>			305	120	5		
<i>Truncatellina cylindrica</i>			4				
<i>Chondrina clienta</i>			16				
<i>Pupilla triplicata</i>			12	10	31		
<i>Pupilla sterri</i>				3	67		
<i>Pupilla muscorum</i>				15	103		
<i>Orcula doliolum</i>			1				
<i>Vallonia costata</i>			51	38	184		
<i>Vallonia tenuilabris</i>				90	121		
<i>Chondrula tridens</i>			20	6			
<i>Cochlodina laminata</i>			79		1		
<i>Iphigena plicatula</i>			64	7			
<i>Clausilia dubia</i>			1	19	9		
<i>Clausilia pumila</i>			58	7			
<i>Laciniaria plicata</i>			60	4			
<i>Clausiliidae</i> indet.			894	50			
<i>Discus ruderatus</i>			1				
<i>Discus rotundatus</i>			8	4			
<i>Punctum pygmaeum</i>			2				
<i>Vitrea crystallina</i>	7	1					
<i>Vitrea contracta</i>		2					
<i>Oxychilus inopinatus</i>	1						
<i>Aegopinella minor</i>	2	4					
<i>Euconulus fulvus</i>		20	109				
<i>Phenacolinax annularis</i>	2	2					
<i>Limax</i> cf. <i>maximus</i>	8	1					
<i>Limacidae</i> indet.	134	7	3				
<i>Bradybaena fruticum</i>	1						
<i>Helicella hungarica</i>	24	12	5				
<i>Trichia hispida</i>		15	35				

	Süttő 6/0	Süttő 6/1	Süttő 6/2-3	Süttő 6/4-5	Süttő 6/6-10	Süttő 6/12	Süttő 6/13
<i>Trichia striolata</i>		8	259				
<i>Trichia</i> sp. iuv. (<i>hispida</i> + <i>striolata</i>)			1274				
<i>Helicidae</i> indet.	37						
<i>Perforatella incarnata</i>	5						
<i>Euomphalia strigella</i>	22						
<i>Helicodonta obvilata</i>	6						
<i>Helix pomatia</i>	12	1					
<i>Cepaea vindobonensis</i>	28	7					
<i>Arianta arbustorum</i>					143		

*note that pollen percentages are approximate values read from pollen diagram

	Süttő 3	Süttő 4/1	Süttő 4/2	Süttő 5	Süttő 7/L	Süttő 7/M	Süttő 7/U	Süttő 8/1	Süttő 8/2	Süttő 8/3	Süttő 9	Süttő 12/A	Süttő 12/B	Süttő 13
<i>Mus</i> sp.											1			
<i>Mus</i> cf. <i>musculus</i>					3									
<i>Apodemus sylvaticus</i>	5	1	2		107		5			1	2	9	6	
<i>Arvicola terrestris</i>					3		1	1			2	1		
<i>Microtus (Terricola) subterraneus</i>			2		8					1		1		
<i>Microtus (Stenocranius) gregaloides</i>							1			1				
<i>Lagurus</i> cf. <i>lagurus</i>					1				1		3	2	1	
<i>Microtus arvalis</i>	8				38		5			1	32	11	2	
<i>Microtus nivalis</i>					1									1
<i>Microtus oeconomus</i>							1				8	1		
<i>Microtus gregalis</i>					1		1				1	3	2	
<i>Microtus</i> sp.		5												
<i>Myodes glareolus</i>										2		2	1	
<i>Myodes</i> sp.	3				58		1				6			
<i>Ochotona pusilla</i>					3		1					6	2	
<i>Lepus</i> aff. <i>praetimidus</i>					2		1				4	5	1	
<i>Ursus arctos</i>					5									
<i>Canis lupus</i>														1
<i>Mustela</i> cf. <i>nivalis</i>					2									
<i>Sus scrofa</i>					2									
<i>Equus</i> sp. (smaller)											2			
<i>Equus</i> sp. (larger)														1
<i>Dama</i> sp.	*				3									
<i>Capreolus capreolus</i>					5									
<i>Cervus elaphus</i>											3			1
<i>Bos</i> sive <i>Bison</i>					1						3			
cf. <i>Coelodonta antiquitatis</i>														1
molluscs														
<i>Cochlicopa lubrica</i>			3								2			
<i>Cochlicopa lubricella</i>			*								2			
<i>Succinea oblonga</i>			6	158		*		22	16		29	1	3	
<i>Cochlicopa lubrica</i>						3	2		*		4	1		
<i>Cochlicopa lubricella</i>					1	1					*			
<i>Columella edentula</i>												4	3	
<i>Columella columella</i>			2						*					
<i>Granaria frumentum</i>			10		37	103	6	*	3	*	259	40	14	
<i>Vertigo pusilla</i>										2				
<i>Vertigo alpestris</i>					2	1	5	8		*	1		1	
<i>Truncatellina cylindrica</i>					2	5						2		
<i>Truncatellina callicratis</i>												2		
<i>Chondrina clienta</i>					*	1								
<i>Pupilla triplicata</i>			5	3	9	2	1	2	2	1	12	25	12	
<i>Pupilla sterri</i>			2	2	*	1					7	11	13	
<i>Pupilla muscorum</i>				141			1	3	2	3	25	7	1	
<i>Orcula dolium</i>			1				*	1	*		1		*	
<i>Orcula doliolum</i>					3	2			*					
<i>Vallonia pulchella</i>				2							258	33	13	
<i>Vallonia costata</i>				18	65	146	9	9	4	80	245	35	25	

	Süttő 3	Süttő 4/1	Süttő 4/2	Süttő 5	Süttő 7/L	Süttő 7/M	Süttő 7/U	Süttő 8/1	Süttő 8/2	Süttő 8/3	Süttő 9	Süttő 12/A	Süttő 12/B	Süttő 13
<i>Vallonia tenuilabris</i>			3	2	3	26	70	35	18	1	12	11	11	
<i>Acanthinula aculeata</i>			3									2	1	
<i>Chondrula tridens</i>	39		5	*	*		1	*			479	22	10	
<i>Pyramidula rupestris</i>						2								
<i>Cochlodina laminata</i>					3	12	4		1	12	1	1	1	
<i>Iphigena plicatula</i>					31	286	38		6	180		1	1	
<i>Iphigena ventricosa</i>						1								
<i>Clausilia dubia</i>			1		1	3	4	28	9		22	1		
<i>Clausilia pumila</i>					1	8	25	6	3	2	2			
<i>Laciniaria plicata</i>			1		10	145	15	1	3	54	1			
<i>Neostyriaca cf. corynodes</i>			2											
<i>Clausiliidae</i> indet.			4		*	*	*	16		359	19	4	1	
<i>Discus ruderatus</i>						5	10	15	1		28			
<i>Discus rotundatus</i>					10	16	2		1	65		2	3	
<i>Punctum pygmaeum</i>			1		1						1	4		
<i>Aegopsis verticillus</i>					*									
<i>Vitrea crystallina</i>			2		2			1	1		5			
<i>Vitrea subrimata</i>					1							8	10	
<i>Vitrea contracta</i>												10	7	
<i>Oxychilus draparnaudi</i>					1	13	1					12	20	
<i>Oxychilus inopinatus</i>			1		2	11	1					11	8	
<i>Aegopinella minor</i>					3	6	4			*	1	9	7	
<i>Nesovitrea hammonis</i>					1		1	1	1		18			
<i>Zonitidae</i> indet.			1			11						66	30	
<i>Euconulus fulvus</i>			1								1	1		
<i>Vitrina pellucida</i>					3	1								
<i>Phenacolimax annularis</i>			3		4	5				*	45	7	2	
<i>Semilimax semilimax</i>							1	1			1			
<i>Limax cf. maximus</i>						25				15	24	4	6	
<i>Limacidae</i> indet.			1		22	99	18	1	11	28	41	33	9	
<i>Bradybaena fruticum</i>											4		1	
<i>Helicopsis striata</i>	24				*	13	2		2	2	390	72	23	
<i>Thricia unidentata</i>					1							2	1	
<i>Trichia hispida</i>				143		1	2	5	7					
<i>Trichia striolata</i>			5								58	1	1	
<i>Helicidae</i> indet.			2		5	1		4			10	12		
<i>Perforatella bidentata</i>											3			
<i>Monachoides incarnata</i>					6	28	*	1		*		10	1	
<i>Monachoides umbrosa</i>						1								
<i>Euomphalia strigella</i>					2	8	1	1	1	11		2		
<i>Helicodonta obvulata</i>					6	29	4		*	1		15	7	
<i>Soosia diodonta</i>												2		
<i>Helix pomatia</i>					1	2	1		*	*		2		
<i>Cepaea vindobonensis</i>	1				1	6	*			*	17	6	2	
<i>Arianta arbustorum</i>				2					3		8	1		

	Süttő 16/6	Süttő 16/5	Süttő 16/4	Süttő 16/3	Süttő 16/2	Süttő 16/1	Süttő 17/C	Süttő 17/S	Süttő 19	Süttő 20
plants										
<i>Betula</i>	3		16							
cf. <i>Alnus</i>			1							
<i>Pinus</i>			7	3						
cf. <i>Pinus</i>					1					
cf. <i>Picea</i>			2							
<i>Larix/Picea</i>			1	1						
Coniferous indet.				1	1					
vertebrates										
<i>Osteichthyes</i> indet.	1									
<i>Limax</i>		1								
<i>Bufo bufo</i>	2		2		1					
<i>Bufo</i> cf. <i>viridis</i>							2			
<i>Bufo</i> sp.		1								
<i>Rana</i> cf. <i>arvalis</i>	1									
<i>Rana</i> cf. <i>temporaria</i>					1					
<i>Rana</i> sp. 1			1		1					
<i>Rana</i> sp. 2			1							
<i>Pelobates</i> cf. <i>fuscus</i>					2					
<i>Pelobates</i> sp.							1	1		
<i>Pseudopus</i> cf. <i>panonicus</i>									*	
<i>Hierophis</i> cf. <i>viridiflavus</i>									*	
cf. <i>Zamenis longissimus</i>										1
<i>Anura</i>	3	4	1							
<i>Lacertilia</i> indet.				6	1					
<i>Lacerta</i> sp.	8		6		25					
<i>Lacerta</i> cf. <i>agilis</i>		6								
<i>Natrix</i> sp.					5					
<i>Coronella</i> sp.					2					
<i>Vipera</i> sp.		5	2		2					1
Aves indet.				6	12					
Chiroptera indet.	1			1			7	5	1	
<i>Talpa europaea</i>	9	6	7	4	9		1	2		
<i>Sorex araneus</i>	11	10	13	4	15		2			
<i>Sorex minutus</i>	1	1	1	3				1		
<i>Beremendia fissidens</i>									2	
<i>Sicista</i> sp.					9					
<i>Spalax</i> cf. <i>leucodon</i>									1	
<i>Cricetus cricetus</i>	1	1	1		2	2	1	1		
<i>Allocricetus</i> sp.								1		
<i>Sciurus vulgaris</i>							1			
<i>Spermophilus citellus</i>								1		
<i>Apodemus sylvaticus</i>		1			1					
<i>Arvicola terrestris</i>	3	3	6	4	4	2				
<i>Microtus (Stenocranius) gregaloides</i>							5	3		
<i>Lagurus</i> cf. <i>lagurus</i>			1							
<i>Microtus arvalis</i>	13	10	9	9	36					
<i>Microtus arvalinus</i>							1			


	Süttő 6/2-3	Süttő 6/4-5
mammals		
<i>Microtus arvalis</i>	48.305	43.59
<i>Sicista cf. subtilis</i>	1.6949	5.1282
<i>Spalax leucodon</i>	2.5424	
<i>Spermophilus citelloides</i>	8.4746	7.6923
steppe species (%)	61.017	56.41
<i>Crocidura leucodon</i> group		2.5641
<i>Crocidura cf. suaveolens</i>	0.8475	
<i>Glis glis</i>	1.6949	2.5641
<i>Dryomys nitedula</i>	0.8475	
<i>Mus</i> sp.	0.8475	
<i>Apodemus sylvaticus</i>	11.017	15.385
<i>Myodes</i> sp.	5.0847	12.821
<i>Dama</i> sp.	0.8475	
forest species (%)	21.186	33.333
molluscs		
<i>Chondrula tridens</i>	1.0735	1.0601
<i>Cochlicopa lubrica</i>		0.1767
<i>Granaria frumentum</i>	16.371	21.201
<i>Pupilla triplicata</i>	0.6441	1.7668
<i>Euconulus fulvus</i>		3.5336
xerophilous species (%)	18.089	27.739
<i>Vallonia costata</i>	2.7375	6.7138
<i>Clausilia dubia</i>	0.0537	3.3569
<i>Clausilia pumila</i>	3.1133	1.2367
<i>Discus ruderatus</i>	0.0537	
hygrophilous species (%)	5.9581	11.307
<i>Pupilla sterri</i>		0.53
<i>Pupilla muscorum</i>		2.6502
<i>Succinea oblonga</i>		16.961
<i>Vallonia tenuilabris</i>		15.901
<i>Trichia hispida</i>		2.6502
<i>Trichia striolata</i>		1.4134
oligotherm species (%)	0	40.106
<i>Laciniaria plicata</i>	3.2206	0.7067
<i>Aegopinella minor</i>	0.1074	0.7067
<i>Phenacolimax amularis</i>	0.1074	0.3534
<i>Cepaea vindobonensis</i>	1.503	1.2367
thermophilous species (%)	4.9383	3.0035

	Süttő 6/2-3	Süttő 6/4-5
Süttő 3	56.949	68.205
Süttő 7/L	49.67	50.38
Süttő 7/U	60.023	62.587
Süttő 9	60.26	61.71
Süttő 12/A	51.57	49
Süttő 12/B	35.36	35.58


ACCEPTED MANUSCRIPT


	Süttő 3	Süttő 7/L	Süttő 7/U	Süttő 9	Süttő 12/A	Süttő 12/B
small mammals						
<i>Sicista cf. subtilis</i>		1.961	4.545			
<i>Spalax cf. leucodon</i>		1.307		5.263	3.704	8.333
<i>Cricetus cricetus</i>				7.895		
<i>Cricetulus sp.</i>					1.852	
<i>Spermophilus citelloides</i>		1.307	9.091			
<i>Microtus arvalis</i>	47.06	12.42	22.73	42.11	20.37	8.333
steppe species (%)	47.06	16.99	36.36	55.26	25.93	16.67
<i>Crocidura leucodon</i> group		0.654		3.947	9.259	4.167
<i>Crocidura cf. suaveolens</i>	5.882					25
<i>Glis glis</i>		6.209			1.852	
<i>Dromomys nitedula</i>		0.327				
<i>Mus sp.</i>				1.316		
<i>Mus cf. musculus</i>		0.98				
<i>Apodemus sylvaticus</i>	29.41	34.97	22.73	2.632	16.67	25
<i>Myodes glareolus</i>	17.65	18.95	4.545	7.895	3.704	4.167
forest species (%)	52.94	62.09	27.27	15.79	31.48	58.33
molluscs						
<i>Cochlicopa lubrica</i>			0.873	0.197	0.202	
<i>Granaria frumentum</i>		15.42	2.62	12.75	8.081	5.645
<i>Pupilla triplicata</i>		3.75	0.437	0.591	5.051	4.839
<i>Chondrula tridens</i>	60.94		0.437	23.57	4.444	4.032
<i>Euconulus fulvus</i>				0.049	0.202	
<i>Helicopsis striata</i>	37.5		0.873	19.19	14.55	9.274
xerophilous species (%)	98.44	19.17	5.24	56.35	32.53	23.79
<i>Vallonia costata</i>		27.08	3.93	12.06	7.071	10.08
<i>Clausilia dubia</i>		0.417	1.747	1.083	0.202	
<i>Clausilia pumila</i>		0.417	10.92	0.098		
<i>Discus ruderratus</i>			4.367	1.378		
<i>Nesovitrea hammonis</i>		0.417	0.437	0.886		
<i>Arianta arbustorum</i>				0.394	0.202	
hygrophilous species (%)	0	28.33	21.4	15.9	7.475	10.08
<i>Succinea oblonga</i>				1.427	0.202	1.21
<i>Pupilla sterri</i>				0.344	2.222	5.242
<i>Pupilla muscorum</i>			0.437	1.23	1.414	0.403
<i>Vallonia tenuilabris</i>		1.25	30.57	0.591	2.222	4.435
<i>Trichia hispida</i>			0.873	2.854		
<i>Trichia striolata</i>					0.202	0.403
oligotherm species (%)	0	1.25	31.88	6.447	6.263	11.69
<i>Acanthinula aculeata</i>					0.404	0.403
<i>Laciniaria plicata</i>		4.167	6.55	0.049		
<i>Aegopsis verticillus</i>						
<i>Aegopinella minor</i>		1.25	1.747	0.049	1.818	2.823
<i>Phenacolimax annularis</i>		1.667		2.215	1.414	0.806
<i>Soosia diodonta</i>					0.404	
<i>Cepaea vindobonensis</i>	1.563	0.417		0.837	1.212	0.806
thermophilous species (%)	1.563	7.5	8.297	3.15	5.253	4.839

	Süttő 16/6	Süttő 16/5	Süttő 16/4	Süttő 16/3	Süttő 16/2
small mammals					
<i>Sicista</i> sp.					10
<i>Cricetus cricetus</i>	1.818	2.632	1.923		2.222
<i>Microtus arvalis</i>	23.64	26.32	17.31	25.71	40
steppe species (%)	25.45	28.95	19.23	25.71	52.22
<i>Apodemus sylvaticus</i>		2.632			1.111
<i>Myodes glareolus</i>	23.64	10.53	25	25.71	8.889
forest species (%)	23.64	13.16	25	25.71	10
molluscs					
<i>Cochlicopa lubrica</i>					9.091
<i>Granaria frumentum</i>	12.12				
<i>Pupilla triplicata</i>	3.03				
<i>Euconulus fulvus</i>	3.03				
<i>Helicopsis striata</i>	3.03				
xerophilous species (%)	21.21	0	0	0	9.091
<i>Vallonia costata</i>	27.27		50	10	63.64
<i>Clausilia dubia</i>	15.15				
<i>Clausilia pumila</i>	6.061		50		
<i>Nesovitrea hammonis</i>				10	
hygrophilous species (%)	48.48	0	100	20	63.64
<i>Pupilla muscorum</i>	9.091	33.33		40	
<i>Vallonia tenuilabris</i>	9.091				
<i>Trichia hispida</i>	12.12				
oligotherm species (%)	30.3	33.33	0	40	0




ACCEPTED






ACCEPTED MANUSCRIPT



ACCEPTED MANUSCRIPT



ACCEPTED

