





The edge of two worlds: A new review and synthesis on Eurasian forest-steppes

László Erdős¹  | Didem Ambarlı² | Oleg A. Anenkhonov³ | Zoltán Bátori⁴ |
 Dániel Cserhalmi⁵ | Márton Kiss^{1,6} | György Kröel-Dulay¹  | Hongyan Liu⁷ |
 Martin Magnes⁸ | Zsolt Molnár¹ | Alireza Naqinezhad⁹ | Yury A. Semenishchenkov¹⁰ |
 Csaba Tölgyesi⁴  | Péter Török¹¹ 

¹Institute of Ecology and Botany, MTA Centre for Ecological Research, Vácrátót, Hungary

²Faculty of Agriculture and Natural Sciences, Düzce University, Konuralp, Turkey

³Institute of General and Experimental Biology SB RAS, Ulan-Ude, Russia

⁴Department of Ecology, University of Szeged, Szeged, Hungary

⁵Department of Botany, University of Veterinary Medicine, Budapest, Hungary

⁶Department of Climatology and Landscape Ecology, University of Szeged, Szeged, Hungary

⁷College of Urban and Environmental Sciences, Peking University, Beijing, China

⁸Institute of Plant Sciences, University of Graz, Graz, Austria

⁹Department of Biology, Faculty of Basic Sciences, University of Mazandaran, Mazandaran, Iran

¹⁰Department of Biology, I. G. Petrovsky Bryansk State University, Bryansk, Russia

¹¹MTA-DE Lendület Functional and Restoration Ecology Research Group, Debrecen, Hungary

Correspondence

Péter Török, MTA-DE Lendület Functional and Restoration Ecology Research Group, Debrecen, Hungary.
 Email: molinia@gmail.com

Funding information

Országos Tudományos Kutatási Alapprogramok, Grant/Award Number: K 119 225 and PD 116114; Hungarian Government, Grant/Award Number: EFOP-3.6.1-16-2016-00014; National Youth Excellence Scholarship, Grant/Award Number: NTP-NFTÖ-16-0623; NKFIH, Grant/Award Number: K 124796 and K 119225; Russian Federal Budget, Grant/Award Number: AAAA-A17-117011810036-3

Co-ordinating Editor: Borja Jiménez-Alfaro

Abstract

Aims: Eurasian forest-steppes are among the most complex non-tropical terrestrial ecosystems. Despite their considerable scientific, ecological and economic importance, knowledge of forest-steppes is limited, particularly at the continental scale. Here we provide an overview of Eurasian forest-steppes across the entire zone: (a) we propose an up-to-date definition of forest-steppes, (b) give a short physiogeographic outline, (c) delineate and briefly characterize the main forest-steppe regions, (d) explore forest-steppe biodiversity and conservation status, and (e) outline forest-steppe prospects under predicted climate change.

Location: Eurasia (29°–56°N, 16°–139°E).

Results and Conclusions: Forest-steppes are natural or near-natural vegetation complexes of arboreal and herbaceous components (typically distributed in a mosaic pattern) in the temperate zone, where the co-existence of forest and grassland is enabled primarily by the semi-humid to semi-arid climate, complemented by complex interactions of biotic and abiotic factors operating at multiple scales. This new definition includes lowland forest–grassland macromosaics (e.g. in Eastern Europe), exposure-related mountain forest-steppes (e.g. in Inner Asia), fine-scale forest–grassland

Nomenclature: Catalogue of Life (www.catalogueoflife.org; accessed 15 Apr 2017)

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2018 The Authors. *Applied Vegetation Science* published by John Wiley & Sons Ltd on behalf of International Association for Vegetation Science.

mosaics (e.g. in the Carpathian Basin) and open woodlands (e.g. in the Middle East). Using criteria of flora, physiognomy, relief and climate, nine main forest-steppe regions are identified and characterized. Forest-steppes are not simple two-phase systems, as they show a high level of habitat diversity, with forest and grassland patches of varying types and sizes, connected by a network of differently oriented edges. Species diversity and functional diversity may also be exceptionally high in forest-steppes. Regarding conservation, we conclude that major knowledge gaps exist in determining priorities at the continental, regional, national and local levels, and in identifying clear target states and optimal management strategies. When combined with other threats, climate change may be particularly dangerous to forest-steppe survival, possibly resulting in compositional changes, rearrangement of the landscape mosaic or even the latitudinal or altitudinal shift of forest-steppes.

KEYWORDS

habitat complexity, landscape heterogeneity, meadow steppe, prairie, semi-arid vegetation, steppe, vegetation mosaic, wooded-steppe, woodland

1 | INTRODUCTION

Mosaic vegetation complexes consisting of woody and herbaceous patches are in the spotlight of current ecological research (e.g. Breshears, 2006; Innes, Anand, & Bauch, 2013; Prevedello, Almeida-Gomes & Lindenmayer, 2018). Forest-steppes belong to the most complex ecosystems outside the tropics in terms of composition, structure and function (Erdős et al., 2014; Walter & Breckle, 1989). While also present in North America (e.g. Leach & Givnish, 1999) and South America (e.g. Kitzberger, 2012), the largest forest-steppes are found in Eurasia.

Eurasian forest-steppes have outstanding ecological and conservation importance. They occupy large areas and appear in a wide variety of types and sub-types on various terrains (plains, hills, mountain ranges, plateaus) from the sea level up to 3,500 m a.s.l. and from sub-Mediterranean to ultracontinental to monsoon climates (Berg, 1958; Walter & Breckle, 1989; Wesche et al., 2016). Forest-steppes have a very high net primary production compared to other non-tropical systems (Pfadenhauer & Klötzli, 2014; Schultz, 2005; Zlotin, 2002), as well as a considerable biomass and C sequestration capacity (Müller, 1981; Schultz, 2005). Species diversity is also high, with many taxa of special conservation interest such as endemics, endangered species and wild relatives of cultivated plants (Bannikova, 1998; Chibilyov, 2002; Olson & Dinerstein, 1998; Zlotin, 2002). Furthermore, forest-steppes are important from an economic perspective, as they are often used as pastures and provide livelihoods for many people (e.g. Ambarlı et al., 2016; Chibilyov, 2002; Pfadenhauer & Klötzli, 2014; Smelansky & Tishkov, 2012). Unfortunately, forest-steppes are among the most threatened ecosystems due to habitat loss, fragmentation and an inadequate network of protected areas (Hoekstra, Boucher, Ricketts, & Roberts, 2005).

The scientific knowledge on Eurasian forest-steppes is relatively scattered (Bone, Johnson, Kelaidis, Kintgen, & Vickerman,

2015). Although the number of studies has increased recently, syntheses are scarce, with several limitations we outline here. First, most reviews have been conducted at national (e.g. Korotchenko & Peregrym, 2012; Molnár, Biró, Bartha & Fekete, 2012) or regional scales (e.g. Berg, 1958; Chibilyov, 2002; Dokhman, 1968; Golubev, 1965; Krashennnikov, 1954; Lavrenko, 1980; Makunina, 2016a; Milkov, 1950, 1951, 1977), while continental-scale studies are almost entirely lacking (but see Walter & Breckle, 1989). Second, most syntheses have focused on the steppe biome, discussing forest-steppes only as a marginal topic (e.g. Lavrenko, 1980; Lavrenko, Karamysheva, & Nikulina, 1991; Nosova, 1973; Rachkovskaya & Bragina, 2012; Wesche et al., 2016). Third, the few regional and continental overviews usually neglect the forest-steppes of the Middle East and the Tian Shan-Pamir ranges (e.g. Lavrenko, 1969; Wendelberger, 1989), resulting in an incomplete view of the ecosystem.

Our aim in this paper was to provide a synthetic overview of Eurasian forest-steppes, by collecting diffuse knowledge of the entire area covered by forest-steppes. First, we provide a formal definition of forest-steppes, identifying inherent difficulties in producing an exact definition and delineation. We briefly discuss spatial extents and gradients. A substantial part of our review focuses on the delineation and brief description of the main forest-steppe regions. Then we review forest-steppe diversity in terms of habitats, species and functional traits, and outline the conservation status of forest-steppes. Finally, we explore the future prospects of forest-steppes under predicted climate change.

2 | DEFINING FOREST-STEPPES

Forest-steppes are known by different names in the literature and across different regions. In most of Eastern Europe and northern

Asia, the terms “forest-steppe” or “wooded-steppe” are used, compared to “steppe forest”, “open woodland” and “sparse arid woodland” in southwestern Asia. In this paper, we treat these terms as synonyms.

The majority of researchers mention one or more of the following points as decisive characteristics in defining forest-steppes: (a) the transitional spatial position (between closed forests and treeless steppes), (b) semi-humid to semi-arid climatic features, and (c) a mosaic-like vegetation pattern. (d) Special soil characteristics as key drivers for vegetation may be considered a fourth criterion (soil is a basic part of nearly all steppe definitions; see for example: Allan, 1946; Berg, 1958; Chibilyov, 2002; Dokuchaev, 1899; Walter & Breckle, 1989). We henceforth discuss the suitability of each of the above four points for defining forest-steppes.

- Standard forest-steppe definitions usually begin with an emphasis on the transitional spatial position of forest-steppes between closed forests (nemoral forests, taiga or Mediterranean forests) and mostly treeless true steppes (e.g. Berg, 1958; Bredenkamp, Spada, & Kazmierczak, 2002; Chibilyov, 2002; Illyés, Bölöni, Kovács, & Kállay-Szerényi, 2007; Kleopov, 1990; Lavrenko, 1980; Magyari et al., 2010; Müller, 1981; Pócs, 2000; Walter & Breckle, 1989), a description which does not apply to all regions. For example, the definition is problematic in both the Carpathian Basin and the Russian Far East, due to the lack of a southern steppe border (Fekete, Molnár, Magyari, Somodi, & Varga, 2014; Ivanov, 2002). Furthermore, forest-steppes occur not only near the northern edge of the steppe zone, but also in the steppe region of the Middle East, without necessarily forming a transition towards the closed forests (Wesche et al., 2016). For instance, in some Iranian and Afghan mountain ranges, open woodlands can be found between low-elevation semi-desert-like steppes and high-mountain thorn cushion communities (Breckle, 2007; Sagheb-Talebi, Sajedi, & Pourhashemi, 2014; Zohary, 1973). In the Qilian Mts, forest-steppes have developed above the lower (arid) timber line, but the closed forest zone is lacking due to the proximity of the upper (cold) timber line (Walter & Breckle, 1989). A simplistic definition of forest-steppe as a transitional zone between treeless steppe and closed forest may therefore be inadequate, and other factors such as topography or soil grain size should be considered.
- Climate is a key defining element for forest-steppes in many scientific publications (e.g. Chibilyov, 2002; Kleopov, 1990; Schultz, 2005; Walter & Breckle, 1989). In the temperate zone, aside from edaphic variations, humid environments are able to support forests, while primary grasslands are typical under more arid conditions (Dengler, Janišová, Török, & Wellstein, 2014). Where climate is transitional (i.e. semi-humid to semi-arid, close to neutral moisture balance), a mosaic of forests and grasslands can develop, as neither of them has a decisive advantage over the other (Berg, 1958; Borhidi, 2002; Bredenkamp et al., 2002; Budyko, 1984; Chibilyov, 2002; Djamali et al., 2011; Kleopov, 1990; Lavrenko & Karamysheva, 1993; Walter & Breckle, 1989).
- An obvious feature of forest-steppes is their mosaic-like pattern. Definitions usually refer to the macromosaic feature, i.e. the spatial alternation of large forest patches and extensive grasslands (e.g. Bredenkamp et al., 2002; Chibilyov, 2002; Müller, 1981; Walter & Breckle, 1989). However, fine-scale mosaics become typical as Mediterranean climatic influences increase (Donitã, 1970; Varga et al., 2000; Wendelberger, 1989; Wesche et al., 2016; Zólyomi & Fekete, 1994). Here, individual patches may be very small. In some cases, the grassland matrix is scattered with solitary trees, which may be regarded as small forest patches (cf. Erdős, Tölgyesi, Cseh, et al., 2015). In sum, forest-steppes may appear as macro- or micromosaics, thus restricting the definition to macromosaics is not justifiable. Another recurring element of forest-steppe definitions is that the grassland component is represented by meadow steppes (e.g. Chibilyov, 2002; Kleopov, 1990; Lavrenko, 1980; Lavrenko & Karamysheva, 1993; Müller, 1981; Zlotin, 2002), i.e. rather mesic tall grasslands with numerous forbs. Several cases, most notably in regions with considerable mediterranean influences, demonstrate that the grassland component is in fact a dry grassland with short grasses, and limited number and cover of forbs. Tragacanthic species are typical, especially in the mountains of the Middle East (Akhani, 1998; Zohary, 1973).
- Generally, the steppe component of forest-steppes frequently grows on chernozem soils, and the forest component on grey forest soil (Knapp, 1979; Rychnovská, 1993; Zamotaev, 2002; Zech, Schad & Hintermaier-Erhard, 2014). Different chernozem varieties may also occur under forest patches (Berg, 1958; Wallis de Vries, Manibazar & Dügerlham, 1996). Solonetz and solonchak soils are quite usual under the steppe component (Lavrenko & Karamysheva, 1993; Müller, 1981; Sochava, 1979; Walter & Breckle, 1989), and solonetz occasionally occurs under forests (Horvat, Glavač, & Ellenberg, 1974; Molnár & Borhidi, 2003). Planosols may support *Betula* stands, while podzols can be found under *Pinus* stands (Berg, 1958; Rachkovskaya & Bragina, 2012). On chestnut soils, both forest patches and steppes can develop (Berg, 1958; Shahgedanova, Mikhailov, Larin & Bredikhin, 2002; Tamura, Asano & Jamsran, 2013; Zhu, 1993). Gley soils are typical of *Larix* forests of Inner Asian mountains with permafrost (Shahgedanova et al., 2002; Walter & Breckle, 1989). Under a strong mediterranean climatic influence, in the Middle East (e.g. Turkey, Iraq, Iran), sierozems are widespread (Kürschner & Parolly, 2012; Singh & Gupta, 1993). For more detail on soil types in the forest-steppes, see also Schultz (2005) and Zech et al. (2014).

Based on the reviewed criteria we argue that a broad yet accurate definition of forest-steppes requires both climatic (semi-humid to semi-arid) and physiognomic (a mosaic of arboreal and herbaceous components) features. In the climatic range where neither closed forest nor treeless grassland is favoured, both have a roughly equal chance to develop. Competition outcomes usually depend on local factors such as aspect, microclimate or soil (e.g. Anenkhonov et al., 2015; Borhidi, 2002; Hais, Chytrý, & Horsák, 2016; Liu et al.,

2012; Mayer, 1984; Walter & Breckle, 1989). When the distribution of forest-steppe is determined primarily by macroclimate, the forest-steppe is zonal. However, forest-steppes may also develop outside this transitional climatic range, provided that local factors modify water availability so that neither component has a competitive advantage. For example, in a region of sufficient humidity to support forests, soils with an extremely low water retention capacity or steep south-facing slopes with a warm microclimate may result in a forest-grassland mosaic. In this case, the forest-steppe is considered extrazonal.

Many additional drivers contribute to the dynamics of the forest-grassland co-existence. The interplay of climate, competition, facilitation, fire, grazing and browsing in maintaining the vegetation mosaic is as yet not fully understood for complex forest-grassland ecosystems (e.g. House, Archer, Breshears, Scholes, & Tree-Grass Interactions Participants, 2003; Sankaran, Ratnam & Hanan, 2004; Scholes & Archer, 1997; Stevens & Fox, 1991).

An exact definition and the accurate delineation of forest-steppes is complicated by inherent ambiguity. The grassland-forest continuum ranges from totally treeless grasslands to closed forests (Breshears, 2006). Based on the physiognomy, forest-steppes lie somewhere between the two extremes, but the proportion of grasslands and forest patches varies widely (Illyés et al., 2007). The middle of the continuum (i.e. 50% arboreal and 50% grassland vegetation) is clearly a forest-steppe, but the designation of lower and upper thresholds is necessarily arbitrary and often difficult (e.g. Berg, 1958; Chibilyov, 2002).

An additional question is whether a mosaic of grasslands and shrubby vegetation should be regarded as forest-steppe. If low shrubs occur only, such as *Prunus tenella*, the complex may be termed shrub-steppe and classified among steppes (Berg, 1958; Lavrenko & Sochava, 1956; Lavrenko, Karamysheva & Nikulina, 1991). In contrast, 2–6-m tall *Pistacia* spp., *Juniperus excelsa* or *Quercus pubescens* individuals or small stands in a grassland matrix are usually classified among forest-steppes.

Considering the arguments outlined above, our definition of forest-steppes is as follows: forest-steppes are natural or near-natural vegetation complexes of arboreal and herbaceous components (typically distributed in a mosaic pattern) in the temperate zone (excluding the Mediterranean), where the co-existence of forest and grassland is enabled primarily by the semi-humid to semi-arid climate, complemented by complex interactions of biotic (e.g. grazing, land use) and abiotic (e.g. soil, topography) factors operating at multiple scales. The arboreal cover (with a minimum height of 2 m) is 10%–70% across the entire landscape mosaic. The vascular vegetation cover within the grassland is at least 10% (corresponding to the grassland definition of Dixon, Faber-Langendoen, Josse, Morrison, & Loucks, 2014; and the steppe definition of Wesche et al., 2016).

Our forest-steppe definition therefore rests on physiognomic features and the underlying environmental factors, the most important of which is climate. This broad understanding of forest-steppes includes lowland forest-grassland macromosaics (e.g. in

Eastern Europe and the southern parts of West Siberia), exposure-related mountain forest-steppes (e.g. in Inner Asia), fine-scale forest-grassland mosaics (e.g. in the Carpathian Basin) and open woodlands (e.g. in the Middle East).

3 | ARE FOREST-STEPPE A BIOME?

Whether forest-steppe is a biome in its own right or only a transition between two neighbouring biomes may be considered a merely semantic question. However, it should be pointed out that forest-steppes differ considerably from both closed forests and treeless steppes in terms of numerous features, including physiognomy, habitat complexity, ecological functions and abiotic parameters, as has been shown for a number of forest-grassland mosaic ecosystems (e.g. Bannikova, 2003; Breshears, 2006; Erdős et al., 2014; Scholes & Archer, 1997; Wendelberger, 1989).

Based on the biogeographic view of Lomolino et al. (2010) and Cox, Moore, and Ladle (2016), who define biomes based on their climate and physiognomy (i.e. vegetation structure), we may conclude that forest-steppes satisfy the criteria to be considered a biome as they have a specific climate and a characteristic physiognomy. Here we have to emphasize that this concept includes latitudinal as well as altitudinal vegetation zones, which fits well with our understanding of forest-steppes. However, the recognition of forest-steppes as a biome is a subject of scientific controversy. Some of the well-known global classification systems treat forest-steppes as a mere contact area between two adjacent biomes or zones (rather than a separate biome or zone in its own right). For example, in the classification of Walter (1979), our forest-steppe definition is equivalent to those of zonoecotone VI/VII (transition between nemoral forest and steppe), zonoecotone VII/VIII (transition between taiga and steppe) and zonoecotone IV/VII (transition between the Mediterranean and steppe), complemented by some parts of the Tibetan subzonobiome (within zonobiome VII) and areas from mountain orobiomes (e.g. Crimean Mts, Caucasus, Kopet Dag, Pamir-Alai, Tian Shan). Regarding the scheme of Schultz (2005), our forest-steppe definition is included in the ecozone “dry midlatitudes” and the contact zone between the ecozones “subtropics with winter rain” and “dry tropics and subtropics”. In the system of Pfadenhauer and Klötzli (2014), our forest-steppes are included mainly in the dry nemoral subzone, but considerable parts belong to the subtropical subzone with winter rain.

4 | PHYSIOGEOGRAPHIC SETTING

Forest-steppes cover vast areas in Eurasia (2.9 million km² according to Wesche et al., 2016; although the figure may be higher, depending on the defining criteria). The altitudinal range of forest-steppes extends from sea level (e.g. Turkey-in-Europe and Crimea) up to some 3,500 m a.s.l. (Qilian Mts), including lowlands, hilly




FIGURE 1 The distribution of Eurasian forest-steppes and the main forest-steppe regions. Region A: Southeast Europe, Region B: East Europe, Region C: North Caucasus and Crimea, Region D: west Siberia and north Kazakhstan, Region E: Inner Asia, Region F: Far East, Region G: Middle East, Region H: Central Asia and southwestern Inner Asia, Region I: eastern Tibetan Plateau. The GIS map (shp file) may be found in Appendix S1. Methods and sources used for delineating forest-steppe areas are given in Appendix S2

areas and mountain ranges. Forest-steppes form two distinct latitudinal belts (Figure 1): northern (ranging from the Carpathian Basin to the Russian Far East), and southern (ranging from Central Anatolia to the Tian Shan). The northernmost reaches of the forest-steppe zone are found in Russia, north of the city of Yekaterinburg in the Ural Mts (56°N) and north of the Kuznetski Alatau Mts (56°N). The southernmost extensions are in Iran, in the Zagros Mts near Shiraz (29°N). The longitudinal extension of the forest-steppe zone is 9,000 km, stretching from the westernmost parts of the Carpathian Basin (near Vienna, Austria, 16°E) to the Amur Lowlands in Russia (139°E).

The most important latitudinal climatic gradient is along the increase in aridity to the south (Zlotin, 2002). Plant species richness usually decreases toward the steppe zone (Liu & Cui, 2009; Zlotin, 2002), although the most obvious change is the reduction of tree abundance (Schultz, 2005). In forest-steppe areas within the proximity of the closed forest zone, steppes are limited to small patches (Walter & Breckle, 1989). As aridity increases towards the south, grasslands become more extensive, while forest patches become smaller. Within the southern forest-steppe belt, forest patches are almost always very small.

Concerning longitudinal gradients, continentality generally increases towards Inner Asia (Chibilyov, 2002; Wesche et al., 2016; Zhu, 1993; Zlotin, 2002). This means that mean annual precipitation decreases (summer precipitation increases, while winter precipitation decreases), mean annual temperature simultaneously decreases, while yearly temperature range increases (summers

remain hot, but winters are long and extremely cold). These changes are accompanied by pronounced changes in cardinal vegetation characteristics (Bannikova, 1998; Berg, 1958; Chibilyov, 2002; Lavrenko, 1942, 1970a,b; Lavrenko et al., 1991; Liu, Cui, Pott, & Speier, 2000; Liu et al., 2012; Zlotin, 2002): with increasing continentality, species richness usually decreases, especially for shrubs and trees, while the root/shoot ratio increases. There are deviations from the described general patterns, depending on the scale of the study and whether it concerns forest or grassland (Lashchinskiy, Korolyuk, Makunina, Anenkhonov, & Liu, 2017; Palpurina et al., 2015) (Box 1).

5 | FOREST-STEPPES ON A COARSE SCALE: MAJOR DIVISIONS

A north-south divide bisects forest-steppes into a western and an eastern part. The transition zone is considered to be either near Lake Baikal (Berg, 1958) or near the Altai Mts and the Yenisei River (Lavrenko, 1969; Lavrenko et al., 1991). Phytogeographic ranges of forest-steppe species lend support to both of these propositions, suggesting a blurred boundary (Hilbig, Jäger, & Knapp, 2004; Nimis et al., 1994; Popov, 1963). However, given that the main floristic and vegetation changes begin in the western part of the Altai Mts, classifying the Altai-Sayan-Baikal area to the eastern forest-steppe section appears well founded. In terms of climate, plant species composition and syntaxa, a major boundary exists at the

Box 1 Eurasian forest-steppes: A fact sheet

Definition: natural or near-natural vegetation complexes of arboreal and herbaceous components (typically distributed in a mosaic pattern) in the temperate zone, where the co-existence of forest and grassland is enabled primarily by the semi-humid to semi-arid climate, complemented by complex interactions of biotic and abiotic factors operating at multiple scales. The arboreal cover (height >2 m) is 10–70% across the entire landscape mosaic, while the vascular vegetation cover within the grassland is at least 10%.

Forest-steppes as a transitional zone or a separate biome: as biome definitions usually rest on climate and physiognomy, it may be concluded that forest-steppes satisfy the criteria to be considered a biome. However, this is a subject of scientific controversy, and some well-known global vegetation classification schemes treat forest-steppes as a mere contact area between two adjacent biomes, rather than a separate biome in its own right.

Geographic extent: 29°–56°N, 16°–139°E.

Main regions: Southeast Europe, East Europe, north Caucasus and Crimea, west Siberia and north Kazakhstan, Inner Asia, Far East, Middle East, Central Asia and southwestern Inner Asia, Eastern Tibetan Plateau.

Dominant life forms: mainly phanerophytes and hemicryptophytes, but also chamaephytes and therophytes, in places many geophytes.

Dominant taxa: Anacardiaceae, Apiaceae, Asteraceae, Betulaceae, Cupressaceae, Cyperaceae, Fabaceae, Fagaceae, Lamiaceae, Pinaceae, Poaceae, Ranunculaceae, Rosaceae, Salicaceae.

northwestern foothills of the Altai Mts (Lashchinskiy et al., 2017), which also appears for edaphic grasslands in the forest-steppe biome (Ermakov, Chytrý, & Valachovič, 2006). Hilbig and Knapp (1983) and Lavrenko and Karamysheva (1993) subsequently place the border to the western foothills of the Altai Mts. Similarly, Pott (2005), in agreement with Wesche et al. (2016), regards the Altai Mts as forming the boundary between western and eastern steppes and forest-steppes.

Another major division must be made between northern and southern forest-steppes, the border running from the Sea of Marmara along the main ridge of the Caucasus and through the arid lands east of the Caspian Sea to the Tian Shan. Generally, northern forest-steppes are relatively mesic, steppes are typically closed and forest patches are often large, although exceptions do exist, especially in extrazonal situations. Southern forest-steppes are more arid, with open grasslands and usually solitary and widely spaced trees (Memariani, Zarrinpour & Akhani, 2016; Zohary, 1973). The position of forest-steppes between the neighbouring vegetation belts differs strongly between northern and southern forest-steppes. While northern forest-steppes occupy space between mesic steppes

and forests, southern forest-steppes usually appear in a transitional zone (a) between forests and semi-desert-like steppes, (b) between forests and alpine communities, or (c) between steppes and alpine/sub-alpine communities.

6 | MAIN FOREST-STEPPE REGIONS

We here provide a basic description of the main regions (Figure 1). Our delineation rests on a combination of floristic and physiognomic characteristics, as well as relief and climate features. We relied on previously published material and expert knowledge, complemented by climatic data of selected stations located within forest-steppe areas. Climate data, as well as information about the remaining forest-steppe areas and current land-use practices are given in Table 1.

6.1 | Region A – Southeast Europe

Carpathian Basin, Lower Danube Plain and Inner Thrace (northeast Austria, southeast Czech Republic, Hungary, south Slovakia, north-east Croatia, Romania, north and northeast Serbia, south Moldova, southwest Ukraine, north and southeast Bulgaria, northeast Greece, Turkey-in-Europe; Bodrozközy, 1957; Zólyomi, 1957; Niklfeld, 1964; Szodfridt, 1969; Donitá, 1970; Horvat et al., 1974; Mayer, 1984; Wallnöfer, 2003; Tzonev, Dimitrov & Roussakova, 2006; Bölöni, Molnár, Biró, & Horváth, 2008; Chytrý, 2012; Molnár et al., 2012) (Figure 2a).

Forest-steppes in this region are under considerable mediterranean climatic influences, with increasing continental effects towards the northeast. Thrace is transitional towards the Anatolian forest-steppes (Region G). Forest-steppes typically occupy plains (from sea level to 250 m a.s.l.), but some hills and mountains (often on south-facing slopes) also host similar forest-grassland mosaics. Mean annual temperature is 9.0–12.5°C (up to 13.5°C in Thrace). Summers are hot, winters are mild. Mean annual precipitation is 420–600 mm, with a maximum in early summer, a secondary maximum in autumn and a semi-arid period in between.

Both forest and grassland patches are mostly xeric. Forest patches are usually small and have an open canopy, with a high number of oak species (among others: *Quercus cerris*, *Quercus frainetto*, *Quercus petraea*, *Q. pubescens*, *Quercus robur*). Other tree species such as *Acer tataricum*, *Carpinus orientalis*, *Fraxinus ornus*, *Populus alba* and *Tilia tomentosa* are also typical. Grasslands are usually characterized by *Chrysopogon gryllus*, *Festuca rupicola*, *Festuca valesiaca*, *Festuca vaginata*, *Stipa capillata*, *S. pennata* and *S. pulcherrima*. Important herbs include *Astragalus austriacus*, *Astragalus dasyanthus*, *Astragalus onobrychis*, *Fragaria viridis*, *Salvia austriaca*, *Salvia nemorosa* and *Salvia nutans*.

6.2 | Region B – East Europe

Southern part of the East European Plain (northeast Romania, Moldova, southeast Poland, Ukraine, southwest Russia; Milkov,

TABLE 1 Basic climatic parameters (to the nearest 0.5°C and 10 mm), remaining areas (+: small, ++: medium, +++: large) and current land-use practices (–: absent or very rare, +: rare, ++ moderately widespread, +++: widespread) of forest-steppes

Region	A	B	C	D	E	F	G	H	I
Mean annual temperature (°C)	9–13.5	3–9	9.5–12	1–4.5	–6 to 5	–1 to 14	10.5–17	0–12	–3 to 7
Mean January temperature (°C)	–3 to 3	–15 to –3	–4 to 1	–20 to –14	–28 to –12	–26 to 0	–5 to 4.5	–24 to –3	–12 to –2
Mean July temperature (°C)	19–25	18–22	21–24	19–22	14–20	20–27	22–31	18–26	8–16
Mean annual precipitation (mm)	420–600	400–660	300–770	270–610	210–550	360–650	270–860	380–600	300–700
Mean summer precipitation (mm)	100–210	160–260	160–230	100–240	150–290	220–400	1–170	40–400	230–490
Proportion of summer precipitation (%)	17–36	28–43	25–40	37–44	36–72	40–70	0.5–34	10–70	47–76
Remaining forest-steppe area	+	+	+	++	+++	++	+	++	+++
Land use of near-natural forest-steppes									
Grazing	++	+++	++	++	++	+++	+++	+++	+++
Mowing	+	+	+	++	+	++	+	+	–
Burning	+	+	–	+	++	++	–	–	–
Traditional crop cultivation	–	–	–	–	–	–	++	–	–
Abandoned	+++	+	++	++	+	+	+	–	+

1950; Krashennikov, 1954; Berg, 1958; Soó, 1957; Jakucs, Fekete, & Gergely, 1959; Borhidi, 1966; Dokhman, 1968; Lavrenko, 1980; Walter & Breckle, 1989; Kleopov, 1990; Chibilyov, 2002; Molnár, Türke & Csathó, 2007; Safronova, 2010; Korotchenko & Peregrym, 2012; Kuzemko et al., 2014; Semenishchenkov, 2015) (Figure 2b).

Stretching from Podolia and the eastern foothills of the Carpathians to the southern foothills of the Ural Mts, forest-steppes of this region occupy lowlands and hilly areas between ca. 90 and 500 m a.s.l. Climate is temperate continental, with some mediterranean influence in the westernmost parts. Mean annual temperature is approximately 9°C in the west, and ca. 3°C in the east. Summers are warm, winters are moderately cold. Mean annual precipitation varies between 400–600 mm (up to 660 mm in Podolia), with a peak in June (–July) and a semi-arid period in late summer.

Large and mesic forest patches are formed mainly by broad-leaved deciduous trees (*Acer platanoides*, *Fraxinus excelsior*, *Q. robur*, *Tilia cordata*, *Ulmus glabra*), although *Populus tremula* and *Betula pendula* are also common. The grassland patches are mesic, hence the names “meadow steppe” and “steppified meadow” (the two differ regarding the role of xeric species, although the distinction is used mainly by Russian and Ukrainian authors; e.g. Kuzemko, 2009; Semenishchenkov, 2009; Averinova, 2010). Important species of the grassland patches include *F. valesiaca*, *Filipendula vulgaris*, *Fragaria viridis*, *Koeleria macrantha*, *Phlomis tuberosa*, *Poa angustifolia*, *Ranunculus polyanthemus*, *Salvia pratensis*, *S. nutans*,

Stipa capillata, *S. pennata*, *S. pulcherrima*, *S. zalesskii*, *Teucrium chamaedrys* and *Trifolium montanum*, as well as different *Tulipa* and *Iris* species.

6.3 | Region C – North Caucasus and Crimea

North Caucasus, Crimea (Southwest Russia, Crimea; Berg, 1958; Walter & Breckle, 1989; Serebryanny, 2002; Volodicheva, 2002) (Figure 2c).

Forest-steppes occupy substantial areas from sea level up to ca. 600 m a.s.l. The whole region is under marked mediterranean climatic influence. Mean annual temperature is 9.5–12.0°C. Mean annual precipitation varies from 300 to 600 (–770) mm, with the maximum in summer.

In the North Caucasus, mesic forest patches are composed of *Acer campestre*, *Carpinus betulus*, *Q. petraea*, *Q. robur* and *Tilia dasystyla*. In the Crimea, forest-steppes are more xeric and show remarkable similarities with those of the Middle East (Region G) and the Lower Danube Plain (in Region A) (Donitã, 1970). In the northwestern part of the Crimean Mts, the most characteristic tree species are *Pyrus communis*, *P. elaeagnifolia*, *Q. petraea*, *Q. pubescens*, *Q. robur* and *Ulmus procera*, while *Arbutus andrachne*, *J. excelsa*, *Pistacia atlantica* and *Q. pubescens* are typical in the southern parts of the Crimean Mts. Some of the most common and characteristic species of the grassland patches in the region are *Adonis vernalis*, *F. rupicola*, *Paeonia tenuifolia*, *Phleum phleoides*, *Stipa capillata*, *S. pennata*, *S. pontica* and *S. pulcherrima*.




FIGURE 2 Eurasian forest-steppe landscapes. (a) *Quercus robur* forest with grasslands of *Festuca rupicola*, *F. wagneri* and *Stipa capillata* in the Kiskunság sand region of the Carpathian Basin, HU (photo Á. Molnár). (b) *Betula pendula*-*Quercus robur* forest and *Stipa pennata* steppe patch with *Salvia pratensis* in the Kulikovo Polye reserve, Tula region, RU (photo Yu. A. Semenishchenkov). (c) *Quercus pubescens* forest and *Stipa pontica* steppe in the Crimean Peninsula (photo Y. P. Didukh). (d) *Betula pendula* patches and *Festuca*-*Stipa* grasslands in the Kostanay Region, north Kazakhstan (photo Z. Bátori). (e) *Betula platyphylla* individuals and groves embedded in a *Leymus chinensis*-*Filifolium sibiricum* grassland in the Ulan Buton area of Inner Mongolia, CN (photo H. Liu). (f) Forest-steppe landscape dominated by *Betula platyphylla* and *Stipa baicalensis* in the southern part of the Greater Khingan Range, China (photo H. Liu). (g) *Quercus brantii* woodland with *Bromus* spp. on calcareous and gypsiferous alluvium deposits in the Zagros Mts, Iran (photo A. Daneshi). (h) Mosaic of *Picea schrenkiana* forests and *Stipa capillata* steppes in the Tian Shan, Xinjiang Uygur Autonomous Region, CN (photo H. Liu)

6.4 | Region D – West Siberia and north Kazakhstan

West Siberia, north Kazakhstan (South Russia, North Kazakhstan; Berg, 1958; Lavrenko & Karamysheva, 1993; Rachkovskaya & Bragina, 2012; Makunina, 2016a; Korolyuk & Yamalov, 2015; Mathar et al., 2016; Bátori et al., 2017; Lashchinskiy et al., 2017; Lebedeva et al., 2017; Tölgyesi et al., 2017) (Figure 2d).

The majority of the forest-steppes of this region occur in lowlands (100–200 m asl), but some, mainly in Kazakhstan, occupy hills (ca. 300–400 m asl). The climate is continental, with mean annual temperatures of 1–4.5 °C. Summers are warm, winters are very cold. Mean annual precipitation is 270–610 mm. Most precipitation falls during the summer months.

Large mesic to semi-dry forest patches alternate with extensive, mostly mesic grasslands. The forest patches are composed of small-leaved deciduous trees (*Betula pendula*, *B. pubescens*, *Populus tremula*) and *Pinus sylvestris*. Principal steppe species include *Artemisia*

glauca, *A. pontica*, *Filipendula vulgaris*, *Festuca rupicola*, *F. valesiaca*, *Fragaria viridis*, *Gypsophila paniculata*, *Helictotrichon hookeri*, *Lathyrus pisiformis*, *L. pratensis*, *P. angustifolia*, *Phleum phleoides*, *Phlomis tuberosa*, *Pimpinella saxifraga*, *Potentilla incana*, *R. polyanthemus*, *Scorzonera ensifolia*, *Stipa capillata*, *S. pennata*, *S. tirsia*, *S. zaleskii*, *Vicia cracca*. The large amount of various halophytic communities is characteristic within the forest-steppes of this region.

6.5 | Region E – Inner Asia

Altai Mts and their northern foothills, Sayan Mts and their northern foothills, Baikal area, Transbaikial Mts, Tarbagatai Mts, Saur Mts, Khangai Mts, Khentei Mts, Inner Mongolia (south Russia, east Kazakhstan, Mongolia, north and northeast China; Hilbig & Knapp, 1983; Hou, 1983; Karamysheva & Khramtsov, 1995; Wallis de Vries et al., 1996; Korotkov & Krasnoshchekov, 1998; Korolyuk & Makunina, 2000; Liu et al., 2000; Shahgedanova et al., 2002; Bannikova, 2003;

Dulamsuren, Hauck, & Mühlenberg, 2005b; Namzalov & Baskhaeva, 2006; Rachkovskaya & Bragina, 2012; Makunina, 2010, 2013, 2014, 2016b, 2017; Hais et al., 2016) (Figure 2e).

The region includes Inner Asian mountain ranges. (The Tarbagatai-Saur range as well as the westernmost extensions of the Altai Mts are transitional towards Region H and Region D, respectively. They are consequently sometimes treated as belonging either to the northern Tian Shan, or to the west Siberian–north Kazakhstan forest-steppes.) Mountain forest-steppes, extending as high as 2,400 m a.s.l., are typically situated between the steppe and forest elevational belts. Forests are usually found on north-facing slopes (often with permafrost), whereas the steppe component occupies mainly south-facing slopes and intermountain depressions. In more arid parts, only small forest patches occur amid dry grasslands. In valleys, forest-steppes can be found as low as 200 m a.s.l. The climate here is ultracontinental. Mean annual temperature is between -6°C and $+2^{\circ}\text{C}$ (up to $+5^{\circ}\text{C}$ in southern Inner Mongolia). Summers are warm, winters are extremely cold. Mean annual temperature amplitude may exceed 50°C . Mean annual precipitation is 210–550 mm; winters are dry, most precipitation falls during summer (July–August).

Forest patches are composed of *Betula pendula*, *B. platyphylla*, *Larix sibirica*, *L. gmelinii*, *Pinus sylvestris* and *Ulmus pumila*. The most common plant species of the grasslands include *Achnatherum sibiricum*, *Agropyron cristatum*, *Artemisia frigida*, *Carex pediformis*, *Cleistogenes squarrosa*, *Cymbaria daurica*, *Filifolium sibiricum*, *Festuca valesiaca*, *F. lenensis*, *K. macrantha*, *Leymus chinensis*, *Nepeta multifida*, *Poa attenuata*, *Pulsatilla patens*, *Stellera chamaejasme*, *Stipa baicalensis* and *S. krylovii*.

6.6 | Region F – Far East

West Manchuria (=northeast China Plain), southern parts of the Greater Khingan Range, eastern parts of the Chinese Loess Plateau, Amur Lowland, southwest Sihote Alin, Khanka Lowland (northeast China, southeast Russia; Berg, 1958; Hou, 1983; Rychnovská, 1993; Zhu, 1993; Ivanov, 2002; Qian et al., 2003; Liu et al., 2015) (Figure 2f).

Forest-steppes of this region occur across a wide range of terrains and altitudinal gradients. Examples of occurrence at low altitudes include the Amur Lowland (ca. 50 m a.s.l.), and west Manchuria of China (120–150 m a.s.l.). Some forest-steppes have developed on low hills, while forest-steppes of the Chinese Loess Plateau and the Greater Khingan Range are found between 800 and 2,500 m a.s.l. Regional climate is influenced by the monsoonal circulation, particularly in the east, while continental influence positively correlates to increasing distance from the ocean. The northeast–southwest direction of the forest-steppe zone in Manchuria, the greater Khingan Range and the Chinese Loess Plateau can be explained by the diminishing effects of the monsoon; the main vegetation zones run more or less parallel with the coast. Mean annual temperature ranges between -1°C and $+14^{\circ}\text{C}$. Mean annual precipitation is 360–650 mm. Winters are cold and dry, summers are warm (western Manchuria of China, Greater Khingan Range, Chinese Loess Plateau) to cool (Amur and Khanka Lowlands, Sihote Alin). Most precipitation falls during

the summer months. Forest-steppes of the Amur Lowland are also known as “Amur prairies”, while those of the Amur and the Khanka Lowlands are sometimes referred to as “East Asian savannas”.

The most typical tree species of the forest patches is *Quercus mongolica*, although *Betula dahurica*, *B. platyphylla* and *Tilia amurensis* are also frequent. Grasslands are steppe and meadow steppe ecosystems with different subtypes occurring in dry and more mesic environments. Typical species include *Arundinella hirta*, *Bothriochloa ischaemum*, *Calamagrostis epigejos*, *Cymbaria daurica*, *Filifolium sibiricum*, *Leymus chinensis*, *Miscanthus sinensis*, *Poa pratensis*, *Stipa baicalensis*, *S. bungeana*, *S. grandis* and *S. pennata*.

6.7 | Region G – Middle East

The peripheral areas of central Anatolia, east and southeast Anatolia, south and east Caucasus, Abdulaziz Mts, Zagros Mts, Persian Plateau, Alborz Mts, Kopet Dag, Badkhyz Mts, central Afghan Mts (Turkey, South Georgia, Armenia, Azerbaijan, southwest Russia, northeast Syria, northeast Iraq, Iran, south Turkmenistan, Afghanistan; Assadi, 1988; Akhani, 1998; Ambarlı et al., 2016; Breckle, 2007; Çolak & Rotherham, 2006; Kürschner & Parolly, 2012; Memariani et al., 2016; Merzlyakova, 2002; Nakhutsrishvili, 2013; Naqinezhad, Zare-Maivan & Gholizadeh, 2015; Popov, 1994; Ravanbakhsh, HamzeH'Ee, Etemad, Marvie Mohadjer & Assadi, 2016; Ravanbakhsh & Moshki, 2016; Uğurlu, Roleček & Bergmeier, 2012; Volodicheva, 2002; Zohary, 1973) (Figure 2g).

The forest-steppes of the Middle East occur on hills and mountains from ca. 200 m (foothills of south and east Caucasus, southwest Iran) to 3,000 m a.s.l. (central Afghan Mts). Other names include “southern forest-steppes”, “arid open woodlands”, “savannoid vegetation”, “semisavanna”, “pseudosavanna”, “steppe-forests” and “light forests”. “Wild orchards”, i.e. grasslands with scattered wild fruit trees, are structurally similar to forest-steppe landscapes, but have probably developed from oak woodlands through selective cutting (Kramer, 1984; Mayer, 1984; Woldring & Cappers, 2001). Mean annual temperature ranges between 10.5 and 17.0°C . Mean annual precipitation varies between 270 and 860 mm. Summers are hot and arid, winters are cold. Forest-steppes are under considerable mediterranean climatic influences. The central Afghan Mts form a transitional zone towards the Pamir and the Tian Shan (Region H), with scattered trees in a semi-desert-like steppe matrix.

The region has been under human influence for so long that it is very difficult to infer its pre-human vegetation (Asouti & Kabukçu, 2014; Frey & Probst, 1986; Wesche et al., 2016). Trees in the forest-steppes of this region either form small groves or occur as scattered individuals. The most common tree species are *Juniperus excelsa*, *J. foetidissima*, *Pinus nigra*, *P. sylvestris*, *Pistacia atlantica*, *P. vera*, *Prunus dulcis*, *Pyrus elaeagnifolia*, *Quercus brantii*, *Q. infectoria*, *Q. ithaburensis*, *Q. macranthera*, *Q. petraea*, *Q. pubescens*, *Q. robur*. Some of the most common taxa of the grasslands are *Agropyron cristatum*, *Astragalus angustifolius*, *A. lycius*, *B. ischaemum*, *C. gryllus*, *F. valesiaca*, *K. macrantha*, *Poa bulbosa*, *Seriphidium fragrans*, *S. sieberi*, *Stipa arabica*, *S. barbata*, *S. capillata*, *S. Lessingiana* and *S. pulcherrima*.



6.8 | Region H – Central Asia and southwestern Inner Asia

Pamir Mts, Alai Mts, Tian Shan, Qilian Mts, Helan Mts (southeast Kazakhstan, Kyrgyzstan, Tajikistan, east and southeast Uzbekistan, northwest China; Berg, 1958; Bone et al., 2015; Jiang, Kang, Liu, Tian, & Lei, 2000; Merzlyakova, 2002; Pang et al., 2013; Rychnovská, 1993; Sang, 2009; Tian, 1996; Wang et al., 2001; Walter & Breckle, 1989; Wu, 1980) (Figure 2h).

Forest-steppes are present in altitudes between ca. 800–3,500 m a.s.l. Climatic influences in the region vary, resulting in significantly different precipitation and temperature records among both individual mountain chains and among slopes of differing aspects. The climate is mostly continental, with mediterranean influences in the western areas. Mean annual temperature is 0–12°C. Mean yearly precipitation varies between 380 and 600 mm, with a maximum occurring during spring (in the west) or summer (in the east).

In the western part of the region (Pamir and Alai Ranges, western and northwestern Tian Shan) forest-steppes are characterized by scattered fruit trees (*Juglans regia*, *Malus sieversii*, *Pistacia* spp. and *Punica granatum*) and *Juniperus* species, embedded in a dry steppe or even a semi-desert-like matrix (*B. ischaemum*, *Ferula tenuisecta*, *Hordeum bulbosum*, *Poa bulbosa* and *Thinopyrum intermedium*). The complex is also known as “open woodland”, “desert with scattered wooded patches”, “orchard” and “wooded field”. While this type shows clear similarities towards the open woodlands of the Middle East (Region G), in the eastern parts of the region (northern and eastern Tian Shan, Qilian Mts, Helan Mts), forest-steppes are similar to those on the mountains of Region E. Such cases are usually located at higher elevations than the fruit tree woodlands. Here, forest patches are found on north-facing slopes, and are formed primarily by *Picea schrenkiana* and *P. crassifolia*, with additional species such as *Betula pendula*, *Larix sibirica*, *Picea asperata*, *Populus tremula* and *Ulmus glaucescens*. Montane steppes occupy south-facing slopes, with the most common species being *Agropyron cristatum*, *Ajanía fruticulosa*, *Artemisia frigida*, *A. lagopus*, *Cleistogenes squarrosa*, *F. rupicola*, *K. macrantha*, *Medicago falcata*, *Oryzopsis chinensis*, *Ptilagrostis pelliottii*, *P. purpurea*, *Stipa capillata*, *S. breviflora* and *S. przewalskyi*.

6.9 | Region I – Eastern Tibetan Plateau

Eastern parts of the Tibetan Plateau (southwest China; Wu, 1980; Chang, 1981; Zhao, Wu, Yin & Yin, 2011). Forest-grassland mosaics of the eastern areas of the Tibetan Plateau may only tentatively be classified among forest-steppes owing to the ambiguity of the primary cause underlying the mosaic pattern. From the southeastern periphery to the central parts of the Plateau, forests gradually give way to meadows and steppes, with a broad transitional zone. The opening up of the forest is a result of a combination of decreasing temperature and decreasing precipitation, although temperature appears as the primary driver in most cases. The elevation is 3,200–4,000 m a.s.l. Mean annual temperature is between –3°C and +7°C. Mean annual precipitation is 300–700 mm.

Forest patches are composed of *Abies fabri*, *A. fargesii*, *A. recurvata*, *A. squamata*, *Picea asperata*, *P. brachytyla*, *P. likiangensis*, *P. purpurea* and *P. wilsonii*. The most typical grassland species are *Kobresia* species (*Kobresia capillifolia*, *K. humilis*, *K. littledalei*, *K. royleana*, *K. tibetica* and *K. vidua*). Other important species are *Argentina stenophylla*, *Carex atrofusca*, *Gentiana algida* and *Thalictrum alpinum*.

7 | BIODIVERSITY FEATURES

Forest-steppes are characterized by a high level of habitat diversity. Forests, scrubs, and grasslands have strongly different physical environmental conditions, resulting in plant communities that differ in terms of vegetation structure and floristic composition (e.g. Anenkhonov et al., 2015; Bannikova, 1985; Berg, 1958; Erdős et al., 2014; Hais et al., 2016; Hilbig & Knapp, 1983; Hilbig et al., 2004; Walter & Breckle, 1989). Moreover, forest, scrub and grassland patches have a number of different types (usually aligned along micro-topographic gradients), further increasing the habitat diversity of forest-steppes (Bátori et al., 2014; Dulamsuren, Hauck, & Mühlenberg, 2005a; Makunina, 2014, 2017; Namzalov & Baskhaeva, 2006; Namzalov et al., 2012; Tölgyesi, Erdős, Körmöczi & Bátori, 2016; Tölgyesi, Zalatnai, et al., 2016; Wallis de Vries et al., 1996). In addition, differently sized patches of the same type may also possess dissimilar environmental and vegetation characteristics. For example, small, medium and large forest patches differed considerably in tree size class distribution and seedling composition (Erdős, Tölgyesi, Cseh, et al., 2015), while the species composition of grasslands also appears to relate to size (Molnár, 1998). Forest-steppes harbour an extensive network of boundaries between different patches, which may be regarded as distinct plant communities, deviating from the communities of habitat interiors (Erdős, Gallé, Körmöczi, & Bátori, 2013; Erdős, Tölgyesi, Körmöczi, & Bátori, 2015; Molnár, 1998). Edges with different orientations may also represent slightly different habitats, showing dissimilar environmental conditions and vegetation features (Erdős et al., 2013). Consequently, forest-steppes should by no means be conceived as simple two-phase systems. Instead, they are characterized by multi-level spatial heterogeneity, where forest, scrub and grassland patches of many types and different sizes, connected by a network of differently oriented edges, form an intricate and highly complex system. An integrated view of these complex ecosystems, including all components, is a prerequisite for the efficient conservation and sustainable use of forest-steppes (cf. Luza, Carlucci, Hartz & Duarte, 2014).

Forest-steppes have been recognized as important biodiversity hotspots (Bannikova, 1998; Habel et al., 2013; Kamp et al., 2016; Makunina, 2016a; Oprea, Goia, Tănase & Sîrbu, 2010; Zlotin, 2002). Habitat diversity, together with vegetation history, is a key determinant of species diversity in forest-steppes (cf. Dengler et al., 2012; Feurdean et al., 2015; Novenko et al., 2016). The grassland component of the forest-steppe may have very high fine-scale plant species richness (Chytrý et al., 2015; Dengler et al., 2016;

Lashchinskiy et al., 2017). For example, meadow steppes that typically form the grassland component of the forest-steppes in Russia may contain on average 64 plant species per 100 m² (Korolyuk, Egorova, Smelansky, & Filippova, 2008). In the forest component, up to 114 species per 100 m² have been registered within the forest-steppe landscape of the northern Altai Mountains, suggesting that those forests are probably the most species-rich forests in non-tropical Eurasia (Chytrý et al., 2012). Diversity and composition of the shrub and herb layers are influenced by variations in canopy cover. If the canopy is relatively open, many xeric steppe species may survive under the trees (Erdős, Tölgyesi, Körmöczi, et al., 2015). Under a closed canopy, mesic conditions develop, providing suitable habitats for plants adapted to more humid conditions (Walter & Breckle, 1989). Forest edges are typically of higher biodiversity than habitat interiors, and provide habitat for several species that are rare or absent in the patch interiors of the studied forest-steppe system (e.g. *Achillea seidlilii*, *Cervaria rivini*, *Geranium sanguineum*, *Hieracium umbellatum*, *Polygonatum odoratum*, *R. polyanthemos*, *Tragopogon pratensis*; Erdős et al., 2013; Erdős, Tölgyesi, Cseh, et al., 2015; Varga, 1989; Wendelberger, 1986; Molnár, 1998).

Forest-steppes provide habitats for many rare, endemic and threatened plants, including IUCN red-listed species (e.g. *Artemisia panicii*, *Astragalus wolgensis*, *Colchicum arenarium*, *Malus sieversii*, *Pistacia vera* and *P. tenella*; Zlotin, 2002; Oprea et al., 2010; Habel et al., 2013; Kamp et al., 2016).

Because of their high structural heterogeneity, forest-steppes also host a high diversity of life-forms. Forests are dominated by phanerophytes (trees and shrubs). In their herb layer, geophytes, hemicryptophytes and/or therophytes are typical, depending on local site conditions. Shrubs are the most characteristic life form in forest edges and steppe thickets. The steppe component is characterized by hemicryptophytes (both graminoids and forbs) and usually chamaephytes. In the steppes of Europe, west Siberia and the Middle East, geophytes play an important role, while therophytes are frequent in dry areas and around disturbed sites (Berg, 1958; Breckle, 2007; Erdős et al., 2014; Kürschner & Parolly, 2012; Rachkovskaya & Bragina, 2012; Rychnovská, 1993; Schultz, 2005; Tzonev et al., 2006; Walter & Breckle, 1989; Wesche et al., 2016).

The functional diversity of forest-steppes is exceptionally high. Although usually not very tall, forests are multi-layered, with evergreen or deciduous woody species (usually both of them in the same place). The steppe component also has multiple layers, with both tall and short herbs, mosses and sometimes even lichens. Both tussock and rhizomatous graminoids are typical. The amount of N-fixing species is considerable (e.g. *Astragalus*, *Medicago* and *Vicia* species). The flowering time of forbs is variable, starting in early spring and lasting till autumn. Steppe plants have evolved numerous strategies to withstand drought, cold, grazing, fire or other natural disturbances, further enhancing the diversity of functional types (Berg, 1958; Korotchenko & Peregrym, 2012; Kürschner & Parolly, 2012; Schultz, 2005; Walter & Breckle, 1989).

8 | CONSERVATION STATUS

Forest-steppes and steppes have been transformed by human activity more than any other part of Northern Eurasia (Chibilyov, 2002), although there are regional differences concerning the level of anthropogenic impacts (Table 1). The proportion of destroyed or severely degraded forest-steppes generally decreases towards the east, where agriculture began later (Zlotin, 2002). Forest-steppes have largely persisted in many Asian landscapes east of the Ural Mts (Lavrenko & Karamysheva, 1993; Smelansky & Tishkov, 2012). However, the situation is much worse in the western parts of the forest-steppe zone.

A large proportion of the steppe patches has been converted into croplands or plantations of non-native species such as *Robinia pseudoacacia*, while many forest patches have been logged or replaced with plantations (e.g. Ambarlı et al., 2016; Berg, 1958; Molnár et al., 2012; Parnikoza & Vasiluk, 2011; Walter & Breckle, 1989). The remaining forest-steppe areas are usually edaphic ones (e.g. on rocky surfaces) and/or small fragments with varying levels of degradation (Smelansky & Tishkov, 2012). As an extreme case of fragmentation, small areas of anthropogenic habitats, such as field margins (Cizek, Hauck, & Pokluda, 2012), railway embankments (Dudáš, Eliáš, & Mártonfi, 2016), river dykes (Bátori et al., 2016), kurgans (Deák et al., 2016) and road verges (Heneberg, Bogusch, & Řezáč, 2017), may serve as the last refuges for steppe and forest-steppe species.

Forest-steppes are highly sensitive to even small changes in factors determining forest/grassland proportion and distribution (Bartha et al., 2008; Kovács-Láng et al., 2000). In many European forest-steppes, the main threats are the invasion of non-native species (e.g. *Asclepias syriaca*, *Elaeagnus angustifolia*, *Robinia pseudoacacia*) and the effects of current agricultural and forestry practices (Molnár et al., 2008; Protopopova, Shevera & Mosyakin, 2006; Smelansky & Tishkov, 2012). In Turkey, forest-steppe ecosystems are negatively affected by agricultural intensification and conversion to croplands, deep ploughing, choosing non-native trees for afforestation, over-exploitation of wild plants and animals (e.g. collecting plants for firewood, poaching and the illegal collection of bulbous plants), overgrazing and road construction (Ambarlı et al., 2016). Iranian pistachio-almond forest-steppe remnants are severely degraded due to firewood cutting (Djamali et al., 2009), while oak forest-steppes suffer from heavy overgrazing (Sagheb-Talebi et al., 2014). The *Pistacia vera* and *J. excelsa* woodlands of Afghanistan are exploited for charcoal production (Breckle, 2007). The biodiversity of the Kazakh forest-steppe is highly threatened by farming and collection of plants (Rachkovskaya & Bragina, 2012). The rapid increase in the number of grazing livestock (especially goats) and logging negatively affect the flora and fauna of the Mongolian forest-steppes (Liu, Evans, et al., 2013; Wallis de Vries et al., 1996). However, since the regime change in the eastern bloc around 1990, abandonment of former croplands has increased (Alcantara et al., 2013; Schierhorn et al., 2013; Smelansky & Tishkov, 2012), providing a unique opportunity for the spontaneous recovery or planned restoration of

the steppe component (Hölzel, Haub, Ingelfinger, Otte, & Pilipenko, 2002; Sojneková & Chytrý, 2015; Török, Vida, Deák, Lengyel & Tóthmérész, 2011).

Forest-steppes have a long history of human presence, and have historically provided humans with countless ecosystem services such as food sources (including crop progenitors), medicinal plants, grazing areas, as well as material for cooking, heating, construction or leaf fodder (Asouti & Kabukçu, 2014; Mosaddegh, Naghibi, Moazzeni, Pirani & Esmaili, 2012). Some forest-steppes in present-day Turkey, Iraq and Iran are located in the region known as “the cradle of civilization” (Asouti & Kabukçu, 2014; Poschlod, 2015; Zohary, 1973). Nomadic, semi-nomadic and sedentary herders of the forest-steppe belt continue to possess rich traditional ecological knowledge of the steppes, their forage species and the spatial and temporal patterns of forage availability (Fernández-Giménez, 2000; Molnár, 2012, 2014). They have developed complex herding systems that utilize diverse pasture types and adapt to the unpredictability of grazing conditions and extremely harsh winters. At the same time, social rules and cultural taboos sometimes protect steppes and forest patches from destruction and over-exploitation. Herders living in forest-steppes, together with ecologists and conservationists, can effectively co-produce knowledge and develop tradition-based but conservation-oriented management systems (Molnár, 2013; Molnár et al., 2016; Zhang et al., 2007).

Forest-steppe conservation requires addressing certain knowledge gaps. First of all, conservation targets at continental, regional, national and local levels must be identified. Second, more research is needed to decide where the conservation of the status quo is a realistic goal, and where inevitable changes must be accepted or even facilitated. Third, practitioners must be equipped with adequate knowledge to choose between non-intervention and active management strategies. It is not yet fully known where the re-establishment of traditional practices in forest-steppe landscapes is a useful strategy, and where land-use pressure should be reduced. While identifying optimal management is challenging, we believe that a thorough knowledge of local circumstances combined with trial-and-error may be the way to success.

9 | CLIMATE CHANGE

Although climate change is not yet considered the greatest threat to forest-steppes (Ambarlı et al., 2016; Kamp et al., 2016), these ecosystems, where both forests and grasslands are near the margin of their tolerances, may be particularly vulnerable. Although drought-adapted forest types such as those in forest-steppes may be able to withstand short (seasonal) droughts, they are threatened by long (multi-year) droughts (Allen et al., 2010).

In line with global trends, increasing temperatures have been detected in many Eurasian forest-steppe areas, including Eastern Europe (Matveev et al., 2017), Turkey (Ambarlı et al., 2016), Kazakhstan (Kamp et al., 2016), Siberia (Tchebakova, Parfenova & Soja, 2011), the Altai Mts (Lkhagvadorj, Hauck, Dulamsuren & Tsogtbaatar,

2013) and Inner Mongolia (Zhang et al., 2011). Precipitation changes, both observed and predicted, are much more variable (Ambarlı et al., 2016; Angerer, Han, Fujisaki, & Havstad, 2008; Matveev et al., 2017; Tchebakova et al., 2011).

Vegetation responses to changing climate may include (a) changing species composition within patches but sustained patchwork of grassland and forest stands, (b) altered pattern of grassland and forest patches, such as shrinkage or expansion of one patch type at the expense of the other, and (c) complete disappearance of one patch type and thus a shift of biome boundaries.

In Trans-Baikalian and northern Mongolian forest-steppes, *Pinus sylvestris* may replace *Larix sibirica* in a drier climate due to its capacity to cope with drought stress (Anenkhonov et al., 2015; Dulamsuren et al., 2009). In the Carpathian Basin, species diversity in forest-steppe grasslands may decrease with increasing aridity, while interannual variability and the number of annual species may increase (Bartha et al., 2008; Kovács-Láng et al., 2000).

The response of forest patches to future climatic changes may mimic behaviour along climatic gradients, where forest patch size and cover decrease with increasing aridity (Kovács-Láng et al., 2000; Xu et al., 2017). Xu et al. (2017) found that small forest patches had increased mortality and decreased regeneration after disturbances than larger patches. A recent field study in Inner Asia has already revealed widespread tree mortality and decreased tree growth at the most xeric sites in response to increased water deficit (Liu, Williams, et al., 2013). Permafrost melting is likely to affect vegetation, including reducing forest cover (Sharkhuu & Sharkhuu, 2012). Continued warming and drying may lead to broad-scale biome shifts. Northward movement of vegetation belts is predicted for several parts of Eurasia (e.g. Angerer et al., 2008; Ishii & Fujita, 2013; Kamp et al., 2016; Zhang et al., 2011). This would lead to an overall decline of forest-steppes in Mongolia (Angerer et al., 2008). Central parts of the Carpathian Basin may be replaced by treeless steppes in the long term (Hickler et al., 2012), with an increase in the proportion of Mediterranean species (Thuiller, Lavorel, Araújo, Sykes & Prentice, 2005).

The non-linear nature of climate change impacts renders detection difficult; systems may resist certain levels of environmental change, which may then be followed by a sudden and large-scale vegetation shift (Liu & Piao, 2013). Extreme climatic events or disturbances may be catalysts of such changes (Kröel-Dulay et al., 2015).

Changing climate may affect ecosystems not only directly, but also in combination with other factors, such as land use or biological invasion. For example, in the forest steppe of the Mongolian Altai Mts, earlier snow melt resulting from warming climate caused reduced migration of pastoral nomads, which, in turn, led to an intensified use of local forest patches (Lkhagvadorj et al., 2013). Drought and associated insect damage resulted in severe forest mortality in Anatolian forest-steppes (Allen et al., 2010).

All these examples demonstrate that forest-steppe ecosystems are already responding to changing climate. With predicted



further warming and changing precipitation regimes in the 21st century (IPCC 2014), climate change may become one of the most important threats to the biodiversity and integrity of numerous ecosystems (Sala et al., 2000), including forest- steppes. Moreover, it has been suggested that the interaction of climate change and habitat fragmentation may have disastrous consequences for biodiversity (Travis, 2003), worsening forest-steppe prospects, given the high level of habitat loss in the biome (Hoekstra et al., 2005).

ACKNOWLEDGEMENTS

The work of L. Erdős was supported by the OTKA PD 116114 grant and the National Youth Excellence Scholarship (NTP-NFTÖ-16-0623). The research of O.A. Anenkhonov was carried out using the framework of project No. AAAA-A17-117011810036-3 supported by the Russian Federal Budget. Z. Bátori was supported by the NKFIH K 124796 grant. A. Naqinezhad thanks the University of Mazandaran for the research grant to support the project. C. Tölgyesi was supported by the EU-funded Hungarian grant EFOP-3.6.1-16-2016-00014. P. Török was supported by the NKFIH K 119225 grant during manuscript preparation. Emmeline Topp with the support of Laura Sutcliffe kindly improved the English of the manuscript.

ORCID

László Erdős  <http://orcid.org/0000-0002-6750-0961>

György Kröel-Dulay  <http://orcid.org/0000-0002-0695-1232>

Csaba Tölgyesi  <http://orcid.org/0000-0002-0770-2107>

Péter Török  <http://orcid.org/0000-0002-4428-3327>

REFERENCES

- Akhani, H. (1998). Plant biodiversity of Golestan National Park, Iran. *Stapfia*, 53, 1–411.
- Alcantara, C., Kuemmerle, T., Baumann, M., Bragina, E. V., Griffiths, P., Hostert, P., ... Schierhorn, F. (2013). Mapping the extent of abandoned farmland in Central and Eastern Europe using MODIS time series satellite data. *Environmental Research Letters*, 8, 035035.
- Allan, H. H. (1946). Tussock grassland or steppe? *New Zealand Geographer*, 2, 223–234.
- Allen, C. D., Macalady, A. K., Chenchouni, H., Bachelet, D., McDowell, N., Vennetier, M., Kitzberger, T., ... Cobb, N. (2010). A global overview of drought and heat-induced tree mortality reveals emerging climate change risks for forests. *Forest Ecology and Management* 259: 660–684.
- Ambarlı, D., Zeydanlı, U. S., Balkız, Ö., Aslan, S., Karaçetin, E., Sözen, M., ... Vural, M. (2016). An overview of biodiversity and conservation status of steppes of the Anatolian Biogeographical Region. *Biodiversity and Conservation* 25: 2491–2519.
- Anenkhonov, O. A., Korolyuk, A. Y., Sandanov, D. V., Liu, H., Zverev, A. A., & Guo, D. (2015). Soil-moisture conditions indicated by field-layer plants help identify vulnerable forests in the forest-steppe of semi-arid Southern Siberia. *Ecological Indicators*, 57, 196–207.
- Angerer, J., Han, G., Fujisaki, I., & Havstad, K. (2008). Climate change and ecosystems of Asia with emphasis on Inner Mongolia and Mongolia. *Rangelands*, 30, 46–51.
- Asouti, E., & Kabukçu, C. (2014). Holocene semi-arid oak woodlands in the Irano-Anatolian region of Southwest Asia: Natural or anthropogenic? *Quaternary Science Reviews*, 90, 158–182.
- Assadi, M. (1988). *Map of the flora of Iran*. Tehran, Iran: Research Institute of Forests and Rangelands. [In Persian.]
- Averinova, E. A. (2010). Sintaksonomiya stepei Tul'skoi oblasti. *Vestnik Bryanskogo Gosudarstvennogo Universiteta*, 4, 73–81. [In Russian.]
- Bannikova, I. A. (1985). Die Gebirgswaldsteppe des Ost-Changai als Ökosystem. *Feddes Repertorium*, 96, 453–465.
- Bannikova, I. A. (1998). *Lesostep' Evrazii (Otsenka floristicheskogo raznoobraziya)*. Moscow, Russia: A.N. Severtsov Institute of Problems of Ecology and Evolution RAS [In Russian.]
- Bannikova, I. A. (2003). *Lesostep' Vnutrenney Azii: struktura i funktsiya*. Moscow, Russia: Collaborative Russian-Mongolian Integrated Biological Expedition [In Russian.]
- Bartha, S., Campetella, G., Ruprecht, E., Kun, A., Házi, J., Horváth, A., ... Molnár, Z. (2008). Will interannual variability in sand grassland communities increase with climate change? *Community Ecology*, 9, 13–21.
- Bátori, Z., Erdős, L., Kelemen, A., Deák, B., Valkó, O., Gallé, R., ... Tölgyesi, C. (2017). Diversity patterns in sandy forest-steppes – a comparative study from the western and central Palaearctic. *Biodiversity and Conservation*, 27, 1011–1030.
- Bátori, Z., Körmöczy, L., Zalatnai, M., Erdős, L., Ódor, P., Tölgyesi, C., ... Török, P. (2016). River dikes in agricultural landscapes: The importance of secondary habitats in maintaining landscape-scale diversity. *Wetlands*, 36, 251–264.
- Bátori, Z., Lengyel, A., Maróti, M., Körmöczy, L., Tölgyesi, C., Bíró, A., ... Erdős, L. (2014). Microclimate-vegetation relationships in natural habitat islands: Species preservation and conservation perspectives. *Időjárás*, 118, 257–281.
- Berg, L. S. (1958). *Die geographischen Zonen der Sowjetunion I-II*. Leipzig, Germany: Teubner.
- Bodrogekőzy, G. (1957). Die Vegetation der Weisspappel-Haine in dem Reservat "Emlékerdő" bei Szeged-Ásotthalom. *Acta Biologica Szegediensis*, 3, 127–140.
- Bölöni, J., Molnár, Z. S., Bíró, M., & Horváth, F. (2008). Distribution of the (semi-)natural habitats in Hungary II. Woodlands and shrublands. *Acta Botanica Hungarica* 50(Suppl.): 107–148.
- Bone, M., Johnson, D., Kelaidis, P., Kintgen, M., & Vickerman, L. G. (2015). *Steppes: The plants and ecology of the world's semi-arid regions*. Portland, OR: Timber Press.
- Borhidi, A. (1966). Erdőtanulmányok a Szovjetunió erdőössztyepp-övében. *Botanikai Közlemények*, 53, 185–190.
- Borhidi, A. (2002). *Gaia zöld ruhája*. Budapest, Hungary: MTA.
- Breckle, S.-W. (2007). Flora and vegetation of Afghanistan. *Basic and Applied Dryland Research*, 1, 155–194.
- Bredenkamp, G. J., Spada, F., & Kazmierczak, E. (2002). On the origin of northern and southern hemisphere grasslands. *Plant Ecology*, 163, 209–229.
- Breshears, D. D. (2006). The grassland-forest continuum: Trends in ecosystem properties for woody plant mosaics? *Frontiers in Ecology and the Environment*, 4, 96–104.
- Budyko, M. I. (1984). *Evolutsiya biosfery*. Leningrad, Russia: Gidrometeoizdat. [In Russian.]
- Chang, D. H. S. (1981). The vegetation zonation of the Tibetan Plateau. *Mountain Research and Development*, 1, 29–48.
- Chibilyov, A. (2002). Steppe and forest-steppe. In M. Shahgedanova (Ed.), *The physical geography of northern Eurasia* (pp. 248–266). Oxford, UK: Oxford University Press.
- Chytrý, M. (2012). Vegetation of the Czech Republic: Diversity, ecology, history and dynamics. *Preslia*, 84, 427–504.



- Chytrý, M., Dražil, T., Hájek, M., Kalníková, V., Preislerová, Z., Šibík, J., ... Vymazalová, M. (2015). The most species-rich plant communities in the Czech Republic and Slovakia (with new world records). *Preslia* 87: 217–278.
- Chytrý, M., Ermakov, N., Danihelka, J., Hájek, M., Hájková, P., Horsák, M. K., ... Zelený, D. (2012). High species richness in hemiboreal forests of the northern Russian Altai, southern Siberia. *Journal of Vegetation Science* 23: 605–616.
- Cizek, L., Hauck, D., & Pokluda, P. (2012). Contrasting needs of grassland dwellers: Habitat preferences of endangered steppe beetles (Coleoptera). *Journal of Insect Conservation*, 16, 281–293.
- Çolak, A. H., & Rotherham, I. D. (2006). A review of the forest vegetation of Turkey: Its status past and present and its future conservation. *Biology and Environment Proceedings of the Royal Irish Academy*, 106, 343–354.
- Cox, C. B., Moore, P. D., & Ladle, R. J. (2016). *Biogeography: An ecological and evolutionary approach*, 9th ed. Oxford, UK: Wiley.
- Deák, B., Tóthmérész, B., Valkó, O., Sudnik-Wójcikowska, B., Moysiyenko, I. I., Bragina, T. M., ... Török, P. (2016). Cultural monuments and nature conservation: A review of the role of kurgans in the conservation and restoration of steppe vegetation. *Biodiversity and Conservation*, 25, 2473–2490.
- Dengler, J., Becker, T., Ruprecht, E., Szabó, A., Becker, U., Beldean, M., ... Uğurlu, E. (2012). Festuco-Brometea of the Transylvanian Plateau (Romania) – a preliminary overview on syntaxonomy, ecology and biodiversity. *Tuexenia* 32: 319–359.
- Dengler, J., Biurrun, I., Apostolova, I., Baumann, E., Becker, T., Becker, U., ... Weiser, F. (2016). Scale-dependent plant diversity in Palaearctic grasslands: A comparative overview. *Bulletin of the Eurasian Dry Grassland Group* 31: 12–26.
- Dengler, J., Janišová, M., Török, P., & Wellstein, C. (2014). Biodiversity of Palaearctic grasslands: A synthesis. *Agriculture, Ecosystems and Environment*, 182, 1–14.
- Dixon, A. P., Faber-Langendoen, D., Josse, C., Morrison, J., & Loucks, C. J. (2014). Distribution mapping of world grassland types. *Journal of Biogeography*, 41, 2003–2019.
- Djamali, M., Akhani, H., Khoshravesh, R., Andrieu-Ponel, V., Ponel, P., & Brewer, S. (2011). Application of the Global Bioclimatic Classification to Iran: Implications for understanding the modern vegetation and biogeography. *Ecologia Mediterranea*, 37, 91–114.
- Djamali, M., de Beaulieu, J.-L., Miller, N. F., Andrieu-Ponel, V., Ponel, P., Lak, R., ... Fazeli, H. (2009). Vegetation history of the SE section of the Zagros Mountains during the last five millennia; a pollen record from the Maharlou Lake, Fars Province, Iran. *Vegetation History and Archaeobotany*, 18, 123–136.
- Dokhman, G. I. (1968). *Lesostep' evropeyskoy chasti SSSR*. Moscow, Russia: Nauka. [In Russian.].
- Dokuchaev, V. V. (1899). *K ucheniyu o zonakh prirody*. Tipografia SPb. Saint-Petersburg, Russia: Gradonachal'stva Kiadó. [In Russian.].
- Donitá, N. (1970). Submediterranean Einflüsse in der Waldflora und -vegetation der Danubischen Provinz. *Feddes Repertorium*, 81, 269–277.
- Dudáš, M., Eliáš, P. Jr, & Mártonfi, P. (2016). Occurrence of *Taraxacum serotinum* (Waldst. et Kit.) Fisch. (sect. *Dioszegia*) in Slovakia. *Thaiszia*, 26, 1–10.
- Dulamsuren, C., Hauck, M., Bader, M., Oyungerel, S., Osokhjargal, D., Nyambayar, S., & Leuschner, C. (2009). The different strategies of *Pinus sylvestris* and *Larix sibirica* to deal with summer drought in a northern Mongolian forest-steppe ecotone suggest a future superiority of pine in a warming climate. *Canadian Journal of Forest Research*, 39, 2520–2528.
- Dulamsuren, C., Hauck, M., & Mühlenberg, M. (2005a). Ground vegetation in the Mongolian taiga forest-steppe ecotone does not offer evidence for the human origin of grasslands. *Applied Vegetation Science*, 8, 149–154.
- Dulamsuren, C., Hauck, M., & Mühlenberg, M. (2005b). Vegetation at the taiga forest-steppe borderline in the western Khentey Mountains, northern Mongolia. *Annales Botanici Fennici*, 42, 411–426.
- Erdős, L., Gallé, R., Körmöcz, L., & Bátori, Z. (2013). Species composition and diversity of natural forest edges: Edge responses and local edge species. *Community Ecology*, 14, 48–58.
- Erdős, L., Tölgyesi, C., Cseh, V., Tolnay, D., Cserhalmi, D., Körmöcz, L., ... Bátori, Z. (2015). Vegetation history, recent dynamics and future prospects of a Hungarian sandy forest-steppe reserve: Forest-grassland relations, tree species composition and size-class distribution. *Community Ecology*, 16, 95–105.
- Erdős, L., Tölgyesi, C., Horzse, M., Tolnay, D., Hurton, Á., Schulcz, N., ... Bátori, Z. (2014). Habitat complexity of the Pannonian forest-steppe zone and its nature conservation implications. *Ecological Complexity*, 17, 107–118.
- Erdős, L., Tölgyesi, C., Körmöcz, L., & Bátori, Z. (2015). The importance of forest patches in supporting steppe-species: A case study from the Carpathian Basin. *Polish Journal of Ecology*, 63, 213–222.
- Ermakov, N., Chytrý, M., & Valachovič, M. (2006). Vegetation of the rock outcrops and screes in the forest-steppe and steppe belts of the Altai and Western Sayan Mts., southern Siberia. *Phytocoenologia*, 36, 509–545.
- Fekete, G., Molnár, Z., Magyari, E., Somodi, I., & Varga, Z. (2014). A new framework for understanding Pannonian vegetation patterns: Regularities, deviations and uniqueness. *Community Ecology*, 15, 12–26.
- Fernández-Giménez, M. E. (2000). The role of Mongolian nomadic pastoralists' ecological knowledge in rangeland management. *Ecological Applications*, 10, 1318–1326.
- Feurdean, A., Marinova, E., Nielsen, A. B., Liakka, J., Veres, D., Hutchinson, S. M., ... Hickler, T. (2015). Origin of the forest steppe and exceptional grassland diversity in Transylvania (Central-Eastern Europe). *Journal of Biogeography*, 42, 951–963.
- Frey, W., & Probst, W. (1986). A synopsis of the vegetation of Iran. In H. Kürschner (Ed.), *Contributions to the vegetation of Southwest Asia* (pp. 9–43). Wiesbaden, Germany: Ludwig Reichert.
- Golubev, V. N. (1965). *Ekologo-biologicheskoye osobennosti travyanistykh rasteniy i rastitel'nykh soobshchestv lesostepi*. Moscow, Russia: Nauka. [In Russian.].
- Habel, J. C., Dengler, J., Janišová, M., Török, P., Wellstein, C., & Wiezik, M. (2013). European grassland ecosystems: Threatened hotspots of biodiversity. *Biodiversity and Conservation*, 22, 2131–2138.
- Hais, M., Chytrý, M., & Horsák, M. (2016). Exposure-related forest-steppe: A diverse landscape type determined by topography and climate. *Journal of Arid Environments*, 135, 75–84.
- Heneberg, P., Bogusch, P., & Řezáč, M. (2017). Roadside verges can support spontaneous establishment of steppe-like habitats hosting diverse assemblages of bees and wasps (Hymenoptera: Aculeata) in an intensively cultivated Central European landscape. *Biodiversity and Conservation*, 26, 843–864.
- Hickler, T., Vohland, K., Feehan, J., Miller, P. A., Smith, B., Costa, L., ... Sykes, M. T. (2012). Projecting the future distribution of European potential natural vegetation zones with a generalized, tree species-based dynamic vegetation model. *Global Ecology and Biogeography* 21: 50–63.
- Hilbig, W., Jäger, E. J., & Knapp, H. D. (2004). Die Vegetation des Bogduul bei Ulaanbaatar (Mongolei) – Standortsbindung und pflanzengeographische Stellung. *Feddes Repertorium*, 115, 265–342.
- Hilbig, W., & Knapp, H. D. (1983). Vegetationsmosaik und Florenelemente an der Wald-Steppen-Grenze im Chentey-Gebirge (Mongolei). *Flora*, 174, 1–89.
- Hoekstra, J. M., Boucher, T. M., Ricketts, T. H., & Roberts, C. (2005). Confronting a biome crisis: Global disparities of habitat loss and protection. *Ecology Letters*, 8, 23–29.
- Hölzel, N., Haub, C., Ingelfinger, M. P., Otte, A., & Pilipenko, V. N. (2002). The return of the steppe large-scale restoration of degraded land in southern Russia during the post-Soviet era. *Journal for Nature Conservation*, 10, 75–85.
- Horvat, I., Glavač, V., & Ellenberg, H. (1974). *Vegetation Südosteuropas*. Stuttgart, Germany: Gustav Fischer.

- Hou, H.-Y. (1983). Vegetation of China with reference to its geographical distribution. *Annals of the Missouri Botanical Garden*, 70, 509–549.
- House, J. I., Archer, S., Breshears, D. D., Scholes, R. J., & Tree-Grass Interactions Participants, N. C. E. A. S. (2003). Conundrums in mixed woody-herbaceous plant systems. *Journal of Biogeography*, 30, 1763–1777.
- Illyés, E., Bölöni, J., Kovács, G., & Kállay-Szerényi, J. (2007). A száraz gyepek jelentősége, elterjedése, helyük a vegetációmozaikban és termőhelyi viszonyaik Magyarországon. In E. Illyés, & J. Bölöni (Eds.), *Lejtősztyepek, löszgyepek és erdőssztyeprétek Magyarországon* (pp. 12–19). Budapest, Hungary: MTA ÖBKI.
- Innes, C., Anand, M., & Bauch, C. T. (2013). The impact of human-environment interactions on the stability of forest-grassland mosaic ecosystems. *Scientific Reports*, 3, 2689.
- IPCC (2014). *Climate Change 2014: Synthesis Report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Geneva, Switzerland: IPCC.
- Ishii, R., & Fujita, N. (2013). A possible future picture of Mongolian forest-steppe vegetation under climate change and increasing livestock: Results from a new vegetation transition model at the topographic scale. In N. Yamamura, N. Fujita, & A. Maekawa (Eds.), *The Mongolian ecosystem network* (pp. 65–82). Tokyo, Japan: Springer.
- Ivanov, A. (2002). The Far East. In M. Shahgedanova (Ed.), *The physical geography of northern Eurasia* (pp. 422–447). Oxford, UK: Oxford University Press.
- Jakucs, P., Fekete, G., & Gergely, J. (1959). Angaben zur Vegetation der Moldau und der Dobrudscha. *Annales Historico-naturales Musei Nationalis Hungarici*, 51, 211–225.
- Jiang, Y., Kang, M., Liu, S., Tian, L., & Lei, M. (2000). A study on the vegetation in the east side of Helan Mountain. *Plant Ecology*, 149, 119–130.
- Kamp, J., Koshkin, M. A., Bragina, T. M., Katzner, T. E., Milner-Gulland, E. J., Schreiber, D., ... Urazaliev, R. (2016). Persistent and novel threats to the biodiversity of Kazakhstan's steppes and semi-deserts. *Biodiversity and Conservation*, 25, 2521–2541.
- Karamysheva, Z. V., & Khramtsov, V. N. (1995). The steppes of Mongolia. *Braun-Blanquetia*, 17, 1–70.
- Kitzberger, T. (2012). Ecotones as complex arenas of disturbance, climate, and human impacts: The trans-Andean forest-steppe ecotone of northern Patagonia. In R. W. Myster (Ed.), *Ecotones between forest and grassland* (pp. 59–88). New York, NY: Springer.
- Kleopov, Y. D. (1990). *Analiz flory shirokolistvennykh lesov Evropeiskoy chasti SSSR*. Kiev, Russia: Naukova Dumka. [In Russian.].
- Knapp, R. (1979). Climate and soils of grassland areas in Europe. In M. Numata (Ed.), *Ecology of grasslands and bamboos in the world* (pp. 49–56). Jena, Germany: Gustav Fisher.
- Korolyuk, A. Y., Egorova, A. V., Smelansky, I. E., & Filippova, N. V. (2008). Structure of vegetation of steppe hills in the Altai Piedmont. *Contemporary Problems of Ecology*, 1, 49–58.
- Korolyuk, A. Y., & Makunina, N. I. (2000). Lugo-vye stepi Altae-Sayanskoy gornoy oblasti. *Krylovia*, 2, 26–37. [In Russian.].
- Korolyuk, A. Y., & Yamalov, S. M. (2015). Differentiation of ecological groups of species according to their reaction to moisture in differentiation of steppes of the West Siberian Plain and Southern Urals. *Contemporary Problems of Ecology*, 8, 162–172.
- Korotchenko, I., & Peregrym, M. (2012). Ukrainian steppes in the past, at present and in the future. In M. J. A. Werger & van Staalduinen M. A., (Eds.), *Eurasian steppes. Ecological problems and livelihoods in a changing world* (pp. 173–196). Dordrecht, Netherlands: Springer.
- Korotkov, I. A., & Krasnoshchekov, Y. N. (1998). Relationships between forest and steppe in Northern Mongolia. *Russian Journal of Ecology*, 29, 230–235.
- Kovács-Láng, E., Kröel-Dulay, G., Kertész, M., Fekete, G., Bartha, S., Mika, J., ... Hahn, I. (2000). Changes in the composition of sand grasslands along a climatic gradient in Hungary and implications for climate change. *Phytocoenologia*, 30, 385–407.
- Kramer, W. (1984). *Tübinger Atlas des Vorderen Orients: Mittlerer Zagros, Iran, Vegetation*. Wiesbaden, Germany: Ludwig Reichert.
- Krashennikov, I. M. (1954). Vzaimootnosheniye lesa i stepi na yuzhnoy okraïne Ural'skoy vozvysheynosti (materialy k istorii lesostepnogo landshafta). In: Y. G. Saushkin, (ed.) *Geograficheskiye raboty* (pp. 174–213). Moscow, Russia: Geografiz. [In Russian.].
- Kröel-Dulay, G., Ransijn, J., Schmidt, I. K., Beier, C., De Angelis, P., de Dato, G., ... Penuelas, J. (2015). Increased sensitivity to climate change in disturbed ecosystems. *Nature Communications*, 6: 6682.
- Kürschner, H., & Parolly, G. (2012). The central Anatolian steppe. In M. J. A. Werger & van Staalduinen M. A. (Eds.), *Eurasian steppes. Ecological problems and livelihoods in a changing world* (pp. 149–171). Dordrecht, Netherlands: Springer.
- Kuzemko, A. A. (2009). Luchna roslinnist'. Klas Molinio-Arrhenatheretea. In Y. R. Shelyag-Sosonko (Ed.), *Shelyag-Sosonko Roslinnist' Ukraïni* (pp. 1–376) s. Kiïv, Ukraine: Fitosotsiotsentr. [In Ukrainian.].
- Kuzemko, A. A., Becker, T., Didukh, Y. P., Ardelean, I. V., Becker, U., Beldean, M., ... Dengler, J. (2014). Dry grassland vegetation of Central Podolia (Ukraine) – a preliminary overview of its syntaxonomy, ecology and biodiversity. *Tuexenia* 34: 391–430.
- Lashchinskiy, N., Korolyuk, A., Makunina, N., Anenkhonov, O., & Liu, H. (2017). Longitudinal changes in species composition of forests and grasslands across the North Asian forest steppe zone. *Folia Geobotanica*, 52, 175–197.
- Lavrenko, E. M. (1942). O provintsial'nom razdelenii Evraziatskoy stepnoy oblasti. *Botanicheskiy Zhurnal SSSR*, 27, 136–142. [In Russian.].
- Lavrenko, E. M. (1969). Über die Lage des eurasiatischen Steppengebiets in dem System der Pflanzengeographischen Gliederung des aussertropischen Eurasiens. *Vegetatio*, 19, 11–20.
- Lavrenko, E. M. (1970a). Provintsial'noe razdelenie Prichernomorsko-Kazakhstanskoy stepnoy podoblasi Evraziatskoy stepnoy oblasti. *Botanicheskiy Zhurnal*, 55, 609–625. [In Russian.].
- Lavrenko, E. M. (1970b). Provintsial'noe razdelenie Central'noaziatskoy stepnoy podoblasi Evraziatskoy stepnoy oblasti. *Botanicheskiy Zhurnal*, 55, 1734–1747. [In Russian.].
- Lavrenko, E. M. (1980). *Stepi. Rastitel'nost' evropeiskoi chasti SSSR*. Leningrad, Russia: Nauka. [In Russian.].
- Lavrenko, E. M., & Karamysheva, Z. V. (1993). Steppes of the former Soviet Union and Mongolia. In R. T. Coupland (Ed.), *Ecosystems of the world 8B. Natural grasslands. Eastern hemisphere and résumé* (pp. 3–59). Amsterdam, Netherlands: Elsevier.
- Lavrenko, E. M., Karamysheva, Z. V., & Nikulina, R. I. (1991). *Stepi Evrazii*. Leningrad, Russia: Nauka. [In Russian.].
- Lavrenko, E. M., & Sochava, V. B. (Eds.) (1956). *Rastitel'nyi pokrov SSSR*. Moscow, Russia: Akademiya Nauk SSSR [In Russian.].
- Leach, M. K., & Givnish, T. J. (1999). Gradients in the composition, structure, and diversity of remnant oak savannas in southern Wisconsin. *Ecological Monographs*, 69, 353–374.
- Lebedeva, M. V., Yamalov, S. M., & Korolyuk, A. Y. (2017). Ecological cenotic groups of species in Bashkir Trans-Ural steppes in relation to key ecological factors. *Contemporary Problems of Ecology*, 10, 455–463.
- Liu, H., & Cui, H. (2009). Patterns of plant biodiversity in the woodland-steppe ecotone in southeastern Inner Mongolia. *Contemporary Problems of Ecology*, 2, 322–329.
- Liu, H., Cui, H., Pott, R., & Speier, M. (2000). Vegetation of the woodland-steppe transition at the southeastern edge of the Inner Mongolian Plateau. *Journal of Vegetation Science*, 11, 525–532.
- Liu, Y. Y., Evans, J. P., McCabe, M. F., de Jeu, R. A. M., van Dijk, A. I. J. M., Dolman, A. J., & Saizen, I. (2013). Changing climate and overgrazing are decimating Mongolian steppes. *PLoS ONE*, 8, e57599.
- Liu, H., He, S., Anenkhonov, O. A., Hu, G., Sandanov, D. V., & Badmaeva, N. K. (2012). Topography-controlled soil water content and the co-existence of forest and steppe in northern China. *Physical Geography*, 33, 561–573.

- Liu, H., & Piao, S. (2013). Drought threatened semi-arid ecosystems in the Inner Asia. *Agricultural and Forest Meteorology*, 178, 1–2.
- Liu, H., Williams, A. P., Allen, C. D., Guo, D., Wu, X., Anenkhonov, O. A., ... Badmaeva, N. K. (2013). Rapid warming accelerates tree growth decline in semi-arid forests of Inner Asia. *Global Change Biology*, 19, 2500–2510.
- Liu, H., Yin, Y., Wang, Q., & He, S. (2015). Climatic effects on plant species distribution within the forest-steppe ecotone in northern China. *Applied Vegetation Science*, 18, 43–49.
- Lkhagvadorj, D., Hauck, M., Dulamsuren, C., & Tsogtbaatar, J. (2013). Pastoral nomadism in the forest-steppe of the Mongolian Altai under a changing economy and a warming climate. *Journal of Arid Environments*, 88, 82–89.
- Lomolino, M. V., Riddle, B. R., Brown, J. H., & Whittaker, R. J. (2010). *Biogeography*, 4th ed. Sunderland, MA: Sinauer Associates.
- Luza, A. L., Carlucci, M. B., Hartz, S. M., & Duarte, L. D. S. (2014). Moving from forest vs. grassland perspectives to an integrated view towards the conservation of forest-grassland mosaics. *Natureza and Conservacao*, 12, 166–169.
- Magyari, E. K., Chapman, J. C., Passmore, D. G., Allen, J. R. M., Huntley, J. P., & Huntley, B. (2010). Holocene persistence of wooded steppe in the Great Hungarian Plain. *Journal of Biogeography*, 37, 915–935.
- Makunina, N. I. (2010). Struktura rastitel'nosti stepnogo i lesostepnogo poyasov mezhgornyykh kotlovin Khakasii i Tuvy. *Rastitel'nyi mir Aziatskoy Rossii*, 2, 50–57. [In Russian.]
- Makunina, N. I. (2013). Rastitel'nost' stepnogo i lesostepnogo poyasov Central'nogo Altaya. *Rastitel'nost' Rossii*, 23, 9–35. [In Russian.]
- Makunina, N. I. (2014). Gornaya lesostep' Yugo-Vostochnogo Altaya i Yugo-Zapadnoy Tuvy. *Rastitel'nost' Rossii*, 24, 86–100. [In Russian.]
- Makunina, N. I. (2016a). *Rastitel'nost' lesostepi Zapadno-Sibirskoy ravniny i Altae-Sayanskoy gornoj oblasti*. Novosibirsk, Russia: Academic Publishing House "Geo" [In Russian.]
- Makunina, N. I. (2016b). Botanical and geographical characteristics of forest steppe of the Altai-Sayan mountain region. *Contemporary Problems of Ecology*, 9, 342–348.
- Makunina, N. I. (2017). Biodiversity of basic vegetation communities in forest steppes of the Altai-Sayan mountain region. *International Journal of Environmental Studies*, 74, 674–684.
- Mathar, W. P., Kämpf, I., Kleinebecker, T., Kuzmin, I., Tolstikov, A., Tupitsin, S., & Hölzel, N. (2016). Floristic diversity of meadow steppes in the Western Siberian Plain: Effects of abiotic site conditions, management and landscape structure. *Biodiversity and Conservation*, 25, 2361–2379.
- Matveev, S. M., Chendev, Y. G., Lupo, A. R., Hubbart, J. A., & Timashchuk, D. A. (2017). Climatic changes in the East-European forest-steppe and effects on Scots pine productivity. *Pure and Applied Geophysics*, 174, 427–443.
- Mayer, H. (1984). *Wälder Europas*. Stuttgart, Germany: Gustav Fischer.
- Memariani, F., Zarrinpour, V., & Akhiani, H. (2016). A review of plant diversity, vegetation, and phytogeography of the Khorassan-Kopet Dagh floristic province in the Irano-Turanian region (northeastern Iran-southern Turkmenistan). *Phytotaxa*, 249, 8–30.
- Merzlyakova, I. (2002). The mountains of Central Asia and Kazakhstan. In M. Shahgedanova (Ed.), *The physical geography of northern Eurasia* (pp. 377–402). Oxford, UK: Oxford University Press.
- Milkov, F. N. (1950). *Lesostep' Russkoy ravniny: Opyt landshaftnoy kharakteristiki*. Moscow, Russia: Izdatel'stvo AN SSSR. [In Russian.]
- Milkov, F. N. (1951). Lesostepnoy landshaft i ego zonal'noe podrazdelenie. *Izvestiya Akademii Nauk SSSR, Seriya Geograficheskaya*, 5, 3–14. [In Russian.]
- Milkov, F. N. (1977). *Prirodnye zony SSSR*. Moscow, Russia: Mysl'. [In Russian.]
- Molnár, Z. (1998). Interpreting present vegetation features by landscape historical data: An example from a woodland-grassland mosaic landscape (Nagykörös Wood, Kiskunság, Hungary). In K. J. Kirby, & C. Watkins (Eds.), *The ecological history of European forests* (pp. 241–263). Wallingford, UK: CAB International.
- Molnár, Z. (2012). Classification of pasture habitats by Hungarian herders in a steppe landscape (Hungary). *Journal of Ethnobiology and Ethnomedicine*, 8, 28.
- Molnár, Z. (2013). Traditional vegetation knowledge of the Hortobágy salt steppe (Hungary): A neglected source of information for vegetation science and conservation. *Phytocoenologia*, 43, 193–205.
- Molnár, Z. (2014). Perception and management of spatio-temporal pasture heterogeneity by Hungarian herders. *Rangeland Ecology and Management*, 67, 107–118.
- Molnár, Zs., Biró, M., Barthá, S., & Fekete, G. (2012). Past trends, present state and future prospects of Hungarian forest-steppes. In M. J. A. Werger & vanStaalduinen M. A., (Eds.), *Eurasian steppes. Ecological problems and livelihoods in a changing world* (pp. 209–252). Dordrecht, Netherlands: Springer.
- Molnár, Z., Bölöni, J., & Horváth, F. (2008). Threatening factors encountered: Actual endangerment of the Hungarian (semi-)natural habitats. *Acta Botanica Hungarica*, 50(Suppl. 1), 199–217.
- Molnár, Z., & Borhidi, A. (2003). Hungarian alkali vegetation: Origins, landscape history, syntaxonomy, conservation. *Phytocoenologia*, 33, 377–408.
- Molnár, Z., Kis, J., Vadász, C., Papp, L., Sándor, I., Béres, S., ... Varga, A. (2016). Common and conflicting objectives and practices of herders and nature conservation managers: The need for the 'conservation herder'. *Ecosystem Health and Sustainability*, 2, e01215.
- Molnár, C., Türke, I. J., & Csathó, A. I. (2007). Botanikai megfigyelések Dél-Bukovina térségében. *Kanitzia*, 15, 19–34.
- Mosaddegh, M., Naghibi, F., Moazzeni, H., Pirani, A., & Esmaeili, S. (2012). Ethnobotanical survey of herbal remedies traditionally used in Kohgiluyeh va Boyer Ahmad province of Iran. *Journal of Ethnopharmacology*, 141, 80–95.
- Müller, P. (1981). *Arealssysteme und Biogeographie*. Stuttgart, Germany: Ulmer.
- Nakhutsrishvili, G. (2013). *The vegetation of Georgia (South Caucasus)*. Berlin, Germany: Springer.
- Namzalov, B. B., & Baskhaeva, T. G. (2006). *Gornaya lesostep' Barguzinskoy kotloviny (Severnoye Pribaikal'ye)*. Ulan-Ude, Russia: Buryat State University Press. [In Russian.]
- Namzalov, B. B., Cholboeva, S. A., Korolyuk, A. J., Baschaeva, T. G., Tszirenova, M. G., & Mongush, A. M. (2012). Osobennosti struktury lesostepi v ekotonnoy zone Yuzhnoy Sibiri i Tsentral'noy Azii. *Aridnye Ecosystemy*, 18, 17–27. [In Russian.]
- Naqinezhad, A., Zare-Maivan, H., & Gholizadeh, H. (2015). A floristic survey of the Hyrcanian forests in Northern Iran, using two lowland mountain transects. *Journal of Forestry Research*, 26, 187–199.
- Niklfeld, H. (1964). Zur xerothermen Vegetation im Osten Niederösterreichs. *Verhandlungen der Zoologisch-Botanischen Gesellschaft in Österreich*, 103(104), 152–181.
- Nimis, P. L., Malyshev, L. I., & Bolognini, G. (1994). A phytogeographic analysis of birch woodlands in the southern part of West Siberia. *Vegetatio*, 113, 25–39.
- Nosova, L. M. (1973). *Floro-geograficheskii analiz severnoi stepi evropeiskoi chasti SSSR*. Moscow, Russia: Nauka. [In Russian.]
- Novenko, E. Y., Tsyganov, A. N., Rudenko, O. V., Volkova, E. V., Zuyganova, I. S., Babeshko, K. V., ... Mazei, Y. A. (2016). Mid- and late-Holocene vegetation history, climate and human impact in the forest-steppe ecotone of European Russia: New data and a regional synthesis. *Biodiversity and Conservation*, 25, 2453–2472.
- Olson, D. M., & Dinerstein, E. (1998). The Global 200: A representation approach to conserving the Earth's most biologically valuable ecoregions. *Conservation Biology*, 12, 502–515.
- Oprea, A., Goia, I., Tănase, C., & Sirbu, C. (2010). Assessment of species composition: Endemics, relicts and red-listed plants (Tracheophytae,

- Bryophytae, and Fungi) in forest natural habitats of Romania. *Contribuții Botanice*, 45, 13–24.
- Palpurina, S., Chytrý, M., Tzonev, R., Danihelka, J., Axmanová, I., Merunková, K., ... Karakiev, T. (2015). Patterns of fine-scale plant species richness in dry grasslands across the eastern Balkan Peninsula. *Acta Oecologica*, 63, 36–46.
- Pang, Y., Zhang, B., Zhao, F., Yao, Y., Zhang, S., & Qi, W. (2013). Omnidirectional distribution patterns of montane coniferous forest in the Helan Mountains of China. *Journal of Mountain Science*, 10, 724–733.
- Parnikoza, I., & Vasiluk, A. (2011). Ukrainian steppes: Current state and perspectives for protection. *Annales Universitatis Mariae Curie-Skłodowska*, 66, 23–37.
- Pfadenhauer, J. S., & Klötzli, F. A. (2014). *Vegetation der Erde: Grundlagen, Ökologie, Verbreitung*. Berlin, Germany: Springer.
- Pócs, T. (2000). *Növényföldrajz. In T. Hortobágyi, & T. Simon (Eds.), Növényföldrajz, társulástan és ökológia* (pp. 25–166). Budapest, Hungary: Nemzeti Tankönyvkiadó.
- Popov, M. G. (1963). *Osnovy florogenetiki*. Moscow, Russia: Nauka. [In Russian].
- Popov, K. P. (1994). Trees, shrubs, and semishrubs in the mountains of Turkmenistan. In K. Fet, & K. I. Atamuradov (Eds.), *Biogeography and ecology of Turkmenistan* (pp. 173–186). Dordrecht, Netherlands: Kluwer Academic.
- Poschlod, P. (2015). *Geschichte der Kulturlandschaft*. Stuttgart, Germany: Eugen Ulmer.
- Pott, R. (2005). *Allgemeine geobotanik: Biogeosysteme und biodiversität*. Berlin, Germany: Springer.
- Prevedello, J. A., Almeida-Gomes, M., & Lindenmayer, D. B. (2018). The importance of scattered trees for biodiversity conservation: A global meta-analysis. *Journal of Applied Ecology*, 55, 205–214.
- Protopopova, V. V., Shevera, M. V., & Mosyakin, S. L. (2006). Deliberate and unintentional introduction of invasive weeds: A case study of the alien flora of Ukraine. *Euphytica*, 148, 17–33.
- Qian, H., Yuan, X.-Y., & Chou, Y.-L. (2003). Forest vegetation of Northeast China. In J. Kolbek, M. Šrutek, & E. O. Box (Eds.), *Forest vegetation of Northeast Asia* (pp. 181–230). Dordrecht, Netherlands: Kluwer Academic.
- Rachkovskaya, E. I., & Bragina, T. M. (2012). Steppes of Kazakhstan: Diversity and present state. In M. J. A. Werger & van Staalduinen M. A. (Eds.), *Eurasian steppes. Ecological problems and livelihoods in a changing world* (pp. 103–148). Dordrecht, Netherlands: Springer.
- Ravanbakhsh, H., Hamzeh'Ee, B., Etemad, V., Marvie Mohadjer, M. R., & Assadi, M. (2016). Phytosociology of *Juniperus excelsa* M. Bieb. forests in Alborz mountain range in the north of Iran. *Plant Biosystems*, 150, 987–1000.
- Ravanbakhsh, H., & Moshki, A. (2016). The influence of environmental variables on distribution patterns of Irano-Turanian forests in Alborz Mountains. *Iran. Journal of Mountain Science*, 13, 1375–1386.
- Rychnovská, M. (1993). Temperate semi-natural grasslands of Eurasia. In R. T. Coupland, (Ed.), *Ecosystems of the world 8B. Natural grasslands. Eastern hemisphere and résumé*, (pp. 125–166). Amsterdam, Netherlands: Elsevier.
- Safronova, I. N. (2010). O podzonal'noi strukture rastitel'nogo pokrova stepnoi zony v evropeiskoi chasti Rossii. *Botanicheskii Zhurnal*, 95, 1126–1134. [In Russian].
- Sagheb-Talebi, K., Sajedi, T., & Pourhashemi, M. (2014). *Forests of Iran: A treasure from the past, a hope for the future*. Berlin, Germany: Springer.
- Sala, O. E., Chapin, F. S. III, Armesto, J. J., Berlow, R., Bloomfield, J., Dirzo, R., ... Wall, D. H. (2000). Global biodiversity scenarios for the year 2100. *Science*, 287, 1770–1774.
- Sang, W. (2009). Plant diversity patterns and their relationships with soil and climatic factors along an altitudinal gradient in the middle Tianshan Mountain area, Xinjiang, China. *Ecological Research*, 24, 303–314.
- Sankaran, M., Ratnam, J., & Hanan, N. P. (2004). Tree-grass coexistence in savannas revisited: Insights from an examination of assumptions and mechanisms invoked in existing models. *Ecology Letters*, 7, 480–490.
- Schierhorn, F., Müller, D., Beringer, T., Prishchepov, A. V., Kuemmerle, T., & Balmann, A. (2013). Post-Soviet cropland abandonment and carbon sequestration in European Russia, Ukraine, and Belarus. *Global Biogeochemical Cycles*, 27, 1175–1185.
- Scholes, R. J., & Archer, S. R. (1997). Tree-grass interactions in savannas. *Annual Review of Ecology and Systematics*, 28, 517–544.
- Schultz, J. (2005). *The ecozones of the world*. Berlin, Germany: Springer.
- Semenishchenkov, Y. A. (2009). *Fitotsenoticheskoe raznoobrazie Sudost'-Desnyanskogo mezhdurech'ya*. Bryansk, Russia: Bryanskii Gosudarstvennyi Universitet. [In Russian].
- Semenishchenkov, Y. A. (2015). Botaniko-geograficheskoe raionirovanie basseina Verkhnego Dnepra na osnove sintaksonomii lesnoi rastitel'nosti. *Botanicheskii Zhurnal*, 100, 625–657. [In Russian].
- Serebryanny, L. (2002). Mixed and deciduous forests. In M. Shahgedanova (Ed.), *The physical geography of northern Eurasia* (pp. 234–247). Oxford, UK: Oxford University Press.
- Shahgedanova, M., Mikhailov, N., Larin, S., & Bredikhin, A. (2002). The mountains of southern Siberia. In M. Shahgedanova (Ed.), *The physical geography of northern Eurasia* (pp. 314–349). Oxford, UK: Oxford University Press.
- Sharkhuu, N., & Sharkhuu, A. (2012). Effects of climate warming and vegetation cover on permafrost of Mongolia. In M. J. A. Werger & van Staalduinen M. A., (Eds.), *Eurasian steppes. Ecological problems and livelihoods in a changing world* (pp. 445–472). Dordrecht, Netherlands: Springer.
- Singh, J. S., & Gupta, S. R. (1993). Grasslands of southern Asia. In R. T. Coupland, (Ed.), *Ecosystems of the world 8B. Natural grasslands. Eastern hemisphere and résumé* (pp. 83–123). Amsterdam, Netherlands: Elsevier.
- Smelansky, I. E., & Tishkov, A. A. (2012). The steppe biome in Russia: Ecosystem services, conservation status, and actual challenges. In M. J. A. Werger & van Staalduinen M. A. (Eds.), *Eurasian steppes. Ecological problems and livelihoods in a changing world* (pp. 45–101). Dordrecht, Netherlands: Springer.
- Sochava, V. B. (1979). Climate and soils of the grassland distribution area in the USSR. In M. Numata (Ed.), *Ecology of grasslands and bamboos in the world* (pp. 43–48). Jena, Germany: Gustav Fisher.
- Sojneková, M., & Chytrý, M. (2015). From arable land to species-rich semi-natural grasslands: Succession in abandoned fields in a dry region of central Europe. *Ecological Engineering*, 77, 373–381.
- Soó, R. (1957). Összehasonlító vegetációtanulmányok a Szovjetunió erdős-sztyepp övéből. *MTA Biológiai Csoportjának Közleményei*, 1, 209–222.
- Stevens, G. C., & Fox, J. F. (1991). The causes of treeline. *Annual Review of Ecology and Systematics*, 22, 177–191.
- Szodfridt, I. (1969). Borókás-nyárasok Bugac környékén. *Botanikai Közlemények*, 56, 159–165.
- Tamura, K., Asano, M., & Jamsran, U. (2013). Soil diversity in Mongolia. In N. Yamamura, N. Fujita, & A. Maekawa (Eds.), *The Mongolian ecosystem network* (pp. 99–103). Tokyo, Japan: Springer.
- Tchebakova, N. M., Parfenova, E. I., & Soja, A. J. (2011). Climate change and climate-induced hot spots in forest shifts in central Siberia from observed data. *Regional Environmental Change*, 11, 817–827.
- Thuiller, W., Lavorel, S., Araújo, M. B., Sykes, M. T., & Prentice, I. C. (2005). Climate change threats to plant diversity in Europe. *Proceedings of the National Academy of Sciences of the United States of America*, 102, 8245–8250.
- Tian, L. X. (1996). *Vegetation in east slope of Helan Mts*. Huhhot, China: Inner Mongolia University Press. [In Chinese].
- Tölgyesi, C., Bragina, T. M., Valkó, O., Deák, B., Kelemen, A., Gallé, R., & Batori, Z. (2017). Micro-environment-vegetation interactions in the sandy forest-steppe of Naurzum Nature Reserve, Kazakhstan. In E. A. Abil, & T. M. Bragina (Eds.), *Biological diversity of Asian steppe* (pp. 190–194). Kostanay, KZ: Kostanay State Pedagogical Institute.



- Tölgyesi, C., Erdős, L., Körmöczi, L., & Bátori, Z. (2016). Hydrologic fluctuations trigger structural changes in wetland-dry grassland ecotones but have no effect on ecotone position. *Community Ecology*, 17, 188–197.
- Tölgyesi, C., Zalatnai, M., Erdős, L., Bátori, Z., Hupp, N., & Körmöczi, L. (2016). Unexpected ecotone dynamics of a sand dune vegetation complex following water table decline. *Journal of Plant Ecology*, 9, 40–50.
- Török, P., Vida, E., Deák, B., Lengyel, S., & Tóthmérész, B. (2011). Grassland restoration on former croplands in Europe: An assessment of applicability of techniques and costs. *Biodiversity and Conservation*, 20, 2311–2332.
- Travis, J. M. J. (2003). Climate change and habitat destruction: A deadly anthropogenic cocktail. *Proceedings of the Royal Society of London, Series B*, 270, 467–473.
- Tzonev, R., Dimitrov, M., & Roussakova, V. (2006). The Western-Pontic steppe vegetation in Bulgaria. *Haquetia*, 5, 5–23.
- Uğurlu, E., Roleček, J., & Bergmeier, E. (2012). Oak woodland vegetation of Turkey – a first overview based on multivariate statistics. *Applied Vegetation Science*, 15, 590–608.
- Varga, Z. (1989). Die Waldteppen des pannonischen Raumes aus biogeographischer Sicht. *Düsseldorfer Geobotanisches Kolloquium*, 6, 35–50.
- Varga, Z., Borhidi, A., Fekete, G., Debreczy, Zs., Bartha, D., Bölöni, J., ... Király, G. (2000). Az erdőssztyepp fogalma, típusai és jellemzésük. In Zs. Molnár & A. Kun (Eds.), *Alföldi erdőssztyepp-maradványok Magyarországon* (pp. 7–19). Budapest-Vácrtót, Hungary: WWF-MTA ÖBKI
- Volodicheva, N. (2002). The Caucasus. In M. Shahgedanova (Ed.), *The physical geography of northern Eurasia* (pp. 350–376). Oxford, UK: Oxford University Press.
- Wallis de Vries, M. F., Manibazar, N., & Dügerlham, S. (1996). The Vegetation of the Forest-Steppe Region of Hustain Nuruu, Mongolia. *Vegetatio*, 122, 111–127.
- Wallnöfer, S. (2003). Thermophile Eichenwaldgesellschaften im Osten Österreichs. *Verhandlungen der Zoologisch-Botanischen Gesellschaft in Österreich*, 140, 1–16.
- Walter, H. (1979). *Vegetation of the earth*, 2nd ed. Berlin, Germany: Springer.
- Walter, H., & Breckle, S.-W. (1989). *Ecological systems of the geobiosphere 3 – Temperate and polar zonobiomes of northern Eurasia*. Berlin, Germany: Springer.
- Wang, J., Chang, X., Ge, S., Miao, Y., Chang, Z., & Zhang, H. (2001). Vertical distribution of the vegetation and water and heat conditions of Qilian Mountain (northern slope). *Journal of Northwest Forestry University*, 16, 1–3. [In Chinese.].
- Wendelberger, G. (1986). Saum- und Mantelgesellschaften des pannonischen Raumes. *Verhandlungen der Zoologisch-Botanischen Gesellschaft in Österreich*, 124, 41–46.
- Wendelberger, G. (1989). Zur Klärung des Waldsteppen-Begriffes: Ein Versuch. *Verhandlungen der Zoologisch-Botanischen Gesellschaft in Österreich*, 126, 185–195.
- Wesche, K., Ambarlı, D., Kamp, J., Török, P., Treiber, J., & Dengler, J. (2016). The Palaearctic steppe biome: A new synthesis. *Biodiversity and Conservation*, 25, 2197–2231.
- Woldring, H., & Cappers, R. (2001). The origin of the 'wild orchards' of Central Anatolia. *Turkish Journal of Botany*, 25, 1–9.
- Wu, Z. Y. (Ed.) (1980). *Vegetation of China*. Beijing, China: Science Press. [In Chinese.].
- Xu, C., Liu, H., Anenkhonov, O. A., Korolyuk, A. Y., Sandanov, D. V., Balsanova, L. D., ... Wu, X. (2017). Long-term forest resilience to climate change indicated by mortality, regeneration, and growth in semiarid southern Siberia. *Global Change Biology*, 23, 2370–2382.
- Zamotaev, I. (2002). Soils. In M. Shahgedanova (Ed.), *The physical geography of northern Eurasia* (pp. 103–121). Oxford, UK: Oxford University Press.
- Zech, W., Schad, P., & Hintermaier-Erhard, G. (2014). *Böden der Welt*, 2nd ed. Berlin, Germany: Springer.
- Zhang, M. A., Borjigin, E., & Zhang, H. (2007). Mongolian nomadic culture and ecological culture: On the ecological reconstruction in the agro-pastoral mosaic zone in Northern China. *Ecological Economics*, 62, 19–26.
- Zhang, G., Kang, Y., Han, G., & Sakurai, K. (2011). Effect of climate change over the past half century on the distribution, extent and NPP of ecosystems of Inner Mongolia. *Global Change Biology*, 17, 377–389.
- Zhao, D., Wu, S., Yin, Y., & Yin, Z.-Y. (2011). Vegetation distribution on Tibetan Plateau under climate change scenario. *Regional Environmental Change*, 11, 905–915.
- Zhu, T.-C. (1993). Grasslands of China. In R. T. Coupland (Ed.), *Ecosystems of the world 8B. Natural grasslands. Eastern hemisphere and résumé* (pp. 61–82). Amsterdam, Netherlands: Elsevier.
- Zlotin, R. (2002). Biodiversity and productivity of ecosystems. In M. Shahgedanova (Ed.), *The physical geography of northern Eurasia* (pp. 169–190). Oxford, UK: Oxford University Press.
- Zohary, M. (1973). *Geobotanical foundations of the Middle East*. Stuttgart, Germany: Gustav Fischer.
- Zólyomi, B. (1957). Der Tatarenahorn-Eichen-Lösswald der Zonalen Waldsteppe (Acereto tatarici-Quercetum). *Acta Botanica Hungarica*, 3, 401–424.
- Zólyomi, B., & Fekete, G. (1994). The Pannonian loess steppe: Differentiation in space and time. *Abstracta Botanica*, 18, 29–41.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

APPENDIX S1.

APPENDIX S2. Methods and sources used for the delineation of the forest-steppe biome and the main forest-steppe regions as shown in Fig. 1.

How to cite this article: Erdős L, Ambarlı D, Anenkhonov OA, et al. The edge of two worlds: A new review and synthesis on Eurasian forest-steppes. *Appl Veg Sci*. 2018;21:345–362. <https://doi.org/10.1111/avsc.12382>