

**Nodular calcrete from the Lower Permian Korpád Sandstone Formation
(borehole Dinnyeberki 9015, Mecsek Mts, Hungary) and its palaeoenvironmental
significance**

VARGA Andrea¹, RAUSIK Béla¹, BAJNÓCZI Bernadett²

¹University of Pécs, Department of Geology, 7624 Pécs, Ifjúság útja 6., e-mail: andrea.varga.geol@gmail.com

²Institute for Geological and Geochemical Research, Research Centre for Astronomy and Earth Sciences, Hungarian Academy of Sciences, 1112 Budapest, Budaörsi út 45.

Abstract

The Korpád Sandstone in the D 9015 drill core (Mecsek Mts.) consists of red mudstones and interbedded calcrete crusts with sandstones and conglomerates. Calcrete microfabrics reveal micritic mottles, rhizocretions, smaller root casts and *Microcodium*-like aggregates. These features together with the mineralogy suggest a relatively dry climate with low amount of rainfall (100–500 mm/year) during pedogenesis. Calcite cements are interpreted to have precipitated first in an oxidizing meteoric environment; then, after initial burial, under reducing conditions.

Keywords: beta calcrete, rhizoliths, Microcodium, early diagenesis, Permian, Mecsek Mts.

Rövid közlemény

**Nodular calcrete from the Lower Permian Korpád Sandstone Formation
(borehole Dinnyeberki 9015, Mecsek Mts, Hungary) and its
palaeoenvironmental significance**

VARGA Andrea¹, RAUSIK Béla¹, BAJNÓCZI Bernadett²

¹University of Pécs, Department of Geology, 7624 Pécs, Ifjúság útja 6., e-mail:
andrea.varga.geol@gmail.com

²Institute for Geological and Geochemical Research, Research Centre for Astronomy and
Earth Sciences, Hungarian Academy of Sciences, 1112 Budapest, Budaörsi út 45.

Abstract

The Korpád Sandstone in the D 9015 drill core (Mecsek Mts.) consists of red mudstones and interbedded calcrete crusts with sandstones and conglomerates. Calcrete microfabrics reveal micritic mottles, rhizocretions, smaller root casts and *Microcodium*-like aggregates. These features together with the mineralogy suggest a relatively dry climate with low amount of rainfall (100–500 mm/year) during pedogenesis. Calcite cements are interpreted to have precipitated first in an oxidizing meteoric environment; then, after initial burial, under reducing conditions.

Keywords: beta calcrete, rhizoliths, Microcodium, early diagenesis, Permian, Mecsek Mts.

Introduction

In southern Transdanubia, SW Hungary, Permian sequences are represented mostly by continental siliciclastic and volcanoclastic rocks. They were deposited in continental strike-slip and rift-related basins, belonging to the internal part of the Variscan orogenic domain (SZEDERKÉNYI 2001; VOZÁROVÁ et al. 2009). The syn-eruptive Cisuralian volcanoclastic deposits with a complex stratigraphic architecture can be subdivided into three lithostratigraphic units: upper part of the Korpád Sandstone, the Gyűrűfü Rhyolite and lower part of the Cserdi Conglomerate (BARABÁS & BARABÁSNÉ STUHL 1998; VARGA 2009). Among these units, the fine-grained siliciclastic deposits of Korpád Sandstone are distinctive in terms of presence of *Microcodium*-bearing paleosols (VARGA 2009).

The alluvial Korpád Sandstone occurs in subsurface in southern Transdanubia and ranges up to 700 m in thickness, consisting of polymictic conglomerate, breccia, sandstone and mudrocks. This formation contains a sparse Early Permian macroflora (e.g. *Pecopteris*, *Voltzites*) and a lowermost Permian microflora composed of the *Potonieisporites* and *Vittatina* assemblage (BARABÁS & BARABÁSNÉ STUHL 1998). Despite a long history of research (BARABÁS & BARABÁSNÉ STUHL 1998 and reference therein; SZEDERKÉNYI 2001), paleosols were not recognized previously in this unit; however, a large amount of individual dolomite concretions and concretion aggregates together with animal burrows were described by JÁMBOR (1964) from the red siltstone samples (drill core Dinnyeberki D 9015; *Fig. 1, A and B*). Related to a current research, VARGA (2009) reported that these carbonate concretions are, at least partially, of rhizogenic origin, representing nodular horizons of calcrete profiles.

General characteristics and palaeoclimatic significance of the Korpád calcrete

The studied calcrete sample is characterized by the presence of root traces and associated biogenic structures (*Figs. 1 and 2*), showing beta microfibrils (*sensu* WRIGHT 1990). Its microscopic features reveal micritic mottles, rhizcretions, smaller root casts and *Microcodium*-like aggregates (KLAPPA 1980; KABANOV et al. 2008). The rhizcretions are complex tubular structures up to 6–9 mm in diameter, with a wall structure of irregular micritic laminae which form roughly concentric layers around the central hollow filled by drusy calcite spar cement (*Figs. 1, D and 2*). In the pedogenic micritic laminae of rhizcretions, carbonate is replaced by tiny authigenic (non-luminescent) quartz having euhedral crystal terminations. Root cast cements are predominantly equant calcite spar with a typical drusy fabric. Crystal sizes may reach 400–500 µm towards the centre of the pores. A distinct pattern of cathodoluminescence zoning, representing different cement generations is observed (*Fig. 2*). The initial generation of the root cast cements is non-luminescent or has a very dull luminescence, the second generation is brightly luminescent containing thin dull or non-luminescent bands. The third generation has a homogenous bright orange CL. This cement occurs as the last void fill in large pores; additionally, it can be seen also around the large rhizoliths (*Fig. 2, C and D*).

The surroundings of the rhizcretions are enriched in relic structure of *in situ* *Microcodium* aggregates (*Fig. 1, D*). Unfortunately, *Microcodium* appears as partly to totally recrystallized calcite grains, so primary morphology could not be determined but relics of dark finely dispersed inclusions are obvious. The shape of the aggregates is variable, ranging from spherical clusters to larger cylinders ('corn-cobs'-like) around 150–200 µm in diameter, showing very similar characteristics to the typical kind of *Microcodium* ('*Microcodium a*'; KABANOV et al. 2008). Occasionally, some aggregates are replaced by limpid calcite spar with coarse anhedral crystals showing sutured contacts. It should be noted that the Cisuralian calcrete occurs about 70 m below the boundary of the Korpád Sandstone and Gyűrűfü

Rhyolite (*Fig. 1, B*). In the western part of the Mecsek Mountains, the ~200 m thick overlying complex is interpreted by VARGA (2009) as a strongly to moderately welded (high-grade) ignimbrite, so the recrystallization of *Microcodium* might have been caused by the heating effect (?) of the subsequent volcanic activity.

X-ray diffraction (XRD) studies of the calcrete indicate that the mineral assemblage of the parent material (mudstone rich in strongly altered glass-shards) is dominated by calcite and, subordinately, quartz. Albite, hematite, illite±muscovite, dolomite, smectite and chlorite are minor to trace components. Its clay fraction (<2 µm) consists of illite±muscovite (~80%), chlorite (~10%) and mixed-layer illite/smectite (5–10 %). In calcrete nodules, calcite is the dominant mineral; quartz, albite and illite±muscovite are present in small proportions.

Carbonate palaeosols which form only when evaporation exceeds precipitation are generally considered as reliable palaeoenvironmental and palaeoclimatic indicators. Microbial decomposition releases CO₂ that controls the dissolution and precipitation of pedogenic carbonate (ALONSO-ZARZA 2003). Drusy calcite spar commonly develops in phreatic meteoric environment, representing the very early diagenetic events. Meteoric cements are typically non-luminescent with minor bright CL zones, reflecting the generally oxidizing nature of shallow meteoric waters. After initial burial, reducing conditions develop that result in luminescent calcites (ELIASSEN & TALBOT 2005). The observed features in the rhizolith cements are consistent with this model.

Chemical weathering of primary silicates in the parent material such as albite results in the formation of clay minerals like smectite (APPELO & POSTMA 2009). Regarding hydrological conditions, smectite is preferentially formed in relatively dry climates with low amount of rainfall, where the rate of flushing of the soil is low, so that the solute concentrations become higher. Its formation is further enhanced when rapidly dissolving material such as volcanic rock is available (ALONSO-ZARZA 2003; APPELO & POSTMA 2009). When calcretes are associated with soils containing oxidized iron, smectite and illite, corresponding to the composition of the studied Permian calcrete, they probably indicate semiarid climates with rainfall averages of 100–500 mm/year (RETALLACK 1990; ALONSO-ZARZA 2003).

Acknowledgements

This research has been supported by the Hungarian Scientific Research Found (OTKA) to AV (No. PD 83511), and it was additionally supported by the 'Developing Competitiveness of Universities in the South Transdanubian Region (SROP-4.2.1.B-10/2/KONV-2010-0002)' project (BR) as well as the János Bolyai Research Scholarship of the Hungarian Academy of Sciences (AV). Thanks are also extended to Andrea MINDSZENTY for providing a thoughtful review.

References

- ALONSO-ZARZA, A. M. 2003: Palaeoenvironmental significance of palustrine carbonates and calcretes in the geological record. – *Earth-Science Reviews* **60**, 261–298.
- APPELO, C. A. J. & POSTMA, D. 2009: *Geochemistry, groundwater and pollution*, 2nd edition, CRC Press, Fourth corrected reprint, 649 p.
- BARABÁS, A. & BARABÁS-NÉ STUHL, Á. 1998: A Mecsek és környéke perm képződményeinek rétegtana. In: BÉRCZI, I. & JÁMBOR, Á. (Eds): *Magyarország geológiai képződményeinek rétegtana*, MOL Rt. and Hung. Geol. Inst. publ., Budapest, 187–215.

- ELIASSEN, A. & TALBOT, M. R. 2005: Solution-collapse breccias of the Minkinfjellet and Wordiekammen Formations, Central Spitsbergen, Svalbard: a large gypsum palaeokarst system. – *Sedimentology* **52**, 775–794.
- JÁMBOR, Á. 1964: A Mecsek hegység alsóperm képződményei. Unpublished report, MÉV Adattár (J-0074), Budapest, 113 p.
- KABANOV, P., ANADÓN, P. & KRUMBEIN, W. E. 2008: *Microcodium*: An extensive review and a proposed non-rhizogenic biologically induced origin for its formation. – *Sedimentary Geology* **205**, 79–99.
- KLAPPA, C. F. 1980: Rhizoliths in terrestrial carbonates: classification, recognition, genesis and significance. *Sedimentology* **27**, 613–629.
- KONRÁD GY. & SEBE K. 2010: Fialat tektonikai jelenségek új észlelései a Nyugat-Mecsekben és környezetében. – *Földtani Közlemények* **140/2**, 135–162.
- RETALLACK, G. J. 1990: *Soils of the Past*. Unwyn Hyman, Boston, MA, 520 p.
- SZEDERKÉNYI, T. 2001: Tisza mega-unit. In: HAAS, J. (Ed.): *Geology of Hungary*, Eötvös Kiadó, Budapest, 148–168.
- VARGA, A. 2009: A dél-dunántúli paleozoos–alsó-triász sziliciklasztos kőzetek közettani és geokémiai vizsgálatának eredményei. PhD dissertation, ELTE Dept. of Petrology and Geochemistry, Budapest, 150 p.
- VOZÁROVÁ, A., EBNER, F., KOVÁCS, S., KRÄUTNER, H-G., SZEDERKÉNYI, T., KRSTIĆ, B., SREMAC, J., ALJINOVIČ, D., NOVAK, M. & SKABERNE, D. 2009: Late Variscan (Carboniferous to Permian) environments in the Circum Pannonian Region. – *Geologica Carpathica* **60/1**, 71–104.
- WRIGHT, P. V. 1990: A micromorphological classification of fossil and recent calcic and petrocalcic microstructures. In: DOUGLAS, L. A. (Ed.): *Soil micromorphology: A Basic and Applied Science*. – *Developments in Soil Science* **19**, 401–407.

Figure captions

Figure 1. A) Structural framework and generalized geological map of the Mecsek Mts. (KONRÁD & SEBE 2010) and sample locality (sample code: 15/22; sample depth: 305.1 m). 1 = Neogene; 2 = Jurassic and Cretaceous; 3 = Triassic; 4 = Upper Permian – Lower Triassic; 5 = Palaeozoic in general; 6 = observed fault; 7 = compiled fault; 8 = observed reverse fault; 9 = compiled reverse fault; 10 = strike-slip fault; 11 = syncline; 12 = anticline; B) Generalized lithological column of the Korpád Sandstone Formation in core D 9015 (VARGA 2009); C) Red mudrock with pedogenic calcite nodules. Carbonate precipitation took place only in discontinuous areas in close association with roots; D) Photomicrographs of the Permian calcrete sample 15/22/1. The biological components of the soil became calcified forming rhizoliths (rhizocretions, root casts), calcified filaments, and nodules (upper photos). Rhizoliths occur in close vicinity to *Microcodium*-like calcite aggregates with relics of dark finely dispersed inclusions (lower photos).

Figure 2. Optical (left) and cathodoluminescence (CL; right) photomicrographs of the Permian calcrete sample 15/22/2. A–B) Rhizocretion with complex tubular structure (XPL and CL). In the central hollow filled with drusy calcite spar cement, a distinct pattern of luminescence zoning representing different calcite cement generations is observed. Authigenic quartz crystals are marked by arrows; C–D) Rhizolith with calcite spar cement filled voids (PPL and CL). Note the inclusion-rich calcite cement with homogenous bright orange CL around the rhizolith



