


# Selective Catalytic Activation of Trimethylsilylacetylenes: A “One – Pot” Route to Unsymmetrical Acetylenes and Heterocycles

Dániel Lasányi,<sup>a</sup> Ádám Mészáros,<sup>a</sup> Zoltán Novák,<sup>a,\*</sup> Gergely L. Tolnai<sup>a,\*</sup>

<sup>a</sup> Institute of Chemistry, Eötvös University, Pázmány P. stny 1/a, H1117, Budapest, Hungary  
novakz@elte.hu; web: <http://zng.elte.hu>, tolnai@chem.elte.hu, web: <http://tolnai.chem.elte.hu>

Supporting Information Placeholder




One-pot conditions, wide scope    **Selective TMS Activation!**    Unsymmetrical acetylenes and heterocycles

**ABSTRACT:** For the synthesis of unsymmetrical acetylenes, a Sonogashira coupling-deprotection-Sonogashira coupling reaction sequence is often used. Removal of protecting groups requires harsh conditions or an excess of difficult to handle and expensive reagents. Herein we disclose a novel catalytic method for the selective deprotection of trimethylsilylacetylenes in Sonogashira reaction. The reagent hexafluorosilicic acid, an inexpensive non-toxic compound was used to promote the selective desilylation. This method enables the efficient synthesis of unsymmetric acetylenes with other silylated functional groups present. Frontiers of the method was explored by synthesis of heterocycles.

Diaryl acetylenes are widely used compounds not only because of their synthetic utility, but their unique optical and electronic properties.<sup>1</sup> In the art of molecular engineering diaryl acetylenes are often used as linear building blocks.<sup>2</sup> Their synthesis therefore is a useful conquest of organic chemistry and has different solutions. The first synthesis of diaryl acetylenes consists of elimination and/or rearrangement strategies, usually requires strong bases and harsh reaction conditions.<sup>3</sup> The synthesis strategies for aryl acetylenes has changed with the Sonogashira reaction<sup>4</sup> that is one of the most efficient sp-sp<sup>2</sup> carbon-carbon bond forming reactions. The modern synthesis of unsymmetrical acetylenes preliminary use tandem Sonogashira reactions.<sup>5</sup>


Despite their obvious drawbacks, the use of protecting groups has an inevitable positive impact on organic synthesis. For the efficient synthesis of the unsymmetrical acetylenes, the choice of the right C2 source is crucial. Carbinols are one of the cheapest acetylene sources,<sup>5-6</sup> however the deprotection requires harsh alkaline conditions in the Sonogashira coupling. (Scheme 1) Propiolic acid can also be used as the alkyne source.<sup>7</sup> The alkynylcarboxylic acid can be activated by excess tetrabutylammoniumfluoride (TBAF) or DBU. The most extensively used acetylene source in tandem Sonogashira reaction is trimethylsilylacetylene<sup>8,9</sup> due to the ease of deprotection with excess of KOH, or fluoride source, preliminary TBAF. This hygroscopic and expensive fluoride source disables the feasibility of the synthesis of complex substrates, as other silyl groups, or base-sensitive functionalities cannot be present elsewhere on the molecule. Established methods for Sonogashira couplings directly from alkynylsilanes require the addition of sub-stoichiometric amount of silver or copper metal, or the addition of equivalent fluoride source.<sup>10</sup> These conditions often result in side-products,<sup>11</sup> as well as desilylation of other protected functionalities. Another general drawback of these methods are the high temperature and high Pd-loading.



**Scheme 1. Tandem Sonogashira Protocols**

There is an interesting, yet very specialized method of using catalytic TBAF for the desilylation of TMS-acetylenes in the synthesis of steroidal oral contraceptives. The deprotection only happens in the very basic conditions of carbonyl addition, and took advantage of the nearby alkoxide, that –according to the authors– was able to regenerate the fluoride ion from TMSF.<sup>12</sup>


A general mild catalytic desilylation method was published by Philipps, using buffer solutions (Scheme 2, left).<sup>13</sup> The method utilizes TBAF or CsF as a fluoride source. An elegant fluoride ion free alternative has been shown by Han and coworkers (Scheme 2, right).<sup>14</sup> With catalytic amount of KOTMS, wide variety of alkynylsilanes could be deprotected. Both methods are effectively removing TES, TBDMS, TIPS protecting groups from oxygen and alkynes.



**Scheme 2. Catalytic Protodesilylation Strategies**

However these methods serve a useful way to gain terminal acetylenes, those type of compounds are rarely the goal of synthesis. Terminal acetylenes are usually serve as building block for internal acetylenes or heterocycles.<sup>15</sup>

DeShong's studies on deprotection of O benzyl-silyl ethers showed, that hexafluorosilicic acid is a very potent compound in cleavage of Si-O bonds.<sup>16</sup> Later studies (Scheme 3) proved, that using  $\text{H}_2\text{SiF}_6$  in a sub-stoichiometric manner provide selective removal of TBDMS over TIPS from benzyl-ethers.<sup>17</sup>




**Scheme 3. Selective Deprotection of O-Si Bond with  $\text{H}_2\text{SiF}_6$**

These results and our experience on desilylation of alkynylacetamides<sup>18</sup> inspired us to find a general selective catalytic desilylation method in tandem Sonogashira reactions.  $\text{H}_2\text{SiF}_6$  is a very cheap<sup>19</sup> industrial starting material available as an aqueous solution, used in water fluorination of drinking water.<sup>20</sup> The fact, that the compound is able to remove the silyl group in a sub-equivalent fashion, encouraged us to further improve its efficiency to establish a general tandem Sonogashira reaction that is able to generate unsymmetrical acetylenes with a wide scope.

To elucidate the frontiers of the desilylating agent in the Sonogashira conditions, iodobenzene (**1a**) and TMS-phenylacetylene (**2**) were applied as model substrates, and the coupling reactions were performed in the presence of common 3 mol%  $\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$ , and CuI catalyst system. To our delight 1.0 and 0.5 equivalent of the addition of hexafluorosilicic acid resulted in complete conversion by GC under the Sonogashira conditions (Table 1, entries 1-2). The addition of 1 % of this compound resulted in 28 % conversion (entry 3). As the  $\text{H}_2\text{SiF}_6$  is in a 34 % aqueous solution, we assumed, that the lack of nucleophile could have led to the loss of activity in our case. Thus, adding 10 equivalents of water increased the yield to 60 %, but only in the presence of hexafluorosilicic acid (entry 4-5). However the reaction resulted in full conversion in 48 h, it was found, that increasing the temperature sped up the reaction without noticeable side effects. The reaction led to 91 % conversion in 24 h on 60 °C (entry 7). As the presence of water turned out to be an important factor, different amounts of water was added over the aqueous solution (entries 8-10). No loss of activity was observed down to 5 equivalents of  $\text{H}_2\text{O}$ , but either increasing to 20 equivalents, or decreasing to 1 equivalent was inferior to 5 or 10 equivalents in these conditions. With 1% HCl, from a 36 % aqueous solution and 10 equivalents of water, the conversion was 5 % (entry 11). A control experiment omitting hexafluorosilicic acid resulted in only 7% yield in our condition (entry 12). The low efficiency of “sila” Sonogashira coupling under these mild conditions predicts the good efficiency of tandem reaction, as the homocoupling could be suppressed, without a further manipulation with catalysts or temperature.

**Table 1. Optimization of Reaction Conditions**




Entry	$\text{H}_2\text{SiF}_6$ (mol%)	$\text{H}_2\text{O}$ added (equiv.)	T (°C)	Yield <sup>a</sup> (%)
1	100		25	98
2	50		25	98
3	1	-	25	28
4	1	10	25	60 <sup>b</sup>
5	0	10	25	traces

6	0.5	10	60	50
7	<b>1</b>	<b>10</b>	<b>60</b>	<b>91</b>
8	1	5	60	87
9	1	20	60	61
10	1	1	60	39
11	1 <sup>c</sup>	10	60	5
12	0	10	60	7

[a] As determined by GC-FID, on 0.5 mmol scale [b] full conversion in 48 h [c] HCl was used instead of H<sub>2</sub>SiF<sub>6</sub>

With the optimized reaction condition in hand, we turned our attention to establish conditions for a tandem coupling reaction. As predicted by the preliminary results, the deprotection of the acetylene was not likely to occur without the H<sub>2</sub>SiF<sub>6</sub>. Indeed, the first Sonogashira coupling proceeded to complete conversion, enabling a one-pot procedure for the coupling. To establish the synthetic utility of the method, we examined the scope of this reaction with 1 mmol of the starting aryl halides, in the presence of only 1 mol% of the desilylating agent.


First, we ran a series of experiments with different monofunctionalized aryl iodides, TMS-acetylene and iodobenzene. As expected, diaryl acetylenes (Scheme 4, **3a-f**) were obtained with different functionalities, including electron donating and electron withdrawing in medium to excellent yields. For medicinal chemistry applications, the use of heterocyclic compounds is very important. Our one pot method proved to be excellent in this field, as iodo- pyridine, indole, thiophene and quinolone (**3g-j**) provided the corresponding internal acetylenes in very good to excellent yields.




**Scheme 4. Synthesis of Unsymmetric Acetylenes (I)**

Acetylenic thiophenes are present in material science as conducting polymers and nonlinear optical materials.<sup>21</sup> Using the developed conditions, series of unsymmetrical thienyl acetylenes (Scheme 5, **3k-m**) were synthesized in good yields. Continuing the exploration of the substrate scope of acetylene products, series of acetylenes with different aryl groups (**3n-t**) were synthesized.


Multicyclic hydrocarbons **3u-v**, with extended conjugation were also synthesized with good efficiency. Synthesis of the molecular rod **3u** provided 82 % yield, with 2.2 equivalent TMSA and 0.5 % H<sub>2</sub>SiF<sub>6</sub> respectively to the TMS.






**Scheme 5. Synthesis of Unsymmetric Acetylenes (II)**

To further challenge our synthetic methodology, we aimed to explore the stability of different silyl groups. Primary silyl ethers are the most labile silyl ethers due to the lack of steric hindrance. 3-Iodobenzyl alcohol was protected with different silyl protecting groups. As a first example, coupling and selective TMS-acetylene activation was demonstrated with the TES, TBDPS, TBDMS and TIPS groups (Scheme 6, **3w-z**). All of these aryl-iodides provided the unsymmetrical diaryl acetylene. Competing deprotection in a ratio of 1:9 was only observed in the case of O-TES, indicating that the low yield might only be a problem with the workup. Iodoindole, with a TIPS-protected acetylene at position 3<sup>22</sup> was coupled with TMS-acetylene, and one pot coupled with iodopyridine, by the aid of 1%  $\text{H}_2\text{SiF}_6$ . To our delight the expected product **3aa** was isolated in 71% yield, and no sign of TIPS-deprotection was observed. In a further control experiment (Scheme 6, down), the mixture the 1 equivalent of iodobenzene was reacted in our optimized conditions with equimolar TES- and TMS-phenylacetylenes. As a result full conversion of PhI and only ~1% of TMS-phenylacetylene was detected after 24 h. The TES-protected acetylene remained intact.




**Scheme 6. Synthesis of Internal Acetylenes with Various Silyl Protecting Groups Present**

One of the preliminary uses of aromatic acetylenes in organic synthesis is applying them as building blocks for the construction of heterocycles. Utilizing the exact same conditions, a 4 step one-pot synthesis of benzofurans was performed, where the Sonogashira coupling, the deprotection and the second coupling followed by ring closing takes place. A series of complex heterocycles were produced (Scheme 7, **4a-f**). This procedure affects the formation of 3 new C-C or C-O bonds. However the conditions are unoptimized and this lead to some degree of (presumably) polymerized side-products in the case of **4a** and **f**, the yields are ranging up to 93% (**4b-e**)




**Scheme 7. Synthesis of Benzofurans**

An other general use of acetylenes are synthesizing triazoles, therefore the general conditions were challenged to examine to scope of a tandem reaction of this kind. Same condition has been applied to generate a small set of triazoles (Scheme 8, **5a-c**) via CuAAC type of ring construction. To our delight the catalytic coupling-desilylation protocol was useful in synthesizing this class of compounds in one-pot.



**Scheme 8. Synthesis of Triazoles**

The mechanistic aspects of this reaction are not clear at the moment. The need of water for the efficient synthesis could suggest the preliminary formation of a terminal acetylene. No meaningful difference in reaction rate was observed, when H<sub>2</sub>O was compared to D<sub>2</sub>O in the coupling of TMS-phenylacetylene (**2**) with iodobenzene (**1a**), under optimized conditions. Control experiment on coupling with bis(trimethylsilyl)acetylene (Scheme 9, **6**) with 2 equivalents of **1a** led to high yield of **3a** (95%), instead of generating ethyn. This could be due to the direct transmetalation from Si to Cu or Pd or the immediate Sonogashira reaction in this condition therefore avoiding the formation of HCCH.



**Scheme 9. Investigation on Mechanism**

These results were examined more in depth: in a control experiment iodobenzene was omitted from the coupling of TMS-phenylacetylene (Scheme 9). Surprisingly, instead of protodesilylation, no change was observed on the starting materials after 24 h. Another pathway would be the formation of <sup>-</sup>OTMS, from water and the protecting group in situ. This version does not seem likely, as Han's method<sup>14</sup> is a general desilylation process, and there is a high TMS selectivity in our case. As a plausible reaction scheme, hexafluorosilicic acid is first hydrolyzed. The transmetalation to the transition metal can be accelerated by the coordination of the SiF<sub>x</sub> as a Lewis acid. Transmetalation from Si to Cu is suggested in the literature.<sup>23</sup>

In conclusion, a new catalyst, the cheap and easy to handle  $\text{H}_2\text{SiF}_6$  was introduced for the selective desilylation of TMS alkynes in tandem Sonogashira cross-coupling reaction. With the optimized protocol a tandem Sonogashira condition was developed, where great variety of unsymmetrical acetylenes were synthesized, with very low catalyst loading. The new selective trimethylsilane activating agent enabled the coupling of substrates with the generally used Si-O and Si-acetylene protecting groups present. Moreover the synthetic utility was demonstrated in the synthesis of benzofuran and triazole heterocycles.

## ASSOCIATED CONTENT

### Supporting Information

This material is available free of charge via the Internet at <http://pubs.acs.org>.

## AUTHOR INFORMATION

### Corresponding Author

\* novakz@elte.hu, \*tolnai@chem.elte.hu

## ACKNOWLEDGMENT

PPD003/2016 and Janos Bolyai Research Scholarship of Hungarian Academy of Science is gratefully acknowledged for financial support. Ms. M. Woodward is acknowledged for proofreading the manuscript.

## EXPERIMENTAL SECTION

Unless otherwise indicated, all starting materials were obtained from commercial suppliers, and were used without further purification. Analytical thin-layer chromatography (TLC) was performed on Merck DC precoated TLC plates with 0.25 mm Kieselgel 60 F<sub>254</sub>. Visualization was performed with a 254 nm UV lamp. The  $^1\text{H}$ ,  $^{13}\text{C}$  and  $^{19}\text{F}$  NMR spectra were recorded on a Bruker Avance-250 spectrometer and in  $\text{CDCl}_3$ , and  $\text{DMSO}-d_6$ . Chemical shifts are expressed in parts per million ( $\delta$ ) using residual solvent protons as internal standards ( $\delta$  7.26 for  $^1\text{H}$ ,  $\delta$  77.0 for  $^{13}\text{C}$ ) for  $\text{CDCl}_3$  and ( $\delta$  2.50 for  $^1\text{H}$ ,  $\delta$  39.5 for  $^{13}\text{C}$ ) for  $\text{DMSO}-d_6$ . Coupling constants ( $J$ ) are reported in Hertz (Hz). Splitting patterns are designated as *s* (singlet), *d* (doublet), *t* (triplet), *q* (quartet), *m* (multiplet). Combination gas chromatography and low resolution mass spectrometry was obtained on an Agilent 6890N Gas Chromatograph (30 m x 0.25 mm column with 0.25  $\mu\text{m}$  HP-5MS coating, He carrier gas) and Agilent 5973 Mass Spectrometer (Ion source: EI+, 70 eV, 230 °C; interface: 300 °C). IR spectra were obtained on a Bruker IFS55 spectrometer on a single-reflection diamond ATR unit. All melting points were measured on Büchi 501 apparatus and are uncorrected. High-resolution mass spectra were acquired on an Agilent 6230 time-of-flight mass spectrometer equipped with a Jet Stream electrospray ion source in positive ion mode. Injections of 0.1–0.3  $\mu\text{L}$  were directed to the mass spectrometer at a flow rate of 0.5 ml/min (70% acetonitrile–water mixture, 0.1 % formic acid), using an Agilent 1260 Infinity HPLC system. Jet: drying gas ( $\text{N}_2$ ) flow and temperature: 10.0 l/min and 325 °C, respectively; nebulizer gas (Stream parameters  $\text{N}_2$ ) pressure: 10 psi; capillary voltage: 4000 V; sheath gas flow and temperature: 325 °C and 7.5 l/min; TOFMS parameters: fragmentor voltage: 120 V; skimmer potential: 120 V; OCT 1 RF Vpp: 750 V. Full-scan mass spectra were acquired over the *m/z* range 100–2500 at an acquisition rate of 250 ms/spectrum and processed by Agilent MassHunter B.03.01 software.

### General procedure:

Into a 7 mL screw cap vial CuI (5.7 mg, 0.03 mmol, 0.03 equiv.) and  $\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$  (21.1 mg, 0.03 mmol, 0.03 equiv.) was and the starting aryl halogenide (1 mmol, 1 equiv.) if a solid, was filled. The vial was closed, and the atmosphere was changed 3 times to Ar. DIPA (4 mL) was added, and the starting aryl halogenide (1 mmol, 1 equiv.) if a liquid. Trimethylsilylacetylene (154  $\mu\text{L}$ , 1.1 mmol, 1.1 equiv.) was added via a syringe and the mixture was stirred at 60 °C for 1 h. The completion of the first coupling was verified by TLC, then the second aryl halogenide (1 mmol, 1 equiv.) was added. Hexafluorosilicic acid (34 % in aqueous solution, 3  $\mu\text{L}$ , 0.01 mmol, 0.01 equiv.) and water (180  $\mu\text{L}$ ) was added via Hamilton syringes. The mixture was stirred for 24 h at 60 °C. The mixture was diluted with EtOAc (15 mL) and water (15 mL), the pH of the aqueous phase was adjusted to neutral. The layers were separated, then the aqueous phase was extracted with EtOAc (2 x 15 mL). The combined organic layers were dried over  $\text{MgSO}_4$ . The solvent was removed under reduced pressure, and the crude product was purified by column chromatography.

### 1,2-Diphenylethyne<sup>24</sup> 3a

The general procedure was followed. White solid (175.5 mg, yield 98%), Mp.: 54–55 °C (Ref.: 54–55 °C),  $R_f$ : 0.53 (in hexanes:EtOAc 25:1).  $^1\text{H}$  NMR (250 MHz,  $\text{CDCl}_3$ )  $\delta$  = 7.70 – 7.51 (m, 4H), 7.48 – 7.30 (m, 6H) ppm.  $^{13}\text{C}$  NMR (63 MHz,  $\text{CDCl}_3$ )  $\delta$  = 132.1, 128.8, 128.7, 123.7, 89.8 ppm. IR (thin film, ATR)  $\nu_{\text{max}}$  3063, 1599, 1571, 1492, 1441, 1386, 1331, 1312, 1280, 1178, 1156, 1102, 1069, 1025, 997, 985, 916, 850, 753, 686, 534, 507, 465  $\text{cm}^{-1}$ . MS (EI, 70 eV) *m/z* (%): 178 (100, [ $\text{M}^+$ ]), 152 (20), 126 (10), 76 (15).

### 5 mmol Scale Experiment

Into a 50 mL round bottomed flask CuI (28.6 mg, 0.15 mmol, 0.03 equiv.) and  $\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$  (105.3 mg, 0.15 mmol, 0.03 equiv.) was and the starting aryl halogenide (5 mmol, 1 equiv.) if a solid, was filled. The flask was closed, and the atmosphere was changed 3 times to Ar. DIPA (20 mL) was added, and the starting aryl halogenide (5 mmol, 1 equiv.) if a liquid. Trimethylsilylacetylene (780  $\mu\text{L}$ , 5.5 mmol, 1.1 equiv.) was added via a syringe and the mixture was stirred at 60 °C for 1 h. The completion of the first coupling was verified by TLC, then the second aryl halogenide or azide (1 mmol, 1 equiv.) was added. Hexafluorosilicic acid (34 % in aqueous solution, 15.2  $\mu\text{L}$ , 0.01 mmol, 0.01 equiv.) and water (900  $\mu\text{L}$ ) was added via Hamilton syringes. The mixture was stirred for 24 h at 60 °C. The mixture was diluted with EtOAc (75 mL) and water (75 mL), the pH of the aqueous phase was adjusted to neutral. The

layers were separated, then the aqueous phase was extracted with EtOAc (2 x50 mL). The combined organic layers were dried over MgSO<sub>4</sub>. The solvent was removed under vacuum, and the crude was purified by column chromatography. White crystals. (868.6 mg, 98 %) Analytical data is the same as at 1 mmol scale.

#### 1-(4-(Phenylethynyl)phenyl)ethan-1-one<sup>24</sup> 3b

The general procedure was followed. White solid (154.2 mg, yield 70%) Mp.= 96-97 °C (Ref.: 95-96 °C), R<sub>f</sub>: 0.23 (in hexanes: EtOAc 10:1). <sup>1</sup>H NMR (250 MHz, CDCl<sub>3</sub>) δ = 7.86 (d, *J* = 8.5 Hz, 2H), 7.57 – 7.41 (m, 4H), 7.34 – 7.23 (m, 3H), 2.53 (s, 3H) ppm. <sup>13</sup>C NMR (63 MHz, CDCl<sub>3</sub>) δ = 197.7, 136.6, 132.1, 132.1, 129.2, 128.8, 128.7, 128.6, 123.0, 93.1, 89.0, 27.0 ppm. IR (thin film, ATR) ν<sub>max</sub> 1680, 1603, 1486, 1443, 1404, 1361, 1266, 1179, 1109, 1071, 960, 835, 762, 693 cm<sup>-1</sup>. MS (EI, 70 eV) *m/z* (%): 220(75), 205(100, [M<sup>+</sup>]), 176(70), 151(32).

#### 1-Methoxy-4-(phenylethynyl)benzene<sup>24</sup> 3c

The general procedure was followed. White solid (119.2 mg, yield 57%) Mp.= 58-59 °C (Ref.: 59-60 °C), R<sub>f</sub>: 0.29 (in hexanes: EtOAc 10:1). <sup>1</sup>H NMR (250 MHz, CDCl<sub>3</sub>) δ = 7.60 – 7.41 (m, 4H), 7.39 – 7.29 (m, 3H), 6.97 – 6.83 (m, 2H), 3.84 (s, 3H) ppm. <sup>13</sup>C NMR (63 MHz, CDCl<sub>3</sub>) δ = 160.0, 133.4, 131.8, 128.7, 128.3, 124.0, 115.8, 114.4, 89.8, 88.5, 55.7 ppm. IR (thin film, ATR) ν<sub>max</sub> 2212, 1604, 1592, 1266, 1506, 1457, 1439, 1285, 1245, 1173, 1136, 1106, 1068, 1025, 914, 835, 828, 778, 751, 688, 655, 641, 545, 519, 472 cm<sup>-1</sup>. MS (EI, 70 eV) *m/z* (%): 208 (100, [M<sup>+</sup>]), 193 (46), 165 (40), 139 (12), 104 (5). The sample contains 5 % of ((4-methoxyphenyl)ethynyl)trimethylsilane. Yield is corrected with purity.

#### 1-Methyl-4-(phenylethynyl)benzene<sup>24</sup> 3d

The general procedure was followed. White solid (107.0 mg, yield 56%) Mp.= 70-72 °C (Ref.: 70-72 °C), R<sub>f</sub>: 0.60 (in hexanes). <sup>1</sup>H NMR (250 MHz, CDCl<sub>3</sub>) δ = 7.47 – 7.36 (m, 2H), 7.32 (d, *J* = 8.1 Hz, 2H), 7.27 – 7.17 (m, 3H), 7.04 (d, *J* = 7.9 Hz, 2H), 2.25 (s, 3H) ppm. <sup>13</sup>C NMR (63 MHz, CDCl<sub>3</sub>) δ = 138.6, 131.9, 131.9, 129.5, 128.7, 128.5, 123.6, 120.4, 90.1, 89.0, 21.7 ppm. IR (thin film, ATR) ν<sub>max</sub> 1593, 1508, 1483, 1440, 1407, 1106, 1089, 1016, 915, 816, 753, 689, 514, 494, 468 cm<sup>-1</sup>. MS (EI, 70 eV) *m/z* (%): 192 (100, [M<sup>+</sup>]), 191 (50), 165 (20), 152 (5), 115 (5).

#### 3-(Phenylethynyl)aniline<sup>25</sup> 3e

The general procedure was followed. Brown solid (170.1 mg, yield 88%) Mp.= 46-48 °C, R<sub>f</sub>: 0.25 (in hexanes: EtOAc 5:1). <sup>1</sup>H NMR (250 MHz, CDCl<sub>3</sub>) δ = 7.51 – 7.35 (m, 2H), 7.32-7.19 (m, 3H), 7.05 (t, *J* = 7.8 Hz, 1H), 6.95 – 6.82 (m, 1H), 6.78 (t, *J* = 1.9 Hz, 1H), 6.64 – 6.50 (m, 1H), 3.60 (s, 2H) ppm. <sup>13</sup>C NMR (63 MHz, CDCl<sub>3</sub>) δ = 146.52, 132.0, 129.7, 128.7, 128.6, 124.3, 123.7, 122.6, 118.3, 115.8, 90.0, 89.2 ppm. IR (thin film, ATR) ν<sub>max</sub> 1678, 1599, 1494, 1445, 1329, 1293, 1252, 1211, 1166, 1132, 1072, 1027, 994, 944, 915, 865, 781, 754, 689 cm<sup>-1</sup>. MS (EI, 70 eV) *m/z* (%): 263(100, [M<sup>+</sup>]), 227(35), 200(40), 131(10).

#### 3-(Phenylethynyl)benzonitrile<sup>26</sup> 3f

The general procedure was followed. Yellow oil (140.8 mg, yield 90%) R<sub>f</sub>: 0.48 (in hexanes:EtOAc 5:1). <sup>1</sup>H NMR (250 MHz, CDCl<sub>3</sub>) δ = 7.87 – 7.78 (m, 1H), 7.73 (dt, *J* = 7.8, 1.4 Hz, 1H), 7.60 (dt, *J* = 7.8, 1.4 Hz, 1H), 7.57 – 7.31 (m, 6H) ppm. <sup>13</sup>C NMR (63 MHz, CDCl<sub>3</sub>) δ = 136.0, 135.3, 132.1, 131.7, 129.7, 129.4, 128.9, 125.4, 122.7, 118.5, 113.3, 92.2, 87.3 ppm. IR (thin film, ATR) ν<sub>max</sub> 2233, 1601, 1574, 1493, 1443, 1414, 1094, 1070, 1026, 896, 797, 756, 681 cm<sup>-1</sup>. MS (EI, 70 eV) *m/z* (%): 203 (100, [M<sup>+</sup>]), 176 (15), 151 (10), 101 (10).

#### 3-(Phenylethynyl)pyridine<sup>27</sup> 3g

The general procedure was followed. White solid (168.2 mg, yield 93%) Mp.= 50-51 °C (Ref.: 49-51 °C), R<sub>f</sub>: 0.29 (in hexanes: EtOAc 20:1). <sup>1</sup>H NMR (250 MHz, CDCl<sub>3</sub>) δ = 8.72 (s, 1H), 8.50 (dd, *J* = 4.9, 1.7 Hz, 1H), 7.76 (dt, *J* = 7.9, 2.0 Hz, 1H), 7.41-7.58 (m, 2H), 7.39 – 7.13 (m, 4H) ppm. <sup>13</sup>C NMR (63 MHz, CDCl<sub>3</sub>) δ = 152.5, 148.8, 138.9, 132.1, 129.2, 128.8, 123.5, 122.9, 120.9, 93.1, 86.3 ppm. IR (thin film, ATR) ν<sub>max</sub> 1559, 1488, 1472, 1413, 1329, 1312, 1280, 1188, 1175, 1145, 1122, 1101, 1071, 1036, 1020, 997, 935, 920, 814, 756, 708, 689, 626, 538, 513, 464 cm<sup>-1</sup>. MS (EI, 70 eV) *m/z* (%): 179(100, [M<sup>+</sup>]), 151(20), 126(30), 100(15).

#### 2-(Phenylethynyl)thiophene<sup>28</sup> 3h

The general procedure was followed. Pale rose crystals (134.8 mg, yield 73%) Mp.= 52-54 °C, R<sub>f</sub>: 0.58 (in hexanes). <sup>1</sup>H NMR (250 MHz, CDCl<sub>3</sub>) δ = 7.60 – 7.46 (m, 2H), 7.44 – 7.25 (m, 5H), 7.03 (dd, *J* = 5.0, 3.8 Hz, 1H) ppm. <sup>13</sup>C NMR (63 MHz, CDCl<sub>3</sub>) δ = 132.3, 131.8, 128.8, 128.8, 127.6, 127.51, 123.7, 123.3, 93.4, 83.0 ppm. IR (thin film, ATR) ν<sub>max</sub> 1594, 1516, 1484, 1441, 1423, 1356, 1275, 1212, 1176, 1155, 1110, 1069, 1037, 1023, 970, 917, 851, 831, 789, 752, 700, 686, 570, 551, 528, 499, 432 cm<sup>-1</sup>. MS (EI, 70 eV) *m/z* (%): 184(100, [M<sup>+</sup>]), 152(21), 139(33).

#### 5-(Phenylethynyl)-1*H*-indole<sup>29</sup> 3i

The general procedure was followed. Yellow solid (179.1 mg, yield 83%) Mp.= 126-128 °C (Ref.: 124-128 °C), R<sub>f</sub>: 0.27 (in hexanes:EtOAc 10:1). <sup>1</sup>H NMR (250 MHz, CDCl<sub>3</sub>) δ = 8.20 (s, 1H), 7.91 (s, 1H), 7.58 (m, *J* = 4.7, 2.7 Hz, 2H), 7.47 – 7.28 (m, 5H), 7.23 (t, *J* = 2.8 Hz, 1H), 6.58 (t, *J* = 2.6 Hz, 1H) ppm. <sup>13</sup>C NMR (63 MHz, CDCl<sub>3</sub>) δ = 135.8, 131.9, 128.7, 128.2, 128.2, 126.1, 125.51, 125.1, 124.34, 114.8, 111.56, 103.3, 91.5, 87.5 ppm. IR (thin film, ATR) ν<sub>max</sub> 2208, 1595, 1490, 1467, 1441, 1413, 1343, 1323, 1300, 1245, 1088, 1067, 1022, 891, 810, 754, 726, 691 cm<sup>-1</sup>. MS (EI, 70 eV) *m/z* (%): 217(100, [M<sup>+</sup>]), 189(30), 108(15).

#### 6-(Phenylethynyl)quinoline<sup>30</sup> 3j

The general procedure was followed. Yellow solid (201.1 mg, yield 88%) Mp.= 94-96 °C, R<sub>f</sub>: 0.35 (in hexanes: EtOAc 1:1). <sup>1</sup>H NMR (250 MHz, CDCl<sub>3</sub>) δ = 8.89 (d, *J* = 4.3 Hz, 1H), 8.21 – 7.91 (m, 3H), 7.81 (d, *J* = 8.7 Hz, 1H), 7.58 (dd, *J* = 6.8, 3.0 Hz, 2H), 7.44 – 7.28 (m, 4H) ppm. <sup>13</sup>C NMR (63 MHz, CDCl<sub>3</sub>) δ = 151.25, 147.96, 136.16, 132.58, 132.10, 131.47, 129.94, 128.98, 128.83,

128.40, 123.30, 122.12, 122.00, 91.10, 89.38 ppm. IR (thin film, ATR)  $\nu_{\max}$  1590, 1567, 1496, 1441, 1372, 1333, 1129, 1071, 1027, 920, 889, 839, 800, 759, 692  $\text{cm}^{-1}$ . MS (EI, 70 eV)  $m/z$  (%): 229(100,  $[\text{M}^+]$ ), 228(20), 200(10), 114(6).

#### **Methyl 4-(thiophen-2-ylethynyl)benzoate<sup>32</sup> 3k**

The general procedure was followed. Yellow crystals (185 mg, yield 76%) Mp.: 107-109 °C (Ref.: 104-105 °C),  $R_f$ : 0.23 (in hexanes: EtOAc 10:1).  $^1\text{H}$  NMR (250 MHz,  $\text{CDCl}_3$ )  $\delta$  = 7.92 (d,  $J$  = 8.3 Hz, 2H), 7.47 (d,  $J$  = 8.2 Hz, 2H), 7.27 – 7.15 (m, 2H), 6.93 (dd,  $J$  = 4.9, 3.9 Hz, 1H), 3.82 (s, 3H) ppm.  $^{13}\text{C}$  NMR (63 MHz,  $\text{CDCl}_3$ )  $\delta$  = 166.9, 133.0, 131.6, 129.9, 128.4, 128.0, 127.6, 123.0, 100.4, 92.8, 86.1, 52.6 ppm. IR (thin film, ATR)  $\nu_{\max}$  1711, 1603, 1439, 1408, 1276, 1215, 1175, 1104, 866, 771, 705  $\text{cm}^{-1}$ . MS (EI, 70 eV)  $m/z$  (%): 243(16), 242(100,  $[\text{M}^+]$ ), 211(95), 183(20), 139(67), 91(10). HRMS calcd for  $\text{C}_{14}\text{H}_{11}\text{O}_2\text{S}$   $[\text{M}+\text{H}]^+$  243.0480 found 243.0471.

#### **5-(Thiophen-2-ylethynyl)-1H-indole 3l**

The general procedure was followed. Brown solid (135 mg, yield 60%) Mp.: 154-156 °C,  $R_f$ : 0.20 (in hexanes: EtOAc 5:1).  $^1\text{H}$  NMR (250 MHz,  $\text{CDCl}_3$ )  $\delta$  = 8.16 (s, 1H), 7.80 (s, 1H), 7.30 (m, 2H), 7.23 – 7.08 (m, 3H), 6.95 (dd,  $J$  = 5.2, 3.6 Hz, 1H), 6.50 (t,  $J$  = 2.6 Hz, 1H) ppm.  $^{13}\text{C}$  NMR (63 MHz,  $\text{CDCl}_3$ )  $\delta$  = 133.4, 129.1, 125.6, 124.9, 124.4, 123.3, 123.0, 122.4, 122.0, 111.8, 109.1, 100.8, 92.6, 78.1 ppm. IR (thin film, ATR)  $\nu_{\max}$  2931, 1612, 1590, 1469, 1412, 1344, 1316, 1259, 1219, 1182, 1089, 1036, 888, 850, 811, 764, 730, 693  $\text{cm}^{-1}$ . MS (EI, 70 eV)  $m/z$  (%): 224(16), 223(100,  $[\text{M}^+]$ ), 195(12), 152(10), 111(5). HRMS calcd for  $\text{C}_{14}\text{H}_{10}\text{NS}$   $[\text{M}+\text{H}]^+$  224.0534 found 224.0535.

#### **2-((3,5-bis(Trifluoromethyl)phenyl)ethynyl)thiophene 3m**

The general procedure was followed. White crystals (260 mg, yield 81%), Mp.: 43-45 °C  $R_f$ : 0.31 (in hexanes).  $^1\text{H}$  NMR (250 MHz,  $\text{CDCl}_3$ )  $\delta$  = 7.94 (s, 2H), 7.82 (s, 1H), 7.43 – 7.32 (m, 2H), 7.06 (dd,  $J$  = 5.0, 3.8 Hz, 1H) ppm.  $^{13}\text{C}$  NMR (63 MHz,  $\text{CDCl}_3$ )  $\delta$  = 133.6, 132.4 (q,  $J$  = 34.2 Hz), 131.5 (m), 129.0, 127.7, 125.6, 123.3 (d,  $J$  = 273.3 Hz), 122.1, 121.9 (p,  $J$  = 3.8 Hz), 90.4, 86.6 ppm.  $^{19}\text{F}$  NMR (235 MHz,  $\text{CDCl}_3$ )  $\delta$  = -63.21 ppm. IR (thin film, ATR)  $\nu_{\max}$  2214, 1383, 1349, 1277, 1174, 1132, 896, 849, 698  $\text{cm}^{-1}$ . MS (EI, 70 eV)  $m/z$  (%): 320(100,  $[\text{M}^+]$ ), 301(20), 251(10), 206(10), 160(5), 69(10). HRMS cannot be obtained by our facility.

#### **1-((3-Fluorophenyl)ethynyl)-3,5-dimethylbenzene 3n**

The general procedure was followed. White crystals (198 mg, yield 88%), Mp.: 38-40 °C  $R_f$ : 0.37 (in hexanes).  $^1\text{H}$  NMR (250 MHz,  $\text{CDCl}_3$ )  $\delta$  = 7.33 – 7.14 (m, 5H), 7.10 – 6.90 (m, 2H), 2.33 (s, 6H) ppm.  $^{13}\text{C}$  NMR (63 MHz,  $\text{CDCl}_3$ )  $\delta$  = 162.8 (d,  $J$  = 245.6 Hz), 138.4, 130.9, 130.3 (d,  $J$  = 8.4 Hz), 129.7, 127.8 (d,  $J$  = 3.0 Hz), 125.7 (d,  $J$  = 8.6 Hz), 122.8, 118.7 (d,  $J$  = 23.8 Hz), 115.8 (d,  $J$  = 23.8 Hz), 91.1, 87.8, 21.5 ppm.  $^{19}\text{F}$  NMR (235 MHz,  $\text{CDCl}_3$ )  $\delta$  = -113.07 ppm. IR (thin film, ATR)  $\nu_{\max}$  2920, 2863, 2219, 1610, 1580, 1489, 1436, 1379, 1335, 1304, 1261, 1181, 1119, 1074, 1039, 960, 933, 850, 784, 719, 681  $\text{cm}^{-1}$ . MS (EI, 70 eV)  $m/z$  (%): 295(100,  $[\text{M}^+]$ ), 276(10), 233(30), 190(20), 163(15), 138(10), 108(5), 75(10). HRMS cannot be obtained by our facility.

#### **1-Chloro-4-((4-fluorophenyl)ethynyl)benzene<sup>32</sup> 3o**

The general procedure was followed. White solid (121 mg, yield 53%), Mp.: 112-113 °C, (Ref.: 113 °C)  $R_f$ : 0.26 (in hexanes).  $^1\text{H}$  NMR (250 MHz,  $\text{CDCl}_3$ )  $\delta$  = 7.57 – 7.40 (m, 4H), 7.38 – 7.26 (m, 2H), 7.15 – 6.96 (m, 2H) ppm.  $^{13}\text{C}$  NMR (63 MHz,  $\text{CDCl}_3$ )  $\delta$  = 163.0 (d,  $J$  = 250.0 Hz), 134.7, 133.9 (d,  $J$  = 8.5 Hz), 133.2, 129.1, 122.0, 119.4 (d,  $J$  = 3.5 Hz), 116.1 (d,  $J$  = 22.9 Hz), 89.6, 88.3 ppm.  $^{19}\text{F}$  NMR (235 MHz,  $\text{CDCl}_3$ )  $\delta$  = -110.48 ppm. IR (thin film, ATR)  $\nu_{\max}$  1601, 1506, 1400, 1226, 1158, 1088, 1012, 831, 702  $\text{cm}^{-1}$ . MS (EI, 70 eV)  $m/z$  (%): 230(100,  $[\text{M}^+]$ ), 194(32), 175(15), 144(5), 115(5), 97(5).

#### **1-Methoxy-4-((4-nitrophenyl)ethynyl)benzene<sup>33</sup> 3p**

The general procedure was followed. Yellow solid (114 mg, yield 45%), Mp.: 122- 124 °C (Ref.: 122-124 °C),  $R_f$ : 0.25 (in hexanes: EtOAc 10:1).  $^1\text{H}$  NMR (250 MHz,  $\text{CDCl}_3$ )  $\delta$  = 8.09 (d,  $J$  = 8.9 Hz, 2H), 7.52 (d,  $J$  = 8.9 Hz, 2H), 7.40 (d,  $J$  = 8.9 Hz, 2H), 6.81 (d,  $J$  = 8.9 Hz, 2H), 3.75 (s, 3H) ppm.  $^{13}\text{C}$  NMR (63 MHz,  $\text{CDCl}_3$ )  $\delta$  = 160.8, 147.0, 133.8, 132.4, 131.1, 124.0, 114.6, 95.6, 87.0, 78.0, 55.7 ppm. IR (thin film, ATR)  $\nu_{\max}$  2211, 1589, 1511, 1464, 1441, 1339, 1291, 1248, 1176, 1139, 1108, 1030, 857, 836, 749, 688  $\text{cm}^{-1}$ . MS (EI, 70 eV)  $m/z$  (%): 253(100,  $[\text{M}^+]$ ), 223(20), 207(20), 163(80), 152(15), 87(10).

#### **5-Chloro-2-(p-tolylethynyl)aniline<sup>34</sup> 3q**

The general procedure was followed. Off-white solid. (177 mg, yield 79%) Mp.: 135 - 137 °C (Ref.: not given),  $R_f$  (Hex.-EtOAc = 5:1) = 0.50;  $^1\text{H}$  NMR (250 MHz,  $\text{CDCl}_3$ )  $\delta$  = 7.28 (d,  $J$  = 8.0 Hz, 2H), 7.13 (d,  $J$  = 8.3 Hz, 1H), 7.00 (d,  $J$  = 8.1 Hz, 2H), 6.68 – 6.45 (m, 2H), 4.46 (s, 2H), 2.22 (s, 3H) ppm.  $^{13}\text{C}$  NMR (63 MHz,  $\text{CDCl}_3$ )  $\delta$  = 148.0, 138.8, 135.2, 133.1, 131.5, 129.3, 119.9, 118.7, 114.6, 107.3, 95.9, 84.2, 21.7 ppm. IR (thin film, ATR): 1614, 1558, 1486, 1426, 1255, 917, 857, 820, 798  $\text{cm}^{-1}$ . MS (EI, 70 eV)  $m/z$  (% relative intensity, ion): 243(32,  $[\text{M}^+]$ ), 241(100,  $[\text{M}^+]$ ), 204(23), 178(14), 164(6), 120(7), 105(15), 89(16), 76(6).

#### **2-((4-Chlorophenyl)ethynyl)-4-(trifluoromethyl)aniline 3r**

The general procedure was followed. Brown crystals (260 mg, yield 81%), Mp.: 61-63 °C  $R_f$ : 0.21 (in hexanes:EtOAc 5:1).  $^1\text{H}$  NMR (250 MHz,  $\text{CDCl}_3$ )  $\delta$  = 7.62 (s, 1H), 7.51 – 7.41 (m, 2H), 7.40 – 7.31 (m, 3H), 6.74 (d,  $J$  = 8.5 Hz, 1H), 4.57 (s, 2H) ppm.  $^{13}\text{C}$  NMR (63 MHz,  $\text{CDCl}_3$ )  $\delta$  = 150.6, 135.1, 133.1, 130.0 (q,  $J$  = 3.6 Hz), 129.2, 127.2 (q,  $J$  = 3.6 Hz), 124.7 (q,  $J$  = 270.9 Hz), 121.5, 120.3 (q,  $J$  = 33.1 Hz), 114.2, 107.5, 94.8, 85.8 ppm.  $^{19}\text{F}$  NMR (235 MHz,  $\text{CDCl}_3$ )  $\delta$  = -61.46 ppm. IR (thin film, ATR)  $\nu_{\max}$  1623, 1490, 1431, 1398, 1338, 1281, 1149, 1111, 1014, 908, 825, 724  $\text{cm}^{-1}$ . MS (EI, 70 eV)  $m/z$  (%): 295(100,  $[\text{M}^+]$ ), 276(10), 233(30), 190(20), 163(15), 138(10), 108(5), 75(10). HRMS calcd for  $\text{C}_{15}\text{H}_{10}\text{ClF}_3\text{N}$   $[\text{M}+\text{H}]^+$  296.0454 found 296.0458.

#### **3-((3-Bromophenyl)ethynyl)-2-methylquinolin-4-amine 3s**

The general procedure was followed. Light brown solid (211 mg, yield 63%). Mp.: 177 - 179 °C.  $R_f$  (EtOAc) = 0.53;  $^1\text{H}$  NMR (250 MHz,  $\text{DMSO}-d_6$ )  $\delta$  = 8.27 (d,  $J$  = 8.2 Hz, 1H), 7.98 (t,  $J$  = 1.6 Hz, 1H), 7.74 – 7.53 (m, 5H), 7.43 – 7.32 (m, 2H), 7.13 (s, 2H), 2.64



(s, 3H) ppm.  $^{13}\text{C}$  NMR (63 MHz,  $\text{DMSO}-d_6$ )  $\delta$  = 158.9, 152.5, 146.7, 133.3, 130.9, 130.5, 130.0, 129.9, 128.4, 125.5, 123.9, 122.5, 121.6, 116.0, 97.7, 95.0, 86.7, 24.5 ppm. MS (EI, 70 eV)  $m/z$  (% relative intensity, ion): 338(92,  $[\text{M}^+]$ ), 336(100,  $[\text{M}^+]$ ), 281(15), 256(55), 207(61), 193(20), 147(24), 128(41), 114(14), 94(13), 73(59). IR (thin film, ATR): 2200, 1642, 1614, 1570, 1501, 1475, 1450, 1381, 1068, 872, 777, 762, 684  $\text{cm}^{-1}$ . HRMS  $m/z$   $[\text{M}+\text{H}]^+$  calculated for  $\text{C}_{18}\text{H}_{14}\text{N}_2\text{Br}^+$ : 337.0340; found: 337.0339.

### 7-Chloro-4-(phenylethynyl)quinolone 3t

The general procedure was followed. Brown solid (154 mg, yield 70%) Mp.: 95-96 °C,  $R_f$ : 0.28 (in hexanes: EtOAc 10:1).  $^1\text{H}$  NMR (250 MHz,  $\text{CDCl}_3$ )  $\delta$  = 8.89 (d,  $J$  = 4.5 Hz, 1H), 8.29 (d,  $J$  = 8.9 Hz, 1H), 8.13 (d,  $J$  = 2.2 Hz, 1H), 7.74 – 7.50 (m, 4H), 7.49 – 7.32 (m, 3H) ppm.  $^{13}\text{C}$  NMR (63 MHz,  $\text{CDCl}_3$ )  $\delta$  = 151.0, 148.6, 136.4, 132.4, 130.5, 130.0, 129.0, 129.0, 128.6, 127.8, 126.6, 124.0, 122.3, 99.8, 84.9 ppm. IR (thin film, ATR)  $\nu_{\text{max}}$  2217, 1608, 1574, 1496, 1443, 1417, 1376, 1353, 1284, 1197, 1147, 1084, 1029, 897, 856, 825, 756, 689  $\text{cm}^{-1}$ . MS (EI, 70 eV)  $m/z$  (%): 263(100,  $[\text{M}^+]$ ), 227(35), 200(40), 131(10). HRMS calcd for  $\text{C}_{17}\text{H}_{11}\text{ClN}$   $[\text{M}+\text{H}]^+$  264.0580 found 264.0582.

### 1,4-bis(Phenylethynyl)benzene<sup>35</sup> 3u

The general procedure was modified. (1,4-iodobenzene (1 mmol, 330 mg), trimethylsilylacetylene (2.2 equiv., 2.2 mmol) and iodobenzene (2.0 equiv., 2 mmol) were used) Yellow crystals (228 mg, yield 82%) mp.: 175-176 °C (Ref.: 178-179 °C),  $R_f$ : 0.25 (in hexanes).  $^1\text{H}$  NMR (250 MHz,  $\text{CDCl}_3$ )  $\delta$  = 7.55 (dd,  $J$  = 6.7, 3.1 Hz, 4H), 7.53 (s, 4H), 7.37 (dd,  $J$  = 5.0, 1.7 Hz, 6H) ppm.  $^{13}\text{C}$  NMR (63 MHz,  $\text{CDCl}_3$ )  $\delta$  = 132.0, 131.9, 128.9, 128.8, 123.5, 123.5, 91.6, 89.5 ppm. IR (thin film, ATR)  $\nu_{\text{max}}$  1594, 1514, 1441, 1070, 839, 752, 690  $\text{cm}^{-1}$ . MS (EI, 70 eV)  $m/z$  (%): 278(100,  $[\text{M}^+]$ ), 276(15), 139(18).

### 2-(Naphthalen-2-ylethynyl)-9H-fluorene 3v

The general procedure was followed. Orange solid (283 mg, yield 45%),  $R_f$ : 0.20 (in hexanes: EtOAc 100:1). Mp.: 137-139 °C  $^1\text{H}$  NMR (250 MHz,  $\text{CDCl}_3$ )  $\delta$  = 8.40 (d,  $J$  = 8.2 Hz, 1H), 7.93 – 6.94 (m, 13H), 3.82 (d,  $J$  = 7.6 Hz, 2H) ppm.  $^{13}\text{C}$  NMR (63 MHz,  $\text{CDCl}_3$ )  $\delta$  = 144.0, 143.7, 142.4, 141.5, 133.7, 133.6, 131.0, 130.7, 129.0, 128.7, 128.6, 127.6, 127.4, 127.2, 126.8, 126.7, 125.7, 125.5, 121.8, 121.5, 120.6, 120.3, 95.6, 88.0, 37.2 ppm. IR (thin film, ATR)  $\nu_{\text{max}}$  2964, 2897, 2829, 1585, 1453, 1396, 905, 832, 798, 768, 732  $\text{cm}^{-1}$ . MS (EI, 70 eV)  $m/z$  (%): 316 (100,  $[\text{M}^+]$ ), 156(35), 143(10), 73(5). HRMS can not be obtained by our facility.

### Triethyl((3-(phenylethynyl)benzyl)oxy)silane 3w

The general procedure was followed. Yellow oil (131 mg, yield 41%),  $R_f$ : 0.25 (in hexanes: EtOAc 100:1).  $^1\text{H}$  NMR (250 MHz,  $\text{CDCl}_3$ )  $\delta$  = 7.57 – 7.38 (m, 3H), 7.37 – 7.18 (m, 6H), 4.64 (s, 2H), 0.90 (t,  $J$  = 7.8 Hz, 9H), 0.58 (q,  $J$  = 7.9 Hz, 6H) ppm.  $^{13}\text{C}$  NMR (63 MHz,  $\text{CDCl}_3$ )  $\delta$  = 142.0, 132.0, 130.6, 129.7, 128.7, 128.7, 128.6, 126.6, 123.7, 123.5, 90.0, 89.6, 64.7, 7.2, 4.9 ppm. IR (thin film, ATR)  $\nu_{\text{max}}$  2958, 2912, 2878, 1604, 1493, 1459, 1413, 1368, 1239, 1201, 1101, 1078, 1005, 974, 915, 889, 817, 784, 754, 728, 689  $\text{cm}^{-1}$ . MS (EI, 70 eV)  $m/z$  (%): 322(15), 293(95), 263(33), 235(20), 206(20), 191(100,  $[\text{M}^+]$ ), 165(30), 95(15), 59(10). HRMS calcd for  $\text{C}_{21}\text{H}_{26}\text{OSiNa}$   $[\text{M}+\text{Na}]^+$  345.1651 found 345.1650.

### tert-Butyldiphenyl((3-(phenylethynyl)benzyl)oxy)silane 3x

The general procedure was followed. Yellow oil (382 mg, yield 86%),  $R_f$ : 0.27 (in hexanes: EtOAc 100:1).  $^1\text{H}$  NMR (250 MHz,  $\text{CDCl}_3$ )  $\delta$  = 7.68 – 7.57 (m, 4H), 7.54 – 7.15 (m, 15H), 4.67 (s, 2H), 1.02 (s, 9H) ppm.  $^{13}\text{C}$  NMR (63 MHz,  $\text{CDCl}_3$ )  $\delta$  = 141.7, 136.0, 133.7, 132.1, 130.6, 130.2, 129.6, 128.8, 128.7, 128.7, 128.2, 126.5, 123.7, 123.5, 90.0, 89.6, 65.6, 27.3, 19.8 ppm. IR (thin film, ATR)  $\nu_{\text{max}}$  2960, 2933, 2856, 1603, 1493, 1471, 1428, 1364, 1201, 1106, 1077, 1028, 999, 918, 888, 823, 784, 756, 688  $\text{cm}^{-1}$ . MS (EI, 70 eV)  $m/z$  (%): 389(90), 311(83), 189(100,  $[\text{M}^+]$ ), 165(45), 105(15), 77(10). HRMS calcd for  $\text{C}_{31}\text{H}_{31}\text{OSi}$   $[\text{M}+\text{H}]^+$  447.2144 found 447.2143.

### tert-Butyldimethyl((3-(phenylethynyl)benzyl)oxy)silane 3y

The general procedure was followed. Yellow oil (261 mg, yield 82%),  $R_f$ : 0.25 (in hexanes: EtOAc 100:1).  $^1\text{H}$  NMR (250 MHz,  $\text{CDCl}_3$ )  $\delta$  = 7.50 – 7.34 (m, 3H), 7.33 – 7.14 (m, 6H), 4.62 (s, 2H), 0.84 (s, 9H), 0.00 (s, 6H) ppm.  $^{13}\text{C}$  NMR (63 MHz,  $\text{CDCl}_3$ )  $\delta$  = 142.1, 132.0, 130.6, 129.5, 128.7, 128.7, 128.6, 126.5, 123.7, 123.5, 89.9, 89.5, 65.0, 26.4, 18.9, -4.8 ppm. IR (thin film, ATR)  $\nu_{\text{max}}$  2958, 2934, 2883, 2857, 1604, 1493, 1463, 1363, 1254, 1201, 1102, 1079, 1027, 1006, 917, 889, 835, 776, 754, 688  $\text{cm}^{-1}$ . MS (EI, 70 eV)  $m/z$  (%): 322(2), 265(100,  $[\text{M}^+]$ ), 235(55), 191(90), 165(20), 75(5). HRMS calcd for  $\text{C}_{21}\text{H}_{26}\text{OSiNa}$   $[\text{M}+\text{Na}]^+$  345.1651 found 345.1656.

### Triisopropyl((3-(phenylethynyl)benzyl)oxy)silane 3z

The general procedure was followed. Colorless oil (281 mg, yield 77%),  $R_f$ : 0.23 (in hexanes: EtOAc 100:1).  $^1\text{H}$  NMR (250 MHz,  $\text{CDCl}_3$ )  $\delta$  = 7.73 – 7.50 (m, 3H), 7.48 – 7.31 (m, 6H), 4.86 (s, 2H), 1.13 (app. s, 21H) ppm.  $^{13}\text{C}$  NMR (63 MHz,  $\text{CDCl}_3$ )  $\delta$  = 142.3, 132.0, 130.5, 129.3, 128.7, 128.7, 128.6, 126.2, 123.8, 123.4, 90.0, 89.5, 65.0, 18.5, 12.4 ppm. IR (thin film, ATR)  $\nu_{\text{max}}$  2942, 2894, 2886, 1604, 1493, 1462, 1367, 1249, 1201, 1105, 1068, 995, 918, 882, 843, 783, 754, 688  $\text{cm}^{-1}$ . MS (EI, 70 eV)  $m/z$  (%): 364(2), 321(100,  $[\text{M}^+]$ ), 293(10), 249(15), 206(25), 191(90), 165(24), 133(10), 95(10), 59(10). HRMS calcd for  $\text{C}_{24}\text{H}_{33}\text{OSi}$   $[\text{M}+\text{H}]^+$  365.2301 found 365.2299.

### 5-(Pyridin-2-ylethynyl)-3-((triisopropylsilyl)ethynyl)-1H-indole 3aa

The general procedure was followed. Green solid (284 mg, yield 71%). Mp.: 168-170 °C,  $R_f$ : 0.21 (in hexanes: EtOAc 2:1).  $^1\text{H}$  NMR (250 MHz,  $\text{CDCl}_3$ )  $\delta$  = 9.34 (s, 1H), 8.72 (s, 1H), 8.43 (dd,  $J$  = 4.8, 1.4 Hz, 1H), 7.85 (s, 1H), 7.78 (dt,  $J$  = 7.9, 1.8 Hz, 1H), 7.37 (d,  $J$  = 2.6 Hz, 1H), 7.34 – 7.18 (m, 3H), 1.09 (app. s, 21H) ppm.  $^{13}\text{C}$  NMR (63 MHz,  $\text{CDCl}_3$ )  $\delta$  = 152.2, 148.0, 139.1, 135.6, 130.1, 129.2, 126.9, 124.4, 123.7, 121.8, 114.7, 112.2, 100.0, 99.8, 95.0, 92.8, 84.4, 19.2, 11.8 ppm. IR (thin film, ATR)  $\nu_{\text{max}}$  2944, 2865, 2725, 2211, 2150, 1589, 1484, 1408, 1344, 1268, 1237, 1189, 1142, 1073, 1042, 995, 926, 883, 848, 801, 699, 675  $\text{cm}^{-1}$ . MS

(EI, 70 eV)  $m/z$  (%): 398(32), 355(100, [M<sup>+</sup>]), 313(43), 285(50), 149(37), 135(27), 59(10). HRMS calcd for C<sub>26</sub>H<sub>31</sub>N<sub>2</sub>Si [M+H]<sup>+</sup> 399.2257 found 399.2249.

#### 2-(4-Methoxyphenyl)benzofuran<sup>5</sup> 4a

The general procedure was followed. White crystals (0.177 g, 79%). Mp.= 140 – 141 °C (Ref.: 143 – 144 °C), R<sub>f</sub> (Hex.-EtOAc = 5:1) = 0.60; <sup>1</sup>H NMR (250 MHz, CDCl<sub>3</sub>) δ = 7.70 (td, *J* = 10, 2.5 Hz, 2H), 7.49 – 7.37 (m, 3H), 7.21 – 7.06 (m, 2H), 6.88 (td, *J* = 32, 2.0 Hz, 2H), 6.78 (d, *J* = 0.8 Hz, 1H), 3.75 (s, 3H) ppm. <sup>13</sup>C NMR (63 MHz, CDCl<sub>3</sub>) δ = 160.1, 156.2, 154.8, 129.6, 126.5, 123.9, 123.5, 123.0, 120.7, 114.4, 111.1, 99.8, 55.5 ppm. MS (EI, 70 eV)  $m/z$  (% relative intensity, ion): 224(100, [M<sup>+</sup>]), 209(78), 181(53), 152(37), 126(8), 112(10), 63(7). IR (thin film, ATR): 1611, 1506, 1456, 1299, 1253, 1184, 1113, 1040, 1025, 921, 837, 802, 744 cm<sup>-1</sup>.

#### Ethyl 4-(benzofuran-2-yl)benzoate<sup>18</sup> 4b

The general procedure was followed. Light yellow solid (0.067 g, 26%). Mp.= 109 - 111 °C (Ref.: 109 - 111 °C), R<sub>f</sub> (Hex.-EtOAc = 5:1) = 0.55; <sup>1</sup>H NMR (250 MHz, CDCl<sub>3</sub>) δ = 7.98 (dt, *J* = 8.5, 1.6 Hz, 2H), 7.77 (dt, *J* = 8.5, 1.6 Hz, 2H), 7.46 (dd, *J* = 7.3, 1.5 Hz, 1H), 7.40 (d, *J* = 8.2 Hz, 1H), 7.23 – 7.05 (m, 2H), 6.99 (s, 1H), 4.26 (q, *J* = 7.1 Hz, 2H), 1.28 (t, *J* = 7.1 Hz, 3H) ppm. <sup>13</sup>C NMR (63 MHz, CDCl<sub>3</sub>) δ = 166.3, 155.3, 154.8, 134.5, 130.2, 130.2, 129.0, 125.1, 124.7, 123.3, 121.4, 111.5, 103.5, 61.2, 14.5. MS (EI, 70 eV)  $m/z$  (% relative intensity, ion): 266(100, [M<sup>+</sup>]), 238 (55), 221(63), 193(13), 165(62), 163(22), 139(11), 110(10), 82(13). IR (thin film, ATR): 1708, 1614, 1454, 1411, 1368, 1277, 1184, 1102, 1035, 1016, 921, 856, 807, 772, 746, 697 cm<sup>-1</sup>.

#### 2-(4-Nitrophenyl)benzofuran<sup>36</sup> 4c

The general procedure was followed. Orange solid (0.148 g, 62%). Mp.= 169 – 171 °C (Ref.: 181 - 183 °C), R<sub>f</sub> (Hex.-EtOAc = 5:1) = 0.55; <sup>1</sup>H NMR (250 MHz, CDCl<sub>3</sub>) δ = 8.16 (d, *J* = 8.6 Hz, 2H), 7.84 (d, *J* = 8.6 Hz, 2H), 7.50 (d, *J* = 7.5 Hz, 1H), 7.41 (d, *J* = 8.0 Hz, 1H), 7.23 (t, *J* = 7.5 Hz, 1H), 7.14 (t, *J* = 7.5 Hz, 1H), 7.09 (s, 1H) ppm. <sup>13</sup>C NMR (62.5 MHz, CDCl<sub>3</sub>) δ = 155.5, 153.4, 147.3, 136.4, 128.8, 126.0, 125.3, 124.4, 123.7, 121.8, 111.6, 105.2 ppm. MS (EI, 70 eV)  $m/z$  (% relative intensity, ion): 239(100, [M<sup>+</sup>]), 209(25), 181(25), 165(98), 163(35), 152(11), 115(10), 63(10). IR (thin film, ATR): 1601, 1573, 1519, 1452, 1346, 1113, 1031, 852, 809, 751, 693 cm<sup>-1</sup>.

#### Methyl 2-(1*H*-indol-5-yl)benzofuran-6-carboxylate 4d

The general procedure was followed. Light brown solid (0.271 g, 93%). Mp.= 134 - 136 °C, R<sub>f</sub> (Hex.-EtOAc = 5:1) = 0.23; δ = <sup>1</sup>H NMR (250 MHz, DMSO-*d*<sub>6</sub>) δ = 11.30 (s, 1H), 8.12 (s, 1H), 8.05 (s, 1H), 7.78 (dd, *J* = 8.2, 1.3 Hz, 1H), 7.63 (m, 2H), 7.45 (d, *J* = 8.5 Hz, 1H), 7.37 (m, 1H), 7.29 (s, 1H), 6.49 (s, 1H), 3.80 (s, 3H) ppm. <sup>13</sup>C NMR (63 MHz, DMSO-*d*<sub>6</sub>) δ = 166.4, 160.7, 153.4, 136.6, 134.1, 127.9, 126.8, 124.6, 124.1, 120.3, 120.2, 118.8, 117.5, 112.2, 111.6, 102.0, 99.6, 52.1 ppm. MS (EI, 70 eV)  $m/z$  (% relative intensity, ion): 291(37, [M<sup>+</sup>]), IR (thin film, ATR): 1596, 1577, 1497, 1473, 1447, 1413, 1340, 1281, 1221, 1197, 1145, 1074, 984, 956, 923, 874, 796, 761 cm<sup>-1</sup>. HRMS  $m/z$  [M+H]<sup>+</sup> calculated for C<sub>18</sub>H<sub>14</sub>NO<sub>3</sub><sup>+</sup>: 292.0974; found: 292.0975.

#### 6-Chloro-2-(thiophen-2-yl)benzofuran 4e

The general procedure was followed. White powder (0.159 g, 68%). Mp.= 142 - 143 °C, R<sub>f</sub> (Hex.-EtOAc = 5:1) = 0.68; <sup>1</sup>H NMR (250 MHz, CDCl<sub>3</sub>) δ = 7.49 (m, 2H), 7.43 – 7.33 (m, 2H), 7.22 (dd, *J* = 8.7, 2.0 Hz, 1H), 7.11 (t, *J* = 1 Hz, 1H), 6.79 (s, 1H) ppm. <sup>13</sup>C NMR (63 MHz, CDCl<sub>3</sub>) δ = 153.3, 153.1, 133.1, 130.9, 129.1, 128.4, 126.8, 125.5, 124.8, 120.6, 112.4, 100.9 ppm. MS (EI, 70 eV)  $m/z$  (% relative intensity, ion): 236(37, [M<sup>+</sup>]), 234(100, [M<sup>+</sup>]), 205(6), 171(44), 145(6), 126(7), 117(8), 85(8). IR (thin film, ATR): 1586, 1448, 1264, 1200, 1064, 999, 880, 854, 796, 740, 708 cm<sup>-1</sup>. HRMS  $m/z$  [M]<sup>+</sup> calculated for C<sub>12</sub>H<sub>7</sub>ClOS<sup>+</sup>: 233.9901; found: 233.9903.

#### 7-Chloro-4-(5-fluorobenzofuran-2-yl)quinoline 4f

The general procedure was followed. Yellow solid (0.105 g, 36%). Mp.= 208 - 210 °C, R<sub>f</sub> (Hex.-EtOAc = 1:1) = 0.55; <sup>1</sup>H NMR (250 MHz, CDCl<sub>3</sub>) δ = 8.97 (d, *J* = 4.6 Hz, 1H), 8.47 (d, *J* = 9.1 Hz, 1H), 8.20 (d, *J* = 2.1 Hz, 1H), 7.74 (d, *J* = 4.6 Hz, 1H), 7.63 – 7.49 (m, 2H), 7.34 (dd, *J* = 8.3, 2.6 Hz, 1H), 7.26 (s, 1H), 7.13 (td, *J* = 9.0, 2.6 Hz, 1H) ppm. <sup>13</sup>C NMR (63 MHz, CDCl<sub>3</sub>) δ = 161.8(d, *J* = 242 Hz), 158.0, 154.4, 152.0, 150.9, 149.4, 136.3, 136.1, 129.4 (d, *J* = 10.9 Hz), 129.2, 128.0 (d, *J* = 110.4 Hz), 123.4, 120.2, 114.2 (d, *J* = 26.6 Hz), 112.7 (d, *J* = 9.6 Hz), 109.3 (d, *J* = 3.8 Hz), 107.6, 107.2 ppm. <sup>19</sup>F NMR (235 MHz, CDCl<sub>3</sub>) δ -119.5. MS (EI, 70 eV)  $m/z$  (% relative intensity, ion): 299(37, [M<sup>+</sup>]), 297(100, [M<sup>+</sup>]), 270(6), 262(20), 234(13), 207(17), 148(4), 135(6), 117(6), 103(8). IR (thin film, ATR): 1700, 1614, 1577, 1435, 1297, 1230, 1204, 1122, 1085, 887, 789, 766, 740 cm<sup>-1</sup>. HRMS  $m/z$  [M+H]<sup>+</sup> calculated for C<sub>17</sub>H<sub>10</sub>NOCIF<sup>+</sup>: 298.0435; found: 298.0428.

#### 5-(1-(4-Bromobenzyl)-1*H*-1,2,3-triazol-4-yl)-1*H*-indole 5a

The general procedure was followed. Pale yellow solid (0.165 g, 47%). Mp.= 173 - 176 °C, R<sub>f</sub> (Hex.-EtOAc = 1:1) = 0.30; <sup>1</sup>H NMR (250 MHz, DMSO-*d*<sub>6</sub>) δ = 11.18 (s, 1H), 8.51 (s, 1H), 8.03 (s, 1H), 7.60 (d, *J* = 8.4 Hz, 3H), 7.44 (d, *J* = 8.4 Hz, 1H), 7.38 – 7.28 (m, 3H), 6.47 (s, 1H), 5.62 (s, 2H) ppm. <sup>13</sup>C NMR (63 MHz, DMSO-*d*<sub>6</sub>) δ = 148.3, 135.7, 135.6, 131.7, 130.2, 127.9, 126.1, 121.6, 121.4, 120.3, 119.0, 116.9, 111.7, 101.4, 52.2 ppm. MS (EI, 70 eV)  $m/z$  (% relative intensity, ion): 354(9, [M<sup>+</sup>]), 352(10, [M<sup>+</sup>]), 325(33), 281(7), 245(15), 218(16), 207(25), 171(18), 169(39), 155(100), 141 (10), 128(44), 122(11), 101(18), 90(24), 77(11), 73(24), 63(10). IR (thin film, ATR): 1491, 1439, 1417, 1350, 1325, 1219, 1070, 1014, 884, 816, 800, 766, 738 cm<sup>-1</sup>. HRMS  $m/z$  [M+H]<sup>+</sup> calculated for C<sub>17</sub>H<sub>14</sub>BrN<sub>4</sub><sup>+</sup>: 353.0402; found: 353.0399.

#### 4-(3-Bromophenyl)-1-(4-nitrobenzyl)-1*H*-1,2,3-triazole<sup>18</sup> 5b

The general procedure was followed. Yellow powder (0.138 g, 39%). Mp.= 88 - 90 °C (Ref.: 90 – 93 °C), R<sub>f</sub> (Hex.-EtOAc = 1:1) = 0.43; <sup>1</sup>H NMR (250 MHz, DMSO-*d*<sub>6</sub>) δ = 8.79 (s, 1H), 8.25 (d, *J* = 8.6 Hz, 2H), 8.05 (s, 1H), 7.88 (d, *J* = 7.7 Hz, 1H), 7.58 (d, *J* = 8.6 Hz, 2H), 7.53 (d, *J* = 8.9 Hz, 3H), 7.41 (t, *J* = 7.8 Hz, 1H), 5.85 (s, 2H) ppm. <sup>13</sup>C NMR (63 MHz, DMSO-*d*<sub>6</sub>) δ = 147.3, 145.3,

143.2, 132.8, 131.1, 130.7, 129.1, 127.7, 124.1, 124.0, 122.7, 122.3, 52.2 ppm. MS (EI, 70 eV)  $m/z$  (% relative intensity, ion): 360(5,  $[M^+]$ ), 358(6,  $[M^+]$ ), 281(7), 207(24), 196(98), 194(100), 169(13), 167(14), 147(10), 136(15), 121(36), 120(28), 115(26), 106(18), 102(12), 90(30), 89(44), 88(24), 78(40), 73(30), 63(17), 51(10). IR (thin film, ATR): 1607, 1568, 1519, 1346, 1230, 1072, 1048, 859, 783, 746, 688  $cm^{-1}$ .

#### 4-(1-(4-Bromobenzyl)-1H-1,2,3-triazol-4-yl)-7-chloroquinoline 5c

The general procedure was followed. Brown solid (0.264 g, 66%). Mp. = 128 - 130 °C,  $R_f$  (EtOAc) = 0.55;  $^1H$  NMR (250 MHz, DMSO- $d_6$ )  $\delta$  = 9.00 (s, 1H), 8.97 (d,  $J$  = 4.6 Hz, 1H), 8.86 (d,  $J$  = 9.1 Hz, 1H), 8.13 (d,  $J$  = 2.2 Hz, 1H), 7.83 (d,  $J$  = 4.6 Hz, 1H), 7.71 (dd,  $J$  = 9.1, 2.3 Hz, 1H), 7.61 (d,  $J$  = 8.4 Hz, 2H), 7.39 (d,  $J$  = 8.4 Hz, 2H), 5.75 (s, 2H) ppm.  $^{13}C$  NMR (63 MHz, DMSO- $d_6$ )  $\delta$  = 151.6, 149.0, 143.8, 135.6, 135.1, 134.2, 131.8, 130.4, 128.2, 128.1, 127.7, 125.9, 123.2, 121.6, 120.4, 52.5 ppm. MS (EI, 70 eV)  $m/z$  (% relative intensity, ion): 400(10,  $[M^+]$ ), 398(7,  $[M^+]$ ), 371(9), 369(7), 291(10), 281(8), 203(29), 201(100), 182(11), 171(69), 169(80), 166(17), 147(17), 90(54), 89(39), 75(10), 73(24). IR (thin film, ATR): 1590, 1491, 1413, 1338, 1299, 1230, 1120, 1074, 1050, 1014, 954, 880, 738, 697  $cm^{-1}$ . HRMS  $m/z$   $[M+H]^+$  calculated for  $C_{17}H_{14}N_4Br^+$ : 353.0402; found: 353.0397.

## REFERENCES

- (1) (a) Fang, Y.; Zhou, Y.; Rui, Q.; Yao, C., Rhodamine–Ferrocene Conjugate Chemosensor for Selectively Sensing Copper(II) with Multisignals: Chromaticity, Fluorescence, and Electrochemistry and Its Application in Living Cell Imaging, *Organometallics* **2015**, *34*, 2962; (b) Grzybowski, M.; Glodkowska-Mrowka, E.; Hugues, V.; Brutkowski, W.; Blanchard-Desce, M.; Gryko, D. T., Polar Diketopyrrolopyrrole-Imidazolium Salts as Selective Probes for Staining Mitochondria in Two-Photon Fluorescence Microscopy, *Chem. Eur. J.* **2015**, *21*, 9101; (c) Bunz, U. H. F., The Larger Linear N-Heteroacenes, *Acc. Chem. Res.* **2015**, *48*, 1676; (d) Yin, H.; Zhang, B.; Yu, H.; Zhu, L.; Feng, Y.; Zhu, M.; Guo, Q.; Meng, X., Two-Photon Fluorescent Probes for Biological  $Mg^{2+}$  Detection Based on 7-Substituted Coumarin, *J. Org. Chem.* **2015**, *80*, 4306; (e) Stang, P. J.; Diederich, F., *Modern Acetylene Chemistry*; Wiley, 2008; (f) Lavastre, O.; Cabioch, S.; Dixneuf, P. H.; Sedlacek, J.; Vohlidal, J., New route to conjugated polymer networks: synthesis of poly(4-ethynyl)phenylacetylene and its transformation into a conjugated network, *Macromolecules* **1999**, *32*, 4477; (g) Lavastre, O.; Cabioch, S.; Dixneuf, P. H.; Vohlidal, J., Selective and efficient access to ortho, meta and para ring-substituted phenylacetylene derivatives  $R-[C\equiv C-C_6H_4](x)-Y$  ( $Y$ : H, NO<sub>2</sub>, CN, I, NH<sub>2</sub>), *Tetrahedron* **1997**, *53*, 7595; (h) Lavastre, O.; Ollivier, L.; Dixneuf, P. H.; Sibandhit, S., Sequential catalytic synthesis of rod-like conjugated poly-ynes, *Tetrahedron* **1996**, *52*, 5495; (i) Rawson, J.; Stuart, A. C.; You, W.; Therien, M. J., Tailoring Porphyrin-Based Electron Accepting Materials for Organic Photovoltaics, *J. Am. Chem. Soc.* **2014**, *136*, 17561; (j) Chen, J.; Wenger, O. S., Fluoride binding to an organoboron wire controls photoinduced electron transfer, *Chem. Sci.* **2015**, *6*, 3582.
- (2) (a) Dordević, L.; Marangoni, T.; Miletic, T.; Rubio-Magnieto, J.; Mohanraj, J.; Amenitsch, H.; Pasini, D.; Liaros, N.; Couris, S.; Armaroli, N.; Surin, M.; Bonifazi, D., Solvent Molding of Organic Morphologies Made of Supramolecular Chiral Polymers, *J. Am. Chem. Soc.* **2015**, *137*, 8150; (b) Miyagawa, M.; Yamaguchi, M., Helicene-Grafted Silica Nanoparticles Capture Hetero-Double-Helix Intermediates during Self-Assembly Gelation, *Chem. Eur. J.* **2015**, *21*, 8408; (c) Lee, H.; Elumalai, P.; Singh, N.; Kim, H.; Lee, S. U.; Chi, K.-W., Selective Synthesis of Ruthenium(II) Metalla[2]Catenane via Solvent and Guest-Dependent Self-Assembly, *J. Am. Chem. Soc.* **2015**, *137*, 4674; (d) Danjo, H.; Hashimoto, Y.; Kidena, Y.; Nogamine, A.; Katagiri, K.; Kawahata, M.; Miyazawa, T.; Yamaguchi, K., Nestable Tetakis(spiroborate) Nanocycles, *Org. Lett.* **2015**, *17*, 2154; (e) Wang, Q.; Yu, C.; Long, H.; Du, Y.; Jin, Y.; Zhang, W., Solution-Phase Dynamic Assembly of Permanently Interlocked Aryleneethynylene Cages through Alkyne Metathesis, *Angew. Chem., Int. Ed.* **2015**, *54*, 7550; (f) Zheng, Q.-N.; Liu, X.-H.; Chen, T.; Yan, H.-J.; Cook, T.; Wang, D.; Stang, P. J.; Wan, L.-J., Formation of Halogen Bond-Based 2D Supramolecular Assemblies by Electric Manipulation, *J. Am. Chem. Soc.* **2015**, *137*, 6128; (g) Nishioka, S.; Morita, S.; Okada, K.; Suzuki, S.; Kozaki, M., Synthesis and Higher-Order Structure of Linear Dendrimeric Assemblies, *Org. Lett.* **2015**, *17*, 2720; (h) Baba, A.; Kojima, T.; Hiraoka, S., Self-Assembly Process of Dodecanuclear Pt(II)-Linked Cyclic Hexagon, *J. Am. Chem. Soc.* **2015**, *137*, 7664; (i) Kaur, I.; Zhao, X.; Bryce, M. R.; Schauer, P. A.; Low, P. J.; Katak, R., Modification of Electrode Surfaces by Self-Assembled Monolayers of Thiol-Terminated Oligo(Phenyleneethynylene)s, *Chem. Phys. Chem.* **2013**, *14*, 431.
- (3) Brandsma, L.; Heus-Kloos, Y. A.; van der Heiden, R.; Verkruisje, H. D. *Preparative acetylenic chemistry*; Elsevier, 1988.
- (4) (a) Sonogashira, K.; Tohda, Y.; Hagihara, N., A convenient synthesis of acetylenes: catalytic substitutions of acetylenic hydrogen with bromoalkenes, iodoarenes and bromopyridines, *Tetrahedron Lett.* **1975**, *16*, 4467; (b) Chinchilla, R.; Nájera, C., The Sonogashira reaction: A booming methodology in synthetic organic chemistry, *Chem. Rev.* **2007**, *107*, 874; (c) Chinchilla, R.; Nájera, C., Recent advances in Sonogashira reactions, *Chem. Soc. Rev.* **2011**, *40*, 5084; (d) Schilz, M.; Plenio, H., A Guide to Sonogashira Cross-Coupling Reactions: The Influence of Substituents in Aryl Bromides, Acetylenes, and Phosphines, *J. Org. Chem.* **2012**, *77*, 2798; (e) Doucet, H.; Hierso, J. C., Palladium-Based Catalytic Systems for the Synthesis of Conjugated Enynes by Sonogashira Reactions and Related Alkynylations, *Angew. Chem., Int. Ed.* **2007**, *46*, 834.
- (5) Csékei, M.; Novák, Z.; Kotschy, A., Development of a one-pot sequential Sonogashira coupling for the synthesis of benzofurans, *Tetrahedron* **2008**, *64*, 8992.
- (6) (a) Novák, Z.; Nemes, P.; Kotschy, A., Tandem Sonogashira Coupling: An Efficient Tool for the Synthesis of Diarylalkynes, *Org. Lett.* **2004**, *6*, 4917; (b) Katritzky, A. R.; Rogovoy, B. V.; Mitrokhin, A. Y., The preparation of diarylacetylenes via diphenyl (benzotriazol-1-yl)(aryl) methylphosphonates, *Arkivoc* **2002**, *13*, 17; (c) Csékei, M.; Novák, Z.; Kotschy, A., Ethynyl-cyclohexanol: an efficient acetylene surrogate in Sonogashira coupling, *Tetrahedron* **2008**, *64*, 975; (d) Ma, L.; Hu, Q. S.; Pu, L., Synthesis of an optically active poly(aryleneethynylene) containing extended conjugation in the repeat unit, *Tetrahedron Asymmetry* **1996**, *7*, 3103; (e) Melissaris, A. P.; Litt, M. H., A simple and economical synthetic route to p-ethynylaniline and ethynyl-terminated substrates, *J. Org. Chem.* **1994**, *59*, 5818; (f) Carpi, A.; Lessi, A.; Rossi, R., One-Pot Palladium-Catalysed Synthesis of Diarylalkynes, *Synthesis* **1984**, *1984*, 571.
- (7) (a) Vokatá, T.; Kumar, M. R.; Park, K.; Moon, J. H.; Lee, S., Synthesis of Poly(phenylenebutadiynylene)s Using the Decarboxylative Coupling of Propiolic Acid and Aryl Iodides, *Synlett* **2013**, *24*, 1563; (b) Moon, J.; Jeong, M.; Nam, H.; Ju, J.; Moon, J. H.; Jung, H. M.; Lee, S., One-Pot Synthesis of Diarylalkynes Using Palladium-Catalyzed Sonogashira Reaction and Decarboxylative Coupling of sp Carbon and sp<sup>2</sup> Carbon, *Org. Lett.* **2008**, *10*, 945; (c) Park, K.; Kim, W.; Lee, S., Efficient one-pot synthesis of the unsymmetrical Diarylalkynes from two different Aryl Bromides and Propiolic acid by using Pd(PPh<sub>3</sub>)<sub>4</sub> Catalyst, *Bull. Korean Chem. Soc.* **2013**, *34*, 2859.
- (8) Severin, R.; Reimer, J.; Doye, S., One-pot procedure for the synthesis of unsymmetrical diarylalkynes, *J. Org. Chem.* **2010**, *75*, 3518.
- (9) (a) Lindström, S.; Ripa, L.; Hallberg, A., Synthesis of two conformationally constrained analogues of the minor tobacco alkaloid anabasine, *Org. Lett.* **2000**, *2*, 2291; (b) Palimkar, S. S.; More, V. S.; Kumar, P. H.; Srinivasan, K. V., Synthesis of an indole containing KDR kinase inhibitor by tandem Sonogashira coupling-5-endo-dig-cyclization as a key step, *Tetrahedron* **2007**, *63*, 12786; (c) Arcadi, A.; Cacchi, S.; Di Giuseppe, S.; Fabrizi, G.; Marinelli, F., New efficient approaches to functionalized 2-substituted furopyridines, *Synlett* **2002**, 453.

- (10) (a) Buendia, J.; Darses, B.; Dauban, P., Tandem Catalytic C(sp<sup>3</sup>)–H Amination/Sila-Sonogashira–Hagihara Coupling Reactions with Iodine Reagents, *Angew. Chem., Int. Ed.* **2015**, *54*, 5697; (b) Nishihara, Y.; Inoue, E.; Noyori, S.; Ogawa, D.; Okada, Y.; Iwasaki, M.; Takagi, K., Synthesis of unsymmetrically disubstituted ethynes by the palladium/copper(I)-cocatalyzed sila-Sonogashira–Hagihara coupling reactions of alkynylsilanes with aryl iodides, bromides, and chlorides through a direct activation of a carbon–silicon bond, *Tetrahedron* **2012**, *68*, 4869; (c) Halbes, U.; Pale, P., A new mild procedure for the direct coupling of 1-trimethylsilyl acetylenes with vinyl triflates or aryl iodide, *Tetrahedron Lett.* **2002**, *43*, 2039; (d) Sorensen, U. S.; Pombo-Villar, E., Copper-free palladium-catalyzed sonogashira-type coupling of aryl halides and 1-aryl-2-(trimethylsilyl)acetylenes, *Tetrahedron* **2005**, *61*, 2697; (e) Nishihara, Y.; Inoue, E.; Okada, Y.; Takagi, K., Sila-sonogashira cross-coupling reactions of activated aryl chlorides with alkynylsilanes, *Synlett* **2008**, 3041; (f) Nishihara, Y.; Inoue, E.; Noyori, S.; Ogawa, D.; Okada, Y.; Iwasaki, M.; Takagi, K., Synthesis of unsymmetrically disubstituted ethynes by the palladium/copper(I)-cocatalyzed sila-Sonogashira–Hagihara coupling reactions of alkynylsilanes with aryl iodides, bromides, and chlorides through a direct activation of a carbon–silicon bond, *Tetrahedron* **2012**, *68*, 4869; (g) Nishihara, Y.; Inoue, E.; Ogawa, D.; Okada, Y.; Noyori, S.; Takagi, K., Palladium/copper-catalyzed sila-Sonogashira reactions of aryl iodides with alkynylsilanes via a direct C–Si bond activation, *Tetrahedron Lett.* **2009**, *50*, 4643; (h) Nishihara, Y.; Ikegashira, K.; Hirabayashi, K.; Ando, J.; Mori, A.; Hiyama, T., Coupling Reactions of Alkynylsilanes Mediated by a Cu(I) Salt: Novel Syntheses of Conjugate Diynes and Disubstituted Ethynes, *J. Org. Chem.* **2000**, *65*, 1780; (i) Carpita, A.; Mannocci, L.; Rossi, R., Silver(I)-catalysed protodesilylation of 1-(Trimethylsilyl)-1-alkynes, *Eur. J. Org. Chem.* **2005**, 1859; (j) Carpita, A.; Lessi, A.; Rossi, R., One-Pot Palladium-Catalysed Synthesis of Diarylalkynes, *Synthesis* **1984**.
- (11) Bhattacharya, P.; Basak, A., An unexpected one step domino conversion of TMS-alkynes to protected ketones in 4-chromenone system, *Tetrahedron Lett.* **2013**, *54*, 5137.
- (12) Wong, F. F.; Chuang, S. H.; Yang, S. c.; Lin, Y. H.; Tseng, W. C.; Lin, S. K.; Huang, J. J., One-pot ethynylation and catalytic desilylation in synthesis of mestranol and levonorgestrel, *Tetrahedron* **2010**, *66*, 4068.
- (13) DiLauro, A. M.; Seo, W.; Phillips, S. T., Use of Catalytic Fluoride under Neutral Conditions for Cleaving Silicon–Oxygen Bonds, *J. Org. Chem.* **2011**, *76*, 7352.
- (14) Yao, W.; Li, R.; Jiang, H.; Han, D., An Additive-Free, Base-Catalyzed Protodesilylation of Organosilanes, *J. Org. Chem.* **2018**, *83*, 2250.
- (15) (a) Chinchilla, R.; Nájera, C., Chemicals from Alkynes with Palladium Catalysts, *Chem. Rev.* **2014**, *114*, 1783; (b) Schore, N. E., Transition metal-mediated cycloaddition reactions of alkynes in organic synthesis, *Chem. Rev.* **1988**, *88*, 1081; (c) Alonso, F.; Beletskaya, I. P.; Yus, M., Transition-Metal-Catalyzed Addition of Heteroatom–Hydrogen Bonds to Alkynes, *Chem. Rev.* **2004**, *104*, 3079.
- (16) Pilcher, A. S.; Hill, D. K.; Shimshock, S. J.; Waltermire, R. E.; DeShong, P., Selective deprotection of trialkylsilyl ethers using fluorosilicic acid, *J. Org. Chem.* **1992**, *57*, 2492.
- (17) Pilcher, A. S.; DeShong, P., Improved protocols for the selective deprotection of trialkylsilyl ethers using fluorosilicic acid, *J. Org. Chem.* **1993**, *58*, 5130.
- (18) Sinai, Á.; Mészáros, Á.; Balogh, Á.; Zwillinger, M.; Novák, Z., Hexafluorosilicic Acid as a Novel Reagent for the Desilylation of Silylacetylenes: Application in Sequential Sonogashira Coupling and Click Reaction, *Synthesis* **2017**, *49*, 2374.
- (19) Price of TBAF is 1300 EUR/mol, KOTMS is 278 EUR/mol, Hexafluorosilicic acid 18 EUR/mol, 2018 February, Sigma-Aldrich price
- (20) (a) Urbansky, E. T., Fate of Fluorosilicate Drinking Water Additives, *Chem. Rev.* **2002**, *102*, 2837; (b) Whitford, G. M.; Sampaio, F. C.; Pinto, C. S.; Maria, A. G.; Cardoso, V. E.; Buzalaf, M. A., Pharmacokinetics of ingested fluoride: lack of effect of chemical compound, *Arch. Oral Biol.* **2008**, *53*, 1037; (c) Broadbent, J. M.; Wills, R.; McMillan, J.; Drummond, B. K.; Whyman, R., Evaluation of evidence behind some recent claims against community water fluoridation in New Zealand, *J. R. Soc. N. Z.* **2015**, *45*, 161.
- (21) Mishra, A.; Ma, C.-Q.; Segura, J. L.; Bäuerle, P. In *Handbook of Thiophene-Based Materials*; John Wiley & Sons, Ltd: **2009**, p 1.
- (22) The starting material was synthesized by a method of Waser et al. Brand, J. P.; Charpentier, J.; Waser, J. *Angew. Chem., Int. Ed.* **2009**, *48*, 9346.
- (23) Zacuto, M. J.; O'Malle, S. J.; Leighton, J. L., Tandem Intramolecular Silylformylation–Crotylsilylation: Highly Efficient Synthesis of Polyketide Fragments, *J. Am. Chem. Soc.* **2002**, *124*, 7890.
- (24) Komáromi, A.; Novák, Z., Efficient copper-free Sonogashira coupling of aryl chlorides with palladium on charcoal, *Chem. Commun.* **2008**, 4968.
- (25) Chen, H. J.; Lin, Z. Y.; Li, M. Y.; Lian, R. J.; Xue, Q. W.; Chung, J. L.; Chen, S. C.; Chen, Y. J., A new, efficient, and inexpensive copper(II)/salicylic acid complex catalyzed Sonogashira-type cross-coupling of haloarenes and iodoheteroarenes with terminal alkynes, *Tetrahedron* **2010**, *66*, 7755.
- (26) Li, P.; Wang, L.; Li, H., Application of recoverable nanosized palladium(0) catalyst in Sonogashira reaction, *Tetrahedron* **2005**, *61*, 8633.
- (27) Paegle, E.; Belyakov, S.; Petrova, M.; Liepinsh, E.; Arsenyan, P., Cyclization of Diaryl(hetaryl)alkynes under Selenobromination Conditions: Regioselectivity and Mechanistic Studies, *Eur. J. Org. Chem.* **2015**, *2015*, 4389.
- (28) Marset, X.; Khoshnood, A.; Sotorrios, L.; Gómez-Bengoa, E.; Alonso, D. A.; Ramón, D. J., Deep Eutectic Solvent Compatible Metallic Catalysts: Cationic Pyridiniophosphine Ligands in Palladium Catalyzed Cross-Coupling Reactions, *ChemCatChem* **2017**, *9*, 1269.
- (29) Tyrrell, E.; Whiteman, L.; Williams, N., Sonogashira Cross-Coupling Reactions and Construction of the Indole Ring System Using a Robust, Silica-Supported Palladium Catalyst, *Synthesis* **2009**, *2009*, 829.
- (30) Adam, R.; Cabrero-Antonino, J. R.; Spannenberg, A.; Junge, K.; Jackstell, R.; Beller, M., A General and Highly Selective Cobalt-Catalyzed Hydrogenation of N-Heteroarenes under Mild Reaction Conditions, *Angew. Chem., Int. Ed.* **2017**, *56*, 3216.
- (31) Xu, K.; Sun, S.; Zhang, G.; Yang, F.; Wu, Y., One-pot synthesis of unsymmetrical diarylacetylenes via Sonogashira/deacetonation/Sonogashira cross-coupling of two different aryl chlorides with 2-methyl-3-buten-2-ol, *RSC Adv.* **2014**, *4*, 32643.
- (32) Li, T.; Qu, X.; Xie, G.; Mao, J., [Cu(acac)<sub>2</sub>]-H<sub>2</sub>O-Catalyzed Sonogashira-Type Couplings of Aryl Halides and Terminal Alkynes, *Chem. Asian J.* **2011**, *6*, 1325.
- (33) Zhang, G., Easy Copper-, Ligand- and Amine-Free Sonogashira Coupling Reaction Catalyzed by Palladium on Carbon at Low Catalyst Loading and by Exposure to Air, *Synlett* **2005**, *2005*, 619.
- (34) Fager-Jokela, E.; Muuronen, M.; Patzschke, M.; Helaja, J., Electronic regioselectivity of diarylalkynes in cobalt-mediated pauson-khand reaction: An experimental and computational study with para- and meta-substituted diarylalkynes and norbornene, *J. Org. Chem.* **2012**, *77*, 9134.
- (35) Dermenci, A.; Whittaker, R. E.; Gao, Y.; Cruz, F. A.; Yu, Z.-X.; Dong, G., Rh-catalyzed decarbonylation of conjugated ynones via carbon-alkyne bond activation: reaction scope and mechanistic exploration via DFT calculations, *Chem. Sci.* **2015**, *6*, 3201.
- (36) Bosiak, M. J., A Convenient Synthesis of 2-Arylbenzo[b]furans from Aryl Halides and 2-Halophenols by Catalytic One-Pot Cascade Method, *ACS Catal.* **2016**, *6*, 2429.