

ORIGINAL ARTICLE

Geographical and ecological outline of metal(loid) accumulating plants in Italian vascular flora

Giuseppe Bazan and Giuseppe Galizia

Department of Biological, Chemical and Pharmaceutical Sciences and Technologies (STEBICEF), University of Palermo, Via Archirafi 38, 90123 Palermo, Italy

E-mail address: giuseppe.bazan@unipa.it

Abstract – The decontamination of heavy metal polluted soils is one of the major challenges that our industrialized world has to face. Remediation technologies are being developed and employed in order to reduce the potential hazards of metal and metalloid contamination. Plants capable of uptaking metals and metalloids in their tissues can be an effective tool to remove such pollutants from contaminated soils. The use of this plant-driven process (Phytoremediation) requires the knowledge of the right phytoextractors to use when facing different types of contamination. The aim of this paper is to provide an inventory of phytoextractors that can be used in Phytoremediation procedures in Italy. The checklist includes 172 native or non-invasive alien accumulating and hyperaccumulating plants. An ecological outline of the accumulating flora was done by using the Ellenberg indicator values (EIVs). The high ecological plasticity of these species in different environmental conditions offers a wide spectrum of phytoextractors to choose from for any phytoremediation procedure.

Keywords – Accumulating plants, Ellenberg indicator values, heavy metals, metal uptake, phytoextractors.

Received: September 27, 2018

Accepted: October 12, 2018

Introduction

Heavy metal soil contamination is a major environmental issue in both developed and developing countries (Panagos et al., 2013; Su et al., 2014).

In the European Union, heavy metals (HMs) are by far the most frequent soil contaminants, accounting for more than 34% of cases of soil contamination, followed by mineral oil (23,8%), polycyclic aromatic hydrocarbons (10,9%) and others (EEA, 2011). Even though the European Environment Agency indicators reveal that during the past 15 years heavy metal (HM) emissions have been decreasing in most of the member states, the “European Commission In-depth Report on soil contamination” (2013) shows that HM contamination is still a widespread problem. Moreover, in developing world, HM pollution has dramatically increased over the past decades (Hu et al., 2014; Zhai et al., 2013).

Intensive farming, wastewater irrigation, mining, heavy industry, smelting procedures and improper waste disposal are some of the human activities that contribute to HM soil contamination (Sharma et al., 2007; Khan et al., 2008). As heavy metals cannot be degraded by microorganisms and tend to remain in polluted sites for long periods of time, soil contamination by HMs represents a serious threat to human health (Roberts and Goodman, 1973; Järup, 2003). Conventional soil remediation techniques, such as soil

washing, excavation, in situ vitrification, solidification and stabilization, can be effectively employed in order to achieve remediation of HM contaminated soils. Anyway, in the last decades other approaches have been developed. Some plants are known to be able to uptake and accumulate metals in their above-ground tissues from the soil in which they grow. The exploitation of this ability is known as “phytoextraction” (Salt et al., 1995). Phytoextraction is a well known technology that can be used to remove heavy metals and/or metalloids from contaminated soils for bioremediation purposes (Phytoremediation) or to extract valuable metals for monetary return (Phytomining).

Some plants can also uptake organic pollutants and degrade them enzymatically, a process called phytodegradation. Phytoextraction and phytodegradation are green, non-invasive and feasible methods to clean up contaminated lands. Other phytoremediation techniques include phytostabilization and phytovolatilization (Salt et al., 1998).

Over the last decades, there has been a growing interest in phytoremediation technologies. In developing countries, there is urgent need for cheap and viable remediation techniques to reclaim HM contaminated soils. Therefore, in developing world, Phytoextraction has gained increasing consideration as a reliable, sustainable and low-

cost alternative to conventional soil remediation technologies (Rajakaruna et al., 2006).

The idea that plants could be used to extract metals from soil has been fascinating scientists and businessmen for over one century. Lungwitz (1900) was the first to suggest the inspection of plant tissues to locate gold deposits. In the late 1940s, Minguzzi and Vergnano (1948) identified the first nickel accumulating plant (*Alyssum bertolonii* Desv.). Brooks et al. (1977) first used the term hyperaccumulators to describe plants that contained 1000 mg kg⁻¹ (0.1%) Ni in their dried tissues. Later, Warren and Delavault (1950) found gold and silver in some trees and horsetails. Further investigations on gold uptake by plants were carried out in the Soviet Union by Kitayev and Zhukova (1980). In the last 20 years, researchers (especially in China, India and Australia) have discovered a considerable number of plant species that accumulate or hyperaccumulate metals and/or metalloids. To date, more than 400 metal hyperaccumulating plant species are known. (Sheoran et al., 2009).

Publication data on phytoextracting plants have been collected and stored in several databases, such as “Phytoremediation” (Famulari and Witz., 2015) and PHYTOREM Database (McIntyre, 2003), which are designed for ease of use even for non-scientific users.

In Italy, research on phytoextraction is still in the early stages. Nevertheless, some studies have been carried out in Italy as well. For instance, Marchiol et al. (2007) found that *Helianthus annuus* L. and *Sorghum bicolor* L. were able to remove Cu and Zn from a polluted soil. Massa et al. (2010) did a screening among some autochthonous plants of North West Italy and discovered two HM hyperaccumulators. In an abandoned Sardinian mining site, among native species, Barbafieri et al. (2011) identified three new phytoextractors. Brunetti et al. (2011) investigated the uptake of Cr, Cu, Pb and Zn by *Brassica napus* L. Moreover, extremely high concentrations of thallium were detected in the tissues of *Biscutella laevigata* L. (Fellet et al., 2012). Malagoli et al. (2014) evaluated the phytoextraction potential of *Sinapis alba* L. and *Festuca rubra* L. The following year, Concas et al. (2015) determined the metal content of epigeal and hypogean tissues of *Pistacia lentiscus* L.

More recently, Roccotiello et al. (2016) studied the effects of nickel on the physiology of a Mediterranean plant (*Alyssoides utriculata* (L.) Medik.).

To date, the literature lacks a systematic overview of phytoextracting plants growing in Italy. In this paper, we present a first checklist of phytoextractors, either native or naturalized to Italy. To define the habitat niches and their occurrence along environmental gradients, we characterized the ecology of the accumulating plants by using the Ellenberg indicator values (EIVs). The Ellenberg indicator values are bioindicators that describe the most important abiotic environmental traits (L = light; T = temperature; K = continentality; F = moisture; R = soil reaction = N nitrogen) for each plant species. (Ellenberg et al., 1991). The high ecological diversity of these species offers a huge number of phytoextractors that can be used in different scenarios of contaminated soils.

Materials and methods

The data were gathered from published materials. The existing literature was screened to find metal and/or metalloid (hyper)accumulating plants.

According to Baker et al. (2000), the hyperaccumulation threshold for most of the elements is 1000 mg kg⁻¹ dry mass, except for Gold (mg kg⁻¹), Hg (10 mg kg⁻¹), Cd (100 mg kg⁻¹) Tl (500 mg kg⁻¹), Zn and Mn (10000 mg kg⁻¹). Thus, plants were considered as hyperaccumulators if their aerial tissues contained metal(loid)s at concentrations exceeding these thresholds. Instead, plants failing to exceed the hyperaccumulation threshold, but still capable of concentrating metal(loid)s to levels close to the hyperaccumulation limit were considered as “accumulators”.

We selected the plants that strictly belong to the Italian flora from the available published resources on accumulating species. To ascertain the actual presence of the selected plant species in the Italian flora, we used Conti et al. (2005) annotated checklist of Italian vascular flora. It should be noted that even if allochthonous invasive species are all naturalized plants, such plants pose a potential threat to biodiversity (Sala et al., 2000). To keep these species out of the list, an evaluation of invasiveness was performed using checklist of allochthonous vascular plants of Italy by Celesti-Grappo et al. (2010). Hence, all the species listed in this work are either native or non-invasive allochthonous plants growing in Italy. The list is sorted in alphabetical order and nomenclature follows the standards set by The Plant List (2013). The geographical distribution of Metallophytes within the Italian regions is based on the data provided by Conti et al. (2005). For each taxon, Raunkiaer’s life forms (Raunkiaer, 1934) were recorded according to Pignatti (1982).

The ecological relevance in phytoremediation procedures was also taken into account. The ecological characterization of plants was carried out using Ellenberg’s indicator values (EIVs) (Ellenberg et al., 1991) as published by Pignatti et al. (2005) and updated by Guarino et al. (2012) and Domina et al. (2018). For light and temperature, EIVs have a wider range (up to 12), as modified by Pignatti et al. (2005) to better fit the characteristics of the Italian geographical region.

The statistical analysis was performed using the R-Studio software and map was designed under ESRI ArcGis environment.

Results

Metal(loid) accumulation

In our screening, 172 infraspecific and specific taxa of accumulators were found. As shown in table 1, these species accumulate 18 metals and 2 metalloids.

Table 1 List of Phytoextractors of Italian Flora (hyperaccumulated elements in bold; L.F.= Life Forms).

Taxa	L. F.	Metal(loid)s	References
<i>Achillea ageratum</i> L.	H	Sb	(Baroni et al. 2000)
<i>Alternanthera philoxeroides</i> (Mart.) Griseb.	I	Cd, Mn, Zn	(Nan et al. 2013)
<i>Alyssoides utriculata</i> (L.) Medik.	CH	Ni	(Roccotiello et al. 2016)
<i>Alyssum argenteum</i> All.	CH	Ni	(Vergnano Gambi et al. 1979)
<i>Alyssum bertolonii</i> Desv.	CH	Ni	(Álvarez-López et al. 2016)
<i>Alyssum wulfenianum</i> Benth. ex Willd.	CH	Pb, Tl	(Fellet et al. 2012)
<i>Amaranthus caudatus</i> L.	T	Cd	(Bosiacki et al. 2013)
<i>Anarrhinum bellidifolium</i> (L.) Willd.	H	Hg	(Fernández et al. 2017)
<i>Anthemis cretica</i> L.	H	W	(Erdemir et al. 2017)
<i>Arabidopsis halleri</i> (L.) O'Kane & Al-Shehbaz	H	Cd, Zn	(Küpper et al. 2000; Wenzel and Jockwer 1999)
<i>Arabis alpina</i> var. <i>parviflora</i> Franch	H	Pb	(Yanqun et al. 2005)
<i>Arabis sagittata</i> (Bertol.) DC.	H	Ni, Cu	(Alekseeva-Popova et al. 2015)
<i>Arthrocnemum macrostachyum</i> (Moric.) K.Koch	CH	As, Pb, Zn, Fe	(Martínez-Sánchez et al. 2012)
<i>Asplenium adiantum-nigrum</i> L.	H	Hg	(Fernández et al. 2017)
<i>Atriplex halimus</i> L.	P	Cd, Zn	(Amer et al. 2013; Lutts et al. 2004)
<i>Atriplex prostrata</i> Boucher ex DC.	T	Zn	(Moreira et al. 2011)
<i>Avena strigosa</i> Schreb.	T	Cd	(Uraguchi et al. 2006)
<i>Beta vulgaris</i> L.	H	Cd	(Song et al. 2012)
<i>Betula pubescens</i> Ehrh.	P	Cd	(Fernández et al. 2008)
<i>Bidens pilosa</i> L.	T	Cd	(Sun et al. 2009)
<i>Biscutella laevigata</i> L.	H	Pb, Tl	(Fellet et al. 2012; Wenzel and Jockwer 1999)
<i>Blackstonia perfoliata</i> (L.) Huds.	T	Hg	(Fernández et al. 2017)
<i>Boehmeria nivea</i> (L.) Gaudich.	H	Zn	(Nan et al. 2013)
<i>Brachypodium pinnatum</i> (L.) P.Beauv.	H	Hg	(Fernández et al. 2017)
<i>Brassica juncea</i> (L.) Czern.	T	Cd, U*, Au*	(Chang et al. 2005; Ghosh and Singh 2005; Lamb et al. 2001)
<i>Brassica napus</i> L.	T	Cr	(Brunetti et al. 2011)
<i>Brassica oleracea</i> L.	CH	Mn, Fe, Tl	(Radulescu et al. 2013; Ning et al. 2015)
<i>Brassica rapa</i> L.	T	Cd, Fe, Zn	(Ghosh and Singh 2005; Parveen et al. 2015; Purakayastha et al. 2008)
<i>Callitriche hamulata</i> Kütz. ex W.D.J.Koch	I	U	(Favas et al. 2014)
<i>Callitriche stagnalis</i> Scop.	I	U	(Favas et al. 2014)
<i>Cannabis sativa</i> L.	T	Pb*	(Kos et al. 2003)
<i>Carthamus tinctorius</i> L.	T	Pb	(Al Chami et al. 2015)
<i>Celosia argentea</i> L.	T	Mn	(Liu et al. 2014)
<i>Centaurea nigra</i> L.	H	Hg	(Fernández et al. 2017)
<i>Centaurium erythraea</i> Rafn	H	Hg	(Fernández et al. 2017)
<i>Centaurium pulchellum</i> (Sw.) Druce	T	Hg	(Fernández et al. 2017)
<i>Chenopodium giganteum</i> D.Don	T	Cu	(Bhargava et al. 2008)
<i>Cistus salvifolius</i> L.	NP	Cd	(Barbafieri et al. 2011)
<i>Coincya monensis</i> (L.) Greuter & Burdet	H	Zn	(Fernández et al. 2017)
<i>Convolvulus arvensis</i> L.	G	Cu, Fe, Cr	(Gardea-Torresdey et al. 2004; Massa et al. 2010)
<i>Cornus sanguinea</i> L.	P	Hg	(Fernández et al. 2017)
<i>Cucumis sativus</i> L.	T	As	(Hong et al. 2011)
<i>Cucurbita moschata</i> Duchesne	T	Zn	(Nan et al. 2013)
<i>Dactylis glomerata</i> L.	H	Hg	(Fernández et al. 2017)
<i>Daucus carota</i> L.	H	Hg	(Fernández et al. 2017; Massa et al. 2010)
<i>Dittrichia viscosa</i> (L.) Greuter	H	Cd, Pb, Fe	(Barbafieri et al. 2011; Buscaroli et al. 2017; Fernández et al. 2008)
<i>Dysphania botrys</i> (L.) Mosyakin & Clemants	T	Fe	(Malayeri et al. 2013)
<i>Eleocharis acicularis</i> (L.) Roem. & Schult.	G	Pb, Cu, Ag, In	(Ha et al. 2011)
<i>Elymus elongatus</i> (Host) Runemark	H	Zn	(Sipos et al. 2013)
<i>Equisetum ramosissimum</i> Desf	G	Hg	(Fernández et al. 2017)
<i>Equisetum telmateia</i> Ehrh.	G	Hg	(Fernández et al. 2017)
<i>Erica arborea</i> L.	P	Hg	(Fernández et al. 2017)
<i>Erica cinerea</i> L.	CH	Hg	(Fernández et al. 2017)
<i>Erica scoparia</i> L.	P	Fe	(De La Fuente et al. 2010)
<i>Erodium malacoides</i> (L.) L'Hér.	T	Fe	(Baycu et al. 2015)
<i>Euphorbia dendroides</i> L.	NP	Pb, Zn	(Barbafieri et al. 2011)
<i>Euphrasia stricta</i> D. Wolff	T	Hg	(Fernández et al. 2017)

Taxa	L. F.	Metal(loid)s	References
<i>Festuca ovina</i> L.	H	Pb	(Yanqun et al. 2005)
<i>Festuca rubra</i> L.	H	Cu	(Lago-Vila et al. 2015; Malagoli et al. 2014)
<i>Fraxinus excelsior</i> L.	P	Hg	(Fernández et al. 2017)
<i>Galinsoga parviflora</i> Cav.	T	Cd	(Lin et al. 2014)
<i>Galium aparine</i> L.	T	Hg	(Massa et al. 2010)
<i>Glaucium flavum</i> Crantz	H	Pb, Zn	(Martínez-Sánchez et al. 2012)
<i>Glycine max</i> (L.) Merr.	P	As, Zn, Cu	(Fellet et al. 2007)
<i>Helianthus annuus</i> L.	T	Ni, Cd, As, Pb, U*, Cr	(Cutright et al. 2010; Mihalik et al. 2010; Solhi et al. 2005)
<i>Helichrysum italicum</i> (Roth) G.Don	CH	Pb	(Barbafieri et al. 2011)
<i>Heliotropium europaeum</i> L.	T	Fe	(Nematian and Kazemeini 2013)
<i>Hirschfeldia incana</i> (L.) Lagr.-Foss.	H	Pb	(Fahr et al. 2015)
<i>Holcus lanatus</i> L.	H	As	(Karczewska et al. 2013)
<i>Hordeum distichon</i> L.	H	Zn	(Soriano and Fereres 2003)
<i>Hypericum perforatum</i> L.	H	Hg, Zn	(Fernández et al. 2017)
<i>Iberis carnosa</i> Willd.	H	Zn	(Fernández et al. 2017)
<i>Iberis linifolia</i> L.	T	Tl	(Anderson et al. 1999)
<i>Imperata cylindrica</i> (L.) Raeusch.	G	Al, Fe	(Rodríguez et al. 2005)
<i>Koeleria vallesiana</i> (Honck.) Bertol. ex Schult.	H	Zn	(Fernández et al. 2017)
<i>Lactuca serriola</i> L.	H	Sb	(Hajiani et al. 2015)
<i>Leersia oryzoides</i> (L.) Sw.	G	As	(Ampiah-Bonney et al. 2007)
<i>Lemna gibba</i> L.	I	As	(Mkandawire and Dudel 2005)
<i>Leontodon taraxacoides</i> Hoppe & Hornsch.	H	Hg	(Fernández et al. 2017)
<i>Lepidium sativum</i> L.	T	Pt	(Asztemborska et al. 2015)
<i>Linaria alpina</i> (L.) Mill.	H	Ni	(Vergnano Gambi and Gabrielli 1981)
<i>Lolium perenne</i> L.	H	As, Pb*	(Karczewska et al. 2013; Perry et al. 2012)
<i>Lotus corniculatus</i> L.	H	Zn, Fe, Hg	(Fernández et al. 2017; Massa et al. 2010)
<i>Luzula lutea</i> (All.) DC.	H	Ni	(Vergnano Gambi and Gabrielli 1981)
<i>Lythrum salicaria</i> L.	H	Hg, Zn	(Fernández et al. 2017)
<i>Malva neglecta</i> Wallr.	T	Zn, Cu	(Malayeri et al. 2013; Nematian and Kazemeini 2013)
<i>Malva nicaeensis</i> All.	T	Fe	(Baycu et al. 2015)
<i>Medicago lupulina</i> L.	T	Pb	(Amer et al. 2013)
<i>Medicago sativa</i> L.	H	Pb*	(Pajuelo et al. 2007)
<i>Melilotus albus</i> Medik.	T	Pb	(Fernández et al. 2012)
<i>Melilotus officinalis</i> (L.) Pall.	H	Pb	(Fernández et al. 2012)
<i>Minuartia laricifolia</i> (L.) Schinz & Thell	CH	Ni	(Vergnano Gambi and Gabrielli 1981)
<i>Minuartia verna</i> (L.) Hiern	CH	Pb, Zn	(Fernández et al. 2017; Wenzel and Jockwer 1999)
<i>Mirabilis jalapa</i> L.	G	Cd*	(Wang and Liu 2014)
<i>Moricandia arvensis</i> (L.) DC	T	Cd, Pb	(Karimi et al. 2012)
<i>Musa basjoo</i> Siebold & Zucc. ex Iinuma	G	Pb, Zn	(Nan et al. 2013)
<i>Nicotiana glauca</i> Graham	NP	Zn	(Barazani et al. 2004)
<i>Nicotiana tabacum</i> L.	T	U	(Stojanović et al. 2012)
<i>Persicaria hydropiper</i> (L.) Delarbre	T	Mn	(Liu et al. 2010)
<i>Persicaria lapathifolia</i> (L.) Delarbre	T	Mn	(Liu et al. 2016)
<i>Phagnalon rupestre</i> (L.) DC.	CH	Fe	(Baycu et al. 2015)
<i>Phaseolus vulgaris</i> L.	T	Cd, Pb*, Zn, Cu	(Luo et al. 2005; Luo et al. 2008)
<i>Phillyrea angustifolia</i> L.	P	Cd	(Tapia et al. 2011)
<i>Phragmites australis</i> (Cav.) Trin. ex Steud.	G	Hg, Mn	(Bonanno and Giudice 2010)

Taxa	L. F.	Metal(loid)s	References
<i>Phytolacca americana</i> L.	G	Mn	(Peng et al. 2008)
<i>Picris hieracioides</i> Sibth. & Sm.	H	Hg	(Fernández et al. 2017)
<i>Piptatherum miliaceum</i> (L.) Coss.	H	Hg	(Fernández et al. 2017)
<i>Pistacia lentiscus</i> L.	P	Pb, Zn	(Concas et al. 2015)
<i>Pistacia terebinthus</i> L.	P	Fe	(Baycu et al. 2015)
<i>Pisum sativum</i> L.	T	Pb*	(Chen et al. 2004; Huang et al. 1997)
<i>Plantago lanceolata</i> L.	H	Hg	(Fernández et al. 2017)
<i>Poa alpina</i> L.	H	Zn	(Fernández et al. 2017)
<i>Poa annua</i> L.	T	Cd, Pb, Zn	(Barbafieri et al. 2011)
<i>Polygonum amphibium</i> L.	G	Zn	(Nan et al. 2013)
<i>Polygonum aviculare</i> L.	T	Hg, Zn	(González and González-Chávez 2006; Massa et al. 2010)
<i>Polypogon monspeliensis</i> (L.) Desf.	T	Hg	(Su et al. 2008)
<i>Populus × canadensis</i> Moench	P	Zn	(Nan et al. 2013)
<i>Potentilla micrantha</i> Ramond ex DC.	H	Hg	(Fernández et al. 2017)
<i>Pteridium aquilinum</i> (L.) Kuhn	G	Pb, Zn, Hg	(Fernández et al. 2017)
<i>Pteris cretica</i> L.	H	Sb	(Feng et al. 2015)
<i>Pteris multifida</i> Poir.	H	As	(Wang et al. 2007)
<i>Pteris vittata</i> L.	H	As, Sb, Hg	(Su et al. 2008; Tisarum et al. 2014; Tu et al. 2002)
<i>Quercus robur</i> L.	P	Hg	(Fernández et al. 2017)
<i>Ranunculus peltatus</i> subsp. <i>fucooides</i> (Freyn) Muñoz Garm	I	U	(Favas et al. 2014)
<i>Raphanus raphanistrum</i> subsp. <i>sativus</i> (L.) Domin	T	Zn	(Marchiol et al. 2004)
<i>Rapistrum rugosum</i> (L.) All.	T	Pb	(Saghi et al. 2016)
<i>Reseda lutea</i> L.	H	Fe	(Nematian and Kazemeini 2013)
<i>Rhamnus alaternus</i> L.	P	Cd	(Tapia et al. 2011)
<i>Ricinus communis</i> L.	T	Cd, As	(Huang et al. 2011; Melo et al. 2009)
<i>Rosmarinus officinalis</i> L.	NP	Cd	(Tapia et al. 2011)
<i>Rubus ulmifolius</i> Schott	NP	As, Pb, Hg	(Fernández et al. 2017; Marques et al. 2009)
<i>Salix atrocinerea</i> Brot.	P	Hg, Zn	(Fernández et al. 2017)
<i>Salix caprea</i> L.	P	Hg, Zn	(Fernández et al. 2017)
<i>Salsola kali</i> L.	T	Cr	(Gardea-Torresdey et al. 2005)
<i>Saxifraga exarata</i> Vill.	H	Ni	(Vergnano Gambi and Gabrielli 1981)
<i>Saxifraga paniculata</i> Mill	H	Ni	(Vergnano Gambi and Gabrielli 1981)
<i>Senecio vulgaris</i> L.	T	Fe	(Baycu et al. 2015)
<i>Serapias vomeracea</i> (Burm.f.) Briq.	G	Fe	(Baycu et al. 2015)
<i>Silene ciliata</i> Pourr.	H	Zn	(Fernández et al. 2017)
<i>Silene latifolia</i> Poir.	H	Tl	(Escarré et al. 2011)
<i>Silene nutans</i> L.	H	Hg	(Fernández et al. 2017)
<i>Silene vulgaris</i> (Moench) Garcke	H	Sb	(Baroni et al. 2000)
<i>Sinapis alba</i> L.	T	Tl, Cu, Pt	(Asztemborska et al. 2015; Malagoli et al. 2014; Vaněk et al. 2010)
<i>Sinapis arvensis</i> L.	T	Pb	(Saghi et al. 2016)
<i>Smilax aspera</i> L.	NP	Pb, Ba	(Poschenrieder et al. 2012)
<i>Solanum americanum</i> Mill.	T	Cd	(Wei et al. 2005)
<i>Sonchus asper</i> (L.) Hill	T	Pb	(Yanqun et al. 2005)
<i>Sonchus oleraceus</i> (L.) L.	T	Fe	(Baycu et al. 2015)
<i>Sorghum bicolor</i> (L.) Moench	T	Pb*, Zn, Cu	(Al Chami et al. 2015; Marchiol et al. 2007; Zhuang et al. 2009)
<i>Spinacia oleracea</i> L.	T	Fe	(Pathak et al. 2013)
<i>Stachys recta</i> L.	H	Fe	(Dudić et al. 2007)
<i>Stipa barbata</i> Desf.	H	Cu	(Malayeri et al. 2013)
<i>Tagetes erecta</i> L.	T	Cd	(Uraguchi et al. 2006)
<i>Taraxacum campylodes</i> G.E.Haglund	H	Fe, Pb, Zn	(Bech et al. 2016; Giacomino et al. 2016; Keane et al. 2001)

Taxa	L. F.	Metal(loid)s	References
<i>Teucrium flavum</i> subsp. <i>glaucum</i> (Jord. & Fourr.) Ronniger	CH	Pb	(Cao et al. 2009)
<i>Teucrium polium</i> L.	CH	Ni	(Yaman 2014)
<i>Teucrium scorodonia</i> L.	H	Hg	(Fernández et al. 2017)
<i>Thlaspi caerulescens</i> J.Presl & C.Presl	CH	Ni, Cd, Pb, Zn	(Reeves et al. 2001)
<i>Thlaspi rotundifolium</i> subsp. <i>cepaefolium</i> (Wulfen) Rouy & Foucaud	H	Cd, Pb, Zn	(Wenzel and Jockwer 1999)
<i>Thlaspi sylvium</i> Gaudin	CH	Ni, Zn	(Taylor and Macnair 2006)
<i>Tragopogon porrifolius</i> L.	H	Fe	(Baycu et al. 2015)
<i>Trifolium pallescens</i> Schreb.	H	Ni	(Reeves and Brooks 1983)
<i>Trifolium repens</i> L.	CH	Pb, Zn	(Bech et al. 2016)
<i>Trifolium subterraneum</i> L.	T	Pb*	(Pajuelo et al. 2007)
<i>Trifolium thalii</i> Vill.	H	Zn	(Fernández et al. 2017)
<i>Trisetum flavescens</i> (L.) P.Beauv.	H	W	(Erdemir et al. 2017)
<i>Typha angustifolia</i> L.	G	Pb	(Panich-Pat et al. 2010)
<i>Typha latifolia</i> L.	G	Mn	(Salem et al. 2017)
<i>Vicia faba</i> L.	T	Pb	(Karimi et al. 2012)
<i>Xanthium strumarium</i> L.	T	Fe	(Nematian and Kazemeini 2013)
<i>Zea mays</i> L.	T	Pb*, Cu*	(Huang et al. 1997; Luo et al. 2005)
<i>Zygophyllum fabago</i> L.	NP	Fe	(Martínez-Sánchez et al. 2012)

* Chelating agents added.





Figure 1. Number of accumulators and hyperaccumulators for each metal(loid) (A) and regional distribution of Phytoextractors in Italy (B).

Eighty-eight plant species are hyperaccumulators, while the other phytoaccumulators cannot reach or exceed the hyperaccumulation threshold.

Lead and zinc are the best accumulated metals, followed by mercury, cadmium and iron. (Fig. 1A).

For lead, we report that 25 taxa are hyperaccumulators while 17 are accumulators.

The highest lead concentration is reported in the roots of *Medicago lupulina* L. (63500 mg kg⁻¹). A wetland species, *Typha angustifolia* L. (20173 mg kg⁻¹), and two cultivated plants, *Pisum sativum* L. (11000 mg kg⁻¹) and *Zea mays* L. (10000 mg kg⁻¹), are among the most efficient (and therefore health-threatening) lead hyperaccumulators (Table 1).

For zinc, only 2 of the 42 listed accumulators are hyperaccumulators, with *Thlaspi caerulescens* J. & C. Presl (43710 mg kg⁻¹) being the best extractor, followed by *Alternanthera philoxeroides* (Mart.) Griseb. (43254 mg kg⁻¹). Of the 36 mercury accumulators, 12 are hyperaccumulators. Among these, metal concentration values are very high in two species (43254 mg kg⁻¹ in *Polypogon monspeliensis* (L.) Desf. and 43254 mg kg⁻¹ in *Pteris vittata* L.), with other values dropping to values close to the threshold.

A substantial part of cadmium phytoextractors (19 of 27) are hyperaccumulators, with values ranging from 5722 mg kg⁻¹ in *Arabidopsis halleri* (L.) O'Kane & Al-Shehbaz to 107 mg kg⁻¹ in *Phyllirea angustifolia* L. Iron is another element with a considerable number of phytoaccumulators (> 20). For this metal, six species, such as *Reseda lutea* (48116 mg kg⁻¹) are hyperaccumulators, while 18 cannot reach the threshold.

Worldwide, most of the known Hyperaccumulating plants are nickel accumulators (Boyd, 2009; Sheoran, 2009). In our list, given that the hyperaccumulation threshold for nickel has been set at 1000 mg kg⁻¹ (Baker et al., 2000), *Arabis sagittata* (Bertol.) DC. (120 mg kg⁻¹) is the only plant species that does not reach the specified hyperaccumulation limit. Instead, *Teucrium polium* L. (14110 mg kg⁻¹), *Thlaspi caerulescens* (12880 mg kg⁻¹), *Alyssum bertolonii* (8727 mg kg⁻¹), *Alyssoides utriculata* (L.) Medik (1100 mg kg⁻¹), *Thlaspi sylvium* Gaudin. (1064 mg kg⁻¹) and *Helianthus annuus* L. (2944 mg kg⁻¹), all exceed the threshold and are therefore considered as hyperaccumulators (Tab. 1). Astonishingly, the concentration of nickel in *Teucrium polium* L. is fourteen times higher than the hyperaccumulation limit.

Uncommon metals, such as barium, indium and gold can be extracted by *Smilax aspera* (180 mg kg⁻¹), *Eleocharis acicularis* (L.) Roem. & Schult. (353 mg kg⁻¹) and *Brassica juncea* (L.) Czern. (326 mg kg⁻¹), respectively (see Supplementary file Table S01 for details).

Taxonomical, geographical and ecological characterization

The floristic list reports 167 species, 4 subspecies and 1 variety, for a total of 172 taxa belonging to 52 families. The most represented family is *Brassicaceae* with 25 plant species (14.53%), followed by *Poaceae* (12.21%), *Asteraceae* (11.63%), *Leguminosae* (7.56%),

Chenopodiaceae (4.65%), *Caryophyllaceae* (3.49%), *Lamiaceae* (2.91), *Polygonaceae* (2.33%) and others (< 2%).

Considering life forms, most of the phytoextractors are hemicryptophytes (35.47%) followed by therophytes (31.40%), chamaephytes (8.72%), geophytes (8.72%), phanerophytes (8.72%), nanophanerophytes (4.07%) and idrophytes (2.91%).

The phytoextractors were found to be spread in all the regions of Italy. Toscana has the highest number of Phytoextractors (136), followed by Friuli-Venezia Giulia (130), Lazio (129) and Liguria (129), while Marche (104), Trentino-Alto Adige (99) and Valle d'Aosta (88) are regions with a "low" number of taxa (Fig. 1B). Of all the taxa, 139 are native to Italy, while 33 are alien species.

Ellenberg's indicator values range very widely. As shown in Table 2 and Fig. 2, moisture (F) and nitrogen (N) values have the highest Standard Deviations (2.38 respectively). Mean values (Fig. 2) are: 8.02 (N), 6.78 (T), 4.72 (K), 4.50 (F), 5.63 (R), 3.99 (N), 0.17 (SAL). Compared to the theoretical value (6), light (L) and temperature values (T) have a higher median value, while continentality values (K), moisture values (F) and nitrogen values (N) have lower median values (Fig. 2).




Figure 2. Ecograms of Ellenberg Environmental values for the Phytoextractors of Italy. (L = light; T = temperature; K= continentality; F= moisture; R = Soil reaction= N nitrogen).

Table 2. Ellenberg values¹ statistics for Italian Phytoextractors.

	L	T	K	F	R	N	S
Minimum	3	1	0	1	0	0	0
Maximum	11	12	8	12	9	9	9
Mean	8.02	6.78	4.72	4.50	5.63	3.99	0.17
Median	8	7	5	4	6	4	0
Standard Deviation	1.63	2.05	1.01	2.38	2.02	2.38	0.93
Variance	2.63	4.16	1.02	5.62	4.04	5.61	0.86

¹ L = Light value; T = Temperature value; K = Continentality value; F = Moisture value; R= Reaction of soil value (PH); N = Nitrogen value; S= Salinity value.

Discussion

Geographical distribution of metal(loid) accumulating plants in Italy

In Italy, heavy metal soil contamination has a quite heterogeneous and diversified geographical pattern. For instance, it is not uncommon to find different concentrations of various heavy metals in the same land areas. According to Tóth et al. (2016), many countries of Europe have regions with a high percentage of topsoil samples which have concentrations of inorganic pollutants well above the normal values. Many Regions of Italy are affected by heavy metal soil pollution, as well (Tóth et al., 2016). Anyway, all the Italian Regions, have a fairly large number of phytoextractors in their flora to be used for soil remediation needs. In fact, almost all the regions host more than 100 taxa of phytoextractors (Fig. 1B).

A comparison between the whole flora and the number of phytoextractors for each region gave a 0.6 Pearson Index. This indicates a significant correlation between the two variables and also suggests that the number of phytoextractors might be primarily dependent on the floristic richness of regions. Worldwide, the metal(loid) accumulating plant species are usually abundant in areas with a significant presence of serpentine soils, such as Cuba, South Africa and New Caledonia (Callahan et al., 2012; Reeves et al., 2018). In Italy, serpentine geological islands range from the central-western Alps to the northern Apennines, mainly appearing in the region of Toscana, where several outcrops are scattered in different biogeographic sectors (Selvi, 2006; Bini and Maleci, 2014; Bini et al., 2017). As expected, in our analysis the majority of the phytoaccumulators are plants of the Tuscan flora, where serpentine soils are somewhat common. However, of the 87 serpentinophytes observed by Selvi (2006), only *Alyssum bertolonii*, *Cistus salvifolius* L., *Helichrysum italicum* (Roth) G. Don and *Erica arborea* L. are known to be phytoaccumulators (Table 1). Therefore, further investigations might lead to the discovery of more metal accumulating plant species within the Italian serpentinophytes.

The regional distribution of phytoaccumulators for any given metal is shown in figure 3.

The phytoextractors of indium, barium, aluminium, silver and gold have the most limited geographical distribution.

Taxonomical considerations on metal(loid) accumulating plants

Many different families have large numbers of species that tolerate high levels of HMs. The family with the largest number of phytoextractors is *Brassicaceae*, followed by *Asteraceae*, *Caryophyllaceae*, *Poaceae* and *Leguminosae*. Many authors have pointed out that most of the known accumulators belong to the *Brassicaceae* family (Krämer, 2010; Szczygłowska et al., 2011). As already mentioned above, this is consistent with our data. From the placement of the 52 families of phytoextractors into the phylogenetic tree (Angiosperm Phylogeny Group III, 2009) it can be noted that the phytoextractors tend to distribute in different clades. According to Broadley et al. (2001), the phenomenon of hyperaccumulation is more common in certain orders than in others. For instance, Ni hyperaccumulators are very common in Brassicales, Asterales and Malpighiales.

As noted by Baker and Brooks (1989), even if the (hyper)accumulators of Nickel seem to have some phylogenetic relationships at the family level, the phenomenon of Metal hyperaccumulation has evolved in a wide range of apparently unrelated taxa.

The phylogenetic distribution of the accumulators in our list suggests that the ability to accumulate metals may be independent of phylogeny.

Ecological behavior of metal(loid) accumulating plants

The ecological relationship amongst accumulating plants, climate and soil conditions is described using the Ellenberg's indicator values (Eiv). Such indicators are an empirical tool used to describe the ecological response of plants to environment (Marcenò et Guarino 2015). However, because EIVs are ordinal scales, EIV values are generally not considered mathematically rigorous enough. Still, since it has been proved that all kinds of statistical tests based on mean and variance can be used with acceptable confidence levels (Domina et al., 2018), we decided to use the EIVs to gather informations on the ecology of metal(loid) accumulating plants.




Figure 3. Regional distribution of phytoaccumulators for any given metal.

Pignatti's EIVs assessment for the whole Italian native flora includes 5774 taxa (Pignatti 2005; Domina et al., 2018). In our checklist, the native accumulating plants (139 taxa) represent the 2,41% of these taxa. The alien species have a similar percentage, being the 2.07% of the whole assessed alien taxa (1597), which were listed by Domina et al (2018).

The EIVs mean comparison between the accumulating native flora and the whole flora of Italy (Pignatti et al., 2005) indicates that the native phytoextractors have higher nutrient requirements (N= 3.71 vs 3.26), higher soil reaction requirements (R= 5.75 vs 5.45) and higher edaphic humidity requirements (F= 4.48 vs 4.17), while having similar ecological requirements for Light (L= 7.94 vs 7.84), Temperature (T= 6.50 vs 6.36) and Continentality (K= 4.67 vs 4.60). Despite the low number of alien taxa, the EIVs for the alien phytoextractors are in line with the national EIVs distribution for the alien plants. Within the accumulating plants, the alien species are more thermophilous than the native species and this is consistent with the considerations of Domina et al. (2018) for the whole Italian flora. Mean EIVs for the metal(loid) accumulating flora clearly shows that this subset of flora is ecologically representative of the entire Italian flora.

The distribution of R values (Fig. 4A) for the best accumulated metal(loid)s shows that most of the phytoextractors naturally grow in neutral to basic soils. For example, species with high R-numbers have higher Fe-solubilizing capacity than most species with low values (Bartelheimer and Poschod, 2016). On the contrary, *Erica arborea* (an acidophilous species with an R value of 1) contains very low levels of Fe (202 mg kg⁻¹) (Fernández et al., 2017). For Zn and Pb, a basophile plant such as *Thlaspi rotundifolium* subsp. *cepaefolium* (Wulfen) Rouy (R=9) has a very good accumulation capability and its ecological requirements are in line with the observations of Bartelheimer and Poschod (2016). *Trifolium repens* L. (R=5), a plant growing in soils with a near-neutral pH, is also capable of accumulating high concentrations of metal(loid)s in substrates with a great calcium carbonate percentage (Bech et al., 2016). Consistently with these considerations, we did not find any correlation between the R values and accumulation capability (Pearson $r = 0.145$ for Pb; Pearson $r = -0.001$ for Zn). For nitrogen/nutrients (Fig. 4B), values range from N = 2 in species growing on poor soils to N = 7 in plants often found in richly fertile places. The box plot (Fig. 4B) shows that the data have a broad distribution, with low median values (except for As and Cu). Such variability of the ecological requirements offers a wide spectrum of phytoextractors to choose from for any phytoremediation procedure in Italy.

Outcomes and implications

Over the last decades, the phenomenon of hyperaccumulation of metal(loid)s in plants aroused great interest within the scientific community, both from a biological point of view (understanding the mechanisms of metal absorption and translocation) and for the potential use of metal accumulating plants in bioremediation procedures. The research on this topic produced thousands

of publications, resulting in the discovery of a huge number of metal(loid) accumulating taxa, some of which accumulate metal(loid)s at exceptional concentrations. The bibliographic screening in this work allowed us to identify a fairly large number of metal(loid) accumulating species that belong to the Italian flora. Interestingly, the EIVs ecological outline revealed a high ecological plasticity of the entire accumulating flora.

In conclusion, this work wants to provide a first synthesis of what is known about the Italian phytoaccumulating species. Also, we think that it can be useful to have a list of phytoaccumulators to use in various contexts of soil contamination for phytoremediation purposes.




Figure 4. Box plots of the distribution of the Soil Reaction values (A) and Nitrogen/Nutrient values (B) for the most important metals.

Acknowledgements

We thank unknown referees for their comments that significantly improved this paper.

References

- Al Chami, Z., Amer, N., Al Bitar, L., Cavoski I., 2015. Potential use of *Sorghum bicolor* and *Carthamus tinctorius* in phytoremediation of nickel, lead and zinc. *Int J Environ Sci Technol* 12(12): 3957-3970.
DOI: [10.1007/s13762-015-0823-0](https://doi.org/10.1007/s13762-015-0823-0)
- Alekseeva-Popova, N.V., Drozdova, I.V., Kalimova, I.B., 2015. Accumulation of heavy metals by North Caucasian plant species of the Cruciferae family in regards to phytoremediation. *Geochem Int* 53(5): 456-463.
DOI: [10.1134/S0016702915030027](https://doi.org/10.1134/S0016702915030027)
- Alford, É.R., Pilon-Smits, E.A., Paschke, M.W., 2010. Metallophytes—a view from the rhizosphere. *Plant Soil* 337(1-2): 33-50.
DOI: [10.1007/s11104-010-0482-3](https://doi.org/10.1007/s11104-010-0482-3)
- Álvarez-López, V., Prieto-Fernández, Á, Cabello-Conejo, M.I., Kidd, P.S., 2016. Organic amendments for improving biomass production and metal yield of Ni-hyperaccumulating plants. *Sci Total Environ* 548: 370-379.
DOI: [10.1016/j.scitotenv.2015.12.147](https://doi.org/10.1016/j.scitotenv.2015.12.147)
- Amer, N., Chami, Z.A., Bitar, L.A., Mondelli, D., Dumontet, S., 2013. Evaluation of *Atriplex halimus*, *Medicago lupulina* and *Portulaca oleracea* for phytoremediation of Ni, Pb, and Zn. *Int J Phytoremediat* 15(5): 498-512.
DOI: [10.1080/15226514.2012.716102](https://doi.org/10.1080/15226514.2012.716102)
- Ampiah-Bonney, R.J., Tyson, J.F., Lanza, G.R., 2007. Phytoextraction of arsenic from soil by *Leersia oryzoides*. *Int J Phytoremediat* 9(1): 31-40.
DOI: [10.1080/15226510601139383](https://doi.org/10.1080/15226510601139383)
- Anderson, C.W.N., Brooks, R.R., Chiarucci, A., LaCoste, C.J., Leblanc, M., Robinson, B.H, Stewart, R.B., 1999. Phytomining for nickel, thallium and gold. *J Geochem Explor* 67(1-3): 407-415.
DOI: [10.1016/S0375-6742\(99\)00055-2](https://doi.org/10.1016/S0375-6742(99)00055-2)
- Angiosperm Phylogeny Group., 2009. An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG III. *Bot J Linn Soc* 161(2): 105-121.
- Asztemborska, M., Steborowski, R., Kowalska, J., Bystrzejewska-Piotrowska, G., 2015. Accumulation of platinum nanoparticles by *Sinapis alba* and *Lepidium sativum* plants. *Water Air Soil Pollut* 226(4): 126.
DOI: [10.1007/s11270-015-2381-y](https://doi.org/10.1007/s11270-015-2381-y)
- Baker, A.J.M., 2000. Metal hyperaccumulator plants: a review of the ecology and physiology of a biological resource for phytoremediation of metal-polluted soils. *Phytoremediat Contam Soil Water*.
- Baker, A.J.M., Brooks R., 1989. Terrestrial higher plants which hyperaccumulate metallic elements. A review of their distribution, ecology and phytochemistry. *Biorecovery* 1(2): 81-126.
- Barazani, O., Sathiyamoorthy, P., Manandhar, U., Vulkan, R., Golan-Goldhirsh, A., 2004. Heavy metal accumulation by *Nicotiana glauca* Graham in a solid waste disposal site. *Chemosph* 54(7): 867-872.
DOI: [10.1016/j.chemosphere.2003.10.005](https://doi.org/10.1016/j.chemosphere.2003.10.005)
- Barbafieri, M., Dadea, C., Tassi, E., Bretzel, F., Fanfani, L., 2011. Uptake of heavy metals by native species growing in a mining area in Sardinia, Italy: discovering native flora for phytoremediation. *Int J Phytoremediat* 13(10): 985-997.
DOI: [10.1080/15226514.2010.549858](https://doi.org/10.1080/15226514.2010.549858)
- Baroni, F., Boscagli, A., Protano, G., Riccobono, F., 2000. Antimony accumulation in *Achillea ageratum*, *Plantago lanceolata* and *Silene vulgaris* growing in an old Sb-mining area. *Environ Pollut* 109(2): 347-352.
DOI: [10.1016/S0269-7491\(99\)00240-7](https://doi.org/10.1016/S0269-7491(99)00240-7)
- Baycu, G., Tolunay, D., Ozden H, Csatari I, Karadag, S., Agba, T., Rognes, S.E., 2015. An abandoned copper mining site in Cyprus and assessment of metal concentrations in plants and soil. *Int J Phytoremediat* 17(7): 622-631.
DOI: [10.1080/15226514.2014.922929](https://doi.org/10.1080/15226514.2014.922929)
- Bech, J., Roca, N., Tume, P., Ramos-Miras, J., Gil, C., Boluda, R., 2016. Screening for new accumulator plants in potential hazards elements polluted soil surrounding Peruvian mine tailings. *Catena* 136: 66-73.
DOI: [10.1016/j.catena.2015.07.009](https://doi.org/10.1016/j.catena.2015.07.009)
- Bhargava, A., Shukla, S., Srivastava, J., Singh, N., Ohri, D., 2008. *Chenopodium*: a prospective plant for phytoextraction. *Acta Physiol Plant*, 30(1): 111-120.
DOI: [10.1007/s11738-007-0097-3](https://doi.org/10.1007/s11738-007-0097-3)
- Bini, C., Maleci, L., Wahsha, M., 2017. Potentially toxic elements in serpentine soils and plants from Tuscany (Central Italy). A proxy for soil remediation. *Catena* 148: 60-66.
DOI: [10.1016/j.catena.2016.03.014](https://doi.org/10.1016/j.catena.2016.03.014)
- Bonanno, G., Giudice, R.L., 2010. Heavy metal bioaccumulation by the organs of *Phragmites australis* (common reed) and their potential use as contamination indicators. *Ecol Indic* 10(3): 639-645.
DOI: [10.1016/j.ecolind.2009.11.002](https://doi.org/10.1016/j.ecolind.2009.11.002)
- Bosiacki, M., Kleiber, T., Kaczmarek, J., 2013. Evaluation of suitability of *Amaranthus caudatus* L. and *Ricinus communis* L. in phytoextraction of cadmium and lead from contaminated substrates. *Arch Environ Prot* 39(3): 47-59.
DOI: [10.2478/aep-2013-0022](https://doi.org/10.2478/aep-2013-0022)

- Boyd, R.S., 2009. High-nickel insects and nickel hyperaccumulator plants: A review. *Insect Sci* 16(1): 19-31.
DOI: [10.1111/j.1744-7917.2009.00250.x](https://doi.org/10.1111/j.1744-7917.2009.00250.x)
- Broadley, M.R., Willey, N.J., Wilkins, J.C., Baker, A.J., Mead, A., White, P.J., 2001. Phylogenetic variation in heavy metal accumulation in angiosperms. *New Phytolog* 152(1): 9-27.
DOI: [10.1046/j.0028-646x.2001.00238.x](https://doi.org/10.1046/j.0028-646x.2001.00238.x)
- Brooks R.R., Lee, J., Reeves, R.D Jaffre, T., 1977a. Detection of nickeliferous rocks by analysis of herbarium specimens of indicator plants. *J Geochem Explor* 7: 49-77
DOI: [10.1016/0375-6742\(77\)90074-7](https://doi.org/10.1016/0375-6742(77)90074-7)
- Brunetti, G., Farrag, K., Rovira, P.S., Nigro, F., Senesi, N., 2011. Greenhouse and field studies on Cr, Cu, Pb and Zn phytoextraction by *Brassica napus* from contaminated soils in the Apulia region, Southern Italy. *Geoderma* 160(3): 517-523.
DOI: [10.1016/j.geoderma.2010.10.023](https://doi.org/10.1016/j.geoderma.2010.10.023)
- Burlo, F., Guijarro, I., Carbonell-Barrachina, A.A., Valero, D., Martinez-Sanchez, F., 1999. Arsenic species: effects on and accumulation by tomato plants. *J Agric Food Chem*, 47(3): 1247-1253.
DOI: [10.1021/jf9806560](https://doi.org/10.1021/jf9806560)
- Buscaroli, A., Zannoni, D., Menichetti, M., Dinelli, E., 2017. Assessment of metal accumulation capacity of *Dittrichia viscosa* (L.) Greuter in two different Italian mine areas for contaminated soils remediation. *J Geochem Explor* 182: 123-131.
DOI: [10.1016/j.gexplo.2016.10.001](https://doi.org/10.1016/j.gexplo.2016.10.001)
- Callahan, D.L., Roessner, U., Dumontet, V., De Livera A.M., Doronila, A., Baker, A.J., et al. 2012. Elemental and metabolite profiling of nickel hyperaccumulators from New Caledonia. *Phytochem* 81: 80-89.
DOI: [10.1016/j.phytochem.2012.06.010](https://doi.org/10.1016/j.phytochem.2012.06.010)
- Cao, A., Carucci, A., Lai, T., Bacchetta, G., Casti, M., 2009. Use of native species and biodegradable chelating agents in the phytoremediation of abandoned mining areas. *J Chem Technol Biotechnol* 84(6): 884-889.
DOI: [10.1002/jctb.2179](https://doi.org/10.1002/jctb.2179)
- Celesti-Grapow, L., Accogli R., 2010. Flora vascolare alloctona e invasiva delle regioni d'Italia. Università degli Studi di Roma La Sapienza.
- Chang, P., Kim, K.W., Yoshida, S., Kim, S.Y., 2005. Uranium accumulation of crop plants enhanced by citric acid. *Environ Geochem Health* 27(5-6): 529-538.
DOI: [10.1007/s10653-005-8013-5](https://doi.org/10.1007/s10653-005-8013-5)
- Chen Y, Li, X., Shen Z., 2004. Leaching and uptake of heavy metals by ten different species of plants during an EDTA-assisted phytoextraction process. *Chemosph* 57(3): 187-196.
DOI: [10.1016/j.chemosphere.2004.05.044](https://doi.org/10.1016/j.chemosphere.2004.05.044)
- Concas, S., Lattanzi, P., Bacchetta, G., Barbaferi, M., Vacca A., 2015. Zn, Pb and Hg Contents of *Pistacia lentiscus* L. Grown on Heavy Metal-Rich Soils: Implications for Phytostabilization. *Water Air Soil Pollut* 226(10): 1-15.
DOI: [10.1007/s11270-015-2609-x](https://doi.org/10.1007/s11270-015-2609-x)
- Conti, F., Bonacquisti, S., Scassellati, E., 2005. An annotated checklist of the Italian vascular flora. Palombi, Roma
- Cutright, T., Gunda, N., Kurt, F., 2010. Simultaneous hyperaccumulation of multiple heavy metals by *Helianthus annuus* grown in a contaminated sandy-loam soil. *Int J Phytoremediat* 12(6): 562-573.
DOI: [10.1080/15226510903353146](https://doi.org/10.1080/15226510903353146)
- De La Fuente, V., Rufo, L., Rodríguez, N., Amils, R., Zuluaga, J., 2010. Metal accumulation screening of the Río Tinto flora (Huelva, Spain). *Biol Trace Elem Res* 134(3): 318-341.
DOI: [10.1007/s12011-009-8471-1](https://doi.org/10.1007/s12011-009-8471-1)
- Domina, G., Galasso, G., Bartolucci, F., Guarino, R., 2018. Ellenberg Indicator Values for the vascular flora alien to Italy. *Flora Mediterranea* 28(1): 53-61.
DOI: [10.7320/FIMedit28.053](https://doi.org/10.7320/FIMedit28.053)
- Dudić, B., Rakić, T., Šinžar-Sekulić J., Atanacković, V., Stevanović, B., 2007. Differences of metal concentrations and morpho-anatomical adaptations between obligate and facultative serpentinophytes from Western Serbia. *Arch Biol Sci* 59(4): 341-349.
DOI: [10.2298/ABS0704341D](https://doi.org/10.2298/ABS0704341D)
- EEA 2011. Progress in management of contaminated sites. European Environment Agency. <http://www.eea.europa.eu/data-and-maps/indicators/progress-in-management-of-contaminated-sites-3/assessment>
- Ellenberg, H., 1992. Zeigerwerte von pflanzen in Mitteleuropa. *Scripta geobot*, 18, 1-258.
- Erdemir Ü.S., Arslan H, Güteryüz, G., Güçer, Ş., 2017. Elemental Composition of Plant Species from an Abandoned Tungsten Mining Area: Are They Useful for Biogeochemical Exploration and/or Phytoremediation Purposes? *Bull Environ Contam Toxicol* 98(3): 299-303.
DOI: [10.1007/s00128-016-1899-z](https://doi.org/10.1007/s00128-016-1899-z)
- Escarré, J., Lefèbvre, C., Raboyeau, S., Dossantos, A., Gruber, W., Marel, J.C.C., Van Oort, F., 2011. Heavy metal concentration survey in soils and plants of the Les Malines mining district (Southern France): implications

for soil restoration. *Water Air Soil Pollut* 216(1-4): 485-504.

DOI: [10.1007/s11270-010-0547-1](https://doi.org/10.1007/s11270-010-0547-1)

Fahr, M., Laplaze, L., El Mzibri, M., Dumas, P., Bendaou, N., Hoher, V., Smouni, A., 2015. Assessment of lead tolerance and accumulation in metallicolous and non-metallicolous populations of *Hirschfeldia incana*. *Environ Exp Bot*, 109: 186-192.

DOI: [10.1016/j.envexpbot.2014.07.010](https://doi.org/10.1016/j.envexpbot.2014.07.010)

Famulari, S., Witz K., 2015. A User-Friendly Phytoremediation Database: Creating the Searchable Database, the Users, and the Broader Implications. *Int J Phytoremediat* 17(8): 737-744.

DOI: [10.1080/15226514.2014.987369](https://doi.org/10.1080/15226514.2014.987369)

Favas P.J., Pratas, J., Varun, M., D'Souza, R., Paul, M.S., 2014. Accumulation of uranium by aquatic plants in field conditions: prospects for phytoremediation. *Sci Total Environ*, 470: 993-1002.

DOI: [10.1016/j.scitotenv.2013.10.067](https://doi.org/10.1016/j.scitotenv.2013.10.067)

Fellet, G., Marchiol, L., Perosa, D., Zerbi, G., 2007. The application of phytoremediation technology in a soil contaminated by pyrite cinders. *Ecol eng*, 31(3): 207-214.

DOI: [10.1016/j.ecoleng.2007.06.011](https://doi.org/10.1016/j.ecoleng.2007.06.011)

Fellet, G., Pošćić, F., Casolo, V., Marchiol, L., 2012. Metallophytes and thallium hyperaccumulation at the former Raibl lead/zinc mining site (Julian Alps, Italy). *Plant Biosyst - An International Journal Dealing with all Aspects of Plant Biology*, 146(4), 1023-1036.

DOI: [10.1080/11263504.2012.703250](https://doi.org/10.1080/11263504.2012.703250)

Feng, R., Wang, X., Wei, C., Tu S., 2015. The accumulation and subcellular distribution of arsenic and antimony in four fern plants. *Int J Phytoremediat* 17(4): 348-354.

DOI: [10.1080/15226514.2013.773281](https://doi.org/10.1080/15226514.2013.773281)

Fernández, R., Bertrand, A., Casares, A., García, R., González, A., Tamés, R.S., 2008. Cadmium accumulation and its effect on the in vitro growth of woody fleabane and mycorrhized white birch. *Environ Pollut* 152(3): 522-529.

DOI: [10.1016/j.envpol.2007.07.011](https://doi.org/10.1016/j.envpol.2007.07.011)

Fernández, R., Bertrand, A., García J.I., Tamés, R.S., González, A., 2012. Lead accumulation and synthesis of non-protein thiolic peptides in selected clones of *Melilotus alba* and *Melilotus officinalis*. *Environ Exp Bot*: 78, 18-24.

DOI: [10.1016/j.envexpbot.2011.12.016](https://doi.org/10.1016/j.envexpbot.2011.12.016)

Fernández, S., Poschenrieder, C., Marcenò, C., Gallego J.R., Jiménez-Gómez, D., Bueno, A., Afif, E., 2017. Phytoremediation capability of native plant species living on Pb-Zn and Hg-As mining wastes in the Cantabrian range, north of Spain. *J Geochem Explor*, 174: 10-20.

DOI: [10.1016/j.gexplo.2016.05.015](https://doi.org/10.1016/j.gexplo.2016.05.015)

Gardea-Torresdey J.L., De la Rosa, G., Peralta-Videa, J.R., Montes, M., Cruz-Jimenez, G., Cano-Aguilera, I., 2005. Differential uptake and transport of trivalent and hexavalent chromium by tumbleweed (*Salsola kali*). *Arch Environ Contam Toxicol* 48(2): 225-232. DOI: [10.1007/s00244-003-0162-x](https://doi.org/10.1007/s00244-003-0162-x)

Gardea-Torresdey, J.L., Peralta-Videa, J.R., Montes, M., De la Rosa, G., Corral-Diaz, B., 2004. Bioaccumulation of cadmium, chromium and copper by *Convolvulus arvensis* L.: impact on plant growth and uptake of nutritional elements. *Bioresour Technol* 92(3): 229-235.

DOI: [10.1016/j.biortech.2003.10.002](https://doi.org/10.1016/j.biortech.2003.10.002)

Ghosh, M., Singh, S.P., 2005. A comparative study of cadmium phytoextraction by accumulator and weed species. *Environ Pollut* 133(2): 365-371.

DOI: [10.1016/j.envpol.2004.05.015](https://doi.org/10.1016/j.envpol.2004.05.015)

Giacomino, A., Malandrino, M., Colombo M.L., Miaglia, S., Maimone, P., Blancato, S., Abollino, O., 2016. Metal content in dandelion (*Taraxacum officinale*) leaves: influence of vehicular traffic and safety upon consumption as food. *J Chem* 2016.

DOI: [10.1155/2016/9842987](https://doi.org/10.1155/2016/9842987)

Gisbert, C., Clemente, R., Navarro-Avinó, J., Baixauli, C., Ginér, A., Serrano, R., Bernal, M.P., 2006. Tolerance and accumulation of heavy metals by Brassicaceae species grown in contaminated soils from Mediterranean regions of Spain. *Environ Exp Bot* 56(1): 19-27.

DOI: [10.1016/j.envexpbot.2004.12.002](https://doi.org/10.1016/j.envexpbot.2004.12.002)

González, R.C., González-Chávez, M.C.A., 2006. Metal accumulation in wild plants surrounding mining wastes. *Environ Pollut*, 144(1): 84-92.

Gulz, P.A., Gupta, S.K., Schulin, R., 2005. Arsenic accumulation of common plants from contaminated soils. *Plant Soil*, 272(1-2): 337-347.

DOI: [10.1007/s11104-004-5960-z](https://doi.org/10.1007/s11104-004-5960-z)

Ha, N.T.H., Sakakibara, M., Sano, S., 2011. Accumulation of Indium and other heavy metals by *Eleocharis acicularis*: an option for phytoremediation and phytomining. *Bioresour Technol*, 102(3): 2228-2234.

DOI: [10.1016/j.biortech.2010.10.014](https://doi.org/10.1016/j.biortech.2010.10.014)

Hajiani, N.J., Ghaderian, S.M., Karimi, N., Schat, H., 2015. A comparative study of antimony accumulation in plants growing in two mining areas in Iran, Moghanlo, and Patyar. *Environ Sci Pollut Res* 22(21): 16542-16553.

DOI: [10.1007/s11356-015-4852-5](https://doi.org/10.1007/s11356-015-4852-5)

Hong, S.H., Choi, S.A., Yoon, H., Cho, K.S., 2011. Screening of *Cucumis sativus* as a new arsenic-accumulating plant and its arsenic accumulation in hydroponic culture. *Environ Geochem Health*, 33(1): 143-149.

DOI: [10.1007/s10653-010-9350-6](https://doi.org/10.1007/s10653-010-9350-6)

- Hu, H., Jin, Q., Kavan, P., 2014. A study of heavy metal pollution in China: Current status, pollution-control policies and countermeasures. *Sustain* 6(9): 5820-5838.
DOI: [10.3390/su6095820](https://doi.org/10.3390/su6095820)
- Huang, H., Yu, N., Wang, L., Gupta, D.K., He, Z., Wang, K., Yang, X.E., 2011. The phytoremediation potential of bioenergy crop *Ricinus communis* for DDTs and cadmium co-contaminated soil. *Bioresour Technol* 102(23): 11034-11038.
DOI: [10.1016/j.biortech.2011.09.067](https://doi.org/10.1016/j.biortech.2011.09.067)
- Huang J.W., Chen, J., Berti, W.R., Cunningham S.D., 1997. Phytoremediation of lead-contaminated soils: role of synthetic chelates in lead phytoextraction. *Environ Sci Technol* 31(3): 800-805.
DOI: [10.1021/es9604828](https://doi.org/10.1021/es9604828)
- Järup L., 2003. Hazards of heavy metal contamination. *Br med bull* 68(1): 167-182.
- Jolly Y.N., Islam, A., Akbar, S., 2013. Transfer of metals from soil to vegetables and possible health risk assessment. *SpringerPlus*, 2(1): 385.
DOI: [10.1186/2193-1801-2-385](https://doi.org/10.1186/2193-1801-2-385)
- Karczewska, A., Lewińska, K., Gałka, B., 2013. Arsenic extractability and uptake by velvetgrass *Holcus lanatus* and ryegrass *Lolium perenne* in variously treated soils polluted by tailing spills. *J Hazard Mater*, 262: 1014-1021.
DOI: [10.1016/j.jhazmat.2012.09.008](https://doi.org/10.1016/j.jhazmat.2012.09.008)
- Karimi, R., Chorom, M., Solhi, S., Solhi, M., 2012. Potential of *Vicia faba* and *Brassica arvensis* for phytoextraction of soil contaminated with cadmium, lead and nickel. *Afr J Agric Res* 7(22): 3293-3301.
DOI: [10.5897/AJAR12.165](https://doi.org/10.5897/AJAR12.165)
- Keane, B., Collier, M.H, Shann, J.R., Rogsta, S.H., 2001. Metal content of dandelion (*Taraxacum officinale*) leaves in relation to soil contamination and airborne particulate matter. *Sci Total Environ* 281(1-3): 63-78.
DOI: [10.1016/S0048-9697\(01\)00836-1](https://doi.org/10.1016/S0048-9697(01)00836-1)
- Khan, S., Cao, Q., Zheng, Y.M., Huang, Y.Z., Zhu, Y.G., 2008. Health risks of heavy metals in contaminated soils and food crops irrigated with wastewater in Beijing, China. *Environ Pollut* 152(3): 686-692.
DOI: [10.1016/j.envpol.2007.06.056](https://doi.org/10.1016/j.envpol.2007.06.056)
- Kitayev N.A., Zhukova R.I., 1980. Relationship between the concentrations of gold in soil, forest bedding and the bark of trees. *Sov Geol Geophys* 21(12): 118-121.
- Kos, B., Greman, H., Lestan, D., 2003. Phytoextraction of lead, zinc and cadmium from soil by selected plants. *Plant Soil Environ*, 49(12): 548-553.
DOI: [10.17221/4192-PSE](https://doi.org/10.17221/4192-PSE)
- Krämer, U., 2010. Metal hyperaccumulation in plants. *Annu Rev Plant Biol* 61: 517-534.
DOI: [10.1146/annurev-arplant-042809-112156](https://doi.org/10.1146/annurev-arplant-042809-112156)
- Küpfer, H., Lombi, E., Zhao, F.J., McGrath, S.P., 2000. Cellular compartmentation of cadmium and zinc in relation to other elements in the hyperaccumulator *Arabidopsis halleri*. *Planta*, 212(1): 75-84.
DOI: [10.1007/s004250000366](https://doi.org/10.1007/s004250000366)
- Lago-Vila, M., Arenas-Lago, D., Rodríguez-Seijo, A., Couce, M.A., Vega, F.A., 2015. Cobalt, chromium and nickel contents in soils and plants from a serpentinite quarry. *Solid earth* 6(1): 323.
- Lin, L., Jin, Q., Liu, Y., Ning, B., Liao, M.A., Luo, L., 2014. Screening of a new cadmium hyperaccumulator, *Galinsoga parviflora*, from winter farmland weeds using the artificially high soil cadmium concentration method. *Environ Toxicol Chem*, 33(11): 2422-2428.
DOI: [10.1002/etc.2694](https://doi.org/10.1002/etc.2694)
- Liu, J., Shang, W., Zhang, X., Zhu, Y., Yu, K., 2014. Mn accumulation and tolerance in *Celosia argentea* Linn.: A new Mn-hyperaccumulating plant species. *J Hazard Mater* 267: 136-141.
DOI: [10.1016/j.jhazmat.2013.12.051](https://doi.org/10.1016/j.jhazmat.2013.12.051)
- Liu, K., Yu, F., Chen, M., Zhou, Z., Chen, C., Li M.S., Zh, J., 2016. A newly found manganese hyperaccumulator—*Polygonum lapathifolium* Linn. *Int J Phytoremediat* 18(4): 348-353.
DOI: [10.1080/15226514.2015.1109589](https://doi.org/10.1080/15226514.2015.1109589)
- Liu, P., Tang, X., Gong, C., Xu, G., 2010. Manganese tolerance and accumulation in six Mn hyperaccumulators or accumulators. *Plant Soil*, 335(1-2): 385-395.
DOI: [10.1007/s11104-010-0427-x](https://doi.org/10.1007/s11104-010-0427-x)
- Lungwitz, E.E., 1900. The lixiviation of gold deposits by vegetation. *Eng Min J* 69: 500-502.
- Luo, C.L., Shen, Z.G., Li, X.D., 2008. Hot NTA application enhanced metal phytoextraction from contaminated soil. *Water Air Soil Pollut* 188(1-4): 127-137.
DOI: [10.1007/s11270-007-9529-3](https://doi.org/10.1007/s11270-007-9529-3)
- Luo, C., Shen, Z., Li, X., 2005. Enhanced phytoextraction of Cu, Pb, Zn and Cd with EDTA and EDDS. *Chemosph* 59(1): 1-11.
DOI: [10.1016/j.chemosphere.2004.09.100](https://doi.org/10.1016/j.chemosphere.2004.09.100)
- Lutts, S., Lefevre I, Delpérée, C., Kivits, S., Dechamps, C., Robledo, A., Correal, E., 2004. Heavy metal accumulation by the halophyte species *Mediterranean saltbush*. *J Environ Qual*, 33(4): 1271-1279.
DOI: [10.2134/jeq2004.1271](https://doi.org/10.2134/jeq2004.1271)

- Malagoli, M., Rossignolo, V., Salvalaggio, N., Schiavon, M., 2014. Potential for phytoextraction of copper by *Sinapis alba* and *Festuca rubra* cv. Merlin grown hydroponically and in vineyard soils. *Environ Sci Pollut Res* 21(5): 3294-3303.
DOI: [10.1007/s11356-013-2307-4](https://doi.org/10.1007/s11356-013-2307-4)
- Malayeri BE., Chehregani, A., Mohsenzadeh, F., Kazemeini, F., Asgari, M., 2013. Plants growing in a mining area: screening for metal accumulator plants possibly useful for bioremediation. *Toxicol Environl Chem* 95(3): 434-444.
DOI: [10.1080/02772248.2013.788701](https://doi.org/10.1080/02772248.2013.788701)
- Marcenò, C., Guarino R., 2015. A test on Ellenberg indicator values in the Mediterranean evergreen woods (*Quercetea ilicis*). *Rend Lincei* 26(3): 345-356.
DOI: [10.1007/s12210-015-0448-8](https://doi.org/10.1007/s12210-015-0448-8)
- Marchiol, L., Fellet, G., Perosa, D., Zerbi G., 2007. Removal of trace metals by *Sorghum bicolor* and *Helianthus annuus* in a site polluted by industrial wastes: a field experience. *Plant Physiol Biochem*, 45(5): 379-387.
DOI: [10.1016/j.plaphy.2007.03.018](https://doi.org/10.1016/j.plaphy.2007.03.018)
- Marchiol, L., Sacco, P., Assolari, S., Zerbi, G., 2004. Reclamation of polluted soil: phytoremediation potential of crop-related Brassica species. *Water Air Soil Pollut* 158(1): 345-356.
- Marques A.P., Moreira H, Rangel A.O., Castro P.M., 2009. Arsenic, lead and nickel accumulation in *Rubus ulmifolius* growing in contaminated soil in Portugal. *J Hazard Mater* 165(1-3): 174-179.
DOI: [10.1016/j.jhazmat.2008.09.102](https://doi.org/10.1016/j.jhazmat.2008.09.102)
- Martínez-Sánchez M.J., García-Lorenzo M.L., Pérez-Sirvent, C., Bech, J., 2012. Trace element accumulation in plants from an arid area affected by mining activities. *J Geochem Explor* 123: 8-12.
DOI: [10.1016/j.gexplo.2012.01.007](https://doi.org/10.1016/j.gexplo.2012.01.007)
- Massa, N., Andreucci, F., Poli, M., Aceto, M., Barbato, R., Berta, G., 2010. Screening for heavy metal accumulators amongst autochthonous plants in a polluted site in Italy. *Ecotoxicol Environ Saf* 73(8): 1988-1997.
DOI: [10.1016/j.ecoenv.2010.08.032](https://doi.org/10.1016/j.ecoenv.2010.08.032)
- Melo, E.E.C., Costa, E.T.S., Guilherme, L.R.G., Faquin, V., Nascimento, C.W.A., 2009. Accumulation of arsenic and nutrients by castor bean plants grown on an As-enriched nutrient solution. *J Hazardous Mater*, 168(1): 479-483.
DOI: [10.1016/j.jhazmat.2009.02.048](https://doi.org/10.1016/j.jhazmat.2009.02.048)
- Mihalík, J., Tlustoš, P., Szaková J., 2010. Comparison of willow and sunflower for uranium phytoextraction induced by citric acid. *J Radioanal Nucl Chem* 285(2): 279-285.
DOI: [10.1007/s10967-010-0538-0](https://doi.org/10.1007/s10967-010-0538-0)
- Minguzzi, C., Vergnano, O., 1948. Il contenuto di nichel nelle ceneri di *Alyssum bertolonii* Desv. *Mem Soc Tosc Sci Nat Ser A* 55: 49-77.
- Mkandawire, M., Dudel, E.G., 2005. Accumulation of arsenic in *Lemna gibba* L.(duckweed) in tailing waters of two abandoned uranium mining sites in Saxony, Germany. *Sci Total Environ* 336(1-3): 81-89.
DOI: [10.1016/j.scitotenv.2004.06.002](https://doi.org/10.1016/j.scitotenv.2004.06.002)
- Moreira H, Marques A.P., Rangel O., Castro, P.M., 2011. Heavy metal accumulation in plant species indigenous to a contaminated Portuguese site: prospects for phytoremediation. *Water Air Soil Pollut* 221(1-4): 377.
DOI: [10.1007/s11270-011-0797-6](https://doi.org/10.1007/s11270-011-0797-6)
- Nan, H., Jifang, Z., Dexin, D., Guangyue, L., Jie Y., Xin, C., Jia, Y., 2013. Screening of native hyperaccumulators at the Huayuan River contaminated by heavy metals. *Bioremediat J* 17(1): 21-29.
DOI: [10.1080/10889868.2012.703260](https://doi.org/10.1080/10889868.2012.703260)
- Nematian MA., Kazemeini, F., 2013. Accumulation of Pb, Zn, Cu and Fe in plants and hyperaccumulator choice in Galali iron mine area, Iran. *Int J Agric Crop Sci* 5(4): 426.
- Ning, Z., He, L., Xiao, T., Márton, L., 2015. High accumulation and subcellular distribution of thallium in green cabbage (*Brassica oleracea* L. var. capitata L.). *Int J Phytoremediat* 17(11): 1097-1104.
DOI: [10.1080/15226514.2015.1045133](https://doi.org/10.1080/15226514.2015.1045133)
- Pajuelo, E., Carrasco J.A., Romero, L.C., Chamber, M.A., Gotor, C., 2007. Evaluation of the metal phytoextraction potential of crop legumes. Regulation of the expression of O-Acetylserine (Thiol) Lyase under metal stress. *Plant Biol* 9(5): 672-681.
DOI: [10.1055/s-2007-965439](https://doi.org/10.1055/s-2007-965439)
- Panagos, P., Van Liedekerke, M., Yigini, Y., Montanarella, L., 2013. Contaminated sites in Europe: review of the current situation based on data collected through a European network. *J Environ Public Health* 2013.
DOI: [10.1155/2013/158764](https://doi.org/10.1155/2013/158764)
- Panich-Pat, T., Upatham, S., Pokethitiyook, P., Kruatrachue, M., Lanza, G.R., 2010. Phytoextraction of metal contaminants by *Typha angustifolia*: interaction of lead and cadmium in soil-water microcosms. *J Environ Prot* 1(04): 431.
DOI: [10.4236/jep.2010.14050](https://doi.org/10.4236/jep.2010.14050)
- Parveen, T., Hussain, A., Rao M.S., 2015. Growth and accumulation of heavy metals in turnip (*Brassica rapa*) irrigated with different concentrations of treated municipal wastewater. *Hydrol Res* 46(1): 60-71.
DOI: [10.2166/nh.2014.140](https://doi.org/10.2166/nh.2014.140)

- Pathak, C., Chopra AK., Srivastava, S., 2013. Accumulation of heavy metals in *Spinacia oleracea* irrigated with paper mill effluent and sewage. *Environ Monit Assess* 185(9): 7343-7352.
DOI: [10.1007/s10661-013-3104-8](https://doi.org/10.1007/s10661-013-3104-8)
- Peng, K., Luo, C., You, W., Lian, C., Li, X., Shen, Z., 2008. Manganese uptake and interactions with cadmium in the hyperaccumulator—*Phytolacca Americana* L. *Journal Hazard Mater* 154(1-3): 674-681.
DOI: [10.1016/j.jhazmat.2007.10.080](https://doi.org/10.1016/j.jhazmat.2007.10.080)
- Perry VR., Krogstad E.J., El-Mayas H, Greipsson, S., 2012. Chemically enhanced phytoextraction of lead-contaminated soils. *Int J Phytoremediat* 14(7): 703-713.
DOI: [10.1080/15226514.2011.619236](https://doi.org/10.1080/15226514.2011.619236)
- Pignatti, S., Menegoni, P., Pietrosanti S., 2005. Biondificazione attraverso le piante vascolari. Valori di indicazione secondo Ellenberg (Zeigerwerte) per le specie della Flora d'Italia. – *Braun-Blanquetia* 39: 1-97.
- Poschenrieder, C., Llugany, M., Lombini, A., Dinelli, E., Bech, J., Barceló, J., 2012. *Smilax aspera* L. an evergreen Mediterranean climber for phytoremediation. *J Geochem Explor* 123: 41-44.
DOI: [10.1016/j.gexplo.2012.07.012](https://doi.org/10.1016/j.gexplo.2012.07.012)
- Purakayastha, T.J., Viswanath, T., Bhadraray, S., Chhonkar, P.K., Adhikari, P.P., Suribabu, K., 2008. Phytoextraction of zinc, copper, nickel and lead from a contaminated soil by different species of *Brassica*. *Int J Phytoremediat* 10(1): 61-72.
DOI: [10.1080/15226510701827077](https://doi.org/10.1080/15226510701827077)
- Radulescu, C., Stihl, C., Popescu IV., Dulama, I.D., Chelarescu, E.D., Chilian, A., 2013. Heavy metal accumulation and translocation in different parts of *Brassica oleracea* L. *Rom J Phys* 58(9-10): 1337.
- Rajakaruna, N., Tompkins, K.M., Pavicevic, P.G., 2006. Phytoremediation: an affordable green technology for the clean-up of metal contaminated sites in Sri Lanka. *Ceylon J Sci*, 35: 25-39.
- Reeves, RD., Schwartz, C., Morel, J.L., Edmondson, J., 2001. Distribution and metal-accumulating behavior of *Thlaspi caerulescens* and associated metallophytes in France. *Int J Phytoremediat*: 3(2): 145-172.
DOI: [10.1080/15226510108500054](https://doi.org/10.1080/15226510108500054)
- Reeves, R.D., Van Der Ent, A., Baker, A.J., 2018. Global Distribution and Ecology of Hyperaccumulator Plants. In *Agromining: Farming for Metals* (pp. 75-92). Springer, Cham
- Roberts, T.M., Goodman, G.T., 1973. Persistence of heavy metals in soils and natural vegetation following closure of a smelter. In: *Trace Subst Environ Healt (United States)* 7(CONF-730613)
- Roccotiello, E., Serrano, H.C., Mariotti, M.G., Branquinho, C., 2016. The impact of Ni on the physiology of a Mediterranean Ni-hyperaccumulating plant. *Environ Sci Pollut Res* 23(12): 12414-12422. DOI: [10.1007/s11356-016-6461-3](https://doi.org/10.1007/s11356-016-6461-3)
- Rodríguez, N., Menéndez, N., Tornero, J., Amils, R., De La Fuente, V., 2005. Internal iron biomineralization in *Imperata cylindrica*, a perennial grass: chemical composition, speciation and plant localization. *New Phytol* 165(3): 781-789.
DOI: [10.1111/j.1469-8137.2004.01264.x](https://doi.org/10.1111/j.1469-8137.2004.01264.x)
- Saghi, A., Rashed Mohassel, M.H., Parsa, M., Hammami, H., 2016. Phytoremediation of lead-contaminated soil by *Sinapis arvensis* and *Rapistrum rugosum*. *Int J phytoremediat*, 18(4): 387-392.
DOI: [10.1080/15226514.2015.1109607](https://doi.org/10.1080/15226514.2015.1109607)
- Salem, Z.B., Laffray, X., Al-Ashoor, A., Ayadi H, Aleya, L., 2017. Metals and metalloid bioconcentrations in the tissues of *Typha latifolia* grown in the four interconnected ponds of a domestic landfill site. *J Environ Sci*, 54: 56-68.
DOI: [10.1016/j.jes.2015.10.039](https://doi.org/10.1016/j.jes.2015.10.039)
- Salt, D.E., Blaylock, M., Kumar, N.P., Dushenkov, V., Ensley, B.D., Chet, I., et al. 1995. Phytoremediation: a novel strategy for the removal of toxic metals from the environment using plants. *Nat Biotechnol*, 13(5): 468-474.
DOI: [10.1038/nbt0595-468](https://doi.org/10.1038/nbt0595-468)
- Salt, D.E., Smith, R.D., Raskin, I., 1998. Phytoremediation. *Annu Rev Plant Biol*, 49(1): 643-668.
DOI: [10.1146/annurev.arplant.49.1.643](https://doi.org/10.1146/annurev.arplant.49.1.643)
- Science Communication Unit, University of the West of England, Bristol. 2013. Science for environment policy In-depth Report: Soil Contamination: Impacts on Human Health. Report produced for the European Commission DG Environment, September 2013. Available at: http://ec.europa.eu/environment/integration/research/new_salert/pdf/IR5_en.pdf
- Selvi, F., 2007. Diversity, geographic variation and conservation of the serpentine flora of Tuscany (Italy). *Biodiver Conserv* 16(5): 1423-1439. DOI: [10.1007/s10531-006-6931-x](https://doi.org/10.1007/s10531-006-6931-x)
- Sharma R.K., Agrawal, M., Marshall, F., 2007. Heavy metal contamination of soil and vegetables in suburban areas of Varanasi, India. *Ecotoxicol Environ Saf* 66(2): 258-266.
DOI: [10.1016/j.ecoenv.2005.11.007](https://doi.org/10.1016/j.ecoenv.2005.11.007)
- Sheoran, V., Sheoran, A.S., Poonia, P., 2009. Phytomining: a review. *Miner Eng* 22(12): 1007-1019.
DOI: [10.1016/j.mineng.2009.04.001](https://doi.org/10.1016/j.mineng.2009.04.001)

- Sipos, G., Solti Á, Czech, V., Vashegyi I, Tóth, B., Cseh, E., Fodor, F., 2013. Heavy metal accumulation and tolerance of energy grass (*Elymus elongatus* subsp. *ponticus* cv. Szarvasi-1) grown in hydroponic culture. *Plant Physiol Biochem* 68: 96-103. DOI: [10.1016/j.plaphy.2013.04.006](https://doi.org/10.1016/j.plaphy.2013.04.006)
- Solhi, M., Shareatmadari, H., Hajabbasi, M.A., 2005. Lead and zinc extraction potential of two common crop plants, *Helianthus annuus* and *Brassica napus*. *Water Air Soil Pollut* 167(1-4): 59-71.
- Song, X., Hu, X., Ji, P., Li Y., Chi, G., Song, Y., 2012. Phytoremediation of cadmium-contaminated farmland soil by the hyperaccumulator *Beta vulgaris* L. var. *cicla*. *Bull Environ Contam Toxicol*, 88(4): 623-626. DOI: [10.1007/s00128-012-0524-z](https://doi.org/10.1007/s00128-012-0524-z)
- Soriano, M.A., Fereres, E., 2003. Use of crops for in situ phytoremediation of polluted soils following a toxic flood from a mine spill. *Plant Soil* 256(2): 253-264.
- Stojanović, M.D., Mihajlović, M.L., Milojković, J.V., Lopičić, Z.R., Adamović, M., Stanković, S., 2012. Efficient phytoremediation of uranium mine tailings by tobacco. *Environ Chem Lett* 10(4): 377-381. DOI: [10.1007/s10311-012-0362-6](https://doi.org/10.1007/s10311-012-0362-6)
- Su, C., Jiang, L., Zhang, W., 2014. A review on heavy metal contamination in the soil worldwide: Situation, impact and remediation techniques. *Environ Skept Crit* 3(2): 24.
- Su, Y, Han, F.X., Chen, J., Sridhar, B.M., Monts, D.L., 2008. Phytoextraction and accumulation of mercury in three plant species: Indian mustard (*Brassica juncea*), beard grass (*Polypogon monspeliensis*), and Chinese brake fern (*Pteris vittata*). *Int J Phytoremediat* 10(6): 547-560. DOI: [10.1080/15226510802115091](https://doi.org/10.1080/15226510802115091)
- Sun, Y, Zhou, Q., Wang, L., Liu, W., 2009. Cadmium tolerance and accumulation characteristics of *Bidens pilosa* L. as a potential Cd-hyperaccumulator. *J Hazard Mater* 161(2-3): 808-814. DOI: [10.1016/j.jhazmat.2008.04.030](https://doi.org/10.1016/j.jhazmat.2008.04.030)
- Szczygłowska, M., Piekarska, A., Konieczka, P., Namieśnik, J., 2011. Use of brassica plants in the phytoremediation and biofumigation processes. *Int J Mol Sci* 12(11): 7760-7771. DOI: [10.3390/ijms12117760](https://doi.org/10.3390/ijms12117760)
- Tapia, Y, Cala, V., Eymar, E., Frutos I, Gárate, A., Masaguer, A., 2011. Phytoextraction of cadmium by four Mediterranean shrub species. *Int J Phytoremediat* 13(6): 567-579. DOI: [10.1080/15226514.2010.495152](https://doi.org/10.1080/15226514.2010.495152)
- Taylor, S.I, Macnair, M.R., 2006. Within and between population variation for zinc and nickel accumulation in two species of *Thlaspi* (Brassicaceae). *New Phytol* 169(3): 505-514. DOI: [10.1111/j.1469-8137.2005.01625.x](https://doi.org/10.1111/j.1469-8137.2005.01625.x)
- The Plant List – version 1. 2013. <http://www.theplantlist.org>. Accessed 10 Apr 2017.
- Raunkiaer C. 1934. *The Life Forms of Plants and Statistical Plant Geography*. Oxford: Oxford University Press.
- Tisarum, R., Lessl JT., Dong, X., De Oliveira, L.M., Rathinasabapathi, B., Ma, L.Q., 2014. Antimony uptake, efflux and speciation in arsenic hyperaccumulator *Pteris vittata*. *Environ Pollut* 186: 110-114. DOI : [10.1016/j.envpol.2013.11.033](https://doi.org/10.1016/j.envpol.2013.11.033)
- Tóth, G., Hermann, T., Da Silva, M.R., Montanarella, L., 2016. Heavy metals in agricultural soils of the European Union with implications for food safety. *Environ Int* 88: 299-309. DOI: [10.1016/j.envint.2015.12.017](https://doi.org/10.1016/j.envint.2015.12.017)
- Tu, C., Ma, L.Q., Bondada, B., 2002. Arsenic accumulation in the hyperaccumulator Chinese brake and its utilization potential for phytoremediation. *J Environ Qual* 31(5): 1671-1675. DOI: [10.2134/jeq2002.1671](https://doi.org/10.2134/jeq2002.1671)
- Uraguchi, S., Watanabe, I, Yoshitomi, A., Kiyono, M., Kuno, K., 2006. Characteristics of cadmium accumulation and tolerance in novel Cd-accumulating crops, *Avena strigosa* and *Crotalaria juncea*. *J Exp Bot* 57(12): 2955-2965. DOI: [10.1093/jxb/erl056](https://doi.org/10.1093/jxb/erl056)
- Vaněk, A., Komárek, M., Chrástný, V., Bečka, D., Mihaljevič, M., Šebek, O., Schusterová, Z., 2010. Thallium uptake by white mustard (*Sinapis alba* L.) grown on moderately contaminated soils—Agro-environmental implications. *J Hazard Mater* 182(1-3): 303-308. DOI: [10.1016/j.jhazmat.2010.06.030](https://doi.org/10.1016/j.jhazmat.2010.06.030)
- Vara Prasad, M.N., De Oliveira Freitas, H.M., 2003. Metal hyperaccumulation in plants: biodiversity prospecting for phytoremediation technology. *Electron J Biotechnol* 6(3): 285-321. DOI: [10.2225/vol6-issue3-fulltext-6](https://doi.org/10.2225/vol6-issue3-fulltext-6)
- Wang, H.B., Wong, M.H, Lan CY, Baker, A.J.M., Qin, Y.R., Shu, W.S., Ye, Z.H., 2007. Uptake and accumulation of arsenic by 11 *Pteris* taxa from southern China. *Environ Pollut* 145(1): 225-233. DOI: [10.1016/j.envpol.2006.03.015](https://doi.org/10.1016/j.envpol.2006.03.015)
- Wang, S., Liu J., 2014. The effectiveness and risk comparison of EDTA with EGTA in enhancing Cd phytoextraction by *Mirabilis jalapa* L. *Environ Monit Assess* 186(2): 751-759. DOI: [10.1007/s10661-013-3414-x](https://doi.org/10.1007/s10661-013-3414-x)

- Warren, H.V., Delavault, R.E., 1950. Gold and silver content of some trees and horsetails in British Columbia. *Geol Soc Am Bull*, 61(2): 123-128.
DOI: [10.1130/0016-7606\(1950\)61\[123:GASCOS\]2.0.CO;2](https://doi.org/10.1130/0016-7606(1950)61[123:GASCOS]2.0.CO;2)
- Wei, S., Zhou, Q., Wang, X., Zhang, K., Guo, G., Ma, L.Q., 2005. A newly-discovered Cd-hyperaccumulator *Solatum nigrum* L. *Chin Sci Bull* 50(1): 33-38.
DOI: [10.1360/982004-292](https://doi.org/10.1360/982004-292)
- Wenzel, W.W., Jockwer, F., 1999. Accumulation of heavy metals in plants grown on mineralised soils of the Austrian Alps. *Environ Pollut* 104(1): 145-155.
DOI: [10.1016/s0269-7491\(98\)00139-0](https://doi.org/10.1016/s0269-7491(98)00139-0)
- Wilson, B., Pyatt, F.B., 2007. Heavy metal dispersion, persistence, and bioaccumulation around an ancient copper mine situated in Anglesey, UK. *Ecotoxicol Environ Saf* 66(2): 224-231.
DOI: [10.1016/j.ecoenv.2006.02.015](https://doi.org/10.1016/j.ecoenv.2006.02.015)
- Yabe, J., Ishizuka, M., Umemura, T., 2010. Current levels of heavy metal pollution in Africa. *J Vet Med Sci* 72(10): 1257-1263.
- Yaman, M., 2014. *Teucrium* as a Novel Discovered Hyperaccumulator for the Phytoextraction of Ni-contaminated Soils. *Ekol Derg* 23(90).
DOI: [10.5053/ekoloji.2014.9010](https://doi.org/10.5053/ekoloji.2014.9010)
- Yanqun, Z., Yuan, L., Jianjun, C., Haiyan, C., Li, Q., Schwartz, C., 2005. Hyperaccumulation of Pb, Zn and Cd in herbaceous grown on lead-zinc mining area in Yunnan, China. *Environ Int* 31(5): 755-762.
DOI: [10.1016/j.envint.2005.02.004](https://doi.org/10.1016/j.envint.2005.02.004)
- Yu, L., Xin, G., Gang, W., Zhang, Q., Qiong, S., Guoju, X., 2008. Heavy metal contamination and source in arid agricultural soil in central Gansu Province, China. *J Environ Sci* 20(5): 607-612.
DOI: [10.1016/S1001-0742\(08\)62101-4](https://doi.org/10.1016/S1001-0742(08)62101-4)
- Zhai, S., Xiao H, Shu Y, Zhao, Z., 2013. Countermeasures of heavy metal pollution. *Chin J Geochem* 32(4): 446-450.
DOI: [10.1007/s11631-013-0654-y](https://doi.org/10.1007/s11631-013-0654-y)
- Zhu Y.G., Williams P.N., Meharg, A.A., 2008. Exposure to inorganic arsenic from rice: a global health issue? *Environ Pollut* 154(2): 169-171.
DOI: [10.1016/j.envpol.2008.03.015](https://doi.org/10.1016/j.envpol.2008.03.015)
- Zhuang, P., Wensheng, S.H.U., Zhian L.I., Bin, L., Jintian, L.I., Jingsong, S., 2009. Removal of metals by sorghum plants from contaminated land. *J Environ Sci* 21(10): 1432-1437.
DOI: [10.1016/S1001-0742\(08\)62436-5](https://doi.org/10.1016/S1001-0742(08)62436-5)