



## A note on a nonlinear $m$ -point boundary value problem for $p$ -Laplacian differential inclusions

*Mouffak Benchohra, Sotiris K. Ntouyas, and A. Ouahab*



## A NOTE ON A NONLINEAR $m$ -POINT BOUNDARY VALUE PROBLEM FOR $p$ -LAPLACIAN DIFFERENTIAL INCLUSIONS

M. BENCHOHRA, S. K. NTOUYAS, AND A. OUAHAB

[Received: March 15, 2003]

**ABSTRACT.** In this note a selection theorem due to Bressan and Colombo for lower semi-continuous multi-valued operators with nonempty closed decomposable values combined with Schaefer's fixed point theorem is used to investigate the existence of positive solutions for  $m$ -point boundary value problems for one dimensional  $p$ -Laplacian differential inclusions.

*Mathematics Subject Classification:* 34A60, 34B10, 34B15

*Keywords:* Boundary value problem,  $p$ -Laplacian, positive solutions, differential inclusions,  $m$ -point boundary value problem, continuous selection, existence, fixed point

### 1. INTRODUCTION

This note is concerned with the existence of positive solutions for the following class of boundary value problems for  $m$ -point one dimensional  $p$ -Laplacian differential inclusions

$$(\varphi(y'))' \in F(t, y), \quad \text{a. e. } t \in J := [0, 1]; \quad (1.1)$$

$$y'(0) = \sum_{i=1}^{m-2} b_i y'(\xi_i), \quad y(1) = \sum_{i=1}^{m-2} a_i y'(\xi_i), \quad (1.2)$$

where  $\varphi : \mathbb{R}_+^* \rightarrow \mathbb{R}_+$  defined by  $\varphi(v) := |v|^{p-2}v$ ,  $p > 1$ ,  $F : J \times \mathbb{R} \rightarrow \mathcal{P}(\mathbb{R}_+)$  is a multi-valued map,  $\mathcal{P}(\mathbb{R}_+)$  is the family of all subsets of  $\mathbb{R}_+$ ,  $\xi_i \in (0, 1)$ ,  $0 < \xi_1 < \xi_2 < \dots < \xi_{m-2} < 1$ , and  $a_i, b_i$ ,  $i = 1, \dots, m-2$ , are positive and satisfy  $0 < \sum_{i=1}^{m-2} a_i < 1$ ,  $\sum_{i=1}^{m-2} b_i < 1$ . The study of multi-point boundary value problems for linear second order ordinary differential equations was initiated by Il'in and Moiseev [15, 16]. Then Gupta [10, 11] studied three-point boundary value problems for nonlinear ordinary differential equations, the  $m$ -point boundary value problem was studied by Gupta *et al.*, [12, 14], Ma [18]. Very recently, in a series of papers by Benchohra and Ntouyas (see [2–5]) some extensions to multi-point differential inclusions have been proposed with the aid of fixed point arguments in the cases when the right-hand side is convex

as well as nonconvex valued. Some existence results for one dimensional  $p$ -Laplacian differential equations are given in the papers by Bai and Fang [1] and Jian and Guo [17]. Our goal here is to give the existence of at least one positive solution for  $m$ -point boundary value problems for one dimensional  $p$ -Laplacian differential inclusions. Our approach relies on Schaefer's fixed point theorem combined with a selection theorem due to Bressan and Colombo [6] for lower semi-continuous operators with nonempty closed and decomposable values.

## 2. PRELIMINARIES

In this Section, we introduce notations, definitions, and preliminary facts from multi-valued analysis which are used throughout this paper.

$C([0, 1], \mathbb{R})$  is the Banach space of all continuous functions from  $[0, 1]$  into  $\mathbb{R}$  with the norm

$$\|y\|_{\infty} := \sup\{|y(t)| : 0 \leq t \leq 1\}.$$

$AC([0, 1], \mathbb{R})$  is the space of all absolutely continuous functions  $y$  from  $[0, 1]$  into  $\mathbb{R}$ .  $L^1(J, \mathbb{R})$  denotes the Banach space of functions  $y : J \rightarrow \mathbb{R}$  which are Lebesgue integrable normed by

$$\|y\|_{L^1} = \int_0^1 |y(t)| dt.$$

Let  $\mathcal{A}$  be a subset of  $[0, 1] \times \mathbb{R}$ .  $\mathcal{A}$  is  $\mathcal{L} \otimes \mathcal{B}$  measurable if  $\mathcal{A}$  belongs to the  $\sigma$ -algebra generated by all sets of the form  $\mathcal{N} \times D$  where  $\mathcal{N}$  is Lebesgue measurable in  $[0, 1]$  and  $D$  is Borel measurable in  $\mathbb{R}$ . A subset  $\mathcal{I}$  of  $L^1([0, 1], \mathbb{R})$  is decomposable if for all  $u, v \in \mathcal{I}$  and  $\mathcal{N} \subset [0, 1]$  measurable the function  $u\chi_{\mathcal{N}} + v\chi_{[0, 1] - \mathcal{N}} \in \mathcal{I}$ , where  $\chi_{[0, 1]}$  stands for the characteristic function of  $[0, 1]$ .

Let  $E$  be a Banach space,  $X$  a nonempty closed subset of  $E$  and  $G : X \rightarrow \mathcal{P}(E)$  a multi-valued operator with nonempty closed values.  $G$  is lower semi-continuous (l.s.c.) if the set  $\{x \in X : G(x) \cap B \neq \emptyset\}$  is open for any open set  $B$  in  $E$ .  $G$  has a fixed point if there is  $x \in X$  such that  $x \in G(x)$ . For more details on multi-valued maps we refer to the books by Deimling [7], Górniewicz [9] and Hu and Papageorgiou [19].

**Definition 1.** Let  $Y$  be a separable metric space and let  $N : Y \rightarrow \mathcal{P}(L^1([0, b], \mathbb{R}))$  be a multi-valued operator. We say  $N$  has the property (BC) if

- (1)  $N$  is lower semi-continuous (l.s.c.);
- (2)  $N$  has nonempty closed and decomposable values.

Let  $F : J \times \mathbb{R}^+ \rightarrow \mathcal{P}(\mathbb{R}^+)$  be a multi-valued map with nonempty compact values. Assign to  $F$  the multi-valued operator

$$\mathcal{F} : C([0, 1], \mathbb{R}^+) \rightarrow \mathcal{P}(L^1([0, 1], \mathbb{R}^+))$$

by letting

$$\mathcal{F}(y) = \{w \in L^1([0, 1], \mathbb{R}) : w(t) \in F(t, y(t)) \text{ for a. e. } t \in [0, 1]\}.$$

The operator  $\mathcal{F}$  is called the Niemytzki operator associated with  $F$ .

**Definition 2.** Let  $F : J \times \mathbb{R}^+ \rightarrow \mathcal{P}(\mathbb{R}^+)$  be a multi-valued function with nonempty compact values. We say  $F$  is of lower semi-continuous type (l.s.c. type) if its associated Niemytzki operator  $\mathcal{F}$  is lower semi-continuous and has nonempty closed and decomposable values.

Next we state a selection theorem due to Bressan and Colombo.

**Theorem 1** ([6]). *Assume that  $Y$  is a separable metric space and let  $N : Y \rightarrow \mathcal{P}(L^1([0, 1], \mathbb{R}))$  be a multi-valued operator which has the property (BC). Then  $N$  has a continuous selection, i.e. there exists a continuous function (single-valued)  $g : Y \rightarrow L^1(J, \mathbb{R})$  such that  $g(y) \in N(y)$  for every  $y \in Y$ .*

Let us introduce the following hypotheses which are assumed hereafter:

- (H1)  $F : [0, 1] \times \mathbb{R}^+ \rightarrow \mathcal{P}(\mathbb{R}^+)$  is a nonempty compact valued multi-valued map such that:
- (a)  $(t, y) \mapsto F(t, y)$  is  $\mathcal{L} \otimes \mathcal{B}$  measurable;
  - (b)  $y \mapsto F(t, y)$  is lower semi-continuous for a. e.  $t \in [0, 1]$ ;
- (H2) There exists a function  $h \in L^1([0, 1], \mathbb{R}_+)$  such that

$$\|F(t, y)\| := \sup\{|v| : v \in F(t, y)\} \leq h(t) \text{ for a. e. } t \in [0, 1] \text{ and for } y \in \mathbb{R}.$$

### 3. MAIN RESULT

Let us start by defining what we mean by a solution of problem (1.1)–(1.2).

**Definition 3.** A function  $y \in C^1((0, 1), \mathbb{R})$  with  $\varphi(y') \in AC((0, 1), \mathbb{R})$  is said to be a solution of (1.1), (1.2) if there exists  $v(t) \in L^1(J, \mathbb{R})$  such that  $y$  satisfies the equation  $(\varphi(y'))' = v(t)$  a. e. on  $J$  and the condition (1.2).

**Theorem 2.** *Suppose that hypotheses (H1), (H2) are satisfied. Then the  $m$ -point BVP (1.1), (1.2) has at least one positive solution.*

**PROOF.** (H1) and (H2) imply by Lemma 2.2 in Frigon [8] that  $F$  is of the lower semi-continuous type. Then from Theorem 1 there exists a continuous function  $f : C([0, 1], \mathbb{R}) \rightarrow L^1([0, 1], \mathbb{R})$  such that  $f(y) \in \mathcal{F}(y)$  for all  $y \in C([0, 1], \mathbb{R})$ . Consider the following problem

$$(\varphi(y'))' = f(y(t)), \quad \text{a. e. } t \in J, \quad (3.1)$$

$$y'(0) = \sum_{i=1}^{m-2} b_i y'(\xi_i), \quad y(1) = \sum_{i=1}^{m-2} a_i y'(\xi_i). \quad (3.2)$$

Clearly, if  $y$  is a solution of problem (3.1), (3.2), then  $y$  is a solution to problem (1.1), (1.2).

Transform the problem (3.1), (3.2) into a fixed point problem. Consider the operator  $N : C([0, 1], \mathbb{R}^+) \rightarrow C([0, 1], \mathbb{R})$  defined by:

$$\begin{aligned} N(y)(t) = & - \int_0^t \psi \left( \int_0^s f(y(\tau)) d\tau \right) ds - tB \sum_{i=1}^{m-2} b_i \psi \left( \int_0^{\xi_i} f(y(\tau)) d\tau \right) \\ & + A \left\{ \int_0^1 \psi \left( \int_0^s f(y(\tau)) d\tau \right) ds - \sum_{i=1}^{m-2} a_i \int_0^{\xi_i} \psi \left( \int_0^s f(y(\tau)) d\tau \right) ds \right. \\ & \left. + B \sum_{i=1}^{m-2} b_i \psi \left( \int_0^{\xi_i} f(y(\tau)) d\tau \right) \left( 1 - \sum_{i=1}^{m-2} a_i \xi_i \right) \right\} \end{aligned}$$

where  $\psi$  is the inverse of the function  $\varphi$  defined by  $\psi(w) := |w|^{q-2}w$ , with  $q = \frac{p}{p-1} > 1$  and

$$A = \left( 1 - \sum_{i=1}^{m-2} a_i \right)^{-1}, \quad B = \left( 1 - \sum_{i=1}^{m-2} b_i \right)^{-1}.$$

The fixed points of the operator  $N$  are solutions to problem (3.1), (3.2) (see [1]). It is clear that  $N(y)(t) \geq 0$  on  $J$  for any  $y \in C([0, 1], \mathbb{R}^+)$ . We shall first show that  $N$  is completely continuous. The proof will be given in three steps.

**Step 1:**  $N$  is continuous. Let  $\{y_n\}$  be a sequence such that  $y_n \rightarrow y$  in  $C([0, 1], \mathbb{R})$ . Set

$$L(y)(t) := \int_0^t |f(y(s))| ds.$$

Then

$$|L(y_n)(s) - L(y)(s)| \leq \int_0^s |f(y_n(s)) - f(y(s))| ds \leq \int_0^1 |f(y_n(s)) - f(y(s))| ds.$$

Since  $f$  is a continuous function, it follows that

$$\|L(y_n) - L(y)\|_\infty \leq \|f(y_n(\cdot)) - f(y(\cdot))\|_{L^1} \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Then

$$\begin{aligned}
|N(y_n(t)) - N(y(t))| &\leq \int_0^1 |\psi(L(y_n(t))) - \psi(L(y(t)))| ds \\
&\quad + tB \sum_{i=1}^{m-2} b_i |\psi(L(y_n(\xi_i))) - \psi(L(y(\xi_i)))| \\
&\quad + A \int_0^1 |\psi(L(y_n(s))) - \psi(L(y(s)))| ds \\
&\quad + A \sum_{i=1}^{m-2} a_i \int_0^{\xi_i} |\psi(L(y_n(s))) - \psi(L(y(s)))| ds \\
&\quad + AB \sum_{i=1}^{m-2} b_i |\psi(L(y_n(\xi_i))) - \psi(L(y(\xi_i)))| \left( 1 - \sum_{i=1}^{m-2} a_i \xi_i \right).
\end{aligned}$$

Since  $\psi$  is a continuous function, then

$$\begin{aligned}
\|N(y_n) - N(y)\|_\infty &\leq \|\psi(L(y_n)) - \psi(L(y))\|_\infty + B \sum_{i=1}^{m-2} b_i \|\psi(L(y_n)) - \psi(L(y))\|_\infty \\
&\quad + A \|\psi(L(y_n)) - \psi(L(y))\|_\infty + A \sum_{i=1}^{m-2} a_i \|\psi(L(y_n)) - \psi(L(y))\|_\infty \\
&\quad + AB \sum_{i=1}^{m-2} b_i \|\psi(L(y_n)) - \psi(L(y))\|_\infty \left( 1 - \sum_{i=1}^{m-2} a_i \xi_i \right) \text{ as } n \rightarrow \infty.
\end{aligned}$$

**Step 2:**  $N$  maps bounded sets into bounded sets in  $C([0, 1], \mathbb{R})$ . Indeed, it is enough to show that, for any  $q > 0$ , there exists a positive constant  $\ell$  such that, for each  $y \in B_q = \{y \in C([0, 1], \mathbb{R}) : \|y\|_\infty \leq q\}$ , we have  $\|N(y)\|_\infty \leq \ell$ . From (H2), we have

$$\left| \int_0^1 f(y(s)) ds \right| \leq \|h\|_{L^1} := q^*,$$

and

$$\begin{aligned}
|N(y)(t)| &\leq \int_0^1 |\psi(L(y(s)))| ds + tB \sum_{i=1}^{m-2} b_i |\psi(L(y(\xi_i)))| \\
&+ A \int_0^1 |\psi(L(y(s)))| ds + A \sum_{i=1}^{m-2} a_i \int_0^{\xi_i} |\psi(L(y(s)))| ds \\
&+ AB \sum_{i=1}^{m-2} b_i |\psi(L(y(\xi_i)))| \left( 1 - \sum_{i=1}^{m-2} a_i \xi_i \right).
\end{aligned}$$

Then

$$\begin{aligned}
\|N(y)\|_\infty &\leq \sup_{w \in [-q^*, q^*]} |\psi(w)| + B \sum_{i=1}^{m-2} b_i \sup_{w \in [-q^*, q^*]} |\psi(w)| \\
&+ A \sup_{w \in [-q^*, q^*]} |\psi(w)| + A \sum_{i=1}^{m-2} a_i \sup_{w \in [-q^*, q^*]} |\psi(w)| \\
&+ AB \sum_{i=1}^{m-2} b_i \sup_{w \in [-q^*, q^*]} |\psi(w)| \left( 1 - \sum_{i=1}^{m-2} a_i \xi_i \right) := \ell.
\end{aligned}$$

**Step 3:** *N maps bounded sets into equicontinuous sets of  $C([0, 1], \mathbb{R})$ .* Let  $l_1, l_2 \in [0, 1]$ ,  $l_1 < l_2$  and  $B_q$  be a bounded set of  $C([0, 1], \mathbb{R})$  as in Step 2. Let  $y \in B_q$ , then

$$\begin{aligned}
|N(y)(l_2) - N(y)(l_1)| &\leq (l_2 - l_1) \sup_{w \in [-q^*, q^*]} |\psi(w)| \\
&+ B(l_2 - l_1) \sum_{i=1}^{m-2} b_i \left| \psi \left( \int_0^{\xi_i} f(y(\tau)) d\tau \right) \right|.
\end{aligned}$$

As  $l_2 \rightarrow l_1$ , the right-hand side of the above inequality tends to zero. As a consequence of Steps 1 to 3 together with the Arzelá-Ascoli theorem, we can conclude that  $N : C([0, 1], \mathbb{R}) \rightarrow C([0, 1], \mathbb{R})$  is completely continuous.

**Step 4:** *The set*

$$\mathcal{E}(N) := \{y \in C([0, 1], \mathbb{R}) : y = \lambda N(y), \text{ for some } 0 < \lambda < 1\}$$

*is bounded.*

The reasoning used in the proof of Step 2 shows that the set  $\mathcal{E}(N)$  is bounded.

Set  $X := C([0, 1], \mathbb{R})$ . As a consequence of Schaefer's fixed point theorem [20, p. 29] we deduce that  $N$  has a fixed point  $y$  which is a solution to problem (3.1), (3.2), and hence, a solution to problem (1.1), (1.2).

□

## REFERENCES

- [1] BAI, C. Z. AND FANG, J. X.: *Existence of multiple positive solutions for nonlinear  $m$ -point boundary value problems*, Appl. Math. Comp. **140**(2003), 297-305.
- [2] BENCHOHRA, M. AND NTOUYAS, S. K.: *Multi-point boundary value problem for second order differential inclusions*, Math. Vesnik **53**(2001), 51-58.
- [3] BENCHOHRA, M. AND NTOUYAS, S. K.: *A note on a three-point boundary value problem for second order differential inclusions*, Math. Notes (Miskolc) **2**(2001), 39-47.
- [4] BENCHOHRA, M. AND NTOUYAS, S. K.: *On a three and four-point boundary value problem for second order differential inclusions*, Math. Notes (Miskolc) **2**(2001), 93-101.
- [5] BENCHOHRA, M. AND NTOUYAS, S. K.: *Multi-point boundary value problem for lower semi-continuous differential inclusions*, Math. Notes (Miskolc) **3**(2002), 91-99.
- [6] BRESSAN, A. AND COLOMBO, G.: *Extensions and selections of maps with decomposable values*, Studia Math. **90**(1988), 69-86.
- [7] DEIMLING, K.: *Multivalued Differential Equations*, Walter de Gruyter, Berlin-New York, 1992.
- [8] FRIGON, M.: *Théorèmes d'existence de solutions d'inclusions différentielles*, *Topological Methods in Differential Equations and Inclusions* (edited by A. Granas and M. Frigon), NATO ASI Series C, Vol. 472, Kluwer Acad. Publ., Dordrecht, (1995), 51-87.
- [9] GÓRNIOWICZ, L.: *Topological Fixed Point Theory of Multivalued Mappings*, Mathematics and its Applications, 495, Kluwer Academic Publishers, Dordrecht, 1999.
- [10] GUPTA, C. P.: *Solvability of a three-point second order ordinary differential equation*, J. Math. Anal. Appl. **168**(1992), 540-55.
- [11] GUPTA, C. P.: *A generalized multi-point boundary value problem for second order ordinary differential equation*, Appl. Math. Comput. **89**(1998), 133-146.
- [12] GUPTA, C. P., NTOUYAS, S.K. AND TSAMATOS, P. CH.: *On an  $m$ -point boundary value problem for second order ordinary differential equations*, Nonlinear Anal. **23**(1994), 1427-1436.
- [13] GUPTA, C.P., NTOUYAS, S. K. AND TSAMATOS, P. CH.: *Existence results for  $m$ -point boundary value problem for second order ordinary differential equations*, Differential Equations Dynam. Systems **2**(1994), 289-298.
- [14] GUPTA, C. P., NTOUYAS, S. K. AND TSAMATOS, P. CH.: *On the solvability of some  $m$ -point boundary value problems*, Appl. Math. **41**(1996), 1-17.
- [15] IL'IN, V. A. AND MOISEEV, E. I.: *Nonlocal boundary value problem of the second kind for a Sturm-Liouville operator in its differential and finite difference aspects*, Differential Equations **7**(1987), 803-810.
- [16] IL'IN, V. A. AND MOISEEV, E. I.: *Nonlocal boundary value problem of the second kind for a Sturm-Liouville operator*, Differential Equations **8**(1987), 979-987.
- [17] JIANG, D. AND GUO, W. J.: *Upper and lower solution method and a singular boundary value problem for the one-dimensional  $p$ -Laplacian*, J. Math. Anal. Appl. **252**(2000), 631-648.
- [18] MA, R. Y.: *Existence of solutions of nonlinear  $m$ -point boundary value problems*, J. Math. Anal. Appl. **256**(2001), 556-567.
- [19] HU, SH. AND PAPAGEORGIOU, N.: *Handbook of Multivalued Analysis, Volume I: Theory*, Kluwer Academic Publishers, Dordrecht, 1997.
- [20] SMART, D. R.: *Fixed Point Theorems*, Cambridge Univ. Press, Cambridge, 1974.



*Authors' Addresses***M. Benchohra:**

LABORATOIRE DE MATHÉMATIQUES, UNIVERSITÉ DE SIDI BEL ABBÈS, BP 89, 22000 SIDI BEL ABBÈS,  
ALGÉRIE

*E-mail address:* benchohra@univ-sba.dz

**S. K. Ntouyas:**

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF IOANNINA, 451 10 IOANNINA, GREECE

*E-mail address:* sntouyas@cc.uoi.gr

**A. Ouahab:**

LABORATOIRE DE MATHÉMATIQUES, UNIVERSITÉ DE SIDI BEL ABBÈS, BP 89, 22000 SIDI BEL ABBÈS,  
ALGÉRIE